

# From SVD to PCA

Xin Li

April 6, 2009

## SVD Review

The statement of SVD

A closer look at SVD

## PCA from SVD

Principal components

PCA and SVD in Matlab

Application to face recognition

# Recall the SVD

- ▶ Given a matrix  $\mathbf{X}_{m \times n}$  of rank  $n$ .

# Recall the SVD

- ▶ Given a matrix  $\mathbf{X}_{m \times n}$  of rank  $n$ .
- ▶ We can find orthogonal matrices  $\mathbf{U}_{m \times n}$  and  $\mathbf{V}_{n \times n}$ ,

# Recall the SVD

- ▶ Given a matrix  $\mathbf{X}_{m \times n}$  of rank  $n$ .
- ▶ We can find orthogonal matrices  $\mathbf{U}_{m \times n}$  and  $\mathbf{V}_{n \times n}$ ,
- ▶ and diagonal matrix  $\mathbf{D}_{n \times n}$ :

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

# Recall the SVD

- ▶ Given a matrix  $\mathbf{X}_{m \times n}$  of rank  $n$ .
- ▶ We can find orthogonal matrices  $\mathbf{U}_{m \times n}$  and  $\mathbf{V}_{n \times n}$ ,
- ▶ and diagonal matrix  $\mathbf{D}_{n \times n}$ :

$$\mathbf{X} = \mathbf{U} \mathbf{D} \mathbf{V}^T$$

- ▶ Visualize it as:

$$\begin{bmatrix} \mathbf{X} \end{bmatrix}_{m \times n} = \begin{bmatrix} \mathbf{U} \end{bmatrix}_{m \times n} \begin{bmatrix} \diagdown \end{bmatrix}_{n \times n} \begin{bmatrix} \mathbf{V}^T \end{bmatrix}_{n \times n}$$

# Recall the SVD

- ▶ Given a matrix  $\mathbf{X}_{m \times n}$  of rank  $n$ .
- ▶ We can find orthogonal matrices  $\mathbf{U}_{m \times n}$  and  $\mathbf{V}_{n \times n}$ ,
- ▶ and diagonal matrix  $\mathbf{D}_{n \times n}$ :

$$\mathbf{X} = \mathbf{U} \mathbf{D} \mathbf{V}^T$$

- ▶ Visualize it as:

$$\begin{bmatrix} \mathbf{X} \end{bmatrix}_{m \times n} = \begin{bmatrix} \mathbf{U} \end{bmatrix}_{m \times n} \begin{bmatrix} \diagdown \end{bmatrix}_{n \times n} \begin{bmatrix} \mathbf{V}^T \end{bmatrix}_{n \times n}$$

- ▶ Now we derive SVD when the rank of  $\mathbf{X}$  is  $r < n$ .

# Deriving the SVD

- Find the eigen decomposition of  $\mathbf{X}^T \mathbf{X}$ :

$$\mathbf{X}^T \mathbf{X} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T$$

where  $\mathbf{\Lambda}$  is the diagonal matrix with the eigenvalues  $\lambda_i$ 's of  $\mathbf{X}^T \mathbf{X}$  on its diagonal.



# Deriving the SVD

- ▶ Find the eigen decomposition of  $\mathbf{X}^T \mathbf{X}$ :

$$\mathbf{X}^T \mathbf{X} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T$$

where  $\mathbf{\Lambda}$  is the diagonal matrix with the eigenvalues  $\lambda_i$ 's of  $\mathbf{X}^T \mathbf{X}$  on its diagonal.

- ▶ The matrix  $\mathbf{V}$  is orthogonal whose columns  $\mathbf{v}_i$ 's are eigenvectors of  $\mathbf{X}^T \mathbf{X}$ :  $\mathbf{X}^T \mathbf{X} \mathbf{v}_i = \lambda_i \mathbf{v}_i, \quad i = 1 : n$

# Deriving the SVD

- ▶ Find the eigen decomposition of  $\mathbf{X}^T \mathbf{X}$ :

$$\mathbf{X}^T \mathbf{X} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T$$

where  $\mathbf{\Lambda}$  is the diagonal matrix with the eigenvalues  $\lambda_i$ 's of  $\mathbf{X}^T \mathbf{X}$  on its diagonal.

- ▶ The matrix  $\mathbf{V}$  is orthogonal whose columns  $\mathbf{v}_i$ 's are eigenvectors of  $\mathbf{X}^T \mathbf{X}$ :  $\mathbf{X}^T \mathbf{X} \mathbf{v}_i = \lambda_i \mathbf{v}_i$ ,  $i = 1 : n$
- ▶ It is also known that  $\lambda_i \geq 0$  (Homework: Verify it!). If the rank of  $\mathbf{X}$  is  $r$  ( $r \leq n$ ), there are exactly  $r$  positive eigenvalues, say

$$\lambda_1, \lambda_2, \dots, \lambda_r$$

# Deriving the SVD

- ▶ Find the eigen decomposition of  $\mathbf{X}^T \mathbf{X}$ :

$$\mathbf{X}^T \mathbf{X} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T$$

where  $\mathbf{\Lambda}$  is the diagonal matrix with the eigenvalues  $\lambda_i$ 's of  $\mathbf{X}^T \mathbf{X}$  on its diagonal.

- ▶ The matrix  $\mathbf{V}$  is orthogonal whose columns  $\mathbf{v}_i$ 's are eigenvectors of  $\mathbf{X}^T \mathbf{X}$ :  $\mathbf{X}^T \mathbf{X} \mathbf{v}_i = \lambda_i \mathbf{v}_i$ ,  $i = 1 : n$
- ▶ It is also known that  $\lambda_i \geq 0$  (Homework: Verify it!). If the rank of  $\mathbf{X}$  is  $r$  ( $r \leq n$ ), there are exactly  $r$  positive eigenvalues, say

$$\lambda_1, \lambda_2, \dots, \lambda_r$$

- ▶ For  $i = 1 : r$ , define  $\sigma_i = \sqrt{\lambda_i}$  and  $\mathbf{u}_i = \mathbf{X} \mathbf{v}_i / \sigma_i$ .

# Deriving the SVD

- Find the eigen decomposition of  $\mathbf{X}^T \mathbf{X}$ :

$$\mathbf{X}^T \mathbf{X} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T$$

where  $\mathbf{\Lambda}$  is the diagonal matrix with the eigenvalues  $\lambda_i$ 's of  $\mathbf{X}^T \mathbf{X}$  on its diagonal.

- The matrix  $\mathbf{V}$  is orthogonal whose columns  $\mathbf{v}_i$ 's are eigenvectors of  $\mathbf{X}^T \mathbf{X}$ :  $\mathbf{X}^T \mathbf{X} \mathbf{v}_i = \lambda_i \mathbf{v}_i$ ,  $i = 1 : n$
- It is also known that  $\lambda_i \geq 0$  (Homework: Verify it!). If the rank of  $\mathbf{X}$  is  $r$  ( $r \leq n$ ), there are exactly  $r$  positive eigenvalues, say

$$\lambda_1, \lambda_2, \dots, \lambda_r$$

- For  $i = 1 : r$ , define  $\sigma_i = \sqrt{\lambda_i}$  and  $\mathbf{u}_i = \mathbf{X} \mathbf{v}_i / \sigma_i$ .
- Then

$$\mathbf{X}^T \mathbf{u}_i = \sigma_i \mathbf{v}_i$$

$$\mathbf{X} \mathbf{v}_i = \sigma_i \mathbf{u}_i$$

# Deriving the SVD, continued

- Put in matrix notation:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r]$$

$$\mathbf{X}^T[\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_r] = [\sigma_1 \mathbf{v}_1 \ \sigma_2 \mathbf{v}_2 \ \cdots \ \sigma_r \mathbf{v}_r]$$

# Deriving the SVD, continued

- Put in matrix notation:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r]$$

$$\mathbf{X}^T[\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_r] = [\sigma_1 \mathbf{v}_1 \ \sigma_2 \mathbf{v}_2 \ \cdots \ \sigma_r \mathbf{v}_r]$$

- Note that  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$  is an orthogonal set in  $\mathbb{R}^n$ . We can extend it to an orthogonal basis.

# Deriving the SVD, continued

- Put in matrix notation:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r]$$

$$\mathbf{X}^T[\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_r] = [\sigma_1 \mathbf{v}_1 \ \sigma_2 \mathbf{v}_2 \ \cdots \ \sigma_r \mathbf{v}_r]$$

- Note that  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$  is an orthogonal set in  $\mathbb{R}^n$ . We can extend it to an orthogonal basis.
- Solve  $[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r]\mathbf{v} = \mathbf{0}$   
(What is the dimension of the solution space of this homogeneous system?)

# Deriving the SVD, continued

- Put in matrix notation:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r]$$

$$\mathbf{X}^T[\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_r] = [\sigma_1 \mathbf{v}_1 \ \sigma_2 \mathbf{v}_2 \ \cdots \ \sigma_r \mathbf{v}_r]$$

- Note that  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$  is an orthogonal set in  $\mathbb{R}^n$ . We can extend it to an orthogonal basis.
- Solve  $[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r]\mathbf{v} = \mathbf{0}$   
(What is the dimension of the solution space of this homogeneous system?)
- Answer: Since the coefficient matrix  $\mathbf{V}_1 := [\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r]$  has rank  $r$  (Why?), the dimension of the solution space is  $n - r$ .



# Deriving the SVD, continued

- ▶ Put in matrix notation:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r]$$

$$\mathbf{X}^T[\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_r] = [\sigma_1 \mathbf{v}_1 \ \sigma_2 \mathbf{v}_2 \ \cdots \ \sigma_r \mathbf{v}_r]$$

- ▶ Note that  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$  is an orthogonal set in  $\mathbb{R}^n$ . We can extend it to an orthogonal basis.
- ▶ Solve  $[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r]\mathbf{v} = \mathbf{0}$   
(What is the dimension of the solution space of this homogeneous system?)
- ▶ Answer: Since the coefficient matrix  $\mathbf{V}_1 := [\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_r]$  has rank  $r$  (Why?), the dimension of the solution space is  $n - r$ .
- ▶ So, there are exactly  $n - r$  orthogonal vectors  $\mathbf{v}_{r+1}, \mathbf{v}_{r+2}, \dots, \mathbf{v}_n$  such that  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  form an orthogonal basis of  $\mathbb{R}^n$ .

# Deriving the SVD, continued

- Use the extended basis:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{X}\mathbf{v}_{r+1} \ \cdots \ \mathbf{X}\mathbf{v}_n]$$

# Deriving the SVD, continued

- ▶ Use the extended basis:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{X}\mathbf{v}_{r+1} \ \cdots \ \mathbf{X}\mathbf{v}_n]$$

- ▶ Claim:  $\mathbf{X}\mathbf{v}_j = \mathbf{0}$ ,  $j = r + 1 : n$  (Homework: Verify it!).

# Deriving the SVD, continued

- ▶ Use the extended basis:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{X}\mathbf{v}_{r+1} \ \cdots \ \mathbf{X}\mathbf{v}_n]$$

- ▶ Claim:  $\mathbf{X}\mathbf{v}_j = \mathbf{0}$ ,  $j = r + 1 : n$  (Homework: Verify it!).
- ▶ Thus,

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{0} \ \cdots \ \mathbf{0}]$$

# Deriving the SVD, continued

- ▶ Use the extended basis:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{X}\mathbf{v}_{r+1} \ \cdots \ \mathbf{X}\mathbf{v}_n]$$

- ▶ Claim:  $\mathbf{X}\mathbf{v}_j = \mathbf{0}$ ,  $j = r + 1 : n$  (Homework: Verify it!).
- ▶ Thus,

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{0} \ \cdots \ \mathbf{0}]$$

- ▶ Let  $\mathbf{V} := [\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n]$  and  $D = \text{diag}(\sigma_1, \dots, \sigma_r)$ .

# Deriving the SVD, continued

- ▶ Use the extended basis:

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{X}\mathbf{v}_{r+1} \ \cdots \ \mathbf{X}\mathbf{v}_n]$$

- ▶ Claim:  $\mathbf{X}\mathbf{v}_j = \mathbf{0}$ ,  $j = r + 1 : n$  (Homework: Verify it!).
- ▶ Thus,

$$\mathbf{X}[\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n] = [\sigma_1 \mathbf{u}_1 \ \sigma_2 \mathbf{u}_2 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{0} \ \cdots \ \mathbf{0}]$$

- ▶ Let  $\mathbf{V} := [\mathbf{v}_1 \ \mathbf{v}_2 \ \cdots \ \mathbf{v}_n]$  and  $D = \text{diag}(\sigma_1, \dots, \sigma_r)$ .
- ▶ So,  $\mathbf{XV} = [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_r][D; \mathbf{0}_{r \times (n-r)}]$ . Thus

$$\mathbf{X} = [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_r][D; \mathbf{0}_{r \times (n-r)}]\mathbf{V}^T$$

## Deriving the SVD, continued

- ▶ Finally, to make the  $\mathbf{u}_i$  matrix a square matrix, we extend it an orthogonal basis:  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m$ .

# Deriving the SVD, continued

- ▶ Finally, to make the  $\mathbf{u}_i$  matrix a square matrix, we extend it an orthogonal basis:  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m$ .
- ▶ Define  $U := [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_m]$  and fill 0's under  $[D; \mathbf{0}_{r \times (n-r)}]$ :

$$\mathbf{X} = \mathbf{U} \begin{bmatrix} D; \mathbf{0}_{r \times (n-r)} \\ \mathbf{0}_{m-r \times n} \end{bmatrix} \mathbf{V}^T$$



# Deriving the SVD, continued

- ▶ Finally, to make the  $\mathbf{u}_i$  matrix a square matrix, we extend it an orthogonal basis:  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m$ .
- ▶ Define  $U := [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_m]$  and fill 0's under  $[D; \mathbf{0}_{r \times (n-r)}]$ :

$$\mathbf{X} = \mathbf{U} \begin{bmatrix} D; \mathbf{0}_{r \times (n-r)} \\ \mathbf{0}_{m-r \times n} \end{bmatrix} \mathbf{V}^T$$

- ▶ We are done!

# Back to column vector form

- Put  $\mathbf{U}$  and  $\mathbf{V}$  back to column vector form in SVD:

$$\mathbf{X} = \mathbf{U} \begin{bmatrix} D: \mathbf{0}_{r \times (n-r)} \\ \mathbf{0}_{m-r \times n} \end{bmatrix} \mathbf{V}^T$$

# Back to column vector form

- ▶ Put  $\mathbf{U}$  and  $\mathbf{V}$  back to column vector form in SVD:

$$\mathbf{X} = \mathbf{U} \begin{bmatrix} D: \mathbf{0}_{r \times (n-r)} \\ \mathbf{0}_{m-r \times n} \end{bmatrix} \mathbf{V}^T$$

- ▶ we have

$$\mathbf{X} = [\mathbf{u}_1 \ \cdots \ \mathbf{u}_m] \begin{bmatrix} D: \mathbf{0}_{r \times (n-r)} \\ \mathbf{0}_{m-r \times n} \end{bmatrix} \begin{bmatrix} \mathbf{v}_1^T \\ \mathbf{v}_2^T \\ \vdots \\ \mathbf{v}_n^T \end{bmatrix}$$

# Back to column vector form

- Put  $\mathbf{U}$  and  $\mathbf{V}$  back to column vector form in SVD:

$$\mathbf{X} = \mathbf{U} \begin{bmatrix} D: \mathbf{0}_{r \times (n-r)} \\ \mathbf{0}_{m-r \times n} \end{bmatrix} \mathbf{V}^T$$

- we have

$$\mathbf{X} = [\mathbf{u}_1 \ \cdots \ \mathbf{u}_m] \begin{bmatrix} D: \mathbf{0}_{r \times (n-r)} \\ \mathbf{0}_{m-r \times n} \end{bmatrix} \begin{bmatrix} \mathbf{v}_1^T \\ \mathbf{v}_2^T \\ \vdots \\ \mathbf{v}_n^T \end{bmatrix}$$

- So,

$$\mathbf{X} = [\sigma_1 \mathbf{u}_1 \ \cdots \ \sigma_r \mathbf{u}_r \ \mathbf{0} \ \cdots \ \mathbf{0}] \begin{bmatrix} \mathbf{v}_1^T \\ \mathbf{v}_2^T \\ \vdots \\ \mathbf{v}_n^T \end{bmatrix} = \sum_{i=1}^r \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

# Dimension reduction

- ▶ Assume we have a collection data of column vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$  of  $\mathbb{R}^n$ .

# Dimension reduction

- ▶ Assume we have a collection data of column vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$  of  $\mathbb{R}^n$ .
- ▶ When  $n$  is large, we want “reduce” the dimension the data.

# Dimension reduction

- ▶ Assume we have a collection data of column vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$  of  $\mathbb{R}^n$ .
- ▶ When  $n$  is large, we want “reduce” the dimension the data.
- ▶ How could this be possible?

# Dimension reduction

- ▶ Assume we have a collection data of column vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$  of  $\mathbb{R}^n$ .
- ▶ When  $n$  is large, we want “reduce” the dimension the data.
- ▶ How could this be possible?
- ▶ This is possible in the case that all data vectors are actually coming from a subspace  $U$  of small dimension (say,  $r$ ).



# Dimension reduction

- ▶ Assume we have a collection data of column vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$  of  $\mathbb{R}^n$ .
- ▶ When  $n$  is large, we want “reduce” the dimension the data.
- ▶ How could this be possible?
- ▶ This is possible in the case that all data vectors are actually coming from a subspace  $U$  of small dimension (say,  $r$ ).
- ▶ Find an orthogonal set of vectors  $\mathbf{u}_1, \dots, \mathbf{u}_r$  that forms a basis for  $U$ !

# Dimension reduction

- ▶ Assume we have a collection data of column vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$  of  $\mathbb{R}^n$ .
- ▶ When  $n$  is large, we want “reduce” the dimension the data.
- ▶ How could this be possible?
- ▶ This is possible in the case that all data vectors are actually coming from a subspace  $U$  of small dimension (say,  $r$ ).
- ▶ Find an orthogonal set of vectors  $\mathbf{u}_1, \dots, \mathbf{u}_r$  that forms a basis for  $U$ !
- ▶ If we can do so, then

$$\mathbf{a}_i = f_{i1}\mathbf{u}_1 + f_{i2}\mathbf{u}_2 + \dots + f_{ir}\mathbf{u}_r, \quad i = 1 : m$$

for some coefficients  $f_{ij}$ 's.

# Dimension reduction

- ▶ Assume we have a collection data of column vectors  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$  of  $\mathbb{R}^n$ .
- ▶ When  $n$  is large, we want “reduce” the dimension the data.
- ▶ How could this be possible?
- ▶ This is possible in the case that all data vectors are actually coming from a subspace  $U$  of small dimension (say,  $r$ ).
- ▶ Find an orthogonal set of vectors  $\mathbf{u}_1, \dots, \mathbf{u}_r$  that forms a basis for  $U$ !
- ▶ If we can do so, then

$$\mathbf{a}_i = f_{i1}\mathbf{u}_1 + f_{i2}\mathbf{u}_2 + \dots + f_{ir}\mathbf{u}_r, \quad i = 1 : m$$

for some coefficients  $f_{ij}$ 's.

- ▶ Note that  $f_{ij} = \mathbf{a}_i^T \mathbf{u}_j$ , the projection of  $\mathbf{a}_i$  along the direction of  $\mathbf{u}_j$ .

# How to find the first principal component?

- ▶ Let us find  $\mathbf{u}_1$ .

# How to find the first principal component?

- ▶ Let us find  $\mathbf{u}_1$ .
- ▶ A good choice of  $\mathbf{u}_1$ : the data vectors have the greatest total projection along the direction of  $\mathbf{u}_1$ :

$$\max_{\|\mathbf{w}\|=1} \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{w}|^2$$

# How to find the first principal component?

- ▶ Let us find  $\mathbf{u}_1$ .
- ▶ A good choice of  $\mathbf{u}_1$ : the data vectors have the greatest total projection along the direction of  $\mathbf{u}_1$ :

$$\max_{\|\mathbf{w}\|=1} \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{w}|^2$$

- ▶ Note that

$$\begin{aligned} \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{w}|^2 &= \|\mathbf{A}^T \mathbf{w}\|^2 \\ &= (\mathbf{A}^T \mathbf{w})^T (\mathbf{A}^T \mathbf{w}) = \mathbf{w}^T \mathbf{A} \mathbf{A}^T \mathbf{w} \end{aligned}$$

# How to find the first principal component?

- ▶ Let us find  $\mathbf{u}_1$ .
- ▶ A good choice of  $\mathbf{u}_1$ : the data vectors have the greatest total projection along the direction of  $\mathbf{u}_1$ :

$$\max_{\|\mathbf{w}\|=1} \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{w}|^2$$

- ▶ Note that

$$\begin{aligned} \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{w}|^2 &= \|\mathbf{A}^T \mathbf{w}\|^2 \\ &= (\mathbf{A}^T \mathbf{w})^T (\mathbf{A}^T \mathbf{w}) = \mathbf{w}^T \mathbf{A} \mathbf{A}^T \mathbf{w} \end{aligned}$$

- ▶ So,

$$\max_{\|\mathbf{w}\|=1} \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{w}|^2 = \max_{\|\mathbf{w}\|=1} \mathbf{w}^T (\mathbf{A} \mathbf{A}^T) \mathbf{w} = \max_{\mathbf{w} \neq 0} \frac{\mathbf{w}^T (\mathbf{A} \mathbf{A}^T) \mathbf{w}}{\mathbf{w}^T \mathbf{w}}$$

# Here comes the SVD

- ▶ Let us use SVD of **A**:

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$



# Here comes the SVD

- ▶ Let us use SVD of **A**:

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

- ▶ Then,

$$\mathbf{A}\mathbf{A}^T = \mathbf{U}\mathbf{D}\mathbf{V}^T\mathbf{V}\mathbf{D}^T\mathbf{U}^T = \mathbf{U}\mathbf{D}\mathbf{D}^T\mathbf{U}^T$$

# Here comes the SVD

- ▶ Let us use SVD of  $\mathbf{A}$ :

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

- ▶ Then,

$$\mathbf{A}\mathbf{A}^T = \mathbf{U}\mathbf{D}\mathbf{V}^T\mathbf{V}\mathbf{D}^T\mathbf{U}^T = \mathbf{U}\mathbf{D}\mathbf{D}^T\mathbf{U}^T$$

- ▶ and

$$\frac{\mathbf{w}^T(\mathbf{A}\mathbf{A}^T)\mathbf{w}}{\mathbf{w}^T\mathbf{w}} = \frac{(\mathbf{U}^T\mathbf{w})^T\mathbf{D}\mathbf{D}^T(\mathbf{U}^T\mathbf{w})}{(\mathbf{U}^T\mathbf{w})^T(\mathbf{U}^T\mathbf{w})}$$

# Here comes the SVD

- ▶ Let us use SVD of  $\mathbf{A}$ :

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

- ▶ Then,

$$\mathbf{A}\mathbf{A}^T = \mathbf{U}\mathbf{D}\mathbf{V}^T\mathbf{V}\mathbf{D}^T\mathbf{U}^T = \mathbf{U}\mathbf{D}\mathbf{D}^T\mathbf{U}^T$$

- ▶ and

$$\frac{\mathbf{w}^T(\mathbf{A}\mathbf{A}^T)\mathbf{w}}{\mathbf{w}^T\mathbf{w}} = \frac{(\mathbf{U}^T\mathbf{w})^T\mathbf{D}\mathbf{D}^T(\mathbf{U}^T\mathbf{w})}{(\mathbf{U}^T\mathbf{w})^T(\mathbf{U}^T\mathbf{w})}$$

- ▶ Let  $\mathbf{x} := \mathbf{U}^T\mathbf{w}$  and notice that there are only  $r$  non-zero entries  $\sigma_1, \dots, \sigma_r$  in  $\mathbf{D}$ .

# Here comes the SVD

- ▶ Let us use SVD of  $\mathbf{A}$ :

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

- ▶ Then,

$$\mathbf{A}\mathbf{A}^T = \mathbf{U}\mathbf{D}\mathbf{V}^T\mathbf{V}\mathbf{D}^T\mathbf{U}^T = \mathbf{U}\mathbf{D}\mathbf{D}^T\mathbf{U}^T$$

- ▶ and

$$\frac{\mathbf{w}^T(\mathbf{A}\mathbf{A}^T)\mathbf{w}}{\mathbf{w}^T\mathbf{w}} = \frac{(\mathbf{U}^T\mathbf{w})^T\mathbf{D}\mathbf{D}^T(\mathbf{U}^T\mathbf{w})}{(\mathbf{U}^T\mathbf{w})^T(\mathbf{U}^T\mathbf{w})}$$

- ▶ Let  $\mathbf{x} := \mathbf{U}^T\mathbf{w}$  and notice that there are only  $r$  non-zero entries  $\sigma_1, \dots, \sigma_r$  in  $\mathbf{D}$ .
- ▶ We have

$$\frac{\mathbf{w}^T(\mathbf{A}\mathbf{A}^T)\mathbf{w}}{\mathbf{w}^T\mathbf{w}} = \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \dots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \dots + x_m^2}$$

# Finally...

► We have

$$\max_{\mathbf{w} \neq 0} \frac{\mathbf{w}^T (\mathbf{A} \mathbf{A}^T) \mathbf{w}}{\mathbf{w}^T \mathbf{w}} = \max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2}$$

# Finally...

- We have

$$\max_{\mathbf{w} \neq 0} \frac{\mathbf{w}^T (\mathbf{A} \mathbf{A}^T) \mathbf{w}}{\mathbf{w}^T \mathbf{w}} = \max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2}$$

- Assume  $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r$ . Then

$$\max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2} = \sigma_1^2 = \lambda_1$$

This is the largest eigenvalue of  $A$ .

# Finally...

- ▶ We have

$$\max_{\mathbf{w} \neq 0} \frac{\mathbf{w}^T (\mathbf{A} \mathbf{A}^T) \mathbf{w}}{\mathbf{w}^T \mathbf{w}} = \max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2}$$

- ▶ Assume  $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r$ . Then

$$\max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2} = \sigma_1^2 = \lambda_1$$

This is the largest eigenvalue of  $A$ .

- ▶ A vector  $\mathbf{x}$  that makes the maximum is  $x_1 = 1$  and  $x_i = 0$  for  $i = 2 : m$

# Finally...

- ▶ We have

$$\max_{\mathbf{w} \neq 0} \frac{\mathbf{w}^T (\mathbf{A} \mathbf{A}^T) \mathbf{w}}{\mathbf{w}^T \mathbf{w}} = \max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2}$$

- ▶ Assume  $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r$ . Then

$$\max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2} = \sigma_1^2 = \lambda_1$$

This is the largest eigenvalue of  $\mathbf{A}$ .

- ▶ A vector  $\mathbf{x}$  that makes the maximum is  $x_1 = 1$  and  $x_i = 0$  for  $i = 2 : m$
- ▶ This corresponds to  $\mathbf{w} = \mathbf{U} \mathbf{x} = \mathbf{u}_1$ .



# Finally...

- ▶ We have

$$\max_{\mathbf{w} \neq 0} \frac{\mathbf{w}^T (\mathbf{A}\mathbf{A}^T) \mathbf{w}}{\mathbf{w}^T \mathbf{w}} = \max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2}$$

- ▶ Assume  $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r$ . Then

$$\max_{\mathbf{x} \neq 0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots + \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2} = \sigma_1^2 = \lambda_1$$

This is the largest eigenvalue of  $\mathbf{A}$ .

- ▶ A vector  $\mathbf{x}$  that makes the maximum is  $x_1 = 1$  and  $x_i = 0$  for  $i = 2 : m$
- ▶ This corresponds to  $\mathbf{w} = \mathbf{U}\mathbf{x} = \mathbf{u}_1$ .
- ▶ So, the first principal component is indeed achieved by the first eigenvector  $\mathbf{u}_1$  of  $\mathbf{A}\mathbf{A}^T$

# How about the 2nd, 3rd, ... principal components?

- ▶ Using the same idea, the second principal component is the direction: be orthogonal to the first and maximize the total projection:

$$\max_{\|w\|=1, w^T u_1=0} \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{w}|^2$$

# How about the 2nd, 3rd, ... principal components?

- ▶ Using the same idea, the second principal component is the direction: be orthogonal to the first and maximize the total projection:

$$\max_{\|w\|=1, w^T u_1=0} \sum_{i=1}^m |a_i^T w|^2$$

- ▶ This is the same to

$$\begin{aligned} \max_{w \neq 0, w^T u_1=0} \frac{w^T (AA^T) w}{w^T w} &= \max_{x \neq 0, x^T U^T u_1=0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2} \\ &= \max_{x \neq 0, x_1=0} \frac{\sigma_1^2 x_1^2 + \sigma_2^2 x_2^2 + \cdots \sigma_r^2 x_r^2}{x_1^2 + x_2^2 + \cdots + x_m^2} = \sigma_2^2 = \lambda_2 \end{aligned}$$

This is the largest eigenvalue of  $AA^T$ .

# Finding the 2nd, 3rd, ... principal components

- ▶ A vector  $\mathbf{x}$  that makes the maximum is  $x_2 = 1$  and  $x_i = 0$  for  $i = 1, 3 : m$

# Finding the 2nd, 3rd, ... principal components

- ▶ A vector  $\mathbf{x}$  that makes the maximum is  $x_2 = 1$  and  $x_i = 0$  for  $i = 1, 3 : m$
- ▶ This corresponds to  $\mathbf{w} = \mathbf{U}\mathbf{x} = \mathbf{u}_2$ .

# Finding the 2nd, 3rd, ... principal components

- ▶ A vector  $\mathbf{x}$  that makes the maximum is  $x_2 = 1$  and  $x_i = 0$  for  $i = 1, 3 : m$
- ▶ This corresponds to  $\mathbf{w} = \mathbf{U}\mathbf{x} = \mathbf{u}_2$ .
- ▶ So, the second principal component is indeed achieved by the second eigenvector  $\mathbf{u}_2$  of  $\mathbf{A}\mathbf{A}^T$ .

# Finding the 2nd, 3rd, ... principal components

- ▶ A vector  $\mathbf{x}$  that makes the maximum is  $x_2 = 1$  and  $x_i = 0$  for  $i = 1, 3 : m$
- ▶ This corresponds to  $\mathbf{w} = \mathbf{U}\mathbf{x} = \mathbf{u}_2$ .
- ▶ So, the second principal component is indeed achieved by the second eigenvector  $\mathbf{u}_2$  of  $\mathbf{A}\mathbf{A}^T$ .
- ▶ Now you see the pattern:  
 $\mathbf{u}_i$  is the  $i$ th principal component.

## Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).



## Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).
- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;`  
`scatter3(x,y,z,3*ones(1,p),1:p);colorbar;view(-30,10);`

# Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).
- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;`  
`scatter3(x,y,z,3*ones(1,p),1:p);colorbar;view(-30,10);`
- ▶ We want to find the first principal component.

# Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).
- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;`  
`scatter3(x,y,z,3*ones(1,p),1:p);colorbar;view(-30,10);`
- ▶ We want to find the first principal component.
- ▶ First, we center the data by remove the mean of all points.

# Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).
- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;  
scatter3(x,y,z,3*ones(1,p),1:p);colorbar;view(-30,10);`
- ▶ We want to find the first principal component.
- ▶ First, we center the data by remove the mean of all points.
- ▶ `x=x-mean(x);y=y-mean(y);z=z-mean(z);hold  
on;scatter3(x,y,z,3*ones(1,p),1:p);`

# Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).
- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;  
scatter3(x,y,z,3*ones(1,p),1:p);colorbar;view(-30,10);`
- ▶ We want to find the first principal component.
- ▶ First, we center the data by remove the mean of all points.
- ▶ `x=x-mean(x);y=y-mean(y);z=z-mean(z);hold  
on;scatter3(x,y,z,3*ones(1,p),1:p);`
- ▶ Now, we use PCA to find the principal direction of these points

# Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).
- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;`  
`scatter3(x,y,z,3*ones(1,p),1:p);colorbar;view(-30,10);`
- ▶ We want to find the first principal component.
- ▶ First, we center the data by remove the mean of all points.
- ▶ `x=x-mean(x);y=y-mean(y);z=z-mean(z);hold`  
`on;scatter3(x,y,z,3*ones(1,p),1:p);`
- ▶ Now, we use PCA to find the principal direction of these points
- ▶  
`A=[x;y;z];[U,S,V]=svds(A);scatter3(x,y,z,3*ones(1,p),U(:,1)'*A);`

# Sample code

- ▶ Let us randomly choose 20 points on a line in space (3-dimension).
- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;`  
`scatter3(x,y,z,3*ones(1,p),1:p);colorbar;view(-30,10);`
- ▶ We want to find the first principal component.
- ▶ First, we center the data by remove the mean of all points.
- ▶ `x=x-mean(x);y=y-mean(y);z=z-mean(z);hold`  
`on;scatter3(x,y,z,3*ones(1,p),1:p);`
- ▶ Now, we use PCA to find the principal direction of these points
- ▶  
`A=[x;y;z];[U,S,V]=svds(A);scatter3(x,y,z,3*ones(1,p),U(:,1)'*A);`
- ▶ Color is according to x-projection


# The m-file and plot

```
► p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;  
scatter3(x,y,z,3*ones(1,p),2*[1:p]/p-1);view(-30,10);  
x=x-mean(x);y=y-mean(y);z=z-mean(z);  
A=[x;y;z];[U,S,V]=svds(A);  
hold on;  
scatter3(x,y,z,3*ones(1,p),U(:,1)'*A);
```



# The m-file and plot

- ▶ `p=100;r=rand(1,p);x=2*r-1;y=r+2;z=3-r;`  
`scatter3(x,y,z,3*ones(1,p),2*[1:p]/p-1);view(-30,10);`  
`x=x-mean(x);y=y-mean(y);z=z-mean(z);`  
`A=[x;y;z];[U,S,V]=svds(A);`  
`hold on;`  
`scatter3(x,y,z,3*ones(1,p),U(:,1)'*A);`



# Images as column vectors

- ▶ Assume we have a set of  $n$  images of the same size (say,  $p \times q$ ):  $\mathbf{l}_1, \mathbf{l}_2, \dots, \mathbf{l}_n$ .

# Images as column vectors

- ▶ Assume we have a set of  $n$  images of the same size (say,  $p \times q$ ):  $\mathbf{I}_1, \mathbf{I}_2, \dots, \mathbf{I}_n$ .
- ▶ We transform the  $p \times q$  image matrix  $\mathbf{I}_i$  to a column vector of size  $m := pq$  by stacking its columns. For simplicity, we use the same notation to denote the column vector.

# Images as column vectors

- ▶ Assume we have a set of  $n$  images of the same size (say,  $p \times q$ ):  $\mathbf{l}_1, \mathbf{l}_2, \dots, \mathbf{l}_n$ .
- ▶ We transform the  $p \times q$  image matrix  $\mathbf{l}_i$  to a column vector of size  $m := pq$  by stacking its columns. For simplicity, we use the same notation to denote the column vector.
- ▶ Now, as a standard step, we take out the mean image  $\bar{\mathbf{l}} := \frac{1}{n} \sum_{i=1}^n \mathbf{l}_i$  from every image:

$$\tilde{\mathbf{l}}_i := \mathbf{l}_i - \bar{\mathbf{l}}, \quad i = 1 : n$$

This step guarantees  $\tilde{\mathbf{l}}_i$  are centered at the origin (and hence contained in a true **subspace**).

# Images as column vectors

- ▶ Assume we have a set of  $n$  images of the same size (say,  $p \times q$ ):  $\mathbf{l}_1, \mathbf{l}_2, \dots, \mathbf{l}_n$ .
- ▶ We transform the  $p \times q$  image matrix  $\mathbf{l}_i$  to a column vector of size  $m := pq$  by stacking its columns. For simplicity, we use the same notation to denote the column vector.
- ▶ Now, as a standard step, we take out the mean image  $\bar{\mathbf{l}} := \frac{1}{n} \sum_{i=1}^n \mathbf{l}_i$  from every image:

$$\tilde{\mathbf{l}}_i := \mathbf{l}_i - \bar{\mathbf{l}}, \quad i = 1 : n$$

This step guarantees  $\tilde{\mathbf{l}}_i$  are centered at the origin (and hence contained in a true **subspace**).

- ▶ Form the  $m \times n$  data matrix

$$X := [\tilde{\mathbf{l}}_1 \quad \tilde{\mathbf{l}}_2 \quad \cdots \quad \tilde{\mathbf{l}}_n]$$