

# Complete solution to $Ax = b$

Rohit Budhiraja, IITK

Applied Linear Algebra for Wireless Communications

# Recap and agenda for today's class

- Discussed the following in the last lecture
  - Systematically calculated  $N(A)$  of matrix  $A$

# Recap and agenda for today's class

- Discussed the following in the last lecture
  - Systematically calculated  $N(A)$  of matrix  $A$
- Discuss the following today
  - Systematically calculate complete solution of  $A\mathbf{x} = \mathbf{b}$

# Complete Solution to $A\mathbf{x} = \mathbf{b}$ <sup>1</sup>

- Recall while calculating elimination converted  $A\mathbf{x} = \mathbf{b}$  to  $R\mathbf{x} = \mathbf{b}$

---

<sup>1</sup>Chap 3.4 of the book

# Complete Solution to $A\mathbf{x} = \mathbf{b}$ <sup>1</sup>

- Recall while calculating elimination converted  $A\mathbf{x} = 0$  to  $R\mathbf{x} = 0$ 
  - Free variables were given special values (one and zero)

---

<sup>1</sup>Chap 3.4 of the book

# Complete Solution to $A\mathbf{x} = \mathbf{b}$ <sup>1</sup>

- Recall while calculating elimination converted  $A\mathbf{x} = 0$  to  $R\mathbf{x} = 0$ 
  - Free variables were given special values (one and zero)
  - Pivot variables were found by back substitution

---

<sup>1</sup>Chap 3.4 of the book

# Complete Solution to $A\mathbf{x} = \mathbf{b}$ <sup>1</sup>

- Recall while calculating elimination converted  $A\mathbf{x} = \mathbf{0}$  to  $R\mathbf{x} = \mathbf{0}$ 
  - Free variables were given special values (one and zero)
  - Pivot variables were found by back substitution
  - Solution  $\mathbf{x}$  was in the nullspace of  $A$

---

<sup>1</sup>Chap 3.4 of the book

# Complete Solution to $A\mathbf{x} = \mathbf{b}$ <sup>1</sup>

- Recall while calculating elimination converted  $A\mathbf{x} = 0$  to  $R\mathbf{x} = 0$ 
  - Free variables were given special values (one and zero)
  - Pivot variables were found by back substitution
  - Solution  $\mathbf{x}$  was in the nullspace of  $A$
- We paid no attention to the right side  $\mathbf{b}$  because it stayed at zero

---

<sup>1</sup>Chap 3.4 of the book



# Complete Solution to $A\mathbf{x} = \mathbf{b}$ <sup>1</sup>

- Recall while calculating elimination converted  $A\mathbf{x} = 0$  to  $R\mathbf{x} = 0$ 
  - Free variables were given special values (one and zero)
  - Pivot variables were found by back substitution
  - Solution  $\mathbf{x}$  was in the nullspace of  $A$
- We paid no attention to the right side  $\mathbf{b}$  because it stayed at zero
- Now  $\mathbf{b}$  is not zero. Row operations on left side must act also on the right side

---

<sup>1</sup>Chap 3.4 of the book

# Complete Solution to $A\mathbf{x} = \mathbf{b}$ <sup>1</sup>

- Recall while calculating elimination converted  $A\mathbf{x} = 0$  to  $R\mathbf{x} = 0$ 
  - Free variables were given special values (one and zero)
  - Pivot variables were found by back substitution
  - Solution  $\mathbf{x}$  was in the nullspace of  $A$
- We paid no attention to the right side  $\mathbf{b}$  because it stayed at zero
- Now  $\mathbf{b}$  is not zero. Row operations on left side must act also on the right side
- One way is to create an augmented matrix  $[A \ \mathbf{b}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 1 & 3 & 1 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 7 \end{bmatrix} \quad \text{has the augmented matrix} \quad \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 1 & 3 & 1 & 6 & 7 \end{bmatrix} = [A \ \mathbf{b}]$$

---

<sup>1</sup>Chap 3.4 of the book

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3
- Last zero is crucial, third equation has become  $0 = 0$

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3
- Last zero is crucial, third equation has become  $0 = 0$ 
  - Equations are consistent on LHS and RHS, and hence can be solved

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3
- Last zero is crucial, third equation has become  $0 = 0$ 
  - Equations are consistent on LHS and RHS, and hence can be solved
- For a particular solution  $\mathbf{x}_p$ , choose free variables as zero:  $x_2 = x_4 = 0$



# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3
- Last zero is crucial, third equation has become  $0 = 0$ 
  - Equations are consistent on LHS and RHS, and hence can be solved
- For a particular solution  $\mathbf{x}_p$ , choose free variables as zero:  $x_2 = x_4 = 0$ 
  - Two nonzero equations give two pivot variables  $x_1 = 1$  and  $x_3 = 6$

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3
- Last zero is crucial, third equation has become  $0 = 0$ 
  - Equations are consistent on LHS and RHS, and hence can be solved
- For a particular solution  $\mathbf{x}_p$ , choose free variables as zero:  $x_2 = x_4 = 0$ 
  - Two nonzero equations give two pivot variables  $x_1 = 1$  and  $x_3 = 6$
  - Our particular solution to  $A\mathbf{x} = \mathbf{b}$  (and also  $R\mathbf{x} = \mathbf{d}$ ) is  $\mathbf{x}_p = (1, 0, 6, 0)$

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3
- Last zero is crucial, third equation has become  $0 = 0$ 
  - Equations are consistent on LHS and RHS, and hence can be solved
- For a particular solution  $\mathbf{x}_p$ , choose free variables as zero:  $x_2 = x_4 = 0$ 
  - Two nonzero equations give two pivot variables  $x_1 = 1$  and  $x_3 = 6$
  - Our particular solution to  $A\mathbf{x} = \mathbf{b}$  (and also  $R\mathbf{x} = \mathbf{d}$ ) is  $\mathbf{x}_p = (1, 0, 6, 0)$
- Two nullspace solutions to  $R\mathbf{x} = \mathbf{0}$  come from two free columns of  $R$

# One Particular Solution $A\mathbf{x}_p = \mathbf{b}$

- Apply elimination steps to  $[A \ \mathbf{b}]$ , system reduces to  $[R \ \mathbf{x}]$

$$\begin{bmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \text{ has the augmented matrix } \begin{bmatrix} 1 & 3 & 0 & 2 & 1 \\ 0 & 0 & 1 & 4 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} = [R \ \mathbf{d}].$$

- In this example we subtract row 1 from 3, and the row 2 from 3
- Last zero is crucial, third equation has become  $0 = 0$ 
  - Equations are consistent on LHS and RHS, and hence can be solved
- **For a particular solution  $\mathbf{x}_p$** , choose free variables as zero:  $x_2 = x_4 = 0$ 
  - Two nonzero equations give two pivot variables  $x_1 = 1$  and  $x_3 = 6$
  - Our particular solution to  $A\mathbf{x} = \mathbf{b}$  (and also  $R\mathbf{x} = \mathbf{d}$ ) is  $\mathbf{x}_p = (1, 0, 6, 0)$
- **Two nullspace solutions to  $R\mathbf{x} = \mathbf{0}$**  come from two free columns of  $R$ 
  - By reversing signs of 3, 2, and 4 which gives  $(-3, 1, 0, 0)$  and  $(-2, 0, -4, 1)$

# Complete Solution to $Ax = b$

- Two solutions are given as

$x_{\text{particular}}$

*The particular solution solves*

$$Ax_p = b$$

$x_{\text{nullspace}}$

*The  $n - r$  special solutions solve*

$$Ax_n = 0.$$

# Complete Solution to $Ax = b$

- Two solutions are given as

$x_{\text{particular}}$

*The particular solution solves*

$$Ax_p = b$$

$x_{\text{nullspace}}$

*The  $n - r$  special solutions solve*

$$Ax_n = 0.$$

- Complete solution  $x_p + x_n$  to  $Ax = b$

Complete solution  
one  $x_p$   
many  $x_n$

$$x = x_p + x_n = \begin{bmatrix} 1 \\ 0 \\ 6 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -2 \\ 0 \\ -4 \\ 1 \end{bmatrix}.$$

# Example1 to find complete solution to $Ax = b$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

# Example1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable



## Example1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable
- Create augmented matrix  $[A \ \mathbf{b}]$  and apply elimination

## Example 1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable
- Create augmented matrix  $[A \ \mathbf{b}]$  and apply elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix}$$

# Example 1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable
- Create augmented matrix  $[A \ \mathbf{b}]$  and apply elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix}$$

# Example1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable
- Create augmented matrix  $[A \ \mathbf{b}]$  and apply elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

## Example 1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable
- Create augmented matrix  $[A \ \mathbf{b}]$  and apply elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

- i) Subtracted row 1 of  $[A \ \mathbf{b}]$  from row 2; ii) Added 2 times row 1 to row 3

## Example 1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable
- Create augmented matrix  $[A \ \mathbf{b}]$  and apply elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

- i) Subtracted row 1 of  $[A \ \mathbf{b}]$  from row 2; ii) Added 2 times row 1 to row 3
- Last equation is  $0 = 0$  provided  $b_3 + b_1 + b_2$

# Example1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider an example system of equations

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

- Find condition on  $(b_1, b_2, b_3)$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable
- Create augmented matrix  $[A \ \mathbf{b}]$  and apply elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

- i) Subtracted row 1 of  $[A \ \mathbf{b}]$  from row 2; ii) Added 2 times row 1 to row 3
- Last equation is  $0 = 0$  provided  $b_3 + b_1 + b_2$ 
  - $A\mathbf{x} = \mathbf{b}$  will be solvable

## Example1 to find complete solution to $Ax = b$ (2)

- Augmented matrix  $[A \ b]$  after elimination



## Example1 to find complete solution to $Ax = b$ (2)

- Augmented matrix  $[A \ b]$  after elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

## Example1 to find complete solution to $Ax = b$ (2)

- Augmented matrix  $[A \ b]$  after elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

- Example has no free variables as  $n - r = 2 - 2$ , nullspace solution is  $\mathbf{x}_n = 0$

## Example1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (2)

- Augmented matrix  $[A \ \mathbf{b}]$  after elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

- Example has no free variables as  $n - r = 2 - 2$ , nullspace solution is  $\mathbf{x}_n = \mathbf{0}$
- Particular solution to  $A\mathbf{x} = \mathbf{b}$  and  $R\mathbf{x} = \mathbf{d}$  is at the top of the final column  $\mathbf{d}$

## Example1 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (2)

- Augmented matrix  $[A \ \mathbf{b}]$  after elimination

$$\begin{bmatrix} 1 & 1 & b_1 \\ 1 & 2 & b_2 \\ -2 & -3 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & b_1 \\ 0 & 1 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2b_1 - b_2 \\ 0 & 1 & b_2 - b_1 \\ 0 & 0 & b_3 + b_1 + b_2 \end{bmatrix}$$

- Example has no free variables as  $n - r = 2 - 2$ , nullspace solution is  $\mathbf{x}_n = \mathbf{0}$
- Particular solution to  $A\mathbf{x} = \mathbf{b}$  and  $R\mathbf{x} = \mathbf{d}$  is at the top of the final column  $\mathbf{d}$
- Complete solution is

$$\mathbf{x} = \mathbf{x}_p + \mathbf{x}_n = \begin{bmatrix} 2b_1 - b_2 \\ b_2 - b_1 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

## Example 1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

## Example 1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ),

## Example 1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ), every column has a pivot

## Example 1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ), every column has a pivot  $\Rightarrow$  rank is  $r = n$



## Example1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ), every column has a pivot  $\Rightarrow$  rank is  $r = n$
- Every matrix  $A$  with full column rank ( $r = n$ ) has all these properties:

## Example 1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ), every column has a pivot  $\Rightarrow$  rank is  $r = n$
- Every matrix  $A$  with full column rank ( $r = n$ ) has all these properties:
  - All columns of  $A$  are pivot columns

## Example1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ), every column has a pivot  $\Rightarrow$  rank is  $r = n$
- Every matrix  $A$  with full column rank ( $r = n$ ) has all these properties:
  - All columns of  $A$  are pivot columns
  - There are no free variables or special solutions

## Example1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ), every column has a pivot  $\Rightarrow$  rank is  $r = n$
- Every matrix  $A$  with full column rank ( $r = n$ ) has all these properties:
  - All columns of  $A$  are pivot columns
  - There are no free variables or special solutions
  - Nullspace  $N(A)$  contains only the zero vector  $\mathbf{x} = 0$

## Example1 to find complete solution to $Ax = b$ (3)

- Earlier example of  $A$  is extremely important as  $A$  has full column rank

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & -3 \end{bmatrix}$$

- Matrix is tall and thin ( $m > n$ ), every column has a pivot  $\Rightarrow$  rank is  $r = n$
- Every matrix  $A$  with full column rank ( $r = n$ ) has all these properties:
  - All columns of  $A$  are pivot columns
  - There are no free variables or special solutions
  - Nullspace  $N(A)$  contains only the zero vector  $\mathbf{x} = 0$
  - If  $Ax = \mathbf{b}$  has a solution (it might not) then it has only one solution

## Example2 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider another system  $A\mathbf{x} = \mathbf{b}$

$$x + y + z = 3$$

$$x + 2y - z = 4$$

- Has  $n = 3$  unknowns but only  $m = 2$  equations

## Example2 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider another system  $A\mathbf{x} = \mathbf{b}$

$$x + y + z = 3$$

$$x + 2y - z = 4$$

- Has  $n = 3$  unknowns but only  $m = 2$  equations
- Find  $\mathbf{x}_p$  and  $\mathbf{x}_n$  by elimination on  $[A \ \mathbf{b}]$

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

## Example2 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider another system  $A\mathbf{x} = \mathbf{b}$

$$x + y + z = 3$$

$$x + 2y - z = 4$$

- Has  $n = 3$  unknowns but only  $m = 2$  equations
- Find  $\mathbf{x}_p$  and  $\mathbf{x}_n$  by elimination on  $[A \ \mathbf{b}]$

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

- Particular solution has free variable  $x_3 = 0$  and special solution has  $x_3 = 1$



## Example2 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider another system  $A\mathbf{x} = \mathbf{b}$

$$x + y + z = 3$$

$$x + 2y - z = 4$$

- Has  $n = 3$  unknowns but only  $m = 2$  equations
- Find  $\mathbf{x}_p$  and  $\mathbf{x}_n$  by elimination on  $[A \ \mathbf{b}]$

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

- Particular solution has free variable  $x_3 = 0$  and special solution has  $x_3 = 1$ 
  - $\mathbf{x}_p = (2, 1, 0)$  comes directly from  $\mathbf{d}$
  - $\mathbf{x}_n = (-3, 2, 1)$  comes from third (free) column

## Example2 to find complete solution to $A\mathbf{x} = \mathbf{b}$ (1)

- Consider another system  $A\mathbf{x} = \mathbf{b}$

$$x + y + z = 3$$

$$x + 2y - z = 4$$

- Has  $n = 3$  unknowns but only  $m = 2$  equations
- Find  $\mathbf{x}_p$  and  $\mathbf{x}_n$  by elimination on  $[A \ \mathbf{b}]$

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

- Particular solution has free variable  $x_3 = 0$  and special solution has  $x_3 = 1$ 
  - $\mathbf{x}_p = (2, 1, 0)$  comes directly from  $\mathbf{d}$
  - $\mathbf{x}_n = (-3, 2, 1)$  comes from third (free) column

$$\mathbf{x} = \mathbf{x}_p + \mathbf{x}_n$$



## Example2 to find complete solution to $Ax = b$ (2)

- We have the following system of equation from last slide

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

## Example2 to find complete solution to $Ax = b$ (2)

- We have the following system of equation from last slide

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

- Every matrix  $A$  with full row rank ( $r = m$ ) has all these properties:

## Example2 to find complete solution to $Ax = b$ (2)

- We have the following system of equation from last slide

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ d]$$

- Every matrix  $A$  with full row rank ( $r = m$ ) has all these properties:
  - All rows have pivots, and  $R$  has no zero rows

## Example2 to find complete solution to $Ax = b$ (2)

- We have the following system of equation from last slide

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

- Every matrix  $A$  with full row rank ( $r = m$ ) has all these properties:
  - All rows have pivots, and  $R$  has no zero rows
  - $Ax = \mathbf{b}$  has a solution for every right side  $\mathbf{b}$

## Example2 to find complete solution to $Ax = b$ (2)

- We have the following system of equation from last slide

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

- Every matrix  $A$  with full row rank ( $r = m$ ) has all these properties:
  - All rows have pivots, and  $R$  has no zero rows
  - $Ax = \mathbf{b}$  has a solution for every right side  $\mathbf{b}$
  - Column space is the whole space  $\mathbf{R}^m$

## Example2 to find complete solution to $Ax = b$ (2)

- We have the following system of equation from last slide

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & -1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 3 \\ 0 & 1 & -2 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 3 & 2 \\ 0 & 1 & -2 & 1 \end{bmatrix} = [R \ \mathbf{d}]$$

- Every matrix  $A$  with full row rank ( $r = m$ ) has all these properties:
  - All rows have pivots, and  $R$  has no zero rows
  - $Ax = \mathbf{b}$  has a solution for every right side  $\mathbf{b}$
  - Column space is the whole space  $\mathbf{R}^m$
  - There are  $n - r = n - m$  special solutions in the nullspace of  $A$