

# Null Spaces and Solution of Linear Equations

Rohit Budhiraja, IITK

Applied Linear Algebra for Wireless Communications

# Recap and agenda for today's class

- Discussed the following in the last lecture
  - Vector spaces and subspaces
  - Null space of matrix  $A$  i.e.,  $N(A)$  and column space

# Recap and agenda for today's class

- Discussed the following in the last lecture
  - Vector spaces and subspaces
  - Null space of matrix  $A$  i.e.,  $N(A)$  and column space
- Discuss the following today
  - $N(A)$  and solution of  $A\mathbf{x} = \mathbf{b}$

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ .

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

- Both columns have pivots i.e., no special solutions

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

- Both columns have pivots i.e., no special solutions
  - $N(A) = 0$  – A is invertible

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

- Both columns have pivots i.e., no special solutions
  - $N(A) = 0$  – A is invertible
- Rank of a matrix  $r$  is the number of pivots – rank of A is 2



# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

- Both columns have pivots i.e., no special solutions
  - $N(A) = 0$  – A is invertible
- Rank of a matrix  $r$  is the number of pivots – rank of A is 2
- Number of pivots (i.e., rank  $r$ ) gives the number of independent columns

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

- Both columns have pivots i.e., no special solutions
  - $N(A) = 0$  – A is invertible
- Rank of a matrix  $r$  is the number of pivots – rank of A is 2
- Number of pivots (i.e., rank  $r$ ) gives the number of independent columns
- Case of a zero nullspace is of the greatest importance

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

- Both columns have pivots i.e., no special solutions
  - $N(A) = 0$  – A is invertible
- Rank of a matrix  $r$  is the number of pivots – rank of A is 2
- Number of pivots (i.e., rank  $r$ ) gives the number of independent columns
- Case of a zero nullspace is of the greatest importance
  - It says that All columns have pivots, and they are independent

# Null space of matrix

- Find the nullspaces of  $A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$  and  $C = [A \ 2A]$ . Use elimination

$$Ax = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} x_1 = 0 \\ x_2 = 0 \end{bmatrix}$$

- Both columns have pivots i.e., no special solutions
  - $N(A) = 0$  – A is invertible
- Rank of a matrix  $r$  is the number of pivots – rank of A is 2
- Number of pivots (i.e., rank  $r$ ) gives the number of independent columns
- Case of a zero nullspace is of the greatest importance
  - It says that All columns have pivots, and they are independent
  - No combination of columns gives zero vector (except zero combination)

# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3 (row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

↑ ↑ ↑ ↑  
pivot columns free columns

# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$

pivot columns    free columns

- Elimination produces pivots in the first two columns of  $C$  (**Rank of  $C$  is 2**)

# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$

pivot columns    free columns

- Elimination produces pivots in the first two columns of  $C$  (**Rank of  $C$  is 2**)
  - Number of independent columns is 2

# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$

pivot columns    free columns

- Elimination produces pivots in the first two columns of  $C$  (**Rank of  $C$  is 2**)
  - Number of independent columns is 2
- Last two columns of  $C$  and  $U$  are "free" – they don't have pivots



# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$

pivot columns    free columns

- Elimination produces pivots in the first two columns of  $C$  (**Rank of  $C$  is 2**)
  - Number of independent columns is 2
- Last two columns of  $C$  and  $U$  are "free" – they don't have pivots
  - We get two special solutions in  $N(C)$ , correspond to free variables  $x_3$  and  $x_4$

# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$

pivot columns      free columns

- Elimination produces pivots in the first two columns of  $C$  (**Rank of  $C$  is 2**)
  - Number of independent columns is 2
- Last two columns of  $C$  and  $U$  are "free" – they don't have pivots
  - We get two special solutions in  $N(C)$ , correspond to free variables  $x_3$  and  $x_4$
  - To calculate them, we first set  $x_3 = 1, x_4 = 0$

# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$

pivot columns    free columns

- Elimination produces pivots in the first two columns of  $C$  (**Rank of  $C$  is 2**)
  - Number of independent columns is 2
- Last two columns of  $C$  and  $U$  are "free" – they don't have pivots
  - We get two special solutions in  $N(C)$ , correspond to free variables  $x_3$  and  $x_4$
  - To calculate them, we first set  $x_3 = 1, x_4 = 0$  and second set  $x_3 = 0, x_4 = 1$

# Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$

pivot columns    free columns

- Elimination produces pivots in the first two columns of  $C$  (**Rank of  $C$  is 2**)
  - Number of independent columns is 2
- Last two columns of  $C$  and  $U$  are "free" – they don't have pivots
  - We get two special solutions in  $N(C)$ , correspond to free variables  $x_3$  and  $x_4$
  - To calculate them, we first set  $x_3 = 1, x_4 = 0$  and second set  $x_3 = 0, x_4 = 1$
- Pivot variables  $x_1$  and  $x_2$  are determined by **back substituting in**  $U\mathbf{x} = 0$

## Pivot Columns, Free Columns And Special Solutions

- We will find  $N(C)$  where  $C = [A \ 2A]$

Subtract 3(row 1)  
from row 2 of  $C$

$$C = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 3 & 8 & 6 & 16 \end{bmatrix} \text{ becomes } U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow$   
 pivot columns    free columns

- Elimination produces pivots in the first two columns of  $C$  (Rank of  $C$  is 2)
  - Number of independent columns is 2
- Last two columns of  $C$  and  $U$  are "free" – they don't have pivots
  - We get two special solutions in  $N(C)$ , correspond to free variables  $x_3$  and  $x_4$
  - To calculate them, we first set  $x_3 = 1, x_4 = 0$  and second set  $x_3 = 0, x_4 = 1$
- Pivot variables  $x_1$  and  $x_2$  are determined by back substituting in  $U\mathbf{x} = 0$

<p><b>Special solutions</b></p> <p><math>Cs = 0</math></p> <p><math>Us = 0</math></p>	$s_1 = \begin{bmatrix} -2 \\ 0 \\ 1 \\ 0 \end{bmatrix}$	and $s_2 = \begin{bmatrix} 0 \\ -2 \\ 0 \\ 1 \end{bmatrix}$	<p>← <b>pivot</b></p> <p>← <b>variables</b></p> <p>← <b>free</b></p> <p>← <b>variables</b></p>
---	---	---	--

# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$

# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$ 
  - We can make this  $U$  simpler by reducing it to row echelon form  $R$

# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$ 
  - We can make this  $U$  simpler by reducing it to row echelon form  $R$

Reduced  
form  $R$

$$U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix} \quad \text{becomes} \quad R = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}.$$

$\uparrow \quad \uparrow$



# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$ 
  - We can make this  $U$  simpler by reducing it to row echelon form  $R$

Reduced  
form  $R$

$$U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix} \quad \text{becomes} \quad R = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}.$$

$\uparrow \quad \uparrow$

- Those steps don't change the zero vector on the right side of the equation

# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$ 
  - We can make this  $U$  simpler by reducing it to row echelon form  $R$

Reduced  
form  $R$

$$U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix} \quad \text{becomes} \quad R = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}.$$

$\uparrow \quad \uparrow$

- Those steps don't change the zero vector on the right side of the equation
- Pivot columns of  $R$  contain  $I$ . Nullspace reach the  $N(A) = N(U) = N(R)$

1. *Produce zeros above the pivots.* Use pivot rows to eliminate upward in  $R$ .
2. *Produce ones in the pivots.* Divide the whole pivot row by its pivot.

# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$ 
  - We can make this  $U$  simpler by reducing it to row echelon form  $R$

Reduced  
form  $R$

$$U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix} \quad \text{becomes} \quad R = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}.$$

$\uparrow \quad \uparrow$

- Those steps don't change the zero vector on the right side of the equation
- Pivot columns of  $R$  contain  $I$ . Nullspace reach the  $N(A) = N(U) = N(R)$

1. *Produce zeros above the pivots.* Use pivot rows to eliminate upward in  $R$ .
2. *Produce ones in the pivots.* Divide the whole pivot row by its pivot.

- Special solutions are much easier to find from the reduced system  $R\mathbf{x} = 0$

# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$ 
  - We can make this  $U$  simpler by reducing it to row echelon form  $R$

Reduced  
form  $R$

$$U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix} \quad \text{becomes} \quad R = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}.$$

$\uparrow \quad \uparrow$

- Those steps don't change the zero vector on the right side of the equation
- Pivot columns of  $R$  contain  $I$ . Nullspace reach the  $N(A) = N(U) = N(R)$

1. *Produce zeros above the pivots.* Use pivot rows to eliminate upward in  $R$ .
2. *Produce ones in the pivots.* Divide the whole pivot row by its pivot.

- Special solutions are much easier to find from the reduced system  $R\mathbf{x} = 0$ 
  - In each free column of  $R$  change all the signs to find  $\mathbf{s}$

# Reduced Row Echelon Form R

- When  $A$  is rectangular, elimination will not stop at the upper triangular  $U$ 
  - We can make this  $U$  simpler by reducing it to row echelon form  $R$

Reduced  
form  $R$

$$U = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 0 & 4 \end{bmatrix} \text{ becomes } R = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}.$$

$\uparrow \quad \uparrow$

- Those steps don't change the zero vector on the right side of the equation
- Pivot columns of  $R$  contain  $I$ . Nullspace reach the  $N(A) = N(U) = N(R)$

1. *Produce zeros above the pivots.* Use pivot rows to eliminate upward in  $R$ .

2. *Produce ones in the pivots.* Divide the whole pivot row by its pivot.

- Special solutions are much easier to find from the reduced system  $R\mathbf{x} = 0$ 
  - In each free column of  $R$  change all the signs to find  $\mathbf{s}$
  - Two special solutions are  $\mathbf{s}_1 = (-2, 0, 1, 0)$  and  $\mathbf{s}_2 = (0, -2, 0, 1)$

# Dimensionality of nullspace

- Counting the pivots leads to an extremely important theorem

# Dimensionality of nullspace

- Counting the pivots leads to an extremely important theorem

$$A = \begin{bmatrix} p & p & f & p & f \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix} \quad R = \begin{bmatrix} 1 & 0 & a & 0 & c \\ 0 & 1 & b & 0 & d \\ 0 & 0 & 0 & 1 & e \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad s_1 = \begin{bmatrix} -a \\ -b \\ 1 \\ 0 \\ 0 \end{bmatrix} \quad s_2 = \begin{bmatrix} -c \\ -d \\ 0 \\ -e \\ 1 \end{bmatrix}$$

3 pivot columns  $p$        $I$  in pivot columns      special  $Rs_1 = \mathbf{0}$  and  $Rs_2 = \mathbf{0}$   
2 free columns  $f$        $F$  in free columns      take  $-a$  to  $-e$  from  $R$   
to be revealed by  $R$       3 pivots: rank  $r = 3$        $Rs = \mathbf{0}$  means  $As = \mathbf{0}$

# Dimensionality of nullspace

- Counting the pivots leads to an extremely important theorem

$$A = \left[ \begin{array}{c|c|c|c|c} p & p & f & p & f \\ \hline \vdots & \vdots & \vdots & \vdots & \vdots \end{array} \right] \quad R = \left[ \begin{array}{ccccc} 1 & 0 & a & 0 & c \\ 0 & 1 & b & 0 & d \\ 0 & 0 & 0 & 1 & e \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad s_1 = \begin{bmatrix} -a \\ -b \\ 1 \\ 0 \\ 0 \end{bmatrix} \quad s_2 = \begin{bmatrix} -c \\ -d \\ 0 \\ -e \\ 1 \end{bmatrix}$$

3 pivot columns  $p$        $I$  in pivot columns      special  $Rs_1 = \mathbf{0}$  and  $Rs_2 = \mathbf{0}$   
2 free columns  $f$        $F$  in free columns      take  $-a$  to  $-e$  from  $R$   
to be revealed by  $R$       3 pivots: rank  $r = 3$        $Rs = \mathbf{0}$  means  $As = \mathbf{0}$

- Let  $A$  has more columns than rows ( $n > m$ ), which is a short wide matrix



# Dimensionality of nullspace

- Counting the pivots leads to an extremely important theorem

$$A = \begin{bmatrix} p & p & f & p & f \\ | & | & | & | & | \end{bmatrix} \quad R = \begin{bmatrix} 1 & 0 & a & 0 & c \\ 0 & 1 & b & 0 & d \\ 0 & 0 & 0 & 1 & e \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad s_1 = \begin{bmatrix} -a \\ -b \\ 1 \\ 0 \\ 0 \end{bmatrix} \quad s_2 = \begin{bmatrix} -c \\ -d \\ 0 \\ -e \\ 1 \end{bmatrix}$$

3 pivot columns  $p$        $I$  in pivot columns      special  $Rs_1 = \mathbf{0}$  and  $Rs_2 = \mathbf{0}$   
2 free columns  $f$        $F$  in free columns      take  $-a$  to  $-e$  from  $R$   
to be revealed by  $R$       3 pivots: rank  $r = 3$        $Rs = \mathbf{0}$  means  $As = \mathbf{0}$

- Let  $A$  has more columns than rows ( $n > m$ ), which is a short wide matrix
- Must have least  $n - m$  free variables, since number of pivots cannot exceed  $m$

# Dimensionality of nullspace

- Counting the pivots leads to an extremely important theorem

$$A = \begin{bmatrix} p & p & f & p & f \\ | & | & | & | & | \\ | & | & | & | & | \\ | & | & | & | & | \end{bmatrix} \quad R = \begin{bmatrix} 1 & 0 & a & 0 & c \\ 0 & 1 & b & 0 & d \\ 0 & 0 & 0 & 1 & e \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad s_1 = \begin{bmatrix} -a \\ -b \\ 1 \\ 0 \\ 0 \end{bmatrix} \quad s_2 = \begin{bmatrix} -c \\ -d \\ 0 \\ -e \\ 1 \end{bmatrix}$$

3 pivot columns  $p$        $I$  in pivot columns      special  $Rs_1 = \mathbf{0}$  and  $Rs_2 = \mathbf{0}$   
2 free columns  $f$        $F$  in free columns      take  $-a$  to  $-e$  from  $R$   
to be revealed by  $R$       3 pivots: rank  $r = 3$        $Rs = \mathbf{0}$  means  $As = \mathbf{0}$

- Let  $A$  has more columns than rows ( $n > m$ ), which is a short wide matrix
- Must have least  $n - m$  free variables, since number of pivots cannot exceed  $m$ 
  - $A$  only has  $m$  rows, and a row never has two pivots

# Dimensionality of nullspace

- Counting the pivots leads to an extremely important theorem

$$A = \left[ \begin{array}{c|c|c|c|c} p & p & f & p & f \\ \hline \vdots & \vdots & \vdots & \vdots & \vdots \end{array} \right] \quad R = \left[ \begin{array}{ccccc} 1 & 0 & a & 0 & c \\ 0 & 1 & b & 0 & d \\ 0 & 0 & 0 & 1 & e \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad s_1 = \left[ \begin{array}{c} -a \\ -b \\ 1 \\ 0 \\ 0 \end{array} \right] \quad s_2 = \left[ \begin{array}{c} -c \\ -d \\ 0 \\ -e \\ 1 \end{array} \right]$$

3 pivot columns  $p$        $I$  in pivot columns      special  $Rs_1 = 0$  and  $Rs_2 = 0$   
2 free columns  $f$        $F$  in free columns      take  $-a$  to  $-e$  from  $R$   
to be revealed by  $R$       3 pivots: rank  $r = 3$        $Rs = 0$  means  $As = 0$

- Let  $A$  has more columns than rows ( $n > m$ ), which is a short wide matrix
- Must have least  $n - m$  free variables, since number of pivots cannot exceed  $m$ 
  - $A$  only has  $m$  rows, and a row never has two pivots
  - Of course a row might have no pivot which means an extra free variable

# Dimensionality of nullspace

- Counting the pivots leads to an extremely important theorem

$$A = \left[ \begin{array}{c|c|c|c|c} p & p & f & p & f \\ \hline \end{array} \right] \quad R = \left[ \begin{array}{ccccc} 1 & 0 & a & 0 & c \\ 0 & 1 & b & 0 & d \\ 0 & 0 & 0 & 1 & e \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad s_1 = \begin{bmatrix} -a \\ -b \\ 1 \\ 0 \\ 0 \end{bmatrix} \quad s_2 = \begin{bmatrix} -c \\ -d \\ 0 \\ -e \\ 1 \end{bmatrix}$$

3 pivot columns  $p$        $I$  in pivot columns      special  $Rs_1 = 0$  and  $Rs_2 = 0$   
2 free columns  $f$        $F$  in free columns      take  $-a$  to  $-e$  from  $R$   
to be revealed by  $R$       3 pivots: rank  $r = 3$        $Rs = 0$  means  $As = 0$

- Let  $A$  has more columns than rows ( $n > m$ ), which is a short wide matrix
- Must have least  $n - m$  free variables, since number of pivots cannot exceed  $m$ 
  - $A$  only has  $m$  rows, and a row never has two pivots
  - Of course a row might have no pivot which means an extra free variable
- Nullspace is a subspace. Its “dimension” is the number of free variables