## ChatGPT 3.5 ~

仚



## You

The integral of (a^2\*e^(-a^2/(2\*sigma^2))da) over minus infinity to plus infinity?



## **ChatGPT**

The integral you provided is related to the normal distribution and is commonly encountered in probability and statistics. The integral you're asking about is:

$$\int_{-\infty}^{\infty} a^2 e^{-\frac{a^2}{2\sigma^2}} da$$

To solve this integral, you can use the fact that it involves a Gaussian function. One way to solve it is by completing the square. The integral can be rewritten as follows:

$$\int_{-\infty}^{\infty}a^2e^{-rac{a^2}{2\sigma^2}}\,da=\sqrt{2\pi}\sigma^3\int_{-\infty}^{\infty}rac{1}{\sqrt{2\pi}\sigma}a^2e^{-rac{a^2}{2\sigma^2}}\,da$$

Now, if you let  $u=\frac{a}{\sigma}$ , the integral becomes:

$$\sqrt{2\pi}\sigma^3\int_{-\infty}^\infty rac{1}{\sqrt{2\pi}\sigma}a^2e^{-rac{a^2}{2\sigma^2}}\,da=\sqrt{2\pi}\sigma^3\int_{-\infty}^\infty rac{1}{\sqrt{2\pi}}(\sigma u)^2e^{-rac{u^2}{2}}\,du$$

Solving this integral involves standard Gaussian integral results. The final answer is:

$$\int_{-\infty}^{\infty} a^2 e^{-rac{a^2}{2\sigma^2}} \, da = \sqrt{2\pi}\sigma^3$$

So, the integral evaluates to  $\sqrt{2\pi}\sigma^3$ .





