

Path-spread-resolved quantum transition-state theory for proton transfer

Mercel Vubangsi,^{1,2,3,a)} Fadi Al-Turjman,² and Martin Tchoffo⁴

¹⁾*Department of Physics, Maharishi International University, Fairfield, Iowa 52557, USA*

²⁾*Research Center for AI and IoT, Near East University, Nicosia 99138, Cyprus*

³⁾*Department of Physics, University of Bamenda, Bamili, Cameroon*

⁴⁾*Department of Physics, University of Dschang, Dschang, Cameroon*

(Dated: 12 September 2026)

Centroid quantum transition-state theory and ring-polymer molecular dynamics yield thermal rates, but they collapse the imaginary-time path of the transferring proton onto a single collective coordinate. We define the proton-path spread χ_p as the mean-square bead displacement of the ring polymer along the proton-sharing coordinate—the squared radius of gyration projected onto that coordinate—and use it to resolve the centroid dividing-surface ensemble. The associated path-spread-resolved (QRR) formula decomposes the usual centroid flux. When the transmission coefficient does not depend on path class, that integral is identically the centroid QTST rate k_{cTST} . For a harmonic proton mode, $\langle\chi_p\rangle$ has a closed continuum formula and an exact finite-bead counterpart; Lévy-staging path-integral Monte Carlo reproduces the finite-bead result to within 1%. The H/D ratio interpolates between $\sqrt{m_D/m_H}$ as $T \rightarrow 0$ and m_D/m_H as $T \rightarrow \infty$. Exact Eckart rates at 300 K give a tunneling correction $\kappa_H = 1.28\text{--}2.68$ and a kinetic isotope effect of 1.68–2.28 as the barrier rises from 4 to 16 $k_B T$ (classical isotope factor $\sqrt{2}$). Umbrella PIMC centroid QTST, which is QRR at $\kappa = 1$, overestimates those exact rates by 13–40% on converged points while remaining above TST; the Wigner series undershoots for H. The same dividing-surface ensemble has $f_{\text{deloc}}^H \approx 0.22\text{--}0.38$ versus $f_{\text{deloc}}^D \approx 0.02$. An asymmetric double well shows the same isotope contrast (0.18–0.23 versus 0.01–0.04). We do not compute a class-conditional recrossing factor. The estimator can be accumulated in any existing path-integral code.

Keywords: proton transfer, path-integral molecular dynamics, quantum transition-state theory, kinetic isotope effect, ring-polymer molecular dynamics

I. INTRODUCTION

Proton transfer, and proton-coupled electron transfer (PCET) in particular, is a setting in which nuclear quantum effects, solvent reorganization, and, at electrodes, the applied potential jointly control the rate.^{1–4} Kinetic isotope effects (KIEs) and non-Arrhenius temperature dependence are the usual experimental signatures of tunneling,^{4,5} but they do not by themselves identify the imaginary-time path geometry that carries the reactive flux.

Path-integral methods address that geometry at the level of the thermal density matrix.^{6–8} Centroid quantum transition-state theory (QTST) replaces the classical reaction coordinate by the ring-polymer centroid and evaluates a flux through a dividing surface in centroid space.^{9–12} Ring-polymer molecular dynamics (RPMD) propagates the same polymer in real time and supplies a recrossing correction to the centroid flux.^{13–15} Instanton theory, in the complementary deep-tunneling limit, optimizes a single dominant imaginary-time orbit.^{16,17} Transition-path sampling retains full real-time path information, at a cost that is not organized by equilibrium imaginary-time geometry.¹⁸

What these approaches do not provide is a cheap, equilibrium order parameter that distinguishes compact ring

polymers (thermally activated, localized near the donor or acceptor) from delocalized polymers whose beads span the transfer coordinate (tunneling-like imaginary-time geometry). The ring-polymer radius of gyration is already used as a diagnostic of delocalization in condensed-phase proton studies.^{19,20} Here we restrict that diagnostic to the chemical sharing coordinate, give it exact harmonic limits, and use it to *condition* the QTST flux.

The resulting path-spread-resolved rate, which we call QRR, is

$$k_{\text{QRR}} = \int_0^\infty d\chi_p k_{\text{QTST}}(\chi_p) \kappa(\chi_p) p^\ddagger(\chi_p), \quad (1)$$

where $p^\ddagger(\chi_p)$ is the distribution of path spreads at the dividing surface and $\kappa(\chi_p)$ is a class-conditional transmission coefficient. When κ is taken independent of χ_p , Eq. (1) reduces identically to ordinary centroid QTST: $k_{\text{QRR}} = k_{\text{cTST}}$. QRR is therefore a decomposition of a known estimator, not a claim of a more accurate thermal rate. Its value is mechanistic: it reports which imaginary-time geometries carry the flux, how that mix depends on isotope and barrier, and whether path spread couples to a solvent coordinate.

Section II defines χ_p and the conditional free energy, states the solvable limits, and writes the flux decomposition. Section III describes the samplers. Section IV reports harmonic benchmarks, compares k_{cTST} (QRR at $\kappa = 1$) to exact Eckart rates, and resolves the dividing-surface ensemble of an asymmetric double well. Section V discusses scope and limitations. We do not re-

^{a)}Electronic mail: m.vubangsi@miu.edu

port condensed-phase electrochemical simulations; those require a validated reactive potential and are left as an application of the estimator.

II. THEORY

A. Path-spread functional

The P -bead ring-polymer representation of the canonical density is standard.^{8,10} For a proton transferring between donor and acceptor oxygens we use the bead-resolved sharing coordinate

$$\xi_p^{(s)} = r_{\text{ODH}}^{(s)} - r_{\text{OAH}}^{(s)}, \quad \bar{\xi}_p = P^{-1} \sum_{s=1}^P \xi_p^{(s)}. \quad (2)$$

The path-spread functional is the bead variance along that coordinate,

$$\chi_p = \frac{1}{P} \sum_{s=1}^P (\xi_p^{(s)} - \bar{\xi}_p)^2. \quad (3)$$

This is the squared radius of gyration of the polymer projected onto ξ_p . It is identically zero at $P = 1$ and is non-negative for all configurations. Compact paths have χ_p of order the zero-point spread in a well; delocalized paths have χ_p large enough that beads occupy both sides of the barrier.

The joint density $P(\bar{\xi}_p, \chi_p)$ defines the conditional free energy

$$F(\bar{\xi}_p, \chi_p) = -k_B T \ln P(\bar{\xi}_p, \chi_p) + C. \quad (4)$$

Projection onto $\bar{\xi}_p$ recovers the centroid potential of mean force.¹⁰ Slices at fixed $\bar{\xi}_p$ resolve path class, as sketched in Fig. 1. A solvent or energy-gap coordinate λ_s may be added without changing the construction; the three-dimensional surface $F(\bar{\xi}_p, \chi_p, \lambda_s)$ is the natural object for PCET, but the calculations below are two-dimensional unless stated.

B. Flux decomposition

Let $\bar{\xi}_p^\ddagger$ locate the dividing surface. The marginal

$$p^\ddagger(\chi_p) = \frac{e^{-\beta F(\bar{\xi}_p^\ddagger, \chi_p)}}{\int d\chi'_p e^{-\beta F(\bar{\xi}_p^\ddagger, \chi'_p)}} \quad (5)$$

is the probability that a dividing-surface configuration belongs to path class χ_p . The class-conditional QTST flux is the usual positive-velocity centroid flux restricted to that class,

$$k_{\text{QTST}}(\chi_p) = Z_R^{-1} \langle [\dot{\xi}_p] + \delta(\bar{\xi}_p - \bar{\xi}_p^\ddagger) \delta(\chi_p[\mathbf{R}] - \chi_p) \rangle. \quad (6)$$

A Bennett–Chandler transmission coefficient $\kappa(\chi_p)$ may be attached to each class.^{14,21,22} Integrating over χ_p produces Eq. (1). If κ does not depend on χ_p , the integral collapses to $k_{\text{QRR}} = \kappa k_{\text{QTST}}$, and the choice $\kappa = 1$ gives

$$k_{\text{QRR}}(\kappa = 1) \equiv k_{\text{cTST}}. \quad (7)$$

The Eckart calculations below evaluate this identity by computing k_{cTST} from umbrella PIMC and comparing it to the exact scattering rate. We emphasize two further restrictions. First, a Wigner series in $\hbar^2$ is a tunneling correction to a parabolic barrier,^{23,24} not a recrossing factor; the two must not be conflated. Second, a χ_p -dependent κ is not computed: for highly delocalized paths the Wigner series is not a controlled approximation, and a class-conditional RPMD flux-side correlation would be the consistent dynamical correction.¹⁴ Inventing a $\kappa(\chi_p)$ from the centroid potential of mean force would double-count the barrier and cannot repair the QTST overestimate documented below.

C. Solvable limits

Remark 1 (Classical limit) At $P = 1$, Eq. (3) gives $\chi_p = 0$ identically, so $p^\ddagger(\chi_p) = \delta(\chi_p)$ and k_{QRR} reduces to classical transition-state theory. This is a consistency check, not an independent result.

Theorem 1 (Harmonic continuum) A harmonic mode $V = \frac{1}{2}m\omega^2 q^2$ has

$$\langle \chi_p \rangle_\infty = \frac{\hbar}{2m\omega} \coth\left(\frac{\beta\hbar\omega}{2}\right) - \frac{k_B T}{m\omega^2}. \quad (8)$$

The first term is the equal-time imaginary-time correlator $G(0)$; the second is the centroid variance, which is classical for a harmonic oscillator.¹⁰ The difference is non-negative because $\coth x \geq 1/x$ for $x > 0$. As $T \rightarrow 0$, $\langle \chi_p \rangle \rightarrow \hbar/(2m\omega)$. As $T \rightarrow \infty$, the leading term is the Wigner correction $\hbar^2\beta/(12m)$, independent of ω .²³ The Matsubara-mode derivation is given in the supplementary material.

Proposition 1 (Exact finite P) For the same oscillator discretized with P beads,

$$\langle \chi_p \rangle_P = \frac{1}{P} \sum_{k=1}^{P-1} \frac{1}{\beta\lambda_k}, \quad \lambda_k = \frac{4mP}{\hbar^2\beta^2} \sin^2\left(\frac{\pi k}{P}\right) + \frac{m\omega^2}{P}. \quad (9)$$

As $P \rightarrow \infty$, $\langle \chi_p \rangle_P \rightarrow \langle \chi_p \rangle_\infty$.

Equation (9) is the quantity that a primitive path-integral estimator of Eq. (3) actually samples. Comparing Monte Carlo to Eq. (8) at modest P therefore mixes statistical error with Trotter error. We compare to Eq. (9) first.




FIG. 1. Schematic, not computed data. (a) Centroid potential of mean force $F(\bar{\xi}_p)$, in which all ring-polymer shapes at a given centroid are averaged. (b) Conditional free energy $F(\bar{\xi}_p, \chi_p)$. Compact and delocalized dividing-surface geometries that share $\bar{\xi}_p = 0$ are marked.

Corollary 1 (Harmonic isotope scaling) *On a shared harmonic surface with $k = m\omega^2$ held fixed, $\langle \chi_p \rangle_H > \langle \chi_p \rangle_D$ at every $T > 0$. The ratio $\mathcal{R} = \langle \chi_p \rangle_H / \langle \chi_p \rangle_D$ satisfies*

$$\mathcal{R}(T \rightarrow 0) = \sqrt{m_D/m_H}, \quad \mathcal{R}(T \rightarrow \infty) = m_D/m_H. \quad (10)$$

The low-temperature limit is the ratio of zero-point amplitudes. The high-temperature limit is the ratio of leading Wigner corrections. Both follow from Eq. (8) and do not extend automatically to anharmonic barriers.

Proposition 2 (Factorization) *If $F(\bar{\xi}_p, \chi_p, \lambda_s) = F_p(\bar{\xi}_p, \chi_p) + F_s(\lambda_s)$, then $p^\ddagger(\chi_p)$ is independent of λ_s and the χ_p integral in Eq. (1) reduces to ordinary QTST when κ is class-independent.*

A nonzero mixed derivative $\partial^2 F / \partial \chi_p \partial \lambda_s$, or a nonzero dividing-surface correlation $\text{corr}(\chi_p, \lambda_s)$, would then signal proton-solvent coupling beyond a product Marcus picture.^{2,3} We do not claim to have measured that correlation with statistical control in this work.

III. METHODS

Path-integral Monte Carlo uses Lévy staging of the free-particle bridge,^{25,26} which samples the ring-polymer springs exactly, together with a Metropolis decision on the physical potential and a rigid centroid move. The primitive estimator of χ_p is the bead variance. For the harmonic oscillator we also evaluate Eqs. (8) and (9) in closed form.

Asymmetric double-well (ADW) free energies are obtained from one-dimensional umbrella sampling along $\bar{\xi}_p$ ²⁷ with harmonic restraints, reweighted by the

weighted histogram analysis method (WHAM).²⁸ Two-dimensional histograms of $(\bar{\xi}_p, \chi_p)$ use the same WHAM weights. The delocalized fraction at the dividing surface is

$$f_{\text{deloc}} = \int_{\chi_p \geq 2\chi_c} p^\ddagger(\chi_p) d\chi_p, \quad (11)$$

with the threshold $\chi_c = \hbar/(2m\omega^*)$ evaluated at $\omega^* = 1000 \text{ cm}^{-1}$, giving $\chi_c \approx 0.010 \text{ Å}^2$. Compact and intermediate classes use $0.5\chi_c$ and $2\chi_c$ as bookkeeping cuts; the continuous $p^\ddagger(\chi_p)$ is the primary object.

Exact thermal rates for the symmetric Eckart barrier $V(x) = V_0 \text{sech}^2(x/a)$ are

$$k = \frac{1}{2\pi\hbar Q_r} \int_0^\infty dE e^{-\beta E} T(E), \quad (12)$$

with $T(E)$ the exact transmission probability of the symmetric Eckart barrier²⁹ and $Q_r = (m/2\pi\hbar^2\beta)^{1/2}$. Classical TST is $k_{\text{TST}} = \sqrt{k_B T / (2\pi m)} e^{-\beta V_0}$. The Wigner estimate multiplies k_{TST} by $1 + (\beta\hbar\omega_b)^2/24$, where $\omega_b = \sqrt{2V_0/(ma^2)}$. All Eckart integrals use a logarithmically stable evaluation of $T(E)$.²⁹

Centroid QTST on the same barrier is obtained from umbrella PIMC along the centroid,

$$k_{\text{cTST}} = \sqrt{\frac{k_B T}{2\pi m}} e^{-\beta F(0)}, \quad (13)$$

with F the WHAM potential of mean force, zeroed on the mean of the $|\bar{\xi}_p| > 1.6 \text{ Å}$ tails so that $F(\infty) = 0$. Eleven harmonic windows on $[-2, 2] \text{ Å}$ with $k_{\text{umb}} = 60 \text{ kJ mol}^{-1} \text{ Å}^{-2}$ are used, $P = 32$ (H) and $P = 24$ (D), 4×10^4 production steps after 8×10^3 equilibration. Residual tail scatter in F is of order 0.5 kJ mol^{-1} . Deuterium at $V_0 = 12 k_B T$ is excluded from the rate comparison:

the dividing-surface window drifted and produced the unphysical ordering $F_D(0) < F_H(0)$. With $\kappa = 1$, Eq. (1) equals Eq. (13) identically; both values are stored in the accompanying JSON.

Temperatures are 300 K unless noted. Masses are 1 and 2 amu. Random number generators are seeded at 42. Code and JSON outputs are in the repository accompanying this article.

IV. RESULTS

A. Harmonic oscillator

Figure 2 compares Eq. (8) with Lévy-staging Monte Carlo for a 1000 cm^{-1} mode over 100–700 K. At 300 K the continuum values are $\langle \chi_p \rangle_H = 0.01011 \text{ \AA}^2$ and $\langle \chi_p \rangle_D = 0.00572 \text{ \AA}^2$. With $P = 64$ (H) and $P = 32$ (D), the exact finite-bead formula differs from the continuum by 0.12% and 0.29%, respectively, and the Monte Carlo root-mean-square relative error against Eq. (9) is 0.56% (H) and 1.03% (D).

The isotope ratio $\mathcal{R}(T)$ increases from 1.52 at 100 K toward $\sqrt{2}$ as T is lowered, and from 1.77 at 300 K to 1.94 at 700 K, approaching 2 from below, as required by Corollary 1 [Fig. 2(b)]. Figure 2(c) shows that the primitive estimator tracks Eq. (9) from $P = 4$ to $P = 128$ and that the Trotter error relative to the continuum is already below 0.5% at $P = 32$ for this frequency and temperature. A staging move that draws Brownian-bridge *marginals* independently, rather than the sequential Lévy construction, produces a spurious increase of $\langle \chi_p \rangle$ with P and should not be used.

B. Eckart barrier

The symmetric Eckart barrier with width $a = 0.50 \text{ \AA}$ supplies exact thermal rates that do not depend on path-integral sampling, and therefore a test of Eq. (7): k_{QRR} at $\kappa = 1$ is k_{cTST} , and that number can be compared with the exact scattering rate.

Figure 3(a) and Table I compare four estimates of $\kappa = k/k_{\text{TST}}$ at 300 K. The exact κ_H increases from 1.28 at $V_0 = 4 k_B T$ to 2.68 at $V_0 = 16 k_B T$; the corresponding Wigner factors are 1.22 and 1.86. Wigner is a useful first correction at moderate $\hbar\omega_b/k_B T$ and underestimates the exact κ once $\hbar\omega_b/k_B T \gtrsim 4$. Centroid QTST from umbrella PIMC gives $\kappa_H = 1.54, 1.98$, and 2.30 at 4, 8, and $12 k_B T$, versus exact 1.28, 1.60, and 2.04 (overestimates of 20%, 24%, and 13%). Deuterium is overestimated by 40% and 10% at 4 and $8 k_B T$. In every converged case $F(0) < V_0$, so $\kappa_{\text{cTST}} > 1$ and the nuclear quantum effect has the correct sign. Deuterium at $12 k_B T$ is omitted: the dividing-surface umbrella drifted and produced the unphysical ordering $F_D(0) < F_H(0)$.

The exact KIE [Fig. 3(b)] is 1.68 at $4 k_B T$ and 2.28

at $16 k_B T$, against a classical factor $\sqrt{2} \approx 1.414$. Centroid QTST gives 1.44 and 1.98 at 4 and $8 k_B T$: it barely exceeds classical at the lowest barrier, where H and D overshoot by similar factors, and it overshoots the exact KIE at $8 k_B T$. These comparisons are the numerical content of Eq. (1) at $\kappa = 1$. They are not a claim that QRR improves on QTST.

Figure 3(c) is the diagnostic that the integrated rate does not report. At the dividing surface, $f_{\text{deloc}}^H = 0.22, 0.23$, and 0.38 at 4, 8, and $12 k_B T$, while f_{deloc}^D remains 0.017. The jump at $12 k_B T$ for H is sampled with 629 dividing-surface configurations and should be read as a trend, not a precise fraction. The isotope contrast of an order of magnitude is the robust result.

C. Asymmetric double well

The ADW $V(q) = V_0[(q/q_0)^2 - 1]^2 + \alpha q$ with $q_0 = 0.70 \text{ \AA}$ is a bound proton-transfer model. Figure 4(a) shows the WHAM centroid PMF at $V_0 = 8 k_B T$: the H barrier is 15.9 kJ mol^{-1} and the D barrier is 17.6 kJ mol^{-1} . Figure 4(b) is the quantity QRR actually uses, the dividing-surface density $p^\ddagger(\chi_p)$. It is broader for H than for D, with means $\langle \chi_p \rangle_\ddagger^H = 0.015 \text{ \AA}^2$ and $\langle \chi_p \rangle_\ddagger^D = 0.008 \text{ \AA}^2$. With the bookkeeping cut $2\chi_c = 0.020 \text{ \AA}^2$, the delocalized weight is $f_{\text{deloc}}^H = 0.18\text{--}0.23$ across $V_0 = 2\text{--}8 k_B T$, versus $f_{\text{deloc}}^D = 0.014\text{--}0.045$ [Fig. 4(c) and Table II]. For H the fraction is essentially flat in this barrier window; the robust observation is the isotope contrast, not a strong barrier trend.

These are equilibrium path-class statistics. The isolated ADW is a bound well and has no irreversible thermal rate, so we do not convert f_{deloc} into a QRR rate, nor into a claimed improvement over RPMD. Two-dimensional histograms of $F(\xi_p, \chi_p)$ from the same umbrellas are too sparsely sampled in χ_p to publish as free-energy surfaces.

D. What is not shown

A bilinear proton-solvent model $V = V_{\text{ADW}}(q_p) + \frac{1}{2}k_s q_s^2 + \kappa_c q_p q_s$ was sampled without umbrellas. Dividing-surface correlations $\text{corr}(\chi_p, q_s)$ lie between -0.06 and -0.16 with a few hundred crossings and do not trend cleanly with κ_c . That scan is not evidence for or against Proposition 2; a controlled test would require umbrella sampling in $(\xi_p, \chi_p, \lambda_s)$. We therefore omit it from the figures.

V. DISCUSSION

QRR does not replace RPMD, instanton theory, or vibronic PCET rate laws.^{2,14,17,30} RPMD remains the appropriate dynamical correction to centroid QTST in the




FIG. 2. Harmonic path spread for $\omega = 1000 \text{ cm}^{-1}$. (a) $\langle \chi_p \rangle(T)$ from Eq. (8) (lines) and Lévy-staging PIMC (symbols). (b) Isotope ratio, with the $T \rightarrow 0$ limit $\sqrt{2}$ and the $T \rightarrow \infty$ limit 2. (c) Bead convergence of $\langle \chi_p \rangle_H$ at 300 K: PIMC versus the exact finite- P formula, Eq. (9).

TABLE I. Symmetric Eckart barrier at $T = 300 \text{ K}$, $a = 0.50 \text{ \AA}$. $\kappa = k/k_{\text{TST}}$. Centroid QTST is umbrella PIMC; QRR at $\kappa = 1$ equals that column. Dashes mark points that were not computed, or D at $12 k_B T$, which failed the dividing-surface check. Classical KIE is $\sqrt{2}$ at every barrier.

$V_0/k_B T$	κ_H^{ex}	κ_H^{W}	κ_H^{cTST}	κ_D^{ex}	κ_D^{W}	κ_D^{cTST}	KIE ^{ex}	KIE ^W	KIE ^{cTST}
4	1.28	1.22	1.54	1.08	1.11	1.51	1.68	1.55	1.44
8	1.60	1.43	1.98	1.28	1.22	1.41	1.77	1.67	1.98
12	2.04	1.65	2.30	1.36	1.32	—	2.12	1.76	—
16	2.68	1.86	—	1.66	1.43	—	2.28	1.84	—

TABLE II. Asymmetric double well at $T = 300 \text{ K}$. Centroid barriers from 1D WHAM; f_{deloc} from the WHAM-weighted dividing-surface ensemble with $2\chi_c = 0.020 \text{ \AA}^2$.

$V_0/k_B T$	ΔF_H (kJ/mol)	ΔF_D (kJ/mol)	f_{deloc}^H	f_{deloc}^D	$\langle \chi_p \rangle_{\frac{1}{2}}^H$ ($\text{\AA}^2$)	$\langle \chi_p \rangle_{\frac{1}{2}}^D$ ($\text{\AA}^2$)
2	4.25	4.60	0.18	0.014	0.014	0.007
4	8.90	9.02	0.23	0.020	0.015	0.007
6	12.4	14.3	0.22	0.024	0.015	0.007
8	15.9	17.6	0.23	0.045	0.015	0.008

crossover regime; instanton theory remains the appropriate semiclassical description when a single tunneling orbit dominates; Hammes-Schiffer vibronic theory remains the appropriate nonadiabatic framework when electron-proton nonadiabaticity is essential. What χ_p adds is an equilibrium resolution of imaginary-time geometries that those methods otherwise average or optimize.

The present calculations evaluate Eq. (1) only at $\kappa = 1$, where it is identical to k_{TST} . A class-conditional transmission coefficient is not computed. On the Eckart barrier that common rate overestimates the exact scattering result by 13–40% on converged points, as expected for a recrossing-free centroid theory; QRR cannot reduce that error without a genuine $\kappa(\chi_p)$. The ADW section reports $p^{\ddagger}(\chi_p)$ only. Connecting the two models is an analogy:

the isolated double well has no irreversible thermal rate.

Two condensed-phase hypotheses follow, and should be treated as such. First, if an increase of the observed KIE with driving force^{30,31} is carried by tunneling-like paths, $\Delta f_{\text{deloc}} = f_{\text{deloc}}^H - f_{\text{deloc}}^D$ should increase with that driving force. Second, a breakdown of $F = F_p + F_s$ at the dividing surface would be a signature of proton-solvent coupling beyond a product Marcus picture. Neither hypothesis is tested here.

Limitations are sharp. χ_p is an imaginary-time polymer statistic, not a spectroscopic observable. The cut χ_c is bookkeeping; Fig. 4(b) is the primary object. Recrossing is not resolved by class. The models are one-dimensional. Residual WHAM tail scatter of order 0.5 kJ mol^{-1} limits the precision of κ_{cTST} at the lowest Eckart barrier. Finite- P errors in χ_p are under control for the 1000 cm^{-1} mode at $P \geq 32$; stiffer stretches will need more beads or a virial estimator.

VI. CONCLUSIONS

We have defined a projected ring-polymer path spread χ_p for proton transfer, derived its harmonic continuum and finite-bead limits, and used it to resolve the centroid dividing-surface ensemble. Lévy-staging Monte Carlo reproduces the exact finite-bead harmonic formula to about




FIG. 3. Symmetric Eckart barrier, $a = 0.50 \text{ \AA}$, $T = 300 \text{ K}$. (a) Tunneling correction $\kappa = k/k_{\text{TST}}$. Diamonds are umbrella-PIMC centroid QTST, equal to QRR at $\kappa = 1$. Deuterium at $12 k_B T$ is omitted (dividing-surface window not converged). (b) Kinetic isotope effect. Classical TST is $\sqrt{2}$ at every V_0 . (c) Delocalized weight at the dividing surface, $2\chi_c = 0.020 \text{ \AA}^2$.




FIG. 4. Asymmetric double well, $T = 300 \text{ K}$. (a) Centroid PMF at $V_0 = 8 k_B T$. (b) Dividing-surface path-spread density $p^{\ddagger}(\chi_p)$ at the same barrier; the dotted line is the bookkeeping cut $2\chi_c$. (c) Delocalized fraction versus barrier height. The H fraction is nearly flat; the D fraction remains an order of magnitude smaller.

one percent. Exact Eckart rates quantify when tunneling corrections and KIEs depart from TST. Umbrella PIMC centroid QTST, equal to QRR at $\kappa = 1$, overestimates those exact rates while capturing the sign of the nuclear quantum effect and assigning a much larger delocalized weight to H than to D. On an asymmetric double well the dividing-surface path-spread distribution is robustly broader for H than for D. Equation (1) is a bookkeeping decomposition of QTST; it is not shown here to be a better rate. The estimator is a collective variable and can be accumulated in any path-integral molecular dynamics run.

ACKNOWLEDGMENTS

Computational work was performed on local workstations. No external funding is declared.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in Zenodo at <https://doi.org/10.5281/zenodo.22718950> (Ref. 32). The source repository is <https://github.com/vmercel/QRR-PCET>.

¹S. Hammes-Schiffer, J. Am. Chem. Soc. **137**, 8860 (2015).

²A. Soudackov and S. Hammes-Schiffer, J. Chem. Phys. **113**, 2385 (2000).

³R. A. Marcus, J. Chem. Phys. **24**, 966 (1956).

- ⁴J. P. Klinman and A. Kohen, *Annu. Rev. Biochem.* **82**, 471 (2013).
- ⁵F. H. Westheimer, *Chem. Rev.* **61**, 265 (1961).
- ⁶R. P. Feynman, *Rev. Mod. Phys.* **20**, 367 (1948).
- ⁷R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
- ⁸D. Chandler and P. G. Wolynes, *J. Chem. Phys.* **74**, 4078 (1981).
- ⁹G. A. Voth, D. Chandler, and W. H. Miller, *J. Chem. Phys.* **91**, 7749 (1989).
- ¹⁰J. Cao and G. A. Voth, *J. Chem. Phys.* **100**, 5106 (1994).
- ¹¹M. J. Gillan, *J. Phys. C: Solid State Phys.* **20**, 3621 (1987).
- ¹²T. J. H. Hele and S. C. Althorpe, *J. Chem. Phys.* **138**, 084108 (2013).
- ¹³I. R. Craig and D. E. Manolopoulos, *J. Chem. Phys.* **121**, 3368 (2004).
- ¹⁴I. R. Craig and D. E. Manolopoulos, *J. Chem. Phys.* **122**, 084106 (2005).
- ¹⁵S. Habershon, D. E. Manolopoulos, T. E. Markland, and T. F. Miller, *Annu. Rev. Phys. Chem.* **64**, 387 (2013).
- ¹⁶J. O. Richardson and S. C. Althorpe, *J. Chem. Phys.* **131**, 214106 (2009).
- ¹⁷J. O. Richardson, *Int. Rev. Phys. Chem.* **37**, 171 (2018).
- ¹⁸P. G. Bolhuis, D. Chandler, C. Dellago, and P. L. Geissler, *Annu. Rev. Phys. Chem.* **53**, 291 (2002).
- ¹⁹C. Drechsel-Grau and D. Marx, *Phys. Rev. Lett.* **112**, 148302 (2014).
- ²⁰O. Marsalek and T. E. Markland, *J. Phys. Chem. Lett.* **8**, 1545 (2017).
- ²¹C. H. Bennett, *ACS Symp. Ser.* **46**, 63 (1977).
- ²²D. Chandler, *J. Chem. Phys.* **68**, 2959 (1978).
- ²³E. Wigner, *Phys. Rev.* **40**, 749 (1932).
- ²⁴E. Wigner, *Trans. Faraday Soc.* **34**, 29 (1938).
- ²⁵D. M. Ceperley, *Rev. Mod. Phys.* **67**, 279 (1995).
- ²⁶M. E. Tuckerman, B. J. Berne, G. J. Martyna, and M. L. Klein, *J. Chem. Phys.* **99**, 2796 (1993).
- ²⁷G. M. Torrie and J. P. Valleau, *J. Comput. Phys.* **23**, 187 (1977).
- ²⁸S. Kumar, J. M. Rosenberg, D. Bouzida, R. H. Swendsen, and P. A. Kollman, *J. Comput. Chem.* **13**, 1011 (1992).
- ²⁹C. Eckart, *Phys. Rev.* **35**, 1303 (1930).
- ³⁰Z. K. Goldsmith, Y. C. Lam, A. V. Soudackov, and S. Hammes-Schiffer, *J. Am. Chem. Soc.* **141**, 1084 (2019).
- ³¹M. T. M. Koper, *J. Electroanal. Chem.* **660**, 254 (2011).
- ³²M. Vubangsi, F. Al-Turjman, and M. Tchoffo, "Path-spread-resolved quantum transition-state theory for proton transfer," (2026).