Modeling rice prices using spectral analysis -preliminary

Vidal Marvin Gabriel

January 26, 2021

1 Preliminary study - checking for causation using \mathbb{R}^2 and OLS

Since there is abundance in the rice data from the mid 1990s to the current year, the author explored for some cauality in rice prices and rice supply. This is in accordance with theory that when supply plummets, price increases. The author found four time series datasets, i.e. rice prices, total rice stock (TRS), household rice stock (HRS), commercial rice stock (CRS), and NFA rice stock (NRS). However, upon checking for causation through a simple ordinary least squares regression (OLS), the author found no causation in the empirical data. In the first run, the author used the following linear model:

rice price =
$$\alpha_0 + \alpha_1 TRS + \alpha_2 HRS + \alpha_3 CRS + \alpha_4 NRS$$
 (1)

The resulting R^2 shows a very low value of 0.289 which indicates a pretty low correlation to try to draw conclusions. The author further utilised omitted-variable bias (OVB) technique, to check if R^2 improves. However, R^2 failed to improve. Thus, based from the preliminary study, the rice inflation time series can be modelled using a univariate process.

2 Introduction

Inflation is an economic phenomenon in the form of price increase (Mankiw, 2018). This is the weighted sum of the different consumer price index (CPI) commodites. In the Philippine setting, rice prices greatly affect the food commodity in the CPI basket. This is mainly because most Filipinos' main food in the table is rice. Thus, an increase in rice prices correlates to an increase in the food commodity which then affects the aggregate inflation. Thus forecasting rice inflation may greatly affect aggregate inflation outlook.

The current models used to analyze inflation is in the form of time series. Time series is the collection of observation in the time domain (Hamilton, 2020). It may contain trend, seasonality,

cycle, and stochastic component. Thus, a large body of empirical literature is devoted to modeling inflation. With regards to time series data, the simplest and perhaps most commonly used statistical tools to forecast inflation are the auto regressive (AR), vector autoregression (VAR), and ARIMA models (Moser, Rumler and Scharler, 2007), (Baciu, 2015). However, in the Philippine setting, these models were designed to model inflation in the aggregate level. Thus, there is a need to model disaggregated inflation especially rice which affects food commodity greatly.

In recent years, there have been attempts to look inflation in a different domain, the frequency domain (Ftiti, Essaadi et al., 2008), (Iacobucci, 2005). In this study, we try to use frequency domain to model the rice inflation series in the country.

When we look at time sereies in the frequency domain, a stationary series has a wave-like pattern as it has constant mean and constant variance. Waves have a period which is the distance between peaks or time between two peaks of a wave. Frequency is the closely related property of the wave period and is just the proportion of cycle that occurs during one observation. Frequency domain is used to analyze the data with respect to frequency.

The most common way to transform time domain function into a frequency domain function is through Fourier transformation developed by Joseph Fourier in 1822 (Culling, 1960). This frequency domain analysis is also known as spectral analysis or Fourier analysis. In this paper, spectral analysis approach with periodogram analysis is used to model the rice inflation of the Philippines. Further, the resulting model will be compared with autoregressive integrated moving average (ARIMA) which is the commonly used tool by Economic institutions to model inflation.

This paper develops a model based on Fourier analysis to model rice inflation and compares it with the commonly used statistical model for forecasting time series in econometrics - ARIMA, developed by Box and Jenkins (Box and Jenkins, 1976). Their performance will be assessed using two performance metrics, i.e. root mean square error (RMSE) and mean absolute error (MAE). The remainder of the paper is as follows: Section 3 describes briefly discuss the mathematical theory of Fourier analysis, Section 4 gives a brief literature review.

3 Mathematical theory on Fourier analysis development

This section briefly provides mathematical considerations on the Fourier development.

Let a function $f: R \to R$, with f and f' piecewise continous on R and periodic with period T, therefore $f(x+T)=f(x) \forall x \in R$ Considering Fourier series associated with the function $f: F(x)=\frac{a_0}{2}+\sum_{k=1}^{\infty}(a_k\cos\frac{2k\pi x}{T}+b_k\sin\frac{2k\pi x}{T})$ we then have the following,

$$\int_{-\frac{T}{2}}^{\frac{T}{2}} f(x) \cos \frac{2n\pi x}{T} dx = \frac{a_n T}{2}, n \ge 0, \int_{-\frac{T}{2}}^{\frac{T}{2}} f(x) \sin \frac{2n\pi x}{T} dx = \frac{b_n T}{2}, n \ge 1$$
 (2)

From Fourier series expression, it is observed that $F(x+T)=F(x)\forall x\in R$ so its sum is also a periodic function of period T

The Dirichlet's Theorem (Weisstein, 2004) states that in the conditions above, the Fourier series should converge punctually to f in the points of continuity and to $\frac{f(x+0)+f(x-0)}{2}$ in the discontinuity points.

Considering the partial sum of order n corresponding to the series of function F, the n-th Fourier polynomials is then given by

$$F_n(x) = \frac{a_0}{2} + \sum_{k=1}^{n} (a_k \cos \frac{2k\pi x}{T} + b_k \sin \frac{2k\pi x}{T})$$
 (3)

The Fourier polynomials have the property of approximating the function through one periodical with the observation that the absolute error tends to fall (due to the convergence points) with the rise of n.

Due to the existence of an important number of cyclical phenomena in many scientific fields, we intend, below, to approximate their development by means of Fourier polynomials of degree conveniently chosen.

In the case of the discretized phenomenons, we put the problem in the generation of functions that will pass through a series of data points. A very useful tool is the Lagrange interpolation polynomial. Therefore, considering a set of data (x_i, y_i) , i = 1, k+1, the Lagrange interpolation polynomial takes the form given in **Equation 4**. Further, **Equation 4** is the polynomial of minimum degree k passing through the data points.

$$L + n(x) = \sum_{i=1}^{k+1} \frac{(x - x_1)...(x - x_{i-1})(x - x_{i+1})...(x - x_n)}{(x_i - x_1)...(x_i - x_{i-1})(x_i - x_{i+1})...(x_i - x_n)} y_i$$
(4)

4 Literature review

There is a growing body of literature on the empirical application of frequency domain analysis for modeling in the fields of economics and finance. Results thus far have generally been favorable attesting to the usefulness of said techniques for these fields.

One study (Konarasinghe, Abeynayake and Gunaratne, 2016) was conducted to develop a model to forecast Returns of Sri Lankan share market using Fourier transform. The model was named as "Cirucular model (CM)" which forecasts individual company returns. The study revealed that the

CM was successful in forecasting monthly returns of 80% of the companies. They have concluded that CM is a suitable forecasting technique for Sri Lankan share market. However, the authors of the paper have noted that CM is only appropriate if the data shows no trend (detrended).

Another study was conducted by Thomson and Vuuren in 2016 (Thomson and Van Vuuren, 2016) to propose a spectral analysis to determine the duration of South African business cycle which is measured by using log changes in nominal gross domestic product (GDP). Three dominant cycles are used to forecast log monthly nominal GDP and forecasts are compared with historical data. The authors found that spectral analysis is more effective in estimating the business cycle length as well as in determining the position of the economy in the business cycle.

A study by Sella et al. in 2016 (Sella et al., 2010) which applies several advanced spectral analysis methods to analyse GDP fluctuations in Italy, Netherlands, and the United Kingdom. They have noted that these analysis tools allowed them to spectrally decompose, as well as reconstruct GDP time series from the data. They further noted that the models are well adapted to the analysis of short and noisy data like the GDP time series in their particular study.

To go further, a study (de Melo et al., 2015) was also conducted to use wavelet transform which is an extensiion of frequency domain models in economics and finanace. This study considered a generalized wavelet that in a particular case of unit amplitudes reduces to the Morlet wavelet. The introduced wavelet appears to be handy for the wavelet processing as it enables even very small details. The study shows some remarkable properties of frequency domain models which make them powerful tools to be used in economics. Thus, it can be seen that there have been attempts to integrate frequency domain analysis in the field of economics.

5 Research Methodology

The main objective of this study is to develop a significant model to forecast inflation in the Philippines. For this purpose, monthly consumer price index (CPI) data is gathered from the Philippine Statistics Authority (PSA) and covers the period from January 1994 to December 2020.

First, data preparation is used to transform the rice CPI data to rice inflation rates. Then, periodogram analysis is used to find rice inflation cycle as rice inflation may have periodicity in the series. After which, the period or frequency of sereis, significance of selected period is tested by a test developed by Fisher in 1929 (Fisher, 1929). Fourier analysis is then used to model rice inflation by utilizing selected frequency as Fourier frequency. On the other hand, ARIMA methodology will also be used to model the rice inflation data. However, in this study, performing ARIMA is not extensively discussed as this methodology has been well elaborated in several economic literatures.

To assess the performance of the two models, this paper will utilize root mean square error (RMSE) and Mean Absolute Error (MAE) as the performance metrics (see **Equations 5 & 6**). By using these accuracy measures, we can conclude which model is superior based on smaller values of RMSE and MAE.

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^{n} e_i^2} \tag{5}$$

$$MAE = \frac{1}{n} \sum_{i=1}^{n} |e_i| \tag{6}$$

5.1 Data preparation

Rice inflation is defined as the percent difference between the current CPI and the CPI of the previous year as seen in **Equation 7** below:

$$\pi_{year-on-year} = \frac{CPI_t - CPI_{t-1}}{CPI_{t-1}} * 100$$
 (7)

where CPI_t is the CPI of the current month while CPI_{t-1} is the same month's CPI from the previous year.

5.2 Periodogram analysis

Periodogram is a tool that partitions the total variance of a time series into component variances. This method is similar to ANOVA. The longer cycle shares large variance in the series. In general practice, when periods of cycles are not known then Periodogram is utilized to identify dominant cyclic behavior in the series. In periodogram, time series can be viewed as

$$Y_t = T_t + \sum_{i=1}^{N} (a_i \cos w_i t + b_i \sin w_i t) + \epsilon_t$$
(8)

where T_t is trend, N is total number of observations, a_i and b_i are coefficients, w_i is angular frequency in radians and ϵ_t is the error term. The coefficients are calculated which is presented in **Equations** 9 & 10 below.

$$a_i = \frac{2}{N} \sum_{t=1}^{N} (Y_t - \hat{T}_t) \cos w_i t \tag{9}$$

$$b_i = \frac{2}{N} \sum_{t=1}^{N} (Y_t - \hat{T}_t) \sin w_i t \tag{10}$$

The calculated coefficients are then used to obtain Intensity of Periodogram coordinate at frequency f_i as

$$I(f_i) = \frac{2}{N}(a_i^2 + b_i^2) \tag{11}$$

where i = 1, 2, 3, .q. In case of even number of observations N = 2q, q = n/2 and for odd number of observations N = 2q + 1, q = (N - 1)/2. The period against the largest Intensity that is actually the largest sum of squares is selected as cycle period of series.

5.3 Significance test

The variability in the sizes of sum of squares may be due to just sampling error. The largest ordinate must indicate strong periodicity even for white noise series. Therefore it is necessary to test the significance of largest periodic component in white noise. A test developed by Fisher in 1929 (Fisher, 1929) provides a reasonable method for testing significance of such periodic components. To perform Fisher test, *g* statistic is computed which is the ratio of largest sum of squares (or intensity ordinate) to the total sum of squares. A table of reference is given by Russel (Russel, 1985) which is the critical values for the test statistic. The null hypothesis of white noise series is rejected, if the value of *g* statistic is greater than critical value.

5.4 Spectral analysis

The basic idea of spectral analysis is to transform the time domain series into frequency domain series, which determines the importance of each frequency in the original series. This target is achieved by using Fourier transformation.

The general Fourier series model that contain components of time series is given by

$$Y_t = T_t + \sum_{i=1}^k (\alpha_i \cos iwt + \beta_i \sin iwt) + \epsilon_t$$
 (12)

where $w=2\pi f$, k=n/2 in case of even number, k=(n-1)/2 in case of odd number, n is the number of observations per season or cycle length, f is the Fourier frequency of number of peaks in series, k is the harmoninic of w - a more detailed explanation for this is given in the study by Delurgio (Armstrong, 2001), t is the time index, α_i and β_i are amplitudes which are estimated through multiple linear regression analysis.

5.5 Development of control model

The control model for this study is ARIMA presented by Box & Jenkins (Box and Jenkins, 1976). The model requires the time series to be stationarity. Afterwards parameters p, d, and q, the autoregressive terms, times differenced, and moving average terms respectively. This model is governed by a linear equation in which predictors are lags of dependent variable and lags of error terms, presented in **Equation 13** below.

$$\phi_p(B)\Delta^d Y = \theta_q(B)\epsilon_t \tag{13}$$

where Y_t is the present value, d is the difference, B is the backshift operator, and ϵ_t is the error term.

References

- Armstrong, Jon Scott. 2001. *Principles of forecasting: a handbook for researchers and practitioners*. Vol. 30 Springer Science & Business Media.
- Baciu, Ionut-Cristian. 2015. "Stochastic models for forecasting inflation rate. Empirical evidence from Romania." *Procedia Economics and Finance* 20:44–52.
- Box, George EP and Gwilym M Jenkins. 1976. "Time series analysis: Forecasting and control San Francisco." *Calif: Holden-Day* .
- Culling, WEH. 1960. "Analytical theory of erosion." The Journal of Geology 68(3):336–344.
- de Melo, Fernando, Valery V Maslennikov, Elena V Popova, Tatyana L Bezrukova and Irina V Kyksova. 2015. "Quantitative analysis in economics based on wavelet transform: a new approach." *Asian Social Science* 11(20):66.
- Fisher, Ronald Aylmer. 1929. "Tests of significance in harmonic analysis." *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* 125(796):54–59.
- Ftiti, Zied, Essahbi Essaadi et al. 2008. The inflation Targeting effect on the inflation series: ANew Analysis Approach of evolutionary spectral analysis. In *GATE Working Paper Number 08–32*.
- Hamilton, James Douglas. 2020. Time series analysis. Princeton university press.
- Iacobucci, Alessandra. 2005. Spectral analysis for economic time series. In *New Tools of Economic Dynamics*. Springer pp. 203–219.
- Konarasinghe, WGS, NR Abeynayake and LHP Gunaratne. 2016. "Circular model on forecasting returns of Sri Lankan Share market." *International Journal of Novel Research in Physics, Chemistry and Mathematics* 3(1):49–56.
- Mankiw, Gregory N. 2018. Principles of Economics. 8th ed. Cengage Learning.
- Moser, Gabriel, Fabio Rumler and Johann Scharler. 2007. "Forecasting austrian inflation." *Economic Modelling* 24(3):470–480.
- Russel, R. 1985. "Significance tables for the results of Fast Fourier Transforms." *Br J Math Stat Psychol* 38:116–119.

Sella, Lisa, Gianna Vivaldo, Michael Ghil and Andreas Groth. 2010. "Economic cycles and their synchronization: Spectral analysis of macroeconomic series from Italy, The Netherlands, and the UK." *EGUGA* p. 11847.

Thomson, Daniel and Gary Van Vuuren. 2016. "Forecasting the South African business cycle using Fourier analysis." *International Business & Economics Research Journal (IBER)* 15(4):175–192.

Weisstein, Eric W. 2004. "Fourier transform." https://mathworld. wolfram. com/.