

# W203 Lab 3: Reducing Crime by Regression Analysis

*Thomas Drage, Venkatesh Nagapudi, Miguel Jaime*

*November 2018*

## 1. Introduction

This statistical investigation aims to understand the determinants of crime to suggest policies to the local government. The study is based upon development of causal models for crime rate, based on county level demographic and judicial data for 1987. We identified factors which modify the rate and extended this to the development of policy proposals for the incoming administration.

## 2. Review of Source Data

```
rm(list = ls())
crime_data = read.csv("crime_v2.csv")
objects(crime_data)
```

```
## [1] "avgsen" "central" "county" "crmte" "density" "mix"
## [7] "pctmin80" "pctymle" "polpc" "prbarr" "prbconv" "prbpris"
## [13] "taxpc" "urban" "wcon" "west" "wfed" "wfir"
## [19] "wloc" "wmfg" "wser" "wsta" "wtrd" "wtuc"
## [25] "year"
```

Overview of type and number of observations:

```
str(crime_data)

## 'data.frame': 97 obs. of 25 variables:
## $ county : int 1 3 5 7 9 11 13 15 17 19 ...
## $ year : int 87 87 87 87 87 87 87 87 87 87 ...
## $ crmte : num 0.0356 0.0153 0.013 0.0268 0.0106 ...
## $ prbarr : num 0.298 0.132 0.444 0.365 0.518 ...
## $ prbconv : Factor w/ 92 levels "", "\", "0.068376102", ...: 63 89 13 62 52 3 59 78 42 86 ...
## $ prbpris : num 0.436 0.45 0.6 0.435 0.443 ...
## $ avgsen : num 6.71 6.35 6.76 7.14 8.22 ...
## $ polpc : num 0.001828 0.000746 0.001234 0.00153 0.00086 ...
## $ density : num 2.423 1.046 0.413 0.492 0.547 ...
## $ taxpc : num 31 26.9 34.8 42.9 28.1 ...
## $ west : int 0 0 1 0 1 1 0 0 0 0 ...
## $ central : int 1 1 0 1 0 0 0 0 0 0 ...
## $ urban : int 0 0 0 0 0 0 0 0 0 0 ...
## $ pctmin80: num 20.22 7.92 3.16 47.92 1.8 ...
## $ wcon : num 281 255 227 375 292 ...
## $ wtuc : num 409 376 372 398 377 ...
## $ wtrd : num 221 196 229 191 207 ...
## $ wfir : num 453 259 306 281 289 ...
## $ wser : num 274 192 210 257 215 ...
## $ wmfg : num 335 300 238 282 291 ...
## $ wfed : num 478 410 359 412 377 ...
## $ wsta : num 292 363 332 328 367 ...
## $ wloc : num 312 301 281 299 343 ...
## $ mix : num 0.0802 0.0302 0.4651 0.2736 0.0601 ...
## $ pctymle : num 0.0779 0.0826 0.0721 0.0735 0.0707 ...
```

There are 97 of them.

## Data Cleansing

Initially, we examined the data and removed values which were measurement or recording errors and ensured the formatting of the dataset was consistent and able to be processed.

1. “NA” data is removed in some cases (there were 6 empty rows at the end of the dataset) .

```
crime_data_corr = na.omit(crime_data)
```

2. Due to the presence of a random back-tick character in the now removed “NA” rows at the end of the dataset, the Variable prbconv was interpreted as a factor of levels - we can convert it back to numeric data with no loss.

```
crime_data_corr$prbconv_fix = as.numeric(as.character(crime_data_corr$prbconv))
summary(crime_data_corr$prbconv_fix)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
## 0.06838 0.34541 0.45283 0.55128 0.58886 2.12121
```

3. Probability values are > 1 in some cases.

```
sum(crime_data_corr$prbarr > 1)
```

```
## [1] 1
```

```
sum(crime_data_corr$prbconv_fix > 1)
```

```
## [1] 10
```

```
sum(crime_data_corr$prbpris > 1)
```

```
## [1] 0
```

There are 11 such values, which we removed as they indicate faulty data.

```
good_prob_cond =
  !((crime_data_corr$prbarr > 1) |
    (crime_data_corr$prbconv_fix > 1) |
    (crime_data_corr$prbpris > 1))
crime_data_corr2 = subset (crime_data_corr, good_prob_cond)
str(crime_data_corr2)
```

```
## 'data.frame':   81 obs. of  26 variables:
## $ county      : int  1 5 7 9 11 13 15 17 21 23 ...
## $ year        : int  87 87 87 87 87 87 87 87 87 87 ...
## $ crmrte      : num  0.0356 0.013 0.0268 0.0106 0.0146 ...
## $ prbarr      : num  0.298 0.444 0.365 0.518 0.525 ...
## $ prbconv     : Factor w/ 92 levels "", "", "0.068376102", ...: 63 13 62 52 3 59 78 42 23 37 ...
## $ prbpris     : num  0.436 0.6 0.435 0.443 0.5 ...
## $ avgsen      : num  6.71 6.76 7.14 8.22 13 ...
## $ polpc       : num  0.00183 0.00123 0.00153 0.00086 0.00288 ...
## $ density     : num  2.423 0.413 0.492 0.547 0.611 ...
## $ taxpc       : num  31 34.8 42.9 28.1 35.2 ...
## $ west        : int  0 1 0 1 1 0 0 0 1 1 ...
## $ central     : int  1 0 1 0 0 0 0 0 0 0 ...
## $ urban       : int  0 0 0 0 0 0 0 0 1 0 ...
## $ pctmin80    : num  20.22 3.16 47.92 1.8 1.54 ...
## $ wcon        : num  281 227 375 292 250 ...
## $ wtuc        : num  409 372 398 377 401 ...
## $ wtrd        : num  221 229 191 207 188 ...
## $ wfir        : num  453 306 281 289 259 ...
```

```
## $ wser      : num  274 210 257 215 237 ...
## $ wmfg      : num  335 238 282 291 259 ...
## $ wfed      : num  478 359 412 377 391 ...
## $ wsta      : num  292 332 328 367 326 ...
## $ wloc      : num  312 281 299 343 275 ...
## $ mix       : num  0.0802 0.4651 0.2736 0.0601 0.3195 ...
## $ pctymle   : num  0.0779 0.0721 0.0735 0.0707 0.0989 ...
## $ prbconv_fix: num  0.5276 0.2679 0.5254 0.4766 0.0684 ...
```

4. There is a duplicate entry for county #193, which we removed from the dataset.

```
crime_data_corr2[crime_data_corr2$county == 193, 1:6]
```

```
##   county year   crmrte   prbarr   prbconv prbpris
## 88    193   87 0.0235277 0.266055 0.588859022 0.423423
## 89    193   87 0.0235277 0.266055 0.588859022 0.423423
```

```
crime_data_corr3 = crime_data_corr2[!duplicated(crime_data_corr2), ]
```

5. There is a density value of 0.0002 - this is approximately one person in an area the size of Alabama and presumably a measurement error. Therefore, we removed this record from the dataset.

```
good_density = (crime_data_corr3$density > 0.001)
crime_data_corr4 = subset(crime_data_corr3, good_density)
```

After cleansing we have 79 records, which we store as our master dataset.

```
crime_data_clean = crime_data_corr4
```

### 3. Identification of Key Variables

#### Dependent Variable

Crime rate (“crmrte”) is the key dependent variable in this study and represents the number of crimes committed per person in each county.


Summarizing the variable we note a small range of fractional values, centred on a mean of approximately 3.5 crimes per hundred people in the year period.

```
summary(crime_data_clean$crmrte)
```

```
##   Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
## 0.01062 0.02345 0.03059 0.03578 0.04397 0.09897
```


The distribution of crime rate is right-skewed in this dataset but sufficient data is available for modelling.

```
hist(crime_data_clean$crmrte, breaks = 30,
     main = 'Histogram of Crime Rate',
     xlab = 'Crime Rate', cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9)
```



We can take an immediate view of this dependent's correlation with our set of available independent variables, firstly excluding the wage variables:

```
corrplot(cor(crime_data_clean[, c(3,4,26,6,7,8,9,10,14,24,25)]), method="ellipse")
```




We note the cases with particularly high correlation with crmte and examine them below as candidate regression variables.

### Independent Variables - Judicial

1. Probability of Arrest ("prbarr")
2. Probability of Conviction ("prbconv")
3. Probability of Going to Prison ("prbpris")
4. Average Sentence ("avgscn")


It is likely that crime rate will be lower when the probability of getting arrested, convicted or going to prison is higher due to the deterrent effect. These variables are expected to have causal relationships with crime rate ("crmte") and should reveal correlation, which we examine through a scatterplot matrix:

```
scatterplotMatrix(~ log(crmte) + prbarr + prbconv_fix + prbpris, data=crime_data_clean)
```



```
crime_data_clean %>%
  keep(is.numeric) %>%
  gather() %>%
  ggplot(aes(value)) + facet_wrap(~key, scales = "free") + geom_histogram(bins = 30)
```

```
## Warning: Computation failed in `stat_bin()`:  
## `binwidth` must be positive
```



The  $\log(\text{cmrte})$  is negatively correlated with  $\text{prbarr}$  and  $\text{prbconv\_fix}$ , which is intuitive. There is perhaps

a positive correlation to prbpris, the probability of prison sentencing, which is not intuitive, but the direction of the correlation is not clear from the dataset, therefore we excluded this from our key variable set.

Anlyzing the average sentence (“avgsen”):

```
summary(crime_data_clean$avgsen)


##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##      5.450   7.450   8.990   9.441  11.180  17.410

cor(crime_data_clean$crm rte, crime_data_clean$avgsen )

## [1] 0.1195381
```

There is a small correlation, but it is unclear as to whether there will be a causal relationship and which way it would be directed.

```
scatterplotMatrix(~ log(crm rte) + avgsen, data=crime_data_clean)
```



## Independent Variables - Demographic


1. Police per capita (“polpc”)
2. Density (“density”)
3. Tax revenue per capita (“taxpc”)
4. Percentage of Young males (“pctymle”)
5. Percentage of minorities (“pctmin80”)

The second set of independent variables are demographic factors which may lead to changes in crime rate, typically in relation to the affluence of the county. Given that the data is collected at county level, these represent an average and any one county may contain a mix of areas (urban/suburban, wealthy/low-income) with corresponding variations in demographics and crimes, which are not captured in this dataset.

## Policing / Density / Tax Revenue

We examined the effect of police staffing, population density and tax revenue:

```
scatterplotMatrix(~ log(crmrte) + polpc + density + taxpc, data=crime_data_clean)
```



Crime Rate is positively correlated to police per capita. We considered police staffing a lagging indicator: where crime rate is high, more police officers are deployed.

Looking at population density, there is a positive correlation between crime and density. This is not unexpected given high density housing is often associated with lower incomes and, in some cases, social issues. The density distribution is not normal, and might need to be transformed.

```
summary(crime_data_clean$density)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
## 0.3006  0.5727   1.0262   1.5363  1.5962   8.8277
```

```
cor(crime_data_clean$crmrte, crime_data_clean$density)
```

```
## [1] 0.7197649
```

Tax revenue per capita (“taxpc”) can be considered a proxy for the income level of a county. We assume that the higher the tax paid the more likely that the people are, on average, more wealthy. Wealthier counties might be a more attractive target for property crime; though these effects might be tempered by a higher opportunity cost for committing crime, and higher likelihood of having higher security measures (such as alarms, gated communities, etc.)

```
summary(crime_data_clean$taxpc)
```


```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
## 25.69   30.92   34.87   38.17   40.94   119.76
```

```
cor(crime_data_clean$crmrte, crime_data_clean$taxpc)
```

```
## [1] 0.4807509
```

We see a positive correlation between taxpc and crime rate. The distribution of taxpc is not optimal and we may need to examine outliers closely if this is to be used in modelling.


```
hist(crime_data_clean$taxpc, breaks = 50,
     main = 'Histogram of Tax Revenue Per Capita',
     xlab = 'Tax Revenue Per Capita', cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9)
```



## Wage Data

A number of variables are provided with average wages in various sectors in each county. We can first examine these to see which might be correlated with crime rate or our other key variables:

```
corr_w = cor(crime_data_clean[, c(3,4,26,6,7,8,9,10,14,25,15,16,17,18,19,20,21,22,23)])
corrplot(corr_w[seq(1,10),seq(11,19)])
```




## Minorities and Young Males

Here we examine the relationship between the proportion of young males (“pctymle”) and the percentage of minority population (“pctmin80”) with crime rate:

```
scatterplotMatrix(~ crm rte + pctymle + pctmin80, data=crime_data_clean)
```





The crime rate is higher in places with a higher percentage of young males. The crime rate is also higher when the percentage of minority population is higher. Both variables seem to have non-ideal distributions.

Looking at the correlation between the variables:

```
summary(crime_data_clean$pctymle)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
## 0.06356 0.07546 0.07795 0.08475 0.08377 0.24871
```

```
cor(crime_data_clean$crmrte, crime_data_clean$pctymle)
```

```
## [1] 0.2875495
```

```
summary(crime_data_clean$pctmin80)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##  1.541  10.477  25.629  26.030  38.842  61.942
```

```
cor(crime_data_clean$crmrte, crime_data_clean$pctmin80)
```

```
## [1] 0.1747599
```

The correlation is weak in both cases.


### 3. Data Transformation

#### Crime Rate

As discussed in section 2, our main variable of interest, crime rate, is measured in a way that results in small variations between values and a skewed distribution. The histogram below shows this distribution.

```
hist(crime_data_clean$crmrte, breaks = 30,
     main = 'Histogram of Crime Rate',
     xlab = 'Crime Rate', cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9)
```

```
crime_data_clean['log_crmrte'] = log(crime_data_clean$crmrate)
hist(crime_data_clean$log_crmrte, breaks = 30,
     main = 'Histogram of log(Crime Rate)',
     xlab = 'log(Crime Rate)', cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9)
```



We applied a  $\log()$  transformation, as shown above to the variable which addresses both issues well.


This transformation will change our interpretation, since the model coefficients will represent percentage changes for crime rate. Given the intended usage in reducing this rate and small values of the variable in its original units, this change will make the results easier to interpret.

## Density

Density is right-skewed. Variable becomes more normal if we apply a log transformation.


```
hist(crime_data_clean$density, breaks = 30,
     main = 'Histogram of Density',
     xlab = 'Density', cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9)

hist(log(crime_data_clean$density), breaks = 30,
     main = 'Histogram of Density',
     xlab = 'Density', cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9)
```



The variable has high correlation with our target variable, which increases slightly with the log transformation.

```
scatterplotMatrix(~ log_crmrte + density + log(density), data = crime_data_clean)
```



```
cor(crime_data_clean$log_crmrte, crime_data_clean$density)
```

```
## [1] 0.6387235
```

```
cor(crime_data_clean$log_crmrte, log(crime_data_clean$density))
```

```
## [1] 0.6702706
```

```
crime_data_clean['log_density'] = log(crime_data_clean$density)
```

## 4. Regression Modelling

### Model 1 - minimal using the Judicial system variables only

```
model1 = lm(crime_data_clean$log_crmrte ~
  crime_data_clean$prbarr +
  crime_data_clean$prbconv_fix
)
model1$coefficients
```

```
##              (Intercept)      crime_data_clean$prbarr
##              -2.2602993      -2.5738165
## crime_data_clean$prbconv_fix
##              -0.9779867
```

Our hypothesis underlying this simple model is that the crime rate is correlated with the efficiency of the justice system, all other demographic factors being approximately equal as justice deters and controls the proliferation of criminal activity. The negative coefficients above support this with the probability of arrest being a stronger contributor than the probability of conviction.



However, this is not the most complete model, and the  $R^2$  value is relatively poor and reveals scope for a more sophisticated model:

```
summary(model1)$r.square
```

```
## [1] 0.3731378
```

We can then plot diagnostics for Model 1 to evaluate OLS assumptions:

```
plot(model1, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 1)
plot(model1, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 3)
```



```
lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle
```



```
lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle
```

- *MLR1 Linearity*: The model is linear in the parameters given.
- *MLR2 Random Sampling*: The sampling process is not clear but assumed to be a random selection of counties.
- *MLR3 Colinearity*: Inspection of scatterplots above did not reveal any perfect colinearity amongst the chosen variables.
- *MLR4 Zero Conditional Mean*: The fitted vs. residuals plot above shows a violation of zero-conditional mean for this model. This suggests some non-linearity with our chosen independents.
- *MLR5 Homoskedasticity*: This model is heteroskedastic. We confirm this using the Breusch-Pagan test and note marginal confirmation. For this reason, we will use robust standard errors going forward.

```
bptest(model1)
```

```
##
## studentized Breusch-Pagan test
##
## data: model1
## BP = 6.0363, df = 2, p-value = 0.04889
```

```
plot(model1, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 2)
plot(model1, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 5)
```



$\text{lm}(\text{crime\_data\_clean}\$log\_crrmte \sim \text{crime\_data\_clean}\$prbarr + \text{crime\_data\_cle})$   $\text{lm}(\text{crime\_data\_clean}\$log\_crrmte \sim \text{crime\_data\_clean}\$prbarr + \text{crime\_data\_cle})$

- *MLR6 Normality*: The Q-Q plot indicates a good normality of the residuals.

Finally, there are no points with Cook's distance  $> 1$  which means that there are no outliers with excessive influence.



### Model 3 - using judicial and demographic system variables

Model 3 is a more elaborate model that takes into account both judicial and demographic system variables to come up with a better causal explanation of crime rate. In this model, we included all meaningful variables. We decided to leave out wage-related variables since we did not find them to be relevant to our analysis.

```
model3 = lm(crime_data_clean$log_crmrte ~
  crime_data_clean$prbarr +
  crime_data_clean$prbconv_fix +
  crime_data_clean$prbpris +
  crime_data_clean$avgsen +
  crime_data_clean$polpc +
  crime_data_clean$log_density +
  crime_data_clean$taxpc +
  crime_data_clean$pctmin80 +
  crime_data_clean$pctymle)
```

We can then plot Model 3 to evaluate OLS assumptions:

```
plot(model3, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 1)
plot(model3, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 3)
```



lm(crime\_data\_clean\$log\_crmrte ~ crime\_data\_clean\$prbarr + crime\_data\_clean\$prbconv\_fix + crime\_data\_clean\$prbpris + crime\_data\_clean\$avgsen + crime\_data\_clean\$polpc + crime\_data\_clean\$log\_density + crime\_data\_clean\$taxpc + crime\_data\_clean\$pctmin80 + crime\_data\_clean\$pctymle)


lm(crime\_data\_clean\$log\_crmrte ~ crime\_data\_clean\$prbarr + crime\_data\_clean\$prbconv\_fix + crime\_data\_clean\$prbpris + crime\_data\_clean\$avgsen + crime\_data\_clean\$polpc + crime\_data\_clean\$log\_density + crime\_data\_clean\$taxpc + crime\_data\_clean\$pctmin80 + crime\_data\_clean\$pctymle)

- *MLR1-3*: As per Model 1 above.
- *MLR4 Zero Conditional Mean*: The inclusion of more explanatory variables improves the mean residual, making it closer to zero.
- *MLR5 Homoskedasticity*: This model appears to have improved the scale location plot and the Breusch-Pagan test does not reject homoskedasticity.


```
bptest(model3)
```

```
##
## studentized Breusch-Pagan test
##
## data: model3
## BP = 15.403, df = 9, p-value = 0.08045
```

```
plot(model3, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 2)
plot(model3, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 5)
```



Theoretical Quantiles



Leverage

`lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle` `lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle`

- *MLR6 Normality*: The Q-Q plot shows some deviations from normality which we confirm with a Shapiro-Wilks test. This suggests non-linearity in one or more of our model variables, most prominently in lower quartiles.

```
shapiro.test(model3$residuals)
```

```
##
##  Shapiro-Wilk normality test
##
## data:  model3$residuals
## W = 0.96209, p-value = 0.01916
```

However, we note there are a couple of high influence outliers which may be negatively affecting the regression.

We can then display Model 3 using heteroskedastic robust standard errors:

```
# Using robust errors to compensate for heteroskedasticity
robust_se <- function(model) {
  cov <- vcovHC(model)
  sqrt(diag(cov))
}

robust_errors <- list(robust_se(model3))

stargazer(model3,
  star.cutoffs = c(0.05, 0.01, 0.001),
  se = robust_errors,
  type = 'latex',
  column.labels = 'Model 3',
  font.size = 'small',
  float = FALSE)
```

% Table created by stargazer v.5.2.2 by Marek Hlavac, Harvard University. E-mail: hlavac at fas.harvard.edu % Date and time: Sun, Dec 09, 2018 - 21:01:32

	<i>Dependent variable:</i>
	log_crmrte Model 3
prbarr	-1.545*** (0.357)
prbconv_fix	-0.293 (0.238)
prbpris	-0.361 (0.393)
avgsen	-0.017 (0.012)
polpc	301.725* (129.909)
log_density	0.286*** (0.064)
taxpc	0.004 (0.004)
pctmin80	0.014*** (0.002)
pctymle	1.130 (2.259)
Constant	-3.650*** (0.471)
Observations	79
R <sup>2</sup>	0.827
Adjusted R <sup>2</sup>	0.804
Residual Std. Error	0.222 (df = 69)
F Statistic	36.629*** (df = 9; 69)

Note: \*p<0.05; \*\*p<0.01; \*\*\*p<0.001

We can also evaluate the Akaike Information Criterion for both models to check goodness of fit relative to parsimony:

```
AIC(model1, model3)
```

```
##      df      AIC
## model1  4 85.627029
## model3 11 -2.044801
```

As we can see, Model 3 has significantly improved on the AIC, R<sup>2</sup> and Residual SE, but there are some coefficients which are not statistically significant (prbconv\_fix, polpc). There is likely a more optimized model that has fewer coefficients that we can derive out of the Model 1 and Model 3 experiments above.

By looking at the standardised co-efficients, we can evaluate compare the effect of changes of each variable on the crime rate:

```
lm.beta(model3)
```

```
##
## Call:
## lm(formula = crime_data_clean$log_crmrte ~ crime_data_clean$prbarr +
##      crime_data_clean$prbconv_fix + crime_data_clean$prbpris +
```



```
##      crime_data_clean$avgsen + crime_data_clean$polpc + crime_data_clean$log_density +
##      crime_data_clean$taxpc + crime_data_clean$pctmin80 + crime_data_clean$pctymle)
##
## Standardized Coefficients::
##              (Intercept)      crime_data_clean$prbarr
##              0.00000000      -0.32747628
## crime_data_clean$prbconv_fix  crime_data_clean$prbpris
##              -0.10072985      -0.05264131
##      crime_data_clean$avgsen      crime_data_clean$polpc
##              -0.07972709      0.30646668
## crime_data_clean$log_density      crime_data_clean$taxpc
##              0.43990931      0.09349452
##      crime_data_clean$pctmin80      crime_data_clean$pctymle
##              0.45722169      0.05433308
```

Based on the above we may consider removing those with lower (e.g.  $<0.1$ ) gain in addition to those which are not statistically significant.

## Model 2 - with optimized Judicial and Demographic system variables

From the above models, it is clear that some of the variables added to the model such as the density, polpc and pctmin80 show particularly strong contribution to the model. The probconv\_fix variable seems to of lower significance in the Model 3, possible it correlates heavily with other variables and therefore decreasing in significance.

We therefore select the following for our second model:

1. prbarr
2. density
3. pctmin80
4. prbconv\_fix


Note that we elected to remove the police per capita variable (“polpc”) from this model as we believe it is likely an effect rather than a cause, and heavily correlated to other variables in the regression.

Creating Model 2 out of these variables:

```
model2 = lm(crime_data_clean$log_crmrte ~
             crime_data_clean$prbarr +
             crime_data_clean$log_density +
             crime_data_clean$prbconv_fix +
             crime_data_clean$pctmin80)
```


We can then plot diagnostics for Model 3 to evaluate OLS assumptions:

```
plot(model2, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 1)
plot(model2, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 3)
```



Fitted values

`lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle`



Fitted values


`lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle`

- *MLR1-3*: As per Model 1 above.
- *MLR4 Zero Conditional Mean*: Further improvement and no violation of zero conditional mean in this model.
- *MLR5 Homoskedasticity*: This model appears to have further improved the scale location plot and the Breusch-Pagan test does not reject homoskedasticity.

```
bptest(model2)
```

```
##
## studentized Breusch-Pagan test
##
## data: model2
## BP = 12.593, df = 4, p-value = 0.01344
```

```
plot(model2, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 2)
plot(model2, cex.main = 0.9, cex.lab = 0.9, cex.axis = 0.9, cex.sub = 0.5, which = 5)
```



```
lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle
```

```
lm(crime_data_clean$log_crmrte ~ crime_data_clean$prbarr + crime_data_cle
```

- *MLR6 Normality:* The Q-Q plot is more normal than Model 3. So the coefficients are more robust. We see that the p-values are all very significant unlike Model 3's p-values showing that the coefficients are more consistent. The Shapiro-Wilks test does not reject normality:

```
shapiro.test(model2$residuals)
```

```
##
## Shapiro-Wilk normality test
##
## data: model2$residuals
## W = 0.91475, p-value = 6.013e-05
```

Additionally, are no outliers with high influence in this model specification.

We can now compare all three models:

```
robust_errors <- list(robust_se(model1),
                     robust_se(model2),
                     robust_se(model3))

stargazer(model1, model2, model3,
           star.cutoffs = c(0.05, 0.01, 0.001),
           se = robust_errors,
           type = 'latex',
           column.labels = c('Model 1', 'Model 2', 'Model 3'),
           font.size = 'small',
           float = FALSE)
```

% Table created by stargazer v.5.2.2 by Marek Hlavac, Harvard University. E-mail: hlavac at fas.harvard.edu % Date and time: Sun, Dec 09, 2018 - 21:01:33

	<i>Dependent variable:</i>		
		log_crmrte	
	Model 1	Model 2	Model 3
	(1)	(2)	(3)
prbarr	-2.574*** (0.535)	-1.859*** (0.397)	-1.545*** (0.357)
log_density		0.348*** (0.070)	0.286*** (0.064)
taxpc			0.004 (0.004)
prbconv_fix	-0.978** (0.328)	-0.711** (0.274)	-0.293 (0.238)
prbpris			-0.361 (0.393)
avgsen			-0.017 (0.012)
polpc			301.725* (129.909)
pctmin80		0.013*** (0.002)	0.014*** (0.002)
pctymle			1.130 (2.259)
Constant	-2.260*** (0.259)	-2.966*** (0.272)	-3.650*** (0.471)
Observations	79	79	79
R <sup>2</sup>	0.373	0.732	0.827
Adjusted R <sup>2</sup>	0.357	0.718	0.804
Residual Std. Error	0.403 (df = 76)	0.267 (df = 74)	0.222 (df = 69)
F Statistic	22.619*** (df = 2; 76)	50.557*** (df = 4; 74)	36.629*** (df = 9; 69)

*Note:*

\*p<0.05; \*\*p<0.01; \*\*\*p<0.001

**AIC(model1, model2, model3)**

```
##          df          AIC
## model1  4 85.627029
## model2  6 22.466835
## model3 11 -2.044801
```

Model 2 shows significant improvement over Model1 with a better AIC, much better R<sup>2</sup> and lower Residual SE.

## 5. Discussion - Model Specification & Omitted Variables

It is likely that crime rate will be heavily influenced by the following omitted variables:

1. Demographics: There is very little information on demographics other than pctmin80 which is based on dated information about minorities; the nature and proportion of particular minorities is omitted here. If, say, a particular cultural group was more prone to crime we would expect greater crime rate. It could be useful to get additional information on the type of people comprising the

county population, and inclusion of variables expressing religious make-up could, for example show less crime in regions densely populated by religions valuing non-violence.

2. Education level: Typically we would expect higher the education level to lead to lower the crime rate. This is likely related to educated decision making but also to employment outcomes which promote stability.
3. Wages: The more affluent neighborhoods will tend to have lesser crime. We thought this would be reflected by tax revenues per capita, but it does not appear to be the case. This may be due to the nature of the taxation system itself in creating a representative variable.
4. Employment: Gainful employment of citizens is well known to decrease crime rate however, we have no information about the employment rates in each county. The only possibility is to compare the wage rates in each county which could express greater employment when competition has driven rates up. Limited employment may result in the presence of higher-density (social) housing and therefore positively bias the density coefficient.
5. Commercial Sectors: High crime might be associated with regions which have a lot of bars or similar entertainment venues but less crime in rural residential areas. Such an omitted variable (say indication number of nightlife venues) would potentially bias our density dependence by increasing the co-efficient or even increase the dependence of our model probability of arrest due to police presence in entertainment areas.
6. Age distribution: We are provided with a variable indicating the percent of the population who are young males, but it would be equally useful to be provided with data for children and geriatrics - both parties whose presence would decrease the crime rate. Both of these groups typically increase density, but decrease crime and we would expect their presence to bias the density variable negatively.
7. More detailed crime information: Not all crime is equal, and different types of crime (violent crime, property crime, etc.) might be explained by different factors. Such omitted variables bias the probability of arrest, conviction and average sentence variables. Sentence length, for example, would be dependant on a measure of the proportion of non-violent crime and we may find sentence length is also negatively biased by this omission.

## 6. Conclusion

Based on our analysis, the probability of arrest and conviction help drive down crime rates. Increasing the probability of arrest by one unit is correlated with a 186% decrease in crime rate. Increasing the probability of conviction by one unit is correlated with a 71% decrease in crime rate.

Density is positively correlated with crime rate. An increase of one unit in density is correlated with a 35% increase in the crime rate.

Based on these results, our policy recommendations would be to:

1. Increase awareness of the effectiveness of the judicial system in counties that are effective at bringing perpetrators to justice; and increase resources, training, and oversight in those that are not.
2. Further investigation of the link between population density and crime rate. Are there economic factors at play? Demographics? Policing techniques in urban settings?