n/p - SMASH check

Initial conditions (input config)

Used Skyrme and Symmetry potentials in SMASH simulations:


- Skyrme: $U_{Sk}=\mathbf{A}(\rho/\rho_0)+\mathbf{B}(\rho/\rho_0)^{\tau}$
- Symmetry: $U_{Sym}=\pm 2\mathbf{S_{pot}}\, I_3/I\, \rho_{I3}/\rho_0+S(\rho_B)(\rho_{I3}/\rho_B)^2$


500k events for each data set:

- Set 1: K_0=349.5 MeV, $\mathbf{S_{pot}}$**=0 MeV**
- Set 2: K_0=349.5 MeV, $\mathbf{S_{pot}}$**=18 MeV**
- Set 3: K_0=349.5 MeV, $\mathbf{S_{pot}}$**=90 MeV**

* 10 parallel ensembles were used to produce realistic mean field

Xe+Cs @ 2.5A GeV: E_{kin}-dependence







E_{kin} cut: E_{kin}>300 MeV

Sensitivity to S_{pot} is observed


Xe+Cs @ 3.8A GeV: y-dependence with E_{kin}>300 MeV






Sensitivity to S_{pot} is observed

Xe+Cs @ 3.8A GeV: p_T-dependence with E_{kin}>300 MeV







Sensitivity to S_{pot} is observed

Backup


Xe+Cs @ 3.8A GeV: y-dependence






Sensitivity to S_{pot} is observed

Xe+Cs @ 3.8A GeV: p_T-dependence





Sensitivity to S_{pot} is observed