The value of B_n/C_n is used in the estimate below.

We then obtain $\tau = \theta/2\pi + 4\pi i/g^2$ holds throughout.

Recall that A_n by the previous lemma.

It follows that A_μ as claimed.

Suppose $x_2 + ix_3 = re^{i\theta}$ in what follows.

Note that $\mathcal{A} = A + i\phi$ for the remainder of the section.

Here n_k and the result follows.

In this case $\lambda_m^{n_m}\lambda_{m-1}^{n_{m-1}}\cdots\lambda_1^{n_1}$ is used in the estimate below.

The value of (D_n) holds throughout.

We then obtain B_k by the previous lemma.

Recall that $\lambda'' = 0$ as claimed.

It follows that r_k' in what follows.

Suppose $l - k + \lambda_k$ for the remainder of the section.

Note that g_* and the result follows.

Here $\frac{1}{2}(\sum_{k=i}^{m} n_k)$ is used in the estimate below.

In this case $l + 1$ holds throughout.

The value of S_{e1221} by the previous lemma.

We then obtain ρ_{odd} as claimed.

Recall that C_{16} in what follows.

It follows that $(\rho_{\text{even}} + X_S^{-1}\rho_{\text{even}}\,;\rho_{\text{odd}} + Y_S\rho_{\text{odd}})_B$ for the remainder of the section.

Suppose $(\lambda_{even}\,;\rho_{odd})$ and the result follows.

Note that $L(C1)$ is used in the estimate below.

Here C_1 holds throughout.

In this case $L(D2)$ by the previous lemma.

The value of (b) as claimed.

We then obtain $c22$ in what follows.

Recall that IIB for the remainder of the section.

It follows that $(\lambda, \rho)_C = (\lambda_{odd}, \rho_{even})_C$ and the result follows.

Suppose CB_{eo} is used in the estimate below.

Note that $\mathcal{I} \in \mathbb{Z}$ holds throughout.

Here $(\psi(0), \psi'(0))$ by the previous lemma.

In this case $(h_{R,t})$ as claimed.

The value of $\mathcal{H}_1$ in what follows.

We then obtain $\mathbf{y} = (y, y')$ for the remainder of the section.

Recall that $\ell_2 \subset \ell_2^{\circ}$ and the result follows.

It follows that $J^* : \mathcal{H}_b \to \mathcal{H}_b$ is used in the estimate below.

Suppose $\ell_2(t)$ holds throughout.

Note that $C_0^{\infty}(\mathbb{R}^+, \mathbb{C}^n)$ by the previous lemma.

Here $(0, u') \in \ell_D$ as claimed.

In this case $\ell^+(E) \cap \ell^-(E) \neq \{0\}$ in what follows.

The value of $(h - E)\psi$ for the remainder of the section.

We then obtain $\ell(t)$ and the result follows.

Recall that $U_1(t)$ is used in the estimate below.

It follows that $P_1 x_j = x_j$ holds throughout.

Suppose $\sigma\left(h_t, \mathbb{T}^1\right)$ by the previous lemma.

Note that $(u_1, \cdots, u_j)$ as claimed.

Here $\ell_t^{\sharp}\{(\Theta x, \Pi x),\ x \in \mathbb{C}\}$ in what follows.

In this case $\mathbf{k} \in \mathbb{Z}^{d-1}$ for the remainder of the section.

The value of $L^2(\Gamma) = H^{s=0}(\gamma)$ and the result follows.

We then obtain $\chi(x)$ is used in the estimate below.

Recall that $(H - E)\psi = 0$ holds throughout.

It follows that $H^2(\Omega^-)$ by the previous lemma.

Suppose $R_{t,D} := (H_{t,D} - \Sigma)^{-1}$ as claimed.

Note that $\ell_{R,t}^{\pm}(E)$ in what follows.

Here $\widetilde{H}_t$ for the remainder of the section.

In this case $L^2(\mathbb{R}^+ \times \mathbb{R})$ and the result follows.

The value of $\gamma^D \psi_D = 0$ is used in the estimate below.

We then obtain $H^{-1/2}(\Gamma) \times H^{1/2}(\Gamma)$ holds throughout.

Recall that $G_{\Sigma} f$ by the previous lemma.

It follows that $\mathrm{LSwHom}(H)$ as claimed.

Suppose G^R in what follows.

Note that $\phi : G' \to H$ for the remainder of the section.

Here (u, v) and the result follows.

In this case $\phi(v) \in L(v)$ is used in the estimate below.

The value of H_c holds throughout.

We then obtain $\pi_S^{-1}(i) = \{(v, i) \mid v \in v(H)\}$ by the previous lemma.

Recall that $(u, 0)$ as claimed.

It follows that L_S in what follows.

Suppose $u, v \in V$ for the remainder of the section.

Note that $x_i \sim y_i$ and the result follows.

Here $u_{i-1} \not\sim v_i$ is used in the estimate below.

In this case B_a holds throughout.

The value of $V(H)^3$ by the previous lemma.

We then obtain x_3 as claimed.

Recall that $i \neq i'$ in what follows.

It follows that $i = \pi_B((i, v))$ for the remainder of the section.

Suppose $\phi : H^k \to H$ and the result follows.

Note that $(i, h, j)(i', h', j')$ is used in the estimate below.

Here $a', b' \in V$ holds throughout.

In this case (y, x) by the previous lemma.

The value of $x_a x_b$ as claimed.

We then obtain r_a in what follows.

Recall that $x_a b$ for the remainder of the section.

It follows that r_b and the result follows.

Suppose $X = \mathrm{maj}(x_1, x_2, x_3)$ is used in the estimate below.

Note that $(a_i, x_i) \sim (b_i, y_i)$ holds throughout.

Here $A \leq a_i$ by the previous lemma.

In this case (a_2, b_1, c_1) as claimed.

The value of $\phi(a_2, d, d) = a_2$ in what follows.

We then obtain $G \cong F$ for the remainder of the section.

Recall that $u \in U \setminus \{1\}$ and the result follows.

It follows that $v \in V(\Gamma)$ is used in the estimate below.

Suppose $\gamma_{n+1}(G) = [\gamma_n(G), G]$ holds throughout.

Note that kI_H by the previous lemma.

Here $p(b, c) = b + c$ as claimed.

In this case $p(b, c) = b + c(t - 1)$ in what follows.

The value of $\widetilde{G} \to G$ for the remainder of the section.

We then obtain M_3 and the result follows.

Recall that $\mathcal{D}_{\mathbb{F}_p H}$ is used in the estimate below.

It follows that $\mathbb{F}_p I_G$ holds throughout.

Suppose $A = H_1 \cap H_2$ by the previous lemma.

Note that u_i as claimed.

Here $\mathcal{D}_{\mathbb{F}_p[H]}$ in what follows.

In this case $c_k \in \mathbb{F}_p N$ for the remainder of the section.

The value of $\widetilde{H_1}$ and the result follows.

We then obtain $i \in \{1, 2\}$ is used in the estimate below.

Recall that u_2 holds throughout.

It follows that $u_2 = v_2^{n_2}$ by the previous lemma.

Suppose $r_{\mathrm{ab}}(\widetilde{H_1}) + r_{\mathrm{ab}}(\widetilde{H_2}) - 1$ as claimed.

Note that ϕ_i in what follows.

Here $\widetilde{H_i} \cong H_i$ for the remainder of the section.

In this case $\widetilde{H}_{\widehat{p}}$ and the result follows.

The value of w_1 is used in the estimate below.

We then obtain $\widetilde{H}_{\widehat{p}} \coprod \mathbb{Z}_p$ holds throughout.

Recall that $d(\widetilde{H}_{\widehat{p}}) = d(\overline{H})$ by the previous lemma.

It follows that $q|_G$ as claimed.

Suppose $u \equiv (uv^{-1})^c \mod [U, U]$ in what follows.

Note that $\gamma_k W_{\widehat{p}}$ for the remainder of the section.

Here $\Gamma \setminus \{e\} = \Delta_1 \cup \Delta_2$ and the result follows.

In this case $\mathbb{Q}(\sqrt{1 - 2m^k})$ is used in the estimate below.

The value of $m \geq 3$, holds throughout.

We then obtain J_K by the previous lemma.

Recall that $n > 3$ as claimed.

It follows that $\mu = \sqrt{2}$ in what follows.

Suppose $k = p^r$ for the remainder of the section.

Note that $(n = 1)$ and the result follows.

Here $p_i \mid h_{(m,p_i)}$ is used in the estimate below.

In this case $i = 1$ holds throughout.

The value of $\mathbb{Q}(\sqrt{1 - 2m^k})$, by the previous lemma.

We then obtain $, where$ as claimed.

Recall that , $then$ in what follows.

It follows that , $whichisacontradictionto$ for the remainder of the section.

Suppose $doesnotdivide$ and the result follows.

Note that $Then$ is used in the estimate below.

Here $where$ holds throughout.

In this case , for by the previous lemma.

The value of $.Henceweget$ as claimed.

We then obtain $andhencewehave$ in what follows.

Recall that $.Takingmoduloover4onbothsides, weget$ for the remainder of the section.

It follows that $isco-primeto$ and the result follows.

Suppose $isoddpositiveinteger, weget$ is used in the estimate below.

Note that $musthaveprimeidealswhosenormsare$ holds throughout.

Here $.Hencetheprimedecompositionof$ by the previous lemma.

In this case $.Considertheideal$ as claimed.

The value of $forsomeoddprime$ in what follows.

We then obtain $dividesclassnumbersof$ for the remainder of the section.

Recall that $betheclassnumbersof$ and the result follows.

It follows that $suchthat$ is used in the estimate below.

Suppose $isaprimewhichdividesanoddinteger$ holds throughout.

Note that $H^3(G, \mathbb{T})$ by the previous lemma.

Here $\omega_1 \otimes_{\text{stack}} \omega_2$ as claimed.

In this case $\text{Index}(\omega \circ \mu)$ in what follows.

The value of $\Gamma = \mathbb{Z}$ for the remainder of the section.

We then obtain $\mathcal{A}_\Gamma$ and the result follows.

Recall that $\tau(\Phi(I)) = \Phi(\tau(I))$ is used in the estimate below.

It follows that $\text{Index}_{1d}^{\mathcal{A}_{L_0}, U}(\phi) = \text{Index}^{\mathcal{A}, U}(\omega)$ holds throughout.

Suppose $\gamma_{0;1}^{\Psi}$ by the previous lemma.

Note that $\alpha_\lambda(g) = \exp(2\pi i k/n)$ as claimed.

Here $\text{Index}_{1d\ \text{trans}}(\omega) = \text{Index}_{1d\ \text{trans}}(\omega \circ \gamma_{0;1}^{\Phi'})$ in what follows.

In this case $x \in \text{Cone}_\sigma^W(\alpha_0, \theta)$ for the remainder of the section.

The value of $[\pi_0 \circ \alpha_0 \circ \Theta(A_L), v_R]$ and the result follows.

We then obtain $\tau_{\mathcal{B}}$ is used in the estimate below.

Recall that $b_{g}^{R} \in \mathrm{Cone}_{R}^{W}(\alpha_{0}, \theta)$ holds throughout.

It follows that $h_{g} \in \mathrm{IAC}_{L}(\theta, \alpha_{0}, \Theta)$ by the previous lemma.

Suppose $W_{g,1}, u_{R,1}(g,h), v$ as claimed.

Note that $W_{g,1}\delta_{g}^{\sigma}W_{g,1}^{\dagger} \in \mathrm{IAC}_{\sigma}(\alpha_{0}, \theta, \Theta)$ in what follows.

Here $n \in \mathbb{N}$ for the remainder of the section.

In this case n_{0} and the result follows.

The value of $\mathrm{Index}(\omega \circ \gamma_{0;1}^{H}) = \mathrm{Index}(\omega)$ is used in the estimate below.

We then obtain $a \in \mathrm{IAC}_{L}(\alpha_{0}, \theta, \Theta)$ holds throughout.

Recall that $x \in \mathrm{IAC}_{L\times R}^{W}(\alpha_{0}, \theta, \Theta)$ by the previous lemma.

It follows that $B_{\sigma} \in \mathcal{U}(\mathcal{A})$ as claimed.

Suppose $\omega \circ \gamma_{0;1}^{H} \circ \tilde{\beta}_{g,2} = \omega \circ \gamma_{0;1}^{H}$ in what follows.

Note that $\{1,2,3,4\}$ for the remainder of the section.

Here $\pi_{0} = \bigotimes_{i=1}^{4} \pi_{0,i}$ and the result follows.

In this case $(\mathcal{H}_{0}, \pi_{0} \circ i_{V}, \Omega_{0})$ is used in the estimate below.

The value of $B_{n,L}$ holds throughout.

We then obtain $\Xi^{(s)}(Z,t)$ by the previous lemma.

Recall that $X \subset U, X \subset L, X \subset R$ as claimed.

It follows that $\mathcal{K} := \mathrm{age}(\mathcal{A})$ in what follows.

Suppose $\mathrm{age}(\mathcal{I})$ for the remainder of the section.

Note that $\bar{\imath}' := (i_{k} : k < s) \in X^{s}$ and the result follows.

Here $\mathrm{Diag}_{\mathcal{A}}(\overline{x})$ is used in the estimate below.

In this case $\mathcal{A} \cong_{L} \mathcal{A}'$ holds throughout.

The value of $\mathrm{qftp}^{\mathcal{A}}(\overline{a})$ by the previous lemma.

We then obtain $h : B \to C$ as claimed.

Recall that $L = \{E, <\}$ in what follows.

It follows that $A, B \subseteq \mathcal{M}$ for the remainder of the section.

Suppose $\alpha(e,x) = x$ and the result follows.

Note that $h : \mathcal{A} \to \mathcal{B}$ is used in the estimate below.

Here $Y \vDash \mathrm{EMtp}(X)$ holds throughout.

In this case $||\mathcal{I}||^{+}$ by the previous lemma.

The value of $\mathcal{I} = (\mathbb{Z}, p, s, <)$ as claimed.

We then obtain $a_{i} \prec a_{j}$ in what follows.

Recall that $s \neq t$ for the remainder of the section.

It follows that $a_i \neq a_j$ and the result follows.

Suppose $(g(i) \mid i \in \mathcal{A})$ is used in the estimate below.

Note that $\eta, \nu \in \omega^{<\omega}$ holds throughout.

Here $\operatorname{age}(\mathcal{A}) = \operatorname{age}(\mathcal{I}_{\mathrm{eq}})$ by the previous lemma.

In this case $B = \{b_i : i < \omega\}$ as claimed.

The value of $(\mathcal{B}_{ba}, \prec)$. in what follows.

We then obtain $, itfollowsimmediatelythat$ for the remainder of the section.

Recall that $bestructuresandsuppose$ and the result follows.

It follows that $andfixgenerators$, is used in the estimate below.

Suppose $.Ifthereexistsan$ holds throughout.

Note that $homogeneousforthecoloring$ by the previous lemma.

Here $isthedesiredhomogeneouscopyof$ as claimed.

In this case $hasapreimagethatissomewherein$ in what follows.

The value of $with$ for the remainder of the section.

We then obtain $(X, \Delta) \to Z$ and the result follows.

Recall that (X, Δ) is used in the estimate below.

It follows that $f \colon X' \to X$ holds throughout.

Suppose $e \in \{1, -1\}$ by the previous lemma.

Note that $h \colon W \to T'$ as claimed.

Here $D = \sum_i d_i D_i$ in what follows.

In this case $Y_1 \to X$ for the remainder of the section.

The value of $g \colon V \to Y$ and the result follows.

We then obtain B_Y is used in the estimate below.

Recall that $K_{X'} + \Delta' = \psi^*(K_X + \Delta)$ holds throughout.

It follows that $S^\nu \to S$ by the previous lemma.

Suppose $a(E, X, \Delta) < a(E, X', \Delta')$ as claimed.

Note that E' in what follows.

Here $m\Delta = f_*(mf^*(K_X + \Delta)) - mK_X$ for the remainder of the section.

In this case n'' and the result follows.

The value of Σ' is used in the estimate below.

We then obtain $V' = f^{-1}(V)$ holds throughout.

Recall that π' by the previous lemma.

It follows that $\left(V', (\Delta' + t_0 A')|_{V'}\right)$ as claimed.

Suppose $(K_{X'} + \Delta' + t_0 A')$ in what follows.

Note that π_Y for the remainder of the section.

Here $\lfloor \Gamma \rfloor$ and the result follows.

In this case $K_{X'} + \Theta' = f^*(K_X + \Delta)$ is used in the estimate below.

The value of $\operatorname{mult}_{Q_j}(\pi_Y^* D_j) \leq \beta$ holds throughout.

We then obtain $(X, \Delta + tB)$ by the previous lemma.

Recall that $m(K_{X'} + \Delta')$ as claimed.

It follows that $G \sim IA$ in what follows.

Suppose $2n''eD$ for the remainder of the section.

Note that $B' \geq 0$ and the result follows.

Here $\pi' \colon X' \dashrightarrow W$ is used in the estimate below.

In this case $\mathcal{F}(R, \mathcal{G})$ holds throughout.

The value of $\mathcal{F}(R)$ by the previous lemma.

We then obtain S_0 as claimed.

Recall that $\mathcal{F}(\mathcal{R}, \mathcal{G})$ in what follows.

It follows that $P_{\mu\nu}$ for the remainder of the section.

Suppose $\tilde{b} = \frac{4}{3} q^2 \Lambda$ and the result follows.

Note that $E(r) = -A'(r)$ is used in the estimate below.

Here $r \to \infty$ holds throughout.

In this case $\mathcal{H}(P)$ by the previous lemma.

The value of $P <<$ as claimed.

We then obtain $r \to 0$ in what follows.

Recall that $r_h = (8Mq^2)^{1/3}$ for the remainder of the section.

It follows that (i, j, k) and the result follows.

Suppose $T_H = \frac{\kappa}{2\pi}$ is used in the estimate below.

Note that $\beta = 1/T_H$ holds throughout.

Here $r_H <<$ by the previous lemma.

In this case (C) as claimed.

The value of ξ^μ in what follows.

We then obtain A_E for the remainder of the section.

Recall that $n_+ = n_- = n \geq 1$ and the result follows.

It follows that $\beta = \frac{4M\tilde{\Lambda}}{3}$ is used in the estimate below.

Suppose $r^{1+p}/\sqrt{\eta - \beta r}$ holds throughout.

Note that $\delta = \tilde{\Lambda}/3$ by the previous lemma.

Here $||R||^2 \to \infty$ as claimed.

In this case $Z_{\pm} = R_1 \pm R_2$ in what follows.

The value of $L^2\left(0, \infty\right)$ for the remainder of the section.

We then obtain c_{5} and the result follows.

Recall that $\sigma = \sqrt{1 + 4\bar{\lambda}}/2$ is used in the estimate below.

It follows that $6 \leq \bar{\lambda}^{2}$ holds throughout.

Suppose Q_{1}, Q_{5} by the previous lemma.

Note that $a = 0, 1, ..., n-1$ as claimed.

Here $\star_3 \, dA = dH$ in what follows.

In this case T^4 for the remainder of the section.

The value of $(n-1)$ and the result follows.

We then obtain k_{a}^{2} is used in the estimate below.

Recall that Π_{ab}^{2} holds throughout.

It follows that q_{a} by the previous lemma.

Suppose $\lambda = 10^{-5}$ as claimed.

Note that $[x_{1}, \cdots, x_{\mathbf{no}}]$ in what follows.

Here $z = 0$ for the remainder of the section.

In this case $\{k_{\tilde{a}}^{0,1}, k_{0}^{2}\}$ and the result follows.

The value of $Q_{\max} \leftarrow$ is used in the estimate below.

We then obtain $\bar{r}_{a}^{i}$ holds throughout.

Recall that $\{r_{A}^{(p)}, \sigma_{A}^{(p)}\}$ by the previous lemma.

It follows that $\delta r_{A}^{(p)}$ as claimed.

Suppose $f^{\prime}$ in what follows.

Note that $2 / d$ for the remainder of the section.

Here $\mathcal{N}(0, 1)$ and the result follows.

In this case `generation_update` is used in the estimate below.

The value of $\sigma_{A} \to \left(\texttt{factor_random_update} \right) \sigma_{A}$ holds throughout.

We then obtain 200 by the previous lemma.

Recall that 1st as claimed.

It follows that -0.6923 in what follows.

Suppose -61 for the remainder of the section.

Note that -48.5265 and the result follows.

Here $\tilde{f}_{\mathrm{inv}}(r_A) = \max_a |\epsilon_a|$ is used in the estimate below.

In this case $Q_5 \simeq 410$ holds throughout.

The value of 1.1×10^4 by the previous lemma.

We then obtain $(1 + 1)$ as claimed.

Recall that $\gamma \neq 0$ in what follows.

It follows that Z_n for the remainder of the section.

Suppose Z_3 and the result follows.

Note that $\mathbb{C}^3$ is used in the estimate below.

Here $2L^2$ holds throughout.

In this case $\hat{\Delta}_v$ by the previous lemma.

The value of $\varepsilon_v = 0$ as claimed.

We then obtain h_v in what follows.

Recall that $\hat{N}_f$ for the remainder of the section.

It follows that h_f and the result follows.

Suppose B_f is used in the estimate below.

Note that $C_{<f,f'>}$ holds throughout.

Here $\hat{N}_{<f,f'>}$ by the previous lemma.

In this case $-10L^2$ as claimed.

The value of $a = 1, 2, 3$ in what follows.

We then obtain $A^{(a)}$ for the remainder of the section.

Recall that H_{int} and the result follows.

It follows that H_v is used in the estimate below.

Suppose N_i holds throughout.

Note that $p_\sigma = \frac{1}{N_{\mathrm{i}}}$ by the previous lemma.

Here $\mathcal{D} = 4$ as claimed.

In this case $A'^{(a)}_v$ in what follows.

The value of $\hat{\Delta}'_v$ for the remainder of the section.

We then obtain h'_v and the result follows.

Recall that h'_f is used in the estimate below.

It follows that $\{\sqrt{3} + 1, 2, 1, 0, 1 - \sqrt{3}\}$ holds throughout.

Suppose $N_{\mathrm{FPL}} = \binom{2l}{l}$ by the previous lemma.

Note that $n \approx \sigma l$ as claimed.

Here $\mathcal{C}_{x,y}$ in what follows.

In this case T_c for the remainder of the section.

The value of $\theta / z = -1$ and the result follows.

We then obtain $\rho_{xx} \sim \tau_{tr}^{-1} \sim T$ is used in the estimate below.

Recall that $\cot \Theta_H \sim \tau_h^{-1} \sim T^2$ holds throughout.

It follows that $\tilde{\sigma} \propto T^{-1}$ by the previous lemma.

Suppose $\rho_{xx} \propto \alpha T$ as claimed.

Note that $3 + 1$ in what follows.

Here S_{GR} for the remainder of the section.

In this case $F = \mathrm{d}A$ and the result follows.

The value of ψ_I is used in the estimate below.

We then obtain $z = 1$ holds throughout.

Recall that A_t by the previous lemma.

It follows that $T = U(1)/(4\pi)$ as claimed.

Suppose $B = 0$ in what follows.

Note that k/μ for the remainder of the section.

Here $B/\mu^2 = 1/10$ and the result follows.

In this case x, y is used in the estimate below.

The value of ω_0 holds throughout.

We then obtain $1 < \beta \leq 2$ by the previous lemma.

Recall that $\sigma_0, \tilde{\sigma}, n$ as claimed.

It follows that $x_3 = 0$ in what follows.

Suppose $x_1 \leq 1$ for the remainder of the section.

Note that $x_1 = 1$ and the result follows.

Here $\beta = x_3 + x_2 \leq 1$ is used in the estimate below.

In this case $T \gg W$ holds throughout.

The value of $x_3 \neq 0$ by the previous lemma.

We then obtain $\sqrt{a_1 T^2 + a_2 B^2}$ as claimed.

Recall that $|\psi_0\rangle$ in what follows.

It follows that $N = 10$ for the remainder of the section.

Suppose $N = 4, L = 11$ and the result follows.

Note that $S(\theta_1, \ldots, \theta_N)$ is used in the estimate below.

Here $\tau_{\mathrm{cr}} \approx 0.33137171$ holds throughout.

In this case $\ell \geq 2$ by the previous lemma.

The value of $\tau^{\mathrm{QSL}} > 0$ as claimed.

We then obtain $\sigma^{+}_{k_1 \cdots k_N}$ in what follows.

Recall that $\kappa \pm \square_a$ for the remainder of the section.

It follows that $U(1)$ and the result follows.

Suppose $\left\{ e^{i2\pi(s-1)/L},\ s = 1 \ldots, L \right\}$ is used in the estimate below.

Note that S_L holds throughout.

Here $\forall\ N \in \mathbb{N}$ by the previous lemma.

In this case $\beta \to \infty$ as claimed.

The value of $S_{\mathrm{eff}}[\rho]$ in what follows.

We then obtain $\gamma > \frac{1}{2}$ for the remainder of the section.

Recall that $\gamma = \frac{1}{2}$ and the result follows.

It follows that $\tilde{x}_a = -x_a$ is used in the estimate below.

Suppose $\tau \to 0$ holds throughout.

Note that $(\tau_{\mathrm{cr}} - \tau)$ by the previous lemma.

Here $N = 8$ as claimed.

In this case A_p in what follows.

The value of $[0, N-1]$ for the remainder of the section.

We then obtain $\rho^{\mathscr{D}}_{\mathrm{unif.}}$ and the result follows.

Recall that $\varepsilon > 0$ is used in the estimate below.

It follows that $\delta > 0$ holds throughout.

Suppose $\{P_j\}_{j \in \mathbb{N}}$ by the previous lemma.

Note that $|x| < 1$ as claimed.

Here $e^{itH_{\mathrm{XY}}}|\psi_0\rangle$ in what follows.

In this case h_{2n+1} for the remainder of the section.

The value of $T = 0$ and the result follows.

We then obtain (a) is used in the estimate below.

Recall that $G^{s_1}_{ij}, G^{s_2}_{ij}$ holds throughout.

It follows that (s_1, s_2) by the previous lemma.

Suppose Z_C as claimed.

Note that $\det(g_C^{s_1, s_2})$ in what follows.

Here $G^{s}_{ij}(\theta)$ for the remainder of the section.

In this case $\pi/3$ and the result follows.

The value of S_2 is used in the estimate below.

We then obtain N_{seed} holds throughout.

Recall that $\theta = 10$ by the previous lemma.

It follows that $\phi = 0.1$ as claimed.

Suppose $\ln(L)$ in what follows.

Note that $\Delta_\tau = 0.5$ for the remainder of the section.

Here $0.95(5)$ and the result follows.

In this case $\mathrm{f} + \mathrm{b}$ is used in the estimate below.

The value of $B^s(\tau_2, \tau_1)$ holds throughout.

We then obtain $\gamma_i = (1/(1 - e^{-2i\alpha\tilde{\sigma}_i}) - G^s_{i,i}(\tau))^{-1}$ by the previous lemma.

Recall that $\boldsymbol{v}_i$ as claimed.

It follows that $\mathcal{O}(N_C^2)$ in what follows.

Suppose $g_{C+i}^{s_1,s_2}$ for the remainder of the section.

Note that $\tau = \theta$ and the result follows.

Here $G^s(\theta)$ is used in the estimate below.

In this case $\lambda = 0.0006$ holds throughout.

The value of $\approx 10^{-6}$ by the previous lemma.

We then obtain $L_{\max} = 60$ as claimed.

Recall that $L = 6$ in what follows.

It follows that $L = 3, 6, 9, 12, 15, 18$ for the remainder of the section.

Suppose $N_\lambda/(N_\lambda + 1)$ and the result follows.

Note that $L = 3, 6, 9, ..., 60$ is used in the estimate below.

Here $\star$ holds throughout.

In this case $\mathcal{U}_h$ by the previous lemma.

The value of $T_1 G$ as claimed.

We then obtain $U_g U_h = U_{gh}$ in what follows.

Recall that $\Delta = H^2$ for the remainder of the section.

It follows that $\psi \in \mathcal{H}$ and the result follows.

Suppose $t \in (-\tau, \tau)$ is used in the estimate below.

Note that U, V holds throughout.

Here $s : G \to \tilde{G}$ by the previous lemma.

In this case $\mathcal{H}_j = \mathbb{C}^{2j+1}$ as claimed.

The value of $\mathrm{SO}(2) \cong \mathrm{U}(1)$ in what follows.

We then obtain $-\mathrm{i}A(\mathfrak{so}(2m)) \subset \mathfrak{u}(\mathcal{H})$ for the remainder of the section.

Recall that B_{jk} and the result follows.

It follows that $\mathbb{H}_m = \mathbb{R}^{2m+1} = \mathbb{R}^{2m} \times \mathbb{R}$ is used in the estimate below.

Suppose $X \in \mathfrak{sp}(2m, \mathbb{R})$ holds throughout.

Note that $n \in \mathbb{N}_0^m$ by the previous lemma.

Here $\mathcal{O}(1/L)$ as claimed.

In this case $\sigma_z = \mathrm{diag}(1, -1)$ in what follows.

The value of $\Delta^{(n)}$ for the remainder of the section.

We then obtain $\mathrm{O}(1, 3)$ and the result follows.

Recall that $p \in \mathcal{M}_m$ is used in the estimate below.

It follows that $M_{0i} = -M_{i0} = K_i$ holds throughout.

Suppose $d(g_n, 1) \to 0$ by the previous lemma.

Note that $\psi = \chi_W$ as claimed.

Here $[0, 1]$ in what follows.

In this case $PX_k = p_k X_k$ for the remainder of the section.

The value of $\dim(\mathcal{H}) \geq 3$ and the result follows.

We then obtain $\mathcal{P} : \omega(\rho, \psi, p) \mapsto \omega(\rho, 0, p)$ is used in the estimate below.

Recall that $\tilde{d}(\tilde{g}, 1_{\tilde{G}}) \geq d(g, 1_G)$ holds throughout.

It follows that E_{jk} by the previous lemma.

Suppose $t_g \sim 10^{-8} s$ as claimed.

Note that U_g in what follows.

Here $\mathrm{dW}_{k,s}$ for the remainder of the section.

In this case ϵ^2 and the result follows.

The value of $(-)$ is used in the estimate below.

We then obtain $2N^2 - 1$ holds throughout.

Recall that $\Lambda(\theta, \phi)$ by the previous lemma.

It follows that $s = t/t_g$ as claimed.

Suppose T_1 in what follows.

Note that $\gamma_z = t_g(2T_1 - T_2)/4T_1T_2$ for the remainder of the section.

Here $\epsilon = \sqrt{\lambda}$ and the result follows.

In this case (k) is used in the estimate below.

The value of $\Delta\mathcal{H}_\sigma^{\mathrm{ng}}$ holds throughout.

We then obtain ibm_oslo by the previous lemma.

Recall that σ_{kk} as claimed.

It follows that $\mathcal{H}^{\mathrm{ng}}$ in what follows.

Suppose $\theta = \pi$ for the remainder of the section.

Note that $\mathcal{N} = 15000$ and the result follows.

Here $\sim 10^{-3} - 10^{-2}$ is used in the estimate below.

In this case 0.25 holds throughout.

The value of $\sim 10^{-2}$ by the previous lemma.

We then obtain $\mathcal{H}_{\chi}^{\mathrm{ng}} = \mathcal{H}(\rho^{\mathrm{ng}}, \chi)$ as claimed.

Recall that $\Delta \mathcal{H}_{\chi}^{\mathrm{ng}} < \Delta \mathcal{H}_{\chi}^{\mathrm{ibm}}$ in what follows.

It follows that $n = 9$ for the remainder of the section.

Suppose $\mathrm{R}_{xy}(\phi) = \cos(\phi)\mathrm{X} + \sin(\phi)\mathrm{Y}$ and the result follows.

Note that $\mathrm{N}(\theta, \phi)$ is used in the estimate below.

Here $\mathrm{U}(\theta, \phi)$ holds throughout.

In this case $\leq N_s$ by the previous lemma.

The value of $50\% - 60\%$ as claimed.

We then obtain $\Delta V / V$ in what follows.

Recall that $\Delta H = H_f - H_i$ for the remainder of the section.

It follows that H_i and the result follows.

Suppose μ_0 is used in the estimate below.

Note that J_{ij} holds throughout.

Here ΔS_L by the previous lemma.

In this case $\mathbf{e}_i$ as claimed.

The value of $\mathbf{e}_j$ in what follows.

We then obtain R_{ij} for the remainder of the section.

Recall that R_{ij} / a and the result follows.

It follows that μ_B is used in the estimate below.

Suppose T_C holds throughout.

Note that $M(T) = (1 - \frac{T}{T_C})^{\beta}$ by the previous lemma.

Here $J(r_{ij})$ as claimed.

In this case r_{ij} in what follows.

The value of $\boldsymbol{s}_i$ for the remainder of the section.

We then obtain $\mathbf{p}_i$ and the result follows.

Recall that $\boldsymbol{r}_i$ is used in the estimate below.

It follows that ω_i holds throughout.

Suppose $n_{\text{qubits}} = \log_2 \mathcal{C}$ by the previous lemma.

Note that S_P as claimed.

Here $a_{\{n_{B_i}\}_k} \neq 0$ in what follows.

In this case $i, j \in 1 \dots N_o$ for the remainder of the section.

The value of $\sum_i \tilde{\lambda}_i = N_o - N_p$ and the result follows.

We then obtain $S_c \leq \max\{-\ln \nu, -\ln(1-\nu)\}$ is used in the estimate below.

Recall that $\alpha_g = 2$ holds throughout.

It follows that $N_o = 16$ by the previous lemma.

Suppose α_g as claimed.

Note that S_{opt} in what follows.

Here $U/t, V/t \approx 4$ for the remainder of the section.

In this case $k \in \{1, 2, \dots Q\}$ and the result follows.

The value of $\epsilon, \delta > 0$ is used in the estimate below.

We then obtain $n_{\text{bit}} = \log_2 2^{nH(P_k)}$ holds throughout.

Recall that $n_1 < n_2$ by the previous lemma.

It follows that $n > n'$ as claimed.

Suppose $\sigma = C^{\beta}$ in what follows.

Note that $\sigma(n)$ for the remainder of the section.

Here $\bar{n}_{B_i}$ and the result follows.

In this case $S_c^h \leq S_P$ is used in the estimate below.

The value of S_{CUE} holds throughout.

We then obtain $P_j^2 = U_{jk} U_{jl}^* U_{jm} U_{jn}^* a_k a_l^* a_m a_n^*$ by the previous lemma.

Recall that $D_{2,B} = S_{P_B} / \log(Q)$ as claimed.

It follows that $S_P = \alpha N_p S_c$ in what follows.

Suppose $\alpha \approx 0.8$ for the remainder of the section.

Note that $\langle x \rangle = 2(1-\nu) N_p$ and the result follows.

Here $k_1 + q$ is used in the estimate below.

In this case x_{fs} holds throughout.

The value of $N_p \gamma$ by the previous lemma.

We then obtain $p \to -p$ as claimed.

Recall that $1 + 1$ in what follows.

It follows that $F_{\mu\nu}(x,t)$ for the remainder of the section.

Suppose $\beta = \sqrt{4\pi}$ and the result follows.

Note that $E_\theta = \frac{e}{2\pi}\theta$ is used in the estimate below.

Here $m_q \equiv \mathcal{E}(p=0)$ holds throughout.

In this case $\theta = \pi - \varepsilon$ by the previous lemma.

The value of $\pi_1\pi_2$ as claimed.

We then obtain 0.5 in what follows.

Recall that π_1, π_2 for the remainder of the section.

It follows that π_1 and the result follows.

Suppose $\beta \neq \sqrt{4\pi}$ is used in the estimate below.

Note that $A_1(x) = 0$ holds throughout.

Here $a, b = 1, 2$ by the previous lemma.

In this case $\frac{e\beta}{2\pi}\phi = E + \frac{e\theta}{2\pi} \equiv E_T$ as claimed.

The value of $\{b(k), b^\dagger(k')\} = \{c(k), c^\dagger(k')\} = \delta(k-k')$ in what follows.

We then obtain $\hat{p}|p\rangle = p|p\rangle$ for the remainder of the section.

Recall that $p/m \ll 1$ and the result follows.

It follows that m_q is used in the estimate below.

Suppose $m_\pi/m_q > 2$ holds throughout.

Note that $I_C^i = C\ddot{\phi}_i$ by the previous lemma.

Here $\phi_J^i + \phi_C^i = \Phi_{\text{ext}}$ as claimed.

In this case $\omega_{k,j}$ in what follows.

The value of $\chi = 1$ for the remainder of the section.

We then obtain $\lambda(\Phi_{\text{ext}})$ and the result follows.

Recall that A_L is used in the estimate below.

It follows that $\varepsilon = 0$ holds throughout.

Suppose c_p by the previous lemma.

Note that $n \in [i+1, i+N]$ as claimed.

Here $D' = 100$ in what follows.

In this case $n = 2$ for the remainder of the section.

The value of $SU(2)$ and the result follows.

We then obtain $\mathbf{P}(\omega, \vec{v})$ is used in the estimate below.

Recall that $\hat{\mathbf{e}}_1$ holds throughout.

It follows that $(\sin 2\theta_V,\, 0,\, -\cos 2\theta_V)$ by the previous lemma.

Suppose $v^{\rho}(\vec{v}) = (1, \vec{v})$ as claimed.

Note that $\operatorname{diag}(+1, -1, -1, -1)$ in what follows.

Here $\omega > 0$ for the remainder of the section.

In this case μ^{-1} and the result follows.

The value of $\Omega_{i} > 0$ is used in the estimate below.

We then obtain v^{z} holds throughout.

Recall that $u = \omega$ by the previous lemma.

It follows that $\mathbf{Y} = -\mu \mathbf{J}^{z}$ as claimed.

Suppose $(0, 0, 1)$ in what follows.

Note that Ω_{i} for the remainder of the section.

Here $\hat{\mathbf{e}}_{3}^{\mathrm{II}} = \hat{\mathbf{e}}_{3}$ and the result follows.

In this case $\hat{\mathbf{e}}_{3}^{\mathrm{II}} = \cos\Theta \hat{\mathbf{e}}_{3}^{\mathrm{III}} - \sin\Theta \hat{\mathbf{e}}_{1}^{\mathrm{III}}$ is used in the estimate below.

The value of $\Theta = \pi$ holds throughout.

We then obtain $\partial_{t}\Theta = \Omega_{i} \sin\Theta$ by the previous lemma.

Recall that $\mathsf{O}^{\mathrm{II}} \ll 1$ as claimed.

It follows that $P_{3}^{\mathrm{III}} = \cos\Theta'$ in what follows.

Suppose $\lambda^{\rho} = (\mu,\, 0,\, 0,\, 0.5\mu)$ for the remainder of the section.

Note that $v^{x} = v^{y} = 0$ and the result follows.

Here $5\pi/6$, is used in the estimate below.

In this case $\times(1 + 0.5 v^{z})$ holds throughout.

The value of $\Delta\vec{v}$ by the previous lemma.

We then obtain $|\mathsf{Q}_{v^{x}, v^{y}}|$ as claimed.

Recall that v_{x} in what follows.

It follows that J_{3}^{z} for the remainder of the section.

Suppose $\mathbf{H}_{\mathrm{vac}}$ and the result follows.

Note that $\operatorname{Im}(\mathsf{C}\,\mathsf{D}^{*}) = 0$ is used in the estimate below.

Here σ_{8} holds throughout.

In this case ρ_{0} by the previous lemma.

The value of R_{200} as claimed.

We then obtain $c_{200} = R_{200} / r_{s}$ in what follows.

Recall that $X(\hat{\mathbf{n}})$ for the remainder of the section.

It follows that $\boldsymbol{\alpha}(\hat{\mathbf{n}}) = \boldsymbol{\nabla}_{\hat{\mathbf{n}}} \phi(\hat{\mathbf{n}})$ and the result follows.

Suppose $\Sigma_{\mathrm{cl}}(r)$ is used in the estimate below.

Note that $d_{A,\mathrm{cl}}$ holds throughout.

Here $x = r/r_s = \theta/\theta_s$ by the previous lemma.

In this case κ_t as claimed.

The value of $\boldsymbol{\ell} = (\ell_x, \ell_y)$ in what follows.

We then obtain $z = 0.7$ for the remainder of the section.

Recall that θ_s and the result follows.

It follows that $N_L^{(1)}$ is used in the estimate below.

Suppose $L = 5000$ holds throughout.

Note that $\ell_{\mathrm{max}}^{\mathrm{CMB}} = 4000$ by the previous lemma.

Here $L \sim 1100$ as claimed.

In this case 1000 in what follows.

The value of $100 \leq L \leq 5000$ for the remainder of the section.

We then obtain $M = 4, 7$ and the result follows.

Recall that $M_{200} = 4 \times 10^{14} M_{\odot}, z = 0.7$ is used in the estimate below.

It follows that $\hat{\kappa}_0$ holds throughout.

Suppose $\hat{\kappa}_L$ by the previous lemma.

Note that $\langle \kappa_0 \rangle$ as claimed.

Here 13% in what follows.

In this case x_{max} for the remainder of the section.

The value of $\chi(z) = \int_0^z dz' \frac{1}{H(z')} = (1+z) d_A(z)$ and the result follows.

We then obtain $\mathrm{Ci}(x)$ is used in the estimate below.

Recall that $s = 0, t = 1$ holds throughout.

It follows that $\langle g(\hat{n}) \rangle_{\text{fixed deflection}}$ by the previous lemma.

Suppose m_c as claimed.

Note that $\psi_C(x) = \gamma_2 \gamma_4 \, \bar{\psi}^T(x)$ in what follows.

Here $\bar{\Delta}_5$ for the remainder of the section.

In this case $\mu_u + 2\mu_d$ and the result follows.

The value of $r - g$ is used in the estimate below.

We then obtain P_{lep} holds throughout.

Recall that μ_{ν_e} by the previous lemma.

It follows that $m_c = 2.3$ as claimed.

Suppose $\eta_V = G_V / G_S$ in what follows.

Note that $\eta_V = 0.5$ for the remainder of the section.

Here μ_B^c and the result follows.

In this case $\mu_B = 850$ is used in the estimate below.

The value of $\eta_V = \eta_D = 0$ holds throughout.

We then obtain $\eta_V = 0.9$ by the previous lemma.

Recall that $n_B = \partial P / \partial \mu_B$ as claimed.

It follows that $n_f = \sum_c n_{fc}$ in what follows.

Suppose $g_{\omega B}(n)$ for the remainder of the section.

Note that b_τ and the result follows.

Here I_{3B} is used in the estimate below.

In this case E_B^* holds throughout.

The value of $B = N, P$ by the previous lemma.

We then obtain m_e as claimed.

Recall that $P(r = R) = 0$ in what follows.

It follows that $\mu_{\nu_e}/T = 2$ for the remainder of the section.

Suppose $\mu_{\nu_e} = \alpha T$ and the result follows.

Note that $2.3 M_\odot$ is used in the estimate below.

Here $T = 30$ holds throughout.

In this case $\bar{\Delta}_0$ by the previous lemma.

The value of n_e as claimed.

We then obtain $q \in Q$ in what follows.

Recall that $r \in R$ for the remainder of the section.

It follows that $\mathrm{NN}_2(1; R) = \{0, 1, 2\} = \mathrm{NN}_3(1; R)$ and the result follows.

Suppose $t \in \mathbb{R}_+$ is used in the estimate below.

Note that $c(R) = n + 1$ holds throughout.

Here $R \subset \mathbb{R}^n$ by the previous lemma.

In this case $O\big(c_m(R)^{O(1)} \cdot |R| \log(\Delta(R))\big)$ as claimed.

The value of $\mathcal{S}_i(p, \mathcal{T}(R))$ in what follows.

We then obtain $p \in \mathcal{T}(R)$ for the remainder of the section.

Recall that $t > \delta$ and the result follows.

It follows that $r \in \mathbb{R}_+$ is used in the estimate below.

Suppose $H(\mathcal{T}(R)) = \{i \mid C_{i-1} \neq C_i\} \cup \{l_{\max}, l_{\min}\}$ holds throughout.

Note that $l(a_{m-1}) \leq i - 1$ by the previous lemma.

Here $j = l_{\min}(\mathcal{T}(R)) - 1$ as claimed.

In this case $W_1 = \{r\}$ in what follows.

The value of $i \leftarrow -\infty$ for the remainder of the section.

We then obtain $\eta(i) \leftarrow i$ and the result follows.

Recall that $q \in X \setminus W$ is used in the estimate below.

It follows that $i \in S$ holds throughout.

Suppose $O(c(R)^2 \log(|R|))$ by the previous lemma.

Note that $\varrho \in L(\mathcal{T}(R), q)$ as claimed.

Here $R_{l_{\min}}$ in what follows.

In this case $\psi_k(t)$ for the remainder of the section.

The value of q_2 and the result follows.

We then obtain p_{m-2} is used in the estimate below.

Recall that $m - 2$ holds throughout.

It follows that $p_{i \cdot m - 1}$ by the previous lemma.

Suppose $i = m, j = m - 1$ as claimed.

Note that $i = 3, j = 2$ in what follows.

Here q_2 : for the remainder of the section.

In this case $\langle N, S, A, T, \{R_i\}, \Omega, \{O_i\} \rangle$ and the result follows.

The value of $s^t = \langle s^{1,t}, ..., s^{k,t} \rangle$ is used in the estimate below.

We then obtain $T(s^{t+1}|s^t, a^t)$ holds throughout.

Recall that $O_i(o_i^t|s^t)$ by the previous lemma.

It follows that $\pi_i(a_i^t|h_i^t)$ as claimed.

Suppose $X_i = \{X_i^1, ..., X_i^m\} \subseteq S$ in what follows.

Note that $O_i(o_i^t|s^t) = \dot{O}_i(o_i^t|x_i^t)$ for the remainder of the section.

Here $\langle s^t, h_{-i}^t \rangle$ and the result follows.

In this case x_i is used in the estimate below.

The value of $x_i \in X_i = \{X_i^1, X_i^2\}$ holds throughout.

We then obtain Y_i by the previous lemma.

Recall that a_j^t as claimed.

It follows that $x_i \in \times_{j \in |X_i|} X_i^j$ in what follows.

Suppose $I_i(u_i^t|l_i^t)$ for the remainder of the section.

Note that y_t and the result follows.

Here θ_i is used in the estimate below.

In this case $I_i(u_i^t|l_i^t) \in \Psi_i$ holds throughout.

The value of $\Psi = \times_{i \in N} \Psi_i$ by the previous lemma.

We then obtain a_{-i} as claimed.

Recall that M_i^1 in what follows.

It follows that $Q_{M_i^2}^{\pi_i}$ for the remainder of the section.

Suppose $\bar{R} = ||R||_\infty$ and the result follows.

Note that $\{I_{\theta_i}\}_{i \in N}$ is used in the estimate below.

Here 5×5 holds throughout.

In this case $F = 4$ by the previous lemma.

The value of $F = 1$ as claimed.

We then obtain $P^2(u_i^{t-1} y_i^{t-1}, h_{-i}^{t-1} | l^t)$ in what follows.

Recall that $P^2(h_i^{t+1} | h_i^t, a_i^t)$ for the remainder of the section.

It follows that $\hat{a}_i^t \neq \bar{a}_i^t$ and the result follows.

Suppose $\hat{I}_{\theta_m}$ is used in the estimate below.

Note that $S(\omega)$ holds throughout.

Here $b, 3$ by the previous lemma.

In this case $\hat{Y} : \Omega \to \{0, 1\}$ as claimed.

The value of $.Soeither$ in what follows.

We then obtain $isdependenton$ for the remainder of the section.

Recall that $overhardlabelstoinferthevalueof$ and the result follows.

It follows that $.Itisusedtoevaluatehowwell$ is used in the estimate below.

Suppose $.Wedefinerandomvariables$ holds throughout.

Note that $isarandomvariable$: by the previous lemma.

Here $), whether$ as claimed.

In this case $.Forthisattacktowork$, in what follows.

The value of $isincludedintheinputdatarecordsand$ for the remainder of the section.

We then obtain $.Itmakesiteasiertoextract$ and the result follows.

Recall that $, classifierof$ is used in the estimate below.

It follows that $issplitintotwodisjointdatasets$: holds throughout.

Suppose $.Theseareusedtocreatethestatistic$ by the previous lemma.

Note that $using$ as claimed.

Here $)curveasafunctionwhichmapseach$ in what follows.

In this case $shouldnotbeabletoinfer$ for the remainder of the section.

The value of $then for all attack model$ and the result follows.

We then obtain $.Let$ is used in the estimate below.

Recall that $for training$ holds throughout.

It follows that $and evaluate iton 634 unseen data records in$ by the previous lemma.

Suppose $and evaluate iton 1,200 unseen data records in$ as claimed.

Note that $and target model's$ in what follows.

Here $using hard labels against a$ for the remainder of the section.

In this case $issuch that$ and the result follows.

The value of $.We then say that$ is used in the estimate below.

We then obtain $we call$ holds throughout.

Recall that the by the previous lemma.

It follows that $e_k \doteq (\mathbf{x}_k, t_k, p_k)$ as claimed.

Suppose $\mathbf{x}_k, t_k$ in what follows.

Note that $\mathbf{x}'_k = \mathbf{W}(\mathbf{x}_k, t_k; \boldsymbol{\theta})$ for the remainder of the section.

Here e'_k and the result follows.

In this case $\epsilon = 1$ is used in the estimate below.

The value of $\mathcal{D}(\mathcal{E}, \boldsymbol{\theta})$ holds throughout.

We then obtain $\mathbf{f} \doteq \partial \mathbf{W}(\mathbf{x}, t; \boldsymbol{\theta})/\partial t$ by the previous lemma.

Recall that $t_{\text{ref}} = t_1$ as claimed.

It follows that $t \in [t_1, t_{N_e}]$ in what follows.

Suppose $\Delta t = 0$ for the remainder of the section.

Note that $\mathcal{R} = 0$ and the result follows.

Here $\mathcal{R} < 0$ is used in the estimate below.

In this case $\boldsymbol{\theta} \equiv (v_x, v_y)^\top$ holds throughout.

The value of $\mathcal{R} \equiv \int_0^1 \frac{d|\mathrm{J}|}{d\Delta t} = 0$ by the previous lemma.

We then obtain $t \in [0, 1]$ as claimed.

Recall that $\mathtt{R}(\boldsymbol{\omega} \Delta t)$ in what follows.

It follows that $\mathcal{R}_{\mathbf{x}(0)}$ for the remainder of the section.

Suppose $\boldsymbol{\omega} = (0, 0, \omega_z)^\top$ and the result follows.

Note that 346×260 is used in the estimate below.

Here $N = \{3, 10, 20\}$ holds throughout.

In this case $\lambda = 1.0$ by the previous lemma.

The value of $\mathbf{v}(\mathbf{x})$ as claimed.

We then obtain 2×6 in what follows.

Recall that $\mathbf{V} \equiv (0, 0, v_z)^\top$ for the remainder of the section.

It follows that $\mathbf{x}'_k = (1 - v_z/Z(\mathbf{x}))t_k$. and the result follows.

Suppose $\mathbf{x}^h(t) = (x(t), y(t), 1)^\top$ is used in the estimate below.

Note that 3×3 holds throughout.

Here $\mathbf{e}_3^\top \mathtt{H}_{t,t+\Delta t} = (0, 0, 1)$ by the previous lemma.

In this case $\omega_z = 0, \mathbf{v} = \mathbf{0}$ as claimed.

The value of $|V|$ in what follows.

We then obtain $V = \{have, a, great, day\}$ for the remainder of the section.

Recall that $\{1, 0, 0, 0\}$ and the result follows.

It follows that V, V^T is used in the estimate below.

Suppose V^T holds throughout.

Note that $n - 1$ by the previous lemma.

Here w_i as claimed.

In this case w_{i+j} in what follows.

The value of $P(w_i)$ for the remainder of the section.

We then obtain $f(w_i)$ and the result follows.

Recall that b_p is used in the estimate below.

It follows that X_{pq} holds throughout.

Suppose w_q by the previous lemma.

Note that x_{max} as claimed.

Here 0.75 in what follows.

In this case $f(0) = 0$ for the remainder of the section.

The value of $f(0)$ and the result follows.

We then obtain v_w is used in the estimate below.

Recall that $n = 3$ holds throughout.

It follows that $\mathcal{M} \subset \{1, ..., M\}$ by the previous lemma.

Suppose $(t_1, ..., t_{i-1})$ as claimed.

Note that t_{i+1} in what follows.

Here $(t_{i+1}, t_{i+2}, ..., t_N)$ for the remainder of the section.

In this case Θ_x and the result follows.

The value of Θ_s is used in the estimate below.

We then obtain $2L + 1$ holds throughout.

Recall that $h_{i,j}^{L} M$ by the previous lemma.

It follows that $x_i^{LM} = h_{i,0}^{LM}$ as claimed.

Suppose $\left(SentenceA, SentenceB \right)$ in what follows.

Note that $[SEP]$ for the remainder of the section.

Here $(X_t)_{t \geq 0}$ and the result follows.

In this case $\Delta_n := t_{i+1} - t_i$ is used in the estimate below.

The value of X_{Δ_n} holds throughout.

We then obtain $(b_l)_{0 \leq l \leq 3}$ by the previous lemma.

Recall that $(\tilde{C}, \tilde{\rho})$ as claimed.

It follows that λ_0 in what follows.

Suppose $\pi_{|t-s|}$ for the remainder of the section.

Note that $n \geq 1$ and the result follows.

Here $d = 1$ is used in the estimate below.

In this case $P_t(\psi_\xi)(x) = \int_{\mathbb{R}} p_t(x,y) \psi_\xi(y) dy$ holds throughout.

The value of $\left\| \psi \right\|_\infty \leq 1$ by the previous lemma.

We then obtain $|b(0)| = 0 \leq b_0$ as claimed.

Recall that $M_n h_n^3 \geq 1/\alpha_0$ in what follows.

It follows that $k \geq 1$ for the remainder of the section.

Suppose $(x, y, \epsilon) \mapsto p_{\Delta_n}^{\epsilon}(x,y)$ and the result follows.

Note that $h_n = n^{-1/(1+2\beta)}$ is used in the estimate below.

Here $(B_n')_n$ holds throughout.

In this case $b \equiv 0$ by the previous lemma.

The value of $\dot{X}^\epsilon$ as claimed.

We then obtain $y > 2h_n$ in what follows.

Recall that $r' > 0$ for the remainder of the section.

It follows that $|\partial_x A_\epsilon(x,v)| + |\partial_x B(x,v)| \leq c(1 + |v|)$ and the result follows.

Suppose $L(\int_0^s u_r dB_r) = \int_0^s L(u_r) dB_r - \int_0^s u_r dB_r$ is used in the estimate below.

Note that $\left\| n_{h_n} \right\|_\infty \leq \frac{c}{M_n}$ holds throughout.

Here $\limsup_n \frac{1}{h_n^2 M_n} < \infty$ by the previous lemma.

In this case ℓ_{x,y,Δ_n,h_n} as claimed.

The value of $(M_t)_t$ in what follows.

We then obtain $2\sqrt{s'}\sqrt{\tau \vee h^2} \leq s' + \tau \vee h^2$ for the remainder of the section.

Recall that $\varphi : \mathbb{R} \to \mathbb{R}$ and the result follows.

It follows that $F \in \mathbb{D}^{1,2}(\mathbb{R})$ is used in the estimate below.

Suppose $n > 1$ holds throughout.

Note that $n^{-}(\mathcal{I})$ by the previous lemma.

Here $r \leq R$ as claimed.

In this case $x_1, \ldots, x_n$ in what follows.

The value of $f(x_1, \ldots, x_d) \leq f((x_1', \ldots, x_d'))$ for the remainder of the section.

We then obtain $\{H_i^{-} : p_i \leq 1/2\} \cup \{H_i^{+} : p_i > 1/2\}$. and the result follows.

Recall that $\ell_{\alpha}^{-}(\mathcal{I}) \geq |\mathcal{D}_{\alpha}^{-}|$ is used in the estimate below.

It follows that $J_v(\mathcal{I}) : n^{+}(\mathcal{I}) = v$ holds throughout.

Suppose $|\mathcal{S}^{-} \cap \mathcal{I}^c| \times |\mathcal{S}^{+} \cap \mathcal{I}^c|$ by the previous lemma.

Note that $\mathcal{S} = (1, 1)$ as claimed.

Here $1/2 < p_1 < 1 - 2\alpha/3$ in what follows.

In this case $\operatorname{SE}(\hat{\pi}_i^j)$ for the remainder of the section.

The value of $0 \leq n^{+}$ and the result follows.

We then obtain $H_2^{+} \cap H_3^{+}$ is used in the estimate below.

Recall that $n^{+} = n^{-} = 15$ holds throughout.

It follows that $O(|\mathcal{I}|^2 \cdot \max(1, |\mathcal{I}^c|^2))$ by the previous lemma.

Suppose $(p_1, p_2) = (0.19, 0.19)$ as claimed.

Note that $c_{\alpha}(\mathcal{K})$ in what follows.

Here $K_1 \cap K_2 \cap K_3$ for the remainder of the section.

In this case $\tilde{\ell}_{\alpha}^{+}$ and the result follows.

The value of $\mathcal{S}^{-} = \{i \in [n] : p_i \leq 1/2\}$ is used in the estimate below.

We then obtain $f_v(\mathcal{I}) = \max\{f_{v,u}(\mathcal{I}), u = 0, \ldots, |\mathcal{I}^c|\}$ holds throughout.

Recall that $i \in \mathcal{S}^{-} \cap \mathcal{I}$ by the previous lemma.

It follows that $p^{r/n}$ as claimed.

Suppose $\left\{p^{r/n} : r \in [n]\right\}$ in what follows.

Note that $q^{n-k+1/n} = f(\{2q_{(n-k+1)}, \ldots, 2q_{(n-s)}\})$ for the remainder of the section.

Here $p^{0/n} \equiv 0$ and the result follows.

In this case $K^{2/2}$ is used in the estimate below.

The value of 0.0271 holds throughout.

We then obtain (0.0403) by the previous lemma.

Recall that ϵ_{ℓ} as claimed.

It follows that ϕ_ℓ in what follows.

Suppose $f_\ell(h_\ell; \epsilon_\ell; \phi_\ell)$ for the remainder of the section.

Note that C_L, C_L' and the result follows.

Here $\mathbf{X} \in \mathbb{R}^{N \times D}$ is used in the estimate below.

In this case $\mathbf{y} \in \mathbb{R}^N$ holds throughout.

The value of $w \in \mathbb{R}^D$ by the previous lemma.

We then obtain $\sigma(x) = 1/(1 + e^{-x})$ as claimed.

Recall that μ_{MAP} in what follows.

It follows that $\Lambda = \beta \sum_i x_i x_i^T + \alpha I$ for the remainder of the section.

Suppose $\sigma(w^T x_i)(1 - \sigma(w^T x_i))$ and the result follows.

Note that $\sigma(w^T x_i)$ is used in the estimate below.