The value of $\mathbb{A}^2/\mathbb{Q}$, is used in the estimate below.

We then obtain $\mathscr{S}_P$. holds throughout.

Recall that (m, l); by the previous lemma.

It follows that $k^2 < p^n$: as claimed.

Suppose $D_X := D \times_B X$, in what follows.

Note that $B \to \mathbb{P}^1_B$. for the remainder of the section.

Here (b, f); and the result follows.

In this case $f_b(0) = (A + C, C)$: is used in the estimate below.

The value of $\widetilde{s}(b, A, B, C, D) \in W^{0,v}_{d+u} \times W^{0,v}_u$, holds throughout.

We then obtain $W^{0,v}_d$. by the previous lemma.

Recall that $\mathscr{Z}_{n,d}$; as claimed.

It follows that $X_{b\mathbb{C}}$: in what follows.

Suppose $nz = \sum_i n_i P_i$, for the remainder of the section.

Note that $\mathscr{P}_b$. and the result follows.

Here m, ℓ; is used in the estimate below.

In this case $\widetilde{S}_{\mathbb{Z}}$: holds throughout.

The value of $\widetilde{S_{\mathbb{Z},b}}$, by the previous lemma.

We then obtain $U_{v,d}(\bar{\widetilde{S}}_{\mathbb{Z}})$. as claimed.

Recall that $V_{\mathbb{C}}$; in what follows.

It follows that $K_{k,p,n} = \mathbb{Q}(\sqrt{-d})$: for the remainder of the section.

Suppose $k = 1$, and the result follows.

Note that $F_0 = 0$. is used in the estimate below.

Here D_2; holds throughout.

In this case $(1, d, k^2, p) \in \mathcal{S}$: by the previous lemma.

The value of $\varepsilon = -1$, as claimed.

We then obtain $r \geq 1$. in what follows.

Recall that $\alpha = k + m\sqrt{-d}$; for the remainder of the section.

It follows that $a = \pm t$: and the result follows.

Suppose $q = 3$, is used in the estimate below.

Note that $d > 3$. holds throughout.

Here $H\mathbb{F}_p$; by the previous lemma.

In this case $X = \Omega^2 S^3$: as claimed.

The value of $\xi = 0$, in what follows.

We then obtain $\otimes_A^{\mathbb{L}}$. for the remainder of the section.

Recall that $f : A_0 \to B_0$; and the result follows.

It follows that $H\mathbb{Z}_p \simeq H\pi_0(\mathbb{S}_p^\wedge)$: is used in the estimate below.

Suppose $Mf_{k,pu}$, holds throughout.

Note that $a^\sharp := \lim_{n\to\infty} y_n^{p^n}$. by the previous lemma.

Here $W(\kappa)$; as claimed.

In this case $R^\flat \to D/p$: in what follows.

The value of $(\varphi_{D/p}^n(b_n))_{n=0}^\infty$, for the remainder of the section.

We then obtain $\eta_D : D \to HR$. and the result follows.

Recall that $X_n := (\tau_{\geq n}\mathbb{S}_p^\wedge) \otimes_{\mathbb{S}_p^\wedge} W^+(k)$; is used in the estimate below.

It follows that $t_{R,\xi}$: holds throughout.

Suppose $\Omega S^3 \to * \to S^3$, by the previous lemma.

Note that $\tau_{\leq 0} Mf_{\kappa,pu} \simeq H\kappa$. as claimed.

Here $\xi R^\flat$; in what follows.

In this case $e = v(p)$: for the remainder of the section.

The value of $W^+(k)[[u]]$, and the result follows.

We then obtain $W(k)[[u]]$. is used in the estimate below.

Recall that $HA \otimes_{W^+(k)[[u]]} HA$; holds throughout.

It follows that t_{k,a_0}: by the previous lemma.

Suppose $\deg u_1 = \cdots = \deg u_n = 0$, as claimed.

Note that $k = A/\mathfrak{m}$. in what follows.

Here $R := A/I$; for the remainder of the section.

In this case $H(A/I)$: and the result follows.

The value of $n < 0$, is used in the estimate below.

We then obtain $\pi_0(R)$. holds throughout.

Recall that $M_{(x)}^\wedge$; by the previous lemma.

It follows that $I \subseteq R$: as claimed.

Suppose $O(n^{\omega+\epsilon})$, in what follows.

Note that $\omega < 2.3725$. for the remainder of the section.

Here $i \in [k_X], j \in [k_Y], k \in [k_Z]$; and the result follows.

In this case $\Omega(n^2 \log n)$: is used in the estimate below.

The value of $x_i y_j z_k$, holds throughout.

We then obtain $a_{ijk} = b_{\pi_x(i)\pi_y(j)\pi_z(k)}$. by the previous lemma.

Recall that i, j, k; as claimed.

It follows that $b \times c$: in what follows.

Suppose $q \geq 2$, for the remainder of the section.

Note that $Y' \subseteq Y$. and the result follows.

Here $\{x_0, \ldots, x_{q+1}\}$; is used in the estimate below.

In this case $a = b = c$: holds throughout.

The value of $I \subseteq [n]$, by the previous lemma.

We then obtain $\bigoplus_{i=1}^{n} T_{iii}$. as claimed.

Recall that $i, j, k, i', j', k' \in [n]$; in what follows.

It follows that $I \subseteq [N]$: for the remainder of the section.

Suppose $N/2 = n$, and the result follows.

Note that $1/n$. is used in the estimate below.

Here $B_Y = Y_J \times Z_{K'} \times X_{I''}$; holds throughout.

In this case N_T: by the previous lemma.

The value of $h_X(I', J', K')$, as claimed.

We then obtain C_3. in what follows.

Recall that $g \geq 2$; for the remainder of the section.

It follows that $\alpha_{Z_k} = \gamma_{Z_k}$: and the result follows.

Suppose $X^2 = \{x_{q+1}\}$, is used in the estimate below.

Note that T_{200}. holds throughout.

Here $L_{t,I} := \{A \in \{0, 1, \ldots, q+1\}^t \mid \sum_{\ell=1}^{t} \kappa(A_\ell) = I\}$; by the previous lemma.

In this case $I' \in \{0, 1, \ldots, I\}$: as claimed.

The value of $x_{q+1,q+1}$, in what follows.

We then obtain $(i, j, k) \in S'$. for the remainder of the section.

Recall that H_1; and the result follows.

It follows that $A : H_1 \to H_3$: is used in the estimate below.

Suppose x^*, holds throughout.

Note that A^{T}. by the previous lemma.

Here $(\varepsilon, \min\{\frac{1}{\|A\|^2}, \frac{1}{\|B\|^2}\} - \varepsilon)$; as claimed.

In this case $C = \{x \in H_1 : \mathfrak{f}(x) \leq 0\}$: in what follows.

The value of $P_C x = \operatorname{argmin}_{y \in C} \|x - y\|, \ \forall x \in H$, for the remainder of the section.

We then obtain $\{x_n\}$. and the result follows.

Recall that $\gamma > 0$; is used in the estimate below.

It follows that $x \in C$: holds throughout.

Suppose $\{\zeta_n\}$, by the previous lemma.

Note that $\lim_{n\to\infty} \zeta_n = 0.$ as claimed.

Here $\{v_n\} \subset H_2$; in what follows.

In this case $\lim_{n\to\infty} \delta_n = 0$: for the remainder of the section.

The value of $\{v_n\}$, and the result follows.

We then obtain $\Phi_n.$ is used in the estimate below.

Recall that $\{y_{n_k}\}$; holds throughout.

It follows that $\bar{y} \in Q$: by the previous lemma.

Suppose $v_n \equiv v$, as claimed.

Note that $\Psi_n.$ in what follows.

Here $g : H_2 \to H_2$; for the remainder of the section.

In this case $(x^*, y^*) = P_\Omega(f(x^*), g(y^*)) \in \Omega$: and the result follows.

The value of $\lambda_1 \in (0, \frac{\sqrt{2}}{2})$, is used in the estimate below.

We then obtain $Q = \{(y_1, y_2, y_3) \in H_2,\ y_1^2 - y_2 + 5 \leq 0\}.$ holds throughout.

Recall that x_0; by the previous lemma.

It follows that $\delta_n = \frac{1}{n+1}$: as claimed.

Suppose (d), in what follows.

Note that $\lambda_1.$ for the remainder of the section.

Here $g(y) = 0.6y$; and the result follows.

In this case 10^{-4}: is used in the estimate below.

The value of $V(G) = \{v_1, ..., v_n\}$, holds throughout.

We then obtain $N(G).$ by the previous lemma.

Recall that $u \in V(G)$; as claimed.

It follows that $u \in C$: in what follows.

Suppose $\mathbf{x}_W = (x_1, ..., x_n)^t$, for the remainder of the section.

Note that $v_i \in W.$ and the result follows.

Here $\mathbf{x}_{N[v_i]}$; is used in the estimate below.

In this case $N(G)\mathbf{x}_S = \mathbf{0}$: holds throughout.

The value of $Col(N)$, by the previous lemma.

We then obtain $\rho(G) := dim(Col(N(G)).$ as claimed.

Recall that $\rho(G)$; in what follows.

It follows that $\nu(G) = 0$: for the remainder of the section.

Suppose $pr(a) = 0$, and the result follows.

Note that $pr(A)$. is used in the estimate below.

Here $pr(S) = pr(\mathbf{x}_S)$; holds throughout.

In this case S_1: by the previous lemma.

The value of $nd(v) := \nu(G - v) - \nu(G)$, as claimed.

We then obtain $nd(v) = -1$. in what follows.

Recall that $N\mathbf{x}_S = \mathbf{1}$; for the remainder of the section.

It follows that $pr(S) = pr(\rho(G - v))$: and the result follows.

Suppose $\rho(T)$, is used in the estimate below.

Note that $pr(A \cap B) = \mathbf{x}_A^t \mathbf{x}_B$. holds throughout.

Here A^c; by the previous lemma.

In this case $N\mathbf{1} = \mathbf{x}_{D^c}$: as claimed.

The value of $G_1 \oplus ... \oplus G_m$, in what follows.

We then obtain $V(G_1 \oplus ... \oplus G_m) = \cup_{i=1}^m V(G_i)$. for the remainder of the section.

Recall that G_1; and the result follows.

It follows that $\nu(G_1 \oplus G_2) = \nu(G_1) + \nu(G_2) + 1$: is used in the estimate below.

Suppose $\{G_1, ..., G_m\}$, holds throughout.

Note that $j \neq 0$. by the previous lemma.

Here $G = (N, A; \mathbf{w})$; as claimed.

In this case $N_1, \ldots, N_p \subseteq N$: in what follows.

The value of $i \in N$, for the remainder of the section.

We then obtain N_j. and the result follows.

Recall that $(i, j) \in W$; is used in the estimate below.

It follows that $d(\xi)$: holds throughout.

Suppose $\omega \in \Omega(T)$, by the previous lemma.

Note that $\Omega \equiv \Omega(T)$. as claimed.

Here $p^{t\omega}$; in what follows.

In this case $i \in N_k$: for the remainder of the section.

The value of S_i^t, and the result follows.

We then obtain D_i^t. is used in the estimate below.

Recall that $S_i^t + R_i^t - D_i^{(t-1)} - d_i^{t\omega} < 0$; holds throughout.

It follows that $\overline{NCD}$: by the previous lemma.

Suppose $k = i$, as claimed.

Note that Φ_1. in what follows.

Here $\Phi_1, \dots, \Phi_4$; for the remainder of the section.

In this case (s, x, y): and the result follows.

The value of $1 =: t_0 < t_1 < \cdots < t_K := q$, is used in the estimate below.

We then obtain T_{k+1}. holds throughout.

Recall that T_2^+; by the previous lemma.

It follows that $T_K^+ = T_K$: as claimed.

Suppose $i_0 \in N_l, (i, j) \in W, l \in P$, in what follows.

Note that $(\bar{s}, \bar{x}, \bar{y})$. for the remainder of the section.

Here N_1; and the result follows.

In this case r_j^-, r_j^+: is used in the estimate below.

The value of $\tilde{r}_i^-$, holds throughout.

We then obtain $\tilde{d}_i^t = (1 + \tilde{r}_i^+) d_i^t, \quad i \in N_j, j \in P, t \in T.$. by the previous lemma.

Recall that 250; as claimed.

It follows that $t = 33$: in what follows.

Suppose P_θ, for the remainder of the section.

Note that $\operatorname{Fav}(E(r))$. and the result follows.

Here $f_\sharp \mu$; is used in the estimate below.

In this case $\mathbb{R}^n \setminus \{a\}$: holds throughout.

The value of $|\mathcal{K}_n + \Gamma| \sim \mathcal{P}_\Gamma(\mathcal{K}_n)$, by the previous lemma.

We then obtain $\pi_\theta(x)$. as claimed.

Recall that $c > 0$; in what follows.

It follows that $a \in A$: for the remainder of the section.

Suppose $\gamma \in \Gamma$, and the result follows.

Note that $\Omega = \mathbb{R}^n$. is used in the estimate below.

Here $P_a : E \to S^{n-1}$; holds throughout.

In this case $a, x, y \in B(x, R)$: by the previous lemma.

The value of Γ_i, as claimed.

We then obtain s, t. in what follows.

Recall that $m = 1$; for the remainder of the section.

It follows that $|b_1 - a_1|$: and the result follows.

Suppose $b_1 - a_1 < d_\lambda$, is used in the estimate below.

Note that $d_\lambda = \lambda - b_1 - t_0 = \lambda - a_1 - s_0 \geq 0$. holds throughout.

Here $a_1 \neq b_1$; by the previous lemma.

In this case $|b_2 - a_2|$: as claimed.

The value of $x = 0$, in what follows.

We then obtain $\mathbf{x} = (x_1, x_2)$. for the remainder of the section.

Recall that $(\Gamma + \mathbf{b})$; and the result follows.

It follows that $s = m$: is used in the estimate below.

Suppose $\widetilde{\pi_{\alpha\sharp}\mu} \ll h$, holds throughout.

Note that $r \in (0, 1)$. by the previous lemma.

Here $\psi = \mathcal{H}^{n-1}$; as claimed.

In this case $\nu(\mathcal{K}_n) = 1$: in what follows.

The value of $(0, 1/4)$, for the remainder of the section.

We then obtain $n - 2$. and the result follows.

Recall that A/I; is used in the estimate below.

It follows that (A, J): holds throughout.

Suppose $\Gamma_{\mathbb{Z}[X]}(Y)$, by the previous lemma.

Note that $f(C) \in \mathcal{D}'$. as claimed.

Here $(K_0 \to K)^*$; in what follows.

In this case $A = \mathbb{Z}[x_1, \ldots, x_n]$: for the remainder of the section.

The value of $\mathcal{D} \hookrightarrow \mathcal{C}$, and the result follows.

We then obtain $\mathcal{C}' = \mathcal{C}$. is used in the estimate below.

Recall that $(A, I), (B, J)$; holds throughout.

It follows that $x \in I$: by the previous lemma.

Suppose $A \to A''$, as claimed.

Note that $M_{(A,I)} \to N_{(A,I)}$. in what follows.

Here $N \otimes^{\mathbb{L}}_A M' \to N \otimes^{\mathbb{L}}_A M \to N \otimes^{\mathbb{L}}_A M''$; for the remainder of the section.

In this case $\Gamma^*_{A/I}(I/I^2) \to J^{[*]}/J^{[*+1]}$: and the result follows.

The value of $\Gamma^*_{A/I}(I/I^2) \to \bigoplus J^{[*]}/J^{[*+1]}$, is used in the estimate below.

We then obtain $A \to S$. holds throughout.

Recall that $(B, J, \delta) = (A[X], IA[X], \gamma)$; by the previous lemma.

It follows that $B/J \cong A''/I$: as claimed.

Suppose $\Delta^{1} \times \Delta^{n}$, in what follows.

Note that $[a, b]$. for the remainder of the section.

Here $A \to C$; and the result follows.

In this case $A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{Q} \to D \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{Q}$: is used in the estimate below.

The value of $R/p \to R/p$, holds throughout.

We then obtain $(\mathbb{F}_{p}\{x\}, (x))$. by the previous lemma.

Recall that $M\{n\}$; as claimed.

It follows that $Q_{n} \in A[X_{0}, \ldots, X_{n-1}]$: in what follows.

Suppose $\prod_{(r,y) \in E} \delta^{r}(y/d)$, for the remainder of the section.

Note that $(A\{y\}/^{\mathbb{L}}(d, y))[p^{-1}]$. and the result follows.

Here $\deg(\delta^{j}(z_{2})) = p^{j}$; is used in the estimate below.

In this case (C, d): holds throughout.

The value of $S(R)$, by the previous lemma.

We then obtain $\partial \gamma_{R} = \partial R$. as claimed.

Recall that $\mathbb{R}^{2^{n}-1}$; in what follows.

It follows that $n' \gg n$: for the remainder of the section.

Suppose γ_{AB}, and the result follows.

Note that I_{B}'. is used in the estimate below.

Here $\partial I_{B}'$; holds throughout.

In this case 2^{L}: by the previous lemma.

The value of I_{l}, as claimed.

We then obtain $x_{l} = 1$. in what follows.

Recall that $x_{A} = 1$; for the remainder of the section.

It follows that $R_{J_{r}'} \neq R_{J_{r}}$: and the result follows.

Suppose $2 \leftrightarrow B$, is used in the estimate below.

Note that $x^{A} = 110$. holds throughout.

Here J_{r}; by the previous lemma.

In this case σ_{ABC}: as claimed.

The value of $f(111)_{4} = 0$, in what follows.

We then obtain $\gamma_{I_{l}}$. for the remainder of the section.

Recall that $w \in \mathbb{R}^{m}$; and the result follows.

It follows that $J_{1} = A$: is used in the estimate below.

Suppose y_{r}, holds throughout.

Note that $\tilde{f}(x) = f(x)$. by the previous lemma.

Here A, BC, G; as claimed.

In this case ABG, C, D: in what follows.

The value of $n_k \geq k$, for the remainder of the section.

We then obtain $b^T x \geq 0$. and the result follows.

Recall that $\sum_{i=1}^{m} q_i$; is used in the estimate below.

It follows that $I \ni n + 1$: holds throughout.

Suppose $\S A + \S AB \geq \S B$, by the previous lemma.

Note that $I \cup J$. as claimed.

Here $n = 5$; in what follows.

In this case $I \smallsetminus J$: for the remainder of the section.

The value of $E \subset V \times V$, and the result follows.

We then obtain $\partial V \subseteq V$. is used in the estimate below.

Recall that $C(W) \subseteq E$; holds throughout.

It follows that $[n]$: by the previous lemma.

Suppose Sym_{n+1}, as claimed.

Note that $S_{I^{\complement}} = S_I$. in what follows.

Here G^{σ}; for the remainder of the section.

In this case $\sigma(I) = \{\sigma(i) : i \in I\}$: and the result follows.

The value of $qS \geq 0$, is used in the estimate below.

We then obtain q'_I. holds throughout.

Recall that $\tilde{q} = Nq$; by the previous lemma.

It follows that $w = 2, 4, \ldots, n$: as claimed.

Suppose $w = n, n-2, \ldots,$ in what follows.

Note that $l = 2$. for the remainder of the section.

Here $l = 3$; and the result follows.

In this case 5.5: is used in the estimate below.

The value of $n - i + 1$, holds throughout.

We then obtain $I, J \subseteq [n]$. by the previous lemma.

Recall that $\Pi(M)$; as claimed.

It follows that $\Pi(M_n^{\mathrm{SQEC}})$: in what follows.

Suppose 3rd, for the remainder of the section.

Note that $1/(n/2)!$. and the result follows.

Here $l = 4$; is used in the estimate below.

In this case $-\tilde{S}_1 - 5\tilde{S}_2 + 5\tilde{S}_3 \geq 0$: holds throughout.

The value of $\S A + \S BC - \S ABC \geq 0$, by the previous lemma.

We then obtain $\tilde{S}_1 + \tilde{S}_2 - \tilde{S}_3 \geq 0$. as claimed.

Recall that $-\tilde{S}_1 + 2\tilde{S}_2 - \tilde{S}_3 \geq 0$; in what follows.

It follows that $-\S CD + \S BCD + \S ACD - \S ABCD \geq 0$: for the remainder of the section.

Suppose $\epsilon_{\Delta_V \Delta_W}$, and the result follows.

Note that Λ_L. is used in the estimate below.

Here v_B; holds throughout.

In this case $\lambda_L \leq \frac{2\pi}{\beta}$: by the previous lemma.

The value of T_+, as claimed.

We then obtain T_-. in what follows.

Recall that λ_L^-; for the remainder of the section.

It follows that $\beta_\pm = \beta(1 \mp \ell\Omega)$: and the result follows.

Suppose $x^- \to \lambda x^-$, is used in the estimate below.

Note that $x^+ \to \bar{\lambda} x^+$. holds throughout.

Here $SL(2,R) \times U(1)$; by the previous lemma.

In this case $SL(2,R) \times SL(2,R)$: as claimed.

The value of H^2, in what follows.

We then obtain $SL(2,R)$. for the remainder of the section.

Recall that AdS_3; and the result follows.

It follows that $\tilde{h}_\psi = h_\psi - q^2/k$: is used in the estimate below.

Suppose h_ψ, holds throughout.

Note that q_ψ. by the previous lemma.

Here $\tilde{h}_i$; as claimed.

In this case $m = 1.05$: in what follows.

The value of $m = 10$, for the remainder of the section.

We then obtain $c = l = 1$. and the result follows.

Recall that $j^i = \frac{1}{2}(\delta x^- \nabla^i \delta x^+ - \delta x^+ \nabla^i \delta x^-)$; is used in the estimate below.

It follows that 4122: holds throughout.

Suppose $U^*_{\mathscr{J}}$, by the previous lemma.

Note that $\beta^{\uparrow}$. as claimed.

Here $\partial \mathcal{M}_m$; in what follows.

In this case $G = (V, E)$: for the remainder of the section.

The value of $U \subset V$, and the result follows.

We then obtain $U^*_{\mathscr{J}\complement}$. is used in the estimate below.

Recall that $\mathcal{S} = \Gamma(\mathcal{W})$; holds throughout.

It follows that $\mathbf{w} \in \mathbb{W}$: by the previous lemma.

Suppose $(A\sigma, B\sigma, C\sigma, O\sigma)$, as claimed.

Note that $\mathcal{W}_{\mathfrak{m}'} \neq \mathbf{x}_{\mathcal{W}}$. in what follows.

Here $\mathbf{w}' = \mathbf{x}_{\mathcal{W}_{\mathfrak{m}}}$; for the remainder of the section.

In this case $\{A, \sigma\}$: and the result follows.

The value of G^1_2, is used in the estimate below.

We then obtain $\mathbb{P} = \pi(\rho)$. holds throughout.

Recall that $\beta(\partial V \cap U) = \mathscr{I}$; by the previous lemma.

It follows that $\mathscr{I}(f)$: as claimed.

Suppose $Y = \mathscr{C}^*_{\mathscr{J}} \setminus X$, in what follows.

Note that $\mathscr{I}(e) \triangle \mathscr{I}(f) = \mathscr{I}(e) \cup \mathscr{I}(f)$. for the remainder of the section.

Here $\mathscr{L}_2$; and the result follows.

In this case $\operatorname{Ker} \Gamma^{\intercal} = \{\mathbf{0}\}$: is used in the estimate below.

The value of $\mathscr{I}'$, holds throughout.

We then obtain B'. by the previous lemma.

Recall that $\mathcal{W}_{\mathfrak{m}'} \subseteq \mathcal{W}_{\mathfrak{m}}$; as claimed.

It follows that A_1: in what follows.

Suppose $(1, 1, 1)$, for the remainder of the section.

Note that $\Phi_{6 \to 5}$. and the result follows.

Here $\mathscr{R}_{\text{SSA}} = \mathscr{R}$; is used in the estimate below.

In this case $a + b$: holds throughout.

The value of $\{k\}, \{i, j\}$, by the previous lemma.

We then obtain $|\Psi_{AF}\rangle$. as claimed.

Recall that $g \mapsto U_g \left|\Psi_{AF}\right\rangle$; in what follows.

It follows that $M - N \in \mathscr{P}(\mathcal{H})$: for the remainder of the section.

Suppose $\{|x\rangle\}_x$, and the result follows.

Note that $\mathscr{C}(A \to B)$. is used in the estimate below.

Here $\mathrm{H}(A)_\rho$; holds throughout.

In this case U_1, U_2: by the previous lemma.

The value of $\sigma \in \mathscr{D}(\mathcal{H})$, as claimed.

We then obtain $n_{\lambda} > 1.$ in what follows.

Recall that $\mathcal{G}^{\otimes n}$; for the remainder of the section.

It follows that $q_{\widehat{MM}}$: and the result follows.

Suppose $\varepsilon < 1$, is used in the estimate below.

Note that $x \leq y.$ holds throughout.

Here $\mathcal{C} \in (\mathfrak{E}, \mathfrak{D})$; by the previous lemma.

In this case d_A: as claimed.

The value of $\pi \in S(N)$, in what follows.

We then obtain $k = 0, \ldots, \lfloor N/2 \rfloor.$ for the remainder of the section.

Recall that ρ_A'; and the result follows.

It follows that $\Psi_A^{\otimes n}$: is used in the estimate below.

Suppose $x^{\alpha} \leq x$, holds throughout.

Note that $\{|\psi_{A,n}\rangle\}.$ by the previous lemma.

Here $\mathcal{S}^{f,t}$; as claimed.

In this case $s' \in [0,1]$: in what follows.

The value of $\overline{\sigma}_A$, for the remainder of the section.

We then obtain $(1 - \delta^{F,T})^{-s} < 2^{s}.$ and the result follows.

Recall that $\{|\boldsymbol{b}_F^{l'}\rangle\}_{l'=1}^{L}$; is used in the estimate below.

It follows that $-sH(AF)_{\xi} + \frac{s^2}{2} V(AF)_{\xi}$: holds throughout.

Suppose P_A, by the previous lemma.

Note that $\rho_{AF}.$ as claimed.

Here ${\rho'_{AF}}^{\otimes n}$; in what follows.

In this case $a = (a^i)_{i=1}^{k}$: for the remainder of the section.

The value of $F \circ L[V^{-1}]$, and the result follows.

We then obtain $\hat{\eta} = (\bar{\eta}_1, \ldots, \bar{\eta}_{d-l}) \in \mathbb{R}^{d-l}.$ is used in the estimate below.

Recall that $\theta \in \mathcal{E}$; holds throughout.

It follows that $D^F(\theta_0 \| \hat{\theta}) - D^F(\theta_0 \| \theta^*) = Tx + O(x^2) < 0$: by the previous lemma.

Suppose $\Xi(v_1, \ldots, v_l)$, as claimed.

Note that $\phi_{\mathcal{M}}^{(e)}(\bar{\theta}) := U(\bar{\theta}, \kappa(\bar{\theta}))$. in what follows.

Here $\theta^{(n_m)}$; for the remainder of the section.

In this case $D^F(\theta_t \| \theta_0) > D^F(\theta_1 \| \theta_0)$: and the result follows.

The value of $\mathcal{X}_1 \times \mathcal{X}_2 \times \mathcal{X}_3$, is used in the estimate below.

We then obtain $\xi(x_i)_* \in \mathbb{R}^l$. holds throughout.

Recall that $(f_{n-1-j}(x_0))_{j=1}^{l}$; by the previous lemma.

It follows that $f_1, \ldots, f_{n_1 n_2 - 1}$: as claimed.

Suppose ρ_i, in what follows.

Note that $\{\theta_n^*(\mathcal{M}, \mathcal{E})\}$. for the remainder of the section.

Here $\theta_{(1)}$; and the result follows.

In this case $\theta_{(t_2(\lambda)-1),\lambda}$: is used in the estimate below.

The value of Q_Y, holds throughout.

We then obtain $\frac{d}{d\tau}\hat{F}[P_Y](0) < 0$. by the previous lemma.

Recall that $P_Y^{(t)}$; as claimed.

It follows that $g_i(x, y) := \delta_{i,x}$: in what follows.

Suppose $A \subset \{1, \ldots, m\}$, for the remainder of the section.

Note that $\mathbb{R}^{n_1 n_3 (n_2 - 1)}$. and the result follows.

Here $(\mathbb{R}^{n_1(n_2-1)}, \mu, D^{\mu})$; is used in the estimate below.

In this case $\rho_R \otimes \rho_B^{(t+1)}$: holds throughout.

The value of $\mathbb{R}^{d_A^2 d_B^2 d_C - 1}$, by the previous lemma.

We then obtain d_A^2. as claimed.

Recall that $b_{k+1} := b_k$; in what follows.

It follows that $\theta_0 \in \hat{\mathcal{M}}_\lambda \setminus \mathcal{M}_\lambda$: for the remainder of the section.

Suppose $n_1 - 1$, and the result follows.

Note that $\mathcal{X} := \{1, \ldots, n_1\}$. is used in the estimate below.

Here $(W \cdot Q_X)(x, y) := \sum_{x \in \mathcal{X}} W(y|x) Q_X(x)$; holds throughout.

In this case $\widehat{Q}_{X,*}$: by the previous lemma.

The value of $P_{\theta,Y}$, as claimed.

We then obtain $W \cdot \widehat{Q}_{X,*} = Q_{Y,*}$. in what follows.

Recall that $\theta^i = -H(W_i) + H(W_{n_2})$; for the remainder of the section.

It follows that $f_{n_1}, \ldots, f_{n_2 - 1}$: and the result follows.

Suppose x_1, x_2, is used in the estimate below.

Note that $\mathcal{X} = \{1, 2, 3, 4\}$. holds throughout.

Here $P_{j,Y}$; by the previous lemma.

In this case C_4: as claimed.

The value of $\{1, 2, 4\}$, in what follows.

We then obtain $\epsilon \to 0$. for the remainder of the section.

Recall that $\epsilon \geq 0.3972$; and the result follows.

It follows that C_{**}: is used in the estimate below.

Suppose $\{W_j\}_{j=1}^{n_1}$, holds throughout.

Note that $n_2^2 - 1$. by the previous lemma.

Here $\sigma = \rho_\theta$; as claimed.

In this case $n_2^2 \times n_2^2$: in what follows.

The value of $O(n_2^4)$, for the remainder of the section.

We then obtain $(W(y|x))_{x,y}$. and the result follows.

Recall that $y = n_1, \ldots, n_2$; is used in the estimate below.

It follows that $Q_{X,\epsilon}$: holds throughout.

Suppose $x = 1, \ldots, n_1$, by the previous lemma.

Note that $\{(h_{x,j})_{j=1}^{n_1-1}\}_{x=1}^{n_1-1}$. as claimed.

Here $D(W_x \| Q_{Y,*}) = D(W_x \| \overline{Q}_{Y,*}) + D(\overline{Q}_{Y,*} \| Q_{Y,*})$; in what follows.

In this case f_{x_0}: for the remainder of the section.

The value of $Q_{Y,*} = Q_{Y,t_*}, Q_{Y,0}$, and the result follows.

We then obtain $x \in \mathcal{X} \setminus \{x_0\}$. is used in the estimate below.

Recall that $\widehat{Q}_{X,2}$; holds throughout.

It follows that $m_\phi = 10^{-5} m_\chi$: by the previous lemma.

Suppose $|\tilde{\Psi}\rangle$, as claimed.

Note that $\mu = m_\chi m_N / (m_\chi + m_N)$. in what follows.

Here $f(\mathbf{v})$; for the remainder of the section.

In this case $q > 1/\Delta x$: and the result follows.

The value of N_0, is used in the estimate below.

We then obtain σ_V. holds throughout.

Recall that 10^8; by the previous lemma.

It follows that σ_ϕ: as claimed.

Suppose 2.9×10^{-7}, in what follows.

Note that 1.0×10^{-4}. for the remainder of the section.

Here 0.9; and the result follows.

In this case $N = AN_A$: is used in the estimate below.

The value of $R \propto A^2 N_A$, holds throughout.

We then obtain $j_1(x)$. by the previous lemma.

Recall that $\mathbf{v}_e = 0$; as claimed.

It follows that $m_\chi \to \infty$: in what follows.

Suppose $(y_\chi y_\phi)^2$, for the remainder of the section.

Note that $\sin(\mathbf{q} \cdot \mathbf{\Delta x}) \simeq q \, \Delta x \cos\theta$. and the result follows.

Here $\tilde{v}_e$; is used in the estimate below.

In this case $F'_{\mu\nu}$: holds throughout.

The value of g_B, by the previous lemma.

We then obtain 100 GeV. as claimed.

Recall that $T^{\rm dec}_{\nu_L}$; in what follows.

It follows that T_γ: for the remainder of the section.

Suppose $\theta_{\rm e} \approx 42^\circ$, and the result follows.

Note that $\theta_{\rm x} = \pi/2$. is used in the estimate below.

Here $\phi(t)/\phi_0$; holds throughout.

In this case $F(\mathbf{q})$: by the previous lemma.

The value of $\varepsilon^{(r)}$, as claimed.

We then obtain $10 - 100$. in what follows.

Recall that $\phi(x) = \tanh(gx)$; for the remainder of the section.

It follows that $\{\sigma\}$: and the result follows.

Suppose $g > 1$, is used in the estimate below.

Note that $\mu_\sigma = N_a/N$. holds throughout.

Here σ_β^{DMFT}; by the previous lemma.

In this case μ_σ: as claimed.

The value of $g < \sqrt{2}$, in what follows.

We then obtain $G' = QG$. for the remainder of the section.

Recall that $\langle (J_{ij})^2 \rangle = N^{-1}$; and the result follows.

It follows that $\eta = 0.8$: is used in the estimate below.

Suppose $g \to g(1 + \delta^2)$, holds throughout.

Note that $\sigma_\mu = 0$. by the previous lemma.

Here $[-1, 1]^3$; as claimed.

In this case $\phi(x) = \max(0, x)$: in what follows.

The value of $\phi''(x)$, for the remainder of the section.

We then obtain τ_h. and the result follows.

Recall that $\tau_{switch} \sim \alpha^2 \langle \phi(h)^2 \rangle$; is used in the estimate below.

It follows that $\partial_t \mathbf{h}(t) = F[G(\mathbf{h})]$: holds throughout.

Suppose $\Delta\tau = \tau_E - \tau_I > 0$, by the previous lemma.

Note that $\Delta_h = \int Dx \phi(\sqrt{\Delta_h} x)^2$. as claimed.

Here $\Delta = C_\phi(\Delta)$; in what follows.

In this case $Nd = N(1 - \mu_\sigma)$: for the remainder of the section.

The value of $e_\nu^\mu = \delta_\nu^\mu$, and the result follows.

We then obtain $N \to \infty$. is used in the estimate below.

Recall that N_a; holds throughout.

It follows that $\hat{l}_\mu = 0$: by the previous lemma.

Suppose $\phi_\mu^{'} \equiv \phi'(f_\mu)$, as claimed.

Note that $N = 3, g = 2.0$. in what follows.

Here $s_{E/I}$; for the remainder of the section.

In this case $X = (X_\ell)_{\ell=1}^n = (X_1, X_2, \ldots, X_n)$: and the result follows.

The value of $X_\ell = i$, is used in the estimate below.

We then obtain $j \in \partial i$. holds throughout.

Recall that c^*; by the previous lemma.

It follows that $\lambda_{s,N}[\nu^*]$: as claimed.

Suppose ν^*, in what follows.

Note that $\lambda_{s,N}$. for the remainder of the section.

Here $f(X_l) = k_{X_l}$; and the result follows.

In this case $N - 2$: is used in the estimate below.

The value of $C_n \sim 0$, holds throughout.

We then obtain $N = 11$. by the previous lemma.

Recall that $k_3 = N - 2 = 9$; as claimed.

It follows that $k_4 = N - 3 = 8$: in what follows.

Suppose $\nu^* = (\nu_{12}^*, \nu_{23}^*, \nu_{34}^*, \nu_{44}^*)$, for the remainder of the section.

Note that $N = 15$. and the result follows.

Here $\lambda_{s,N}^{\prime}$; is used in the estimate below.

In this case $\lambda_{s,\infty}$: holds throughout.

The value of $s > s_c$, by the previous lemma.

We then obtain $\tilde{\epsilon}_{\tilde{s}}$. as claimed.

Recall that $(-\log 2/2, -\log 2/2$; in what follows.

It follows that $\tilde{s} \gg 0$: for the remainder of the section.

Suppose $b \to \infty$, and the result follows.

Note that $p_{b1} = 1/b.$ is used in the estimate below.

Here $\sum_{i,j \in \{1,b\}} \nu_{ij} = 1$; holds throughout.

In this case ν_{11}: by the previous lemma.

The value of $s_c = -\log 2$, as claimed.

We then obtain $\lambda_{s,b}.$ in what follows.

Recall that $b = 250$; for the remainder of the section.

It follows that $s^- \equiv s < s_c$: and the result follows.

Suppose $b^{-\gamma}$, is used in the estimate below.

Note that $\sim 160meV.$ holds throughout.

Here $C_{2z}T$; by the previous lemma.

In this case N_M: as claimed.

The value of $e_{Y_1} = e_{Y_2}$, in what follows.

We then obtain $|\Delta_{e_Y=+}| = |\Delta_{e_Y=-}| = \Delta_0.$ for the remainder of the section.

Recall that $V_{pp\sigma}$; and the result follows.

It follows that 5.229: is used in the estimate below.

Suppose $\mathbf{d}_{\alpha s}^{r(i)}$, holds throughout.

Note that $i, j \in x, y, z.$ by the previous lemma.

Here $E_{\pm} = \pm v_f k$; as claimed.

In this case $\mathbf{p} = \mathbf{K}_1 + \delta\mathbf{p}$: in what follows.

The value of τ'_{α}, for the remainder of the section.

We then obtain $\eta\mathbf{K}_D^l - \mathbf{Q}.$ and the result follows.

Recall that $\mathbf{k}_1 = -\mathbf{k}, \mathbf{k}'_1 = -\mathbf{k}'$; is used in the estimate below.

It follows that ζ^a: holds throughout.

Suppose D_i, by the previous lemma.

Note that $\eta = -.$ as claimed.

Here $\mathbf{k} = \Gamma_M$; in what follows.

In this case $\mathbf{p} = \mathbf{k} + \eta\mathbf{K}_D - \mathbf{Q}'_{l\eta}$: for the remainder of the section.

The value of z^2, and the result follows.

We then obtain $S_{rel}^-.$ is used in the estimate below.

Recall that $|\lambda_{C_{2z}}| = 1$; holds throughout.

It follows that U_0: by the previous lemma.

Suppose $\mathbf{k} = 0$, as claimed.

Note that Φ_{+-}. in what follows.

Here $\mu = \nu = y$; for the remainder of the section.

In this case $\beta_{\mathbf{k},n}^{\dagger}$: and the result follows.

The value of $\mathcal{S}^{10} = s_x$, is used in the estimate below.

We then obtain $|\Delta_{\mathbf{k},+}| = |\Delta_{\mathbf{k},-}| = \tilde{\Delta}_{0,k}$. holds throughout.

Recall that $\xi \approx 10nm$; by the previous lemma.

It follows that $\beta(\varphi_0)$: as claimed.

Suppose $\beta(\varphi_0) = \beta_0$, in what follows.

Note that -4.5. for the remainder of the section.

Here $\beta_0 \gg 1$; and the result follows.

In this case $\mathcal{O}(\Omega^2)$: is used in the estimate below.

The value of D/r, holds throughout.

We then obtain 10^{16}. by the previous lemma.

Recall that $\sqrt{M/R^3}$; as claimed.

It follows that $\varphi_0 = 10^{-5}$: in what follows.

Suppose $\bar{I} - \bar{Q}$, for the remainder of the section.

Note that 10%. and the result follows.

Here $f(\varphi)$; is used in the estimate below.

In this case φ^a: holds throughout.

The value of $F_{\mu\nu} = \nabla_\mu X_\nu - \nabla_\nu X_\mu$, by the previous lemma.

We then obtain $\mu_X = 1.6 \times 10^{-11}$. as claimed.

Recall that $g^{tt} < 0$; in what follows.

It follows that $g^{tt} > 0$: for the remainder of the section.

Suppose $F^{\rho\mu}$, and the result follows.

Note that $2\mu_X^2 x \to 4V(x)$. is used in the estimate below.

Here $f_{\mu\nu}$; holds throughout.

In this case $\lambda\varphi^3$: by the previous lemma.

The value of $M = 0$, as claimed.

We then obtain $\nabla^{(0)}$. in what follows.

Recall that $X = -\nabla_\mu \varphi \nabla^\mu \varphi / 2$; for the remainder of the section.

It follows that μ_φ^2: and the result follows.

Suppose $\alpha = \lambda$, is used in the estimate below.

Note that $T < 0$. holds throughout.

Here $f(\varphi) = \lambda^2(1 - e^{-3/2\varphi^2})/12$; by the previous lemma.

In this case $\varphi \to -\varphi$: as claimed.

The value of $R = 0$, in what follows.

We then obtain ≥ 300. for the remainder of the section.

Recall that $\hat{n}_a$; and the result follows.

It follows that $t_e = t$: is used in the estimate below.

Suppose $e^A_{ij}(\hat{\Omega})$, holds throughout.

Note that $z(f, \hat{n})$. by the previous lemma.

Here $Q(f, \hat{\Omega})$; as claimed.

In this case ${}_{\pm s}Y_{LM}(\hat{\Omega})$: in what follows.

The value of $c^I_{00} = 1$, for the remainder of the section.

We then obtain $P_- = (Q - iU)$. and the result follows.

Recall that $E(f, \hat{\Omega})$; is used in the estimate below.

It follows that $|Q|^2 + |U|^2 \leq I^2$: holds throughout.

Suppose $X \in \{E, B\}$, by the previous lemma.

Note that $A^{LM}_{\ell\ell'}$. as claimed.

Here $A^L_{\ell\ell'}$; in what follows.

In this case $|\ell - \ell'| \leq L \leq \ell + \ell'$: for the remainder of the section.

The value of $\left\langle A^{LM}_{\ell\ell'}(A^{LM}_{\ell\ell'})^* \right\rangle$, and the result follows.

We then obtain $\langle z^*_{\ell m} z_{\ell' m'} \rangle$. is used in the estimate below.

Recall that $fL \gg 1$); holds throughout.

It follows that $\hat{\Omega} = (\theta, \phi)$: by the previous lemma.

Suppose $I(\hat{\Omega})$, as claimed.

Note that $\mathcal{Z}^{P_\pm}_{\ell\ell' mm'}$. in what follows.

Here $X = E$; for the remainder of the section.

In this case $A^{LM}_{\ell,\ell'}$: and the result follows.

The value of I, Q, U, is used in the estimate below.

We then obtain $\{\ell', \ell\}$. holds throughout.

Recall that ${}^{(ab)}\Gamma^E_{LM}$; by the previous lemma.

It follows that $LM\ell\ell'$: as claimed.

Suppose $c^{E,B}_{4M}$, in what follows.

Note that $F^{\times}(\hat{\Omega})$. for the remainder of the section.

Here $X \in \{I, P_{+}, P_{-}\}$; and the result follows.

In this case $O(\log(k + \Delta))$: is used in the estimate below.

The value of $(\Delta + 2)$, holds throughout.

We then obtain $L(G)$. by the previous lemma.

Recall that $\mathtt{out}_{v}$; as claimed.

It follows that $x \in D(C)$: in what follows.

Suppose $t_0 \geq t^*$, for the remainder of the section.

Note that C_0. and the result follows.

Here $\pi(G, C) = 0$; is used in the estimate below.

In this case $e \in D(C_t) \cap D(C_{t+1})$: holds throughout.

The value of $\mathtt{out}_{u,t-1}(v) = \perp$, by the previous lemma.

We then obtain $H = G_t(B)$. as claimed.

Recall that $e \in \overline{D}(C_{t-1}) \cap D(\hat{C}_{t-1})$; in what follows.

It follows that $\lambda_{t'}(e)$: for the remainder of the section.

Suppose $V(G) - D$, and the result follows.

Note that $C_t(v) = o$. is used in the estimate below.

Here $t - (t^* + \phi + 1)$; holds throughout.

In this case $V(H_{t'})$: by the previous lemma.

The value of $r_u(v)$, as claimed.

We then obtain $\mathrm{d}_H(u)$. in what follows.

Recall that $F_v^g = \{\{u, v\} \in F_v \mid \mathrm{d}_H(u) \leq \mathrm{d}_H(v)\}$; for the remainder of the section.

It follows that $b_{v,u}(i) = 1$: and the result follows.

Suppose c_e, is used in the estimate below.

Note that $\rho_0 = c$. holds throughout.

Here $((1 + \epsilon)\Delta)$; by the previous lemma.

In this case $m_v = \perp$: as claimed.

The value of $isincidentonatleastonevertexin$, in what follows.

We then obtain $inducedbythenodesthatstartaphaseinround$. for the remainder of the section.

Recall that $asitmayhavebeenremoveddu ringtheroundinterval$; and the result follows.

It follows that $.Ifnode$: is used in the estimate below.

Suppose $)where$, holds throughout.

Note that $A \in \mathbb{R}^{n \times n}$. by the previous lemma.

Here $\boldsymbol{d} \in \mathbb{R}^m$; as claimed.

In this case $(0, 2)$: in what follows.

The value of $k = 0, 1, 2, \ldots$, for the remainder of the section.

We then obtain $\boldsymbol{x}^{(k)} \to \boldsymbol{x}^*$. and the result follows.

Recall that $\boldsymbol{x}^{(2)} = \mathbf{0}$; is used in the estimate below.

It follows that $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^m$: holds throughout.

Suppose $P = D^{-1}$, by the previous lemma.

Note that $x_i^{(k)}$. as claimed.

Here $\omega^{(0)} \in (0, 2)$; in what follows.

In this case $\omega^{(k)} \to 0$: for the remainder of the section.

The value of $\rho \in (0, 1)$, and the result follows.

We then obtain ε_ω. is used in the estimate below.

Recall that $c_1 \in (0, 1)$; holds throughout.

It follows that $\omega^{(0)} = 1$: by the previous lemma.

Suppose $h^{(k+1)} = 2$, as claimed.

Note that k'. in what follows.

Here $\overline{d}^{(l')} < -2$; for the remainder of the section.

In this case $\overline{s}_m^{(k-1)} > \overline{s}_m^{(k-2)}$: and the result follows.

The value of $\boldsymbol{x}_0$, is used in the estimate below.

We then obtain $\mathsf{sprandsym}$. holds throughout.

Recall that $\boldsymbol{b} = A\hat{\boldsymbol{x}} - \boldsymbol{y}$; by the previous lemma.

It follows that $\kappa = 10^4$: as claimed.

Suppose $\omega = 1.77...$, in what follows.

Note that $A \in \mathbb{R}^{10000 \times 10000}$. for the remainder of the section.

Here $\sigma = \min(\mathrm{diag}(A)) = 3030.3...$; and the result follows.

In this case $\omega = 0.1, 0.2, \ldots, 1.9$: is used in the estimate below.

The value of $\boldsymbol{x} \geq \mathbf{0}$, holds throughout.

We then obtain $\{\boldsymbol{x}^{(k)}\}_{k=0}^{\infty}$. by the previous lemma.

Recall that $\omega = 1.7$; as claimed.

It follows that (k, M): in what follows.

Suppose $\mathcal{Y} = \{0, 1\}$, for the remainder of the section.

Note that $k_N / N \to 0$. and the result follows.

Here $k = k_N$; is used in the estimate below.

In this case L_2: holds throughout.

The value of $\alpha > 1$, by the previous lemma.

We then obtain $d \leq 10$. as claimed.

Recall that $\mathcal{Y} \subseteq \mathbb{R}$; in what follows.

It follows that $k \leq N$: for the remainder of the section.

Suppose $x \in \operatorname{supp}(\mu)$, and the result follows.

Note that l_Y. is used in the estimate below.

Here $1 - \delta$; holds throughout.

In this case $N \geq 1$: by the previous lemma.

The value of (β, B), as claimed.

We then obtain $M = M_o N^{\frac{2\alpha}{2\alpha+1}} (\log \frac{1}{\delta})^{\frac{1}{2\alpha+1}} \leq N$. in what follows.

Recall that $M = N^{\gamma}$; for the remainder of the section.

It follows that $m_1, \ldots, m_L$: and the result follows.

Suppose $\Theta(N^{\frac{2\alpha}{2\alpha+1} - \gamma})$, is used in the estimate below.

Note that $\mathbf{X} = X_{1:N} = \{X_i\}_{i=1}^{N}$. holds throughout.

Here (σ, S); by the previous lemma.

In this case $\zeta := \frac{\sigma}{d} \wedge 1$: as claimed.

The value of $\zeta = \frac{\sigma}{d}$, in what follows.

We then obtain $|\mathbf{X}_m| = n \to \infty$. for the remainder of the section.

Recall that $\mathbb{E}[\log U_{kn}^{\infty}(x)] = \Psi(k) - \log p(x)$; and the result follows.

It follows that $\phi_k^{(f)}(u)$: is used in the estimate below.

Suppose $\mathcal{L}^{-1}(F(p))(u)$, holds throughout.

Note that $\alpha \in \mathbb{R}$. by the previous lemma.

Here $(k, M, \frac{M}{2})$; as claimed.

In this case $(1, M, \frac{M}{2}, M)$: in what follows.

The value of $k_{\mathsf{base}} \in \{1, 2, 3, 4, 5\}$, for the remainder of the section.

We then obtain $\boldsymbol{F}(\boldsymbol{S}, \boldsymbol{T})$. and the result follows.

Recall that $\boldsymbol{B}_{id}$; is used in the estimate below.

It follows that $\boldsymbol{B}_{exp}$: holds throughout.

Suppose $\boldsymbol{\alpha} \in \mathbb{R}^{80}$, by the previous lemma.

Note that $\boldsymbol{\beta} \in \mathbb{R}^{64}$. as claimed.

Here $\boldsymbol{n}_{i}$; in what follows.

In this case $\boldsymbol{t}_{i}$: for the remainder of the section.

The value of Φ_{b}, and the result follows.

We then obtain $\boldsymbol{p} \in \mathbb{R}^{6}$. is used in the estimate below.

Recall that $\boldsymbol{R} \in SO(3)$; holds throughout.

It follows that $\Pi : \mathbb{R}^{3} \to \mathbb{R}^{2}$: by the previous lemma.

Suppose $\boldsymbol{T}'$, as claimed.

Note that M_{k}. in what follows.

Here $\boldsymbol{C}_{k} = \{(\boldsymbol{I}_{i}, \{l_{n}\}_{i})\}_{i=1}^{M_{k}}$; for the remainder of the section.

In this case $\boldsymbol{I}'$: and the result follows.

The value of ω_{n}, is used in the estimate below.

We then obtain $\omega_{\boldsymbol{\alpha}} = 0.5$. holds throughout.

Recall that $\omega_{\boldsymbol{\beta}} = 2.0$; by the previous lemma.

It follows that $\lambda_{pho} = 1.92$: as claimed.

Suppose $\lambda_{lan} = 1.6e{-}3$, in what follows.

Note that $\boldsymbol{I}'(\hat{\boldsymbol{x}})$. for the remainder of the section.

Here $\boldsymbol{D}'$; and the result follows.

In this case $\boldsymbol{Z}'_{\boldsymbol{S}}$: is used in the estimate below.

The value of $\boldsymbol{Z}_{max} = \max(\boldsymbol{Z}_{\boldsymbol{S}_{i}})$, holds throughout.

We then obtain $K = 50k$. by the previous lemma.

Recall that $1/7$; as claimed.

It follows that $\boldsymbol{\alpha}_{gt}$: in what follows.

Suppose $\boldsymbol{\beta}_{gt}$, for the remainder of the section.

Note that $1.0e{-}4$. and the result follows.

Here $50k$; is used in the estimate below.

In this case $\mathcal{D}_{1}$: holds throughout.

The value of p^{2}, by the previous lemma.

We then obtain $g^{1}, \ldots, g^{\mathsf{y}}$. as claimed.

Recall that $2\mathsf{d}$; in what follows.

It follows that $\iota_{\mathbb{C}^{d}}$: for the remainder of the section.

Suppose $\mathcal{B} \in \mathfrak{B}$, and the result follows.

Note that $\mathcal{A} \subset \mathcal{C} \subset [\mathsf{n}]$. is used in the estimate below.

Here $\mathcal{DEC}[\mathcal{A}]$; holds throughout.

In this case U_D: by the previous lemma.

The value of $\in \mathbb{F}_q^{\times}$, as claimed.

We then obtain $R = GU_S \in \mathbb{F}_q^{\bar{\mathsf{n}}}$. in what follows.

Recall that $\bar{\mathsf{n}} \times \mathsf{n}$; for the remainder of the section.

It follows that $\mathsf{n} \geq \mathsf{r} > \mathsf{t} > 0$: and the result follows.

Suppose $2(\mathsf{r} - \mathsf{t})/\mathsf{n}$, is used in the estimate below.

Note that $, and.$ holds throughout.

Here $, anda$; by the previous lemma.

In this case $canbeconsideredasastabilizercodewiththestabilizer$: as claimed.

The value of $, theusermakesthemeasurementgivenbythePOVM$, in what follows.

We then obtain $, andsendsthe.$ for the remainder of the section.

Recall that $, thestatereduceddensitywithrespectto$; and the result follows.

It follows that $, theprotocoldoesnothavethematrix$: is used in the estimate below.

Suppose $-EAMDScodewhenitis$, holds throughout.

Note that $onthesystem.$ by the previous lemma.

Here $iscolumnorthogonalto$; as claimed.

In this case $-secureEASPIRprotocolwithrate$: in what follows.

The value of $, wehave$, for the remainder of the section.

We then obtain $, andthesecondrelationshowsthattheMMSP.$ and the result follows.

Recall that $componentsby$; is used in the estimate below.

It follows that $.Thecasewith$: holds throughout.

Suppose $, therelationholdsfor$, by the previous lemma.

Note that $\rho \to \infty$. as claimed.

Here $\rho = \rho_0$; in what follows.

In this case $\{\boldsymbol{\theta}_k\}$: for the remainder of the section.

The value of $|\mathcal{I}_k| = b$, and the result follows.

We then obtain $\mathcal{I} \leftarrow$. is used in the estimate below.

Recall that $g = 0$; holds throughout.

It follows that $0.5 < \gamma \leq 1$: by the previous lemma.

Suppose $m = 5c^2 + 2L^2r^2 + 2LGr$, as claimed.

Note that δ_0. in what follows.

Here $\mathbb{E}\|P_C(\boldsymbol{\theta}_k) - \boldsymbol{\theta}_*\|^2$; for the remainder of the section.

In this case ζ_k: and the result follows.

The value of $P_{C_s}(\boldsymbol{\theta})$, is used in the estimate below.

We then obtain 20%. holds throughout.

Recall that $\gamma = \frac{4}{3}$; by the previous lemma.

It follows that C_u: as claimed.

Suppose $p = q = 64$, in what follows.

Note that $\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_*\|^2$. for the remainder of the section.

Here $\mathcal{O}(pb)$; and the result follows.

In this case $\rho_1 = 1 \times 10^{-2}$: is used in the estimate below.

The value of $\boldsymbol{\theta}_* \in C'$, holds throughout.

We then obtain $\|\nabla g(\boldsymbol{x}_k)\| \leq \|\nabla g(\boldsymbol{x}_{k-1})\|$. by the previous lemma.

Recall that $\sum_{i=1}^{+\infty} b_i < +\infty$; as claimed.

It follows that $k = 0, 1, ...$: in what follows.

Suppose $F[P_C(\boldsymbol{\theta}_{i_k - 1})] - F(\boldsymbol{\theta}_*) \to 0$, for the remainder of the section.

Note that $b_n = b_1 n^{-\beta}$. and the result follows.

Here $\beta = 1$; is used in the estimate below.

In this case $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$: holds throughout.

The value of $L_\delta(\tilde{y}_i - \tilde{\boldsymbol{x}}_i^T \boldsymbol{\theta})$, by the previous lemma.

We then obtain $\|\boldsymbol{\theta}_{\text{true}}\|_2 = 2$. as claimed.

Recall that $\boldsymbol{X}_i \in \mathbb{R}^{p \times p}$; in what follows.

It follows that $Bernoulli(p_i)$: for the remainder of the section.

Suppose μ, Σ, and the result follows.

Note that $\theta \in \Theta$. is used in the estimate below.

Here $g_\theta(x) := g(\theta, x)$; holds throughout.

In this case $\theta_0 \in \Theta$: by the previous lemma.

The value of $q = 1$, as claimed.

We then obtain $v \in S$. in what follows.

Recall that $W(q)$; for the remainder of the section.

It follows that $\Delta(v, X)$: and the result follows.

Suppose $g_\theta(y) := y^\top X \theta$, is used in the estimate below.

Note that $X > z_{1-\alpha}$. holds throughout.

Here $\alpha = 0.025$; by the previous lemma.

In this case $f''(\theta) \leq 1$: as claimed.

The value of $2^p - 1$, in what follows.

We then obtain $\{H_i\}_{i=1}^{I}$. for the remainder of the section.

Recall that $b_j = 1$; and the result follows.

It follows that $\hat{\lambda}_i(\alpha)$: is used in the estimate below.

Suppose $\mathcal{F}_\tau$, holds throughout.

Note that S_τ. by the previous lemma.

Here θ_1; as claimed.

In this case (θ_3, θ_4): in what follows.

The value of $\mathrm{Bern}(p_i)$, for the remainder of the section.

We then obtain λ^*. and the result follows.

Recall that $\theta_3 = -1$; is used in the estimate below.

It follows that $(p_3 = 0)$: holds throughout.

Suppose $(\theta, \theta, \theta, -1)$, by the previous lemma.

Note that (μ, σ^2). as claimed.

Here L_X^2; in what follows.

In this case $\mathrm{Beta}(\alpha, \beta)$: for the remainder of the section.

The value of $\hat{u}(\theta)$, and the result follows.

We then obtain $c_1, c_2, \ldots, c_n$.. is used in the estimate below.

Recall that $h_k(x)$; holds throughout.

It follows that $\mu(x)$: by the previous lemma.

Suppose $Y = \min(T_1, T_2, \ldots, T_n)$, as claimed.

Note that c_j. in what follows.

Here $\mathbf{T} = (T_1, \ldots, T_n)'$; for the remainder of the section.

In this case $S_{\mathbf{T}}(0, \cdots, 0) = 1$: and the result follows.

The value of $t_j \in (0, \infty)$, is used in the estimate below.

We then obtain $S_{\mathbf{T}}(t_1, \cdots, t_n)$. holds throughout.

Recall that T_i; by the previous lemma.

It follows that $T_j < T_i$: as claimed.

Suppose $h(t \,|\, j) = -\frac{d}{dt} \log S(t \,|\, j) = f(t \,|\, j)/S(t \,|\, j)$, in what follows.

Note that $S_j(t_j)$. for the remainder of the section.

Here $i \neq j$, $j = 1, \ldots, n$; and the result follows.

In this case $\prod_{j=1}^{n} S_j(t_j)$: is used in the estimate below.

The value of c_1, holds throughout.

We then obtain $Y = \min(T_1, T_2)$. by the previous lemma.

Recall that g_2; as claimed.

It follows that $\frac{1}{b} \ln \frac{c}{a}$: in what follows.

Suppose $\frac{1}{b} \ln \frac{c}{a(1-kc)}$, for the remainder of the section.

Note that $\frac{c}{a_1 e^{-b_1 x} + a_2 e^{b_2 x} + c}$. and the result follows.

Here e_x; is used in the estimate below.

In this case $g_2(x)$: holds throughout.

The value of M_{GM}, by the previous lemma.

We then obtain e_0. as claimed.

Recall that $x > 0$; in what follows.

It follows that $b = 0.1$: for the remainder of the section.

Suppose $b = 0.85$, and the result follows.

Note that $a = 0.4$. is used in the estimate below.

Here $p(x) = c/\left(ae^{bx} + c\right)$; holds throughout.