The value of $\mathbb{A}^2/\mathbb{Q}$ is used in the estimate below.

We then obtain $\mathscr{S}_P$ holds throughout.

Recall that (m, l) by the previous lemma.

It follows that $k^2 < p^n$ as claimed.

Suppose $D_X := D \times_B X$ in what follows.

Note that $B \to \mathbb{P}^1_B$ for the remainder of the section.

Here (b, f) and the result follows.

In this case $f_b(0) = (A + C, C)$ is used in the estimate below.

The value of $\widetilde{s}(b, A, B, C, D) \in W^{0,v}_{d+u} \times W^{0,v}_u$ holds throughout.

We then obtain $W^{0,v}_d$ by the previous lemma.

Recall that $\mathscr{Z}_{n,d}$ as claimed.

It follows that $X_{b\mathbb{C}}$ in what follows.

Suppose $nz = \sum_i n_i P_i$ for the remainder of the section.

Note that $\mathscr{P}_b$ and the result follows.

Here m, ℓ is used in the estimate below.

In this case $\widetilde{S}_{\mathbb{Z}}$ holds throughout.

The value of $\widetilde{S_{\mathbb{Z},b}}$ by the previous lemma.

We then obtain $U_{v,d}(\bar{\widetilde{S}}_{\mathbb{Z}})$ as claimed.

Recall that $V_{\mathbb{C}}$ in what follows.

It follows that $K_{k,p,n} = \mathbb{Q}(\sqrt{-d})$ for the remainder of the section.

Suppose $k = 1$ and the result follows.

Note that $F_0 = 0$ is used in the estimate below.

Here D_2 holds throughout.

In this case $(1, d, k^2, p) \in \mathcal{S}$ by the previous lemma.

The value of $\varepsilon = -1$ as claimed.

We then obtain $r \geq 1$ in what follows.

Recall that $\alpha = k + m\sqrt{-d}$ for the remainder of the section.

It follows that $a = \pm t$ and the result follows.

Suppose $q = 3$ is used in the estimate below.

Note that $d > 3$ holds throughout.

Here $H\mathbb{F}_p$ by the previous lemma.

In this case $X = \Omega^2 S^3$ as claimed.

The value of $\xi = 0$ in what follows.

We then obtain $\otimes_A^{\mathbb{L}}$ for the remainder of the section.

Recall that $f : A_0 \to B_0$ and the result follows.

It follows that $H\mathbb{Z}_p \simeq H\pi_0(\mathbb{S}_p^\wedge)$ is used in the estimate below.

Suppose $Mf_{k,pu}$ holds throughout.

Note that $a^\sharp := \lim_{n\to\infty} y_n^{p^n}$ by the previous lemma.

Here $W(\kappa)$ as claimed.

In this case $R^\flat \to D/p$ in what follows.

The value of $(\varphi_{D/p}^n(b_n))_{n=0}^\infty$ for the remainder of the section.

We then obtain $\eta_D : D \to HR$ and the result follows.

Recall that $X_n := (\tau_{\geq n}\mathbb{S}_p^\wedge) \otimes_{\mathbb{S}_p^\wedge} W^+(k)$ is used in the estimate below.

It follows that $t_{R,\xi}$ holds throughout.

Suppose $\Omega S^3 \to * \to S^3$ by the previous lemma.

Note that $\tau_{\leq 0} Mf_{\kappa,pu} \simeq H\kappa$ as claimed.

Here $\xi R^\flat$ in what follows.

In this case $e = v(p)$ for the remainder of the section.

The value of $W^+(k)[[u]]$ and the result follows.

We then obtain $W(k)[[u]]$ is used in the estimate below.

Recall that $HA \otimes_{W^+(k)[[u]]} HA$ holds throughout.

It follows that t_{k,a_0} by the previous lemma.

Suppose $\deg u_1 = \cdots = \deg u_n = 0$ as claimed.

Note that $k = A/\mathfrak{m}$ in what follows.

Here $R := A/I$ for the remainder of the section.

In this case $H(A/I)$ and the result follows.

The value of $n < 0$ is used in the estimate below.

We then obtain $\pi_0(R)$ holds throughout.

Recall that $M_{(x)}^\wedge$ by the previous lemma.

It follows that $I \subseteq R$ as claimed.

Suppose $O(n^{\omega+\epsilon})$ in what follows.

Note that $\omega < 2.3725$ for the remainder of the section.

Here $i \in [k_X], j \in [k_Y], k \in [k_Z]$ and the result follows.

In this case $\Omega(n^2 \log n)$ is used in the estimate below.

The value of $x_i y_j z_k$ holds throughout.

We then obtain $a_{ijk} = b_{\pi_x(i)\pi_y(j)\pi_z(k)}$ by the previous lemma.

Recall that i, j, k as claimed.

It follows that $b \times c$ in what follows.

Suppose $q \geq 2$ for the remainder of the section.

Note that $Y' \subseteq Y$ and the result follows.

Here $\{x_0, \ldots, x_{q+1}\}$ is used in the estimate below.

In this case $a = b = c$ holds throughout.

The value of $I \subseteq [n]$ by the previous lemma.

We then obtain $\bigoplus_{i=1}^{n} T_{iii}$ as claimed.

Recall that $i, j, k, i', j', k' \in [n]$ in what follows.

It follows that $I \subseteq [N]$ for the remainder of the section.

Suppose $N/2 = n$ and the result follows.

Note that $1/n$ is used in the estimate below.

Here $B_Y = Y_J \times Z_{K'} \times X_{I''}$ holds throughout.

In this case N_T by the previous lemma.

The value of $h_X(I', J', K')$ as claimed.

We then obtain C_3 in what follows.

Recall that $g \geq 2$ for the remainder of the section.

It follows that $\alpha_{Z_k} = \gamma_{Z_k}$ and the result follows.

Suppose $X^2 = \{x_{q+1}\}$ is used in the estimate below.

Note that T_{200} holds throughout.

Here $L_{t,I} := \{A \in \{0, 1, \ldots, q+1\}^t \mid \sum_{\ell=1}^{t} \kappa(A_\ell) = I\}$ by the previous lemma.

In this case $I' \in \{0, 1, \ldots, I\}$ as claimed.

The value of $x_{q+1,q+1}$ in what follows.

We then obtain $(i, j, k) \in S'$ for the remainder of the section.

Recall that H_1 and the result follows.

It follows that $A : H_1 \to H_3$ is used in the estimate below.

Suppose x^* holds throughout.

Note that A^{T} by the previous lemma.

Here $(\varepsilon, \min\{\frac{1}{\|A\|^2}, \frac{1}{\|B\|^2}\} - \varepsilon)$ as claimed.

In this case $C = \{x \in H_1 : \mathfrak{f}(x) \leq 0\}$ in what follows.

The value of $P_C x = \operatorname{argmin}_{y \in C} \|x - y\|, \ \forall x \in H$ for the remainder of the section.

We then obtain $\{x_n\}$ and the result follows.

Recall that $\gamma > 0$ is used in the estimate below.

It follows that $x \in C$ holds throughout.

Suppose $\{\zeta_n\}$ by the previous lemma.

Note that $\lim_{n\to\infty} \zeta_n = 0$ as claimed.

Here $\{v_n\} \subset H_2$ in what follows.

In this case $\lim_{n\to\infty} \delta_n = 0$ for the remainder of the section.

The value of $\{v_n\}$ and the result follows.

We then obtain Φ_n is used in the estimate below.

Recall that $\{y_{n_k}\}$ holds throughout.

It follows that $\bar{y} \in Q$ by the previous lemma.

Suppose $v_n \equiv v$ as claimed.

Note that Ψ_n in what follows.

Here $g : H_2 \to H_2$ for the remainder of the section.

In this case $(x^*, y^*) = P_\Omega(f(x^*), g(y^*)) \in \Omega$ and the result follows.

The value of $\lambda_1 \in (0, \frac{\sqrt{2}}{2})$ is used in the estimate below.

We then obtain $Q = \{(y_1, y_2, y_3) \in H_2,\ y_1^2 - y_2 + 5 \leq 0\}$ holds throughout.

Recall that x_0 by the previous lemma.

It follows that $\delta_n = \frac{1}{n+1}$ as claimed.

Suppose (d) in what follows.

Note that λ_1 for the remainder of the section.

Here $g(y) = 0.6y$ and the result follows.

In this case 10^{-4} is used in the estimate below.

The value of $V(G) = \{v_1, ..., v_n\}$ holds throughout.

We then obtain $N(G)$ by the previous lemma.

Recall that $u \in V(G)$ as claimed.

It follows that $u \in C$ in what follows.

Suppose $\mathbf{x}_W = (x_1, ..., x_n)^t$ for the remainder of the section.

Note that $v_i \in W$ and the result follows.

Here $\mathbf{x}_{N[v_i]}$ is used in the estimate below.

In this case $N(G)\mathbf{x}_S = \mathbf{0}$ holds throughout.

The value of $Col(N)$ by the previous lemma.

We then obtain $\rho(G) := dim(Col(N(G))$ as claimed.

Recall that $\rho(G)$ in what follows.

It follows that $\nu(G) = 0$ for the remainder of the section.

Suppose $pr(a) = 0$ and the result follows.

Note that $pr(A)$ is used in the estimate below.

Here $pr(S) = pr(\mathbf{x}_S)$ holds throughout.

In this case S_1 by the previous lemma.

The value of $nd(v) := \nu(G - v) - \nu(G)$ as claimed.

We then obtain $nd(v) = -1$ in what follows.

Recall that $N\mathbf{x}_S = \mathbf{1}$ for the remainder of the section.

It follows that $pr(S) = pr(\rho(G - v))$ and the result follows.

Suppose $\rho(T)$ is used in the estimate below.

Note that $pr(A \cap B) = \mathbf{x}_A^t \mathbf{x}_B$ holds throughout.

Here A^c by the previous lemma.

In this case $N\mathbf{1} = \mathbf{x}_{D^c}$ as claimed.

The value of $G_1 \oplus ... \oplus G_m$ in what follows.

We then obtain $V(G_1 \oplus ... \oplus G_m) = \cup_{i=1}^{m} V(G_i)$ for the remainder of the section.

Recall that G_1 and the result follows.

It follows that $\nu(G_1 \oplus G_2) = \nu(G_1) + \nu(G_2) + 1$ is used in the estimate below.

Suppose $\{G_1, ..., G_m\}$ holds throughout.

Note that $j \neq 0$ by the previous lemma.

Here $G = (N, A; \mathbf{w})$ as claimed.

In this case $N_1, \ldots, N_p \subseteq N$ in what follows.

The value of $i \in N$ for the remainder of the section.

We then obtain N_j and the result follows.

Recall that $(i, j) \in W$ is used in the estimate below.

It follows that $d(\xi)$ holds throughout.

Suppose $\omega \in \Omega(T)$ by the previous lemma.

Note that $\Omega \equiv \Omega(T)$ as claimed.

Here $p^{t\omega}$ in what follows.

In this case $i \in N_k$ for the remainder of the section.

The value of S_i^t and the result follows.

We then obtain D_i^t is used in the estimate below.

Recall that $S_i^t + R_i^t - D_i^{(t-1)} - d_i^{t\omega} < 0$ holds throughout.

It follows that $\overline{NCD}$ by the previous lemma.

Suppose $k = i$ as claimed.

Note that Φ_1 in what follows.

Here $\Phi_1, \dots, \Phi_4$ for the remainder of the section.

In this case (s, x, y) and the result follows.

The value of $1 =: t_0 < t_1 < \cdots < t_K := q$ is used in the estimate below.

We then obtain T_{k+1} holds throughout.

Recall that T_2^+ by the previous lemma.

It follows that $T_K^+ = T_K$ as claimed.

Suppose $i_0 \in N_l, (i, j) \in W, l \in P$ in what follows.

Note that $(\bar{s}, \bar{x}, \bar{y})$ for the remainder of the section.

Here N_1 and the result follows.

In this case r_j^-, r_j^+ is used in the estimate below.

The value of $\tilde{r}_i^-$ holds throughout.

We then obtain $\tilde{d_i^t} = (1 + \tilde{r}_i^+) d_i^t, \quad i \in N_j, j \in P, t \in T.$ by the previous lemma.

Recall that 250 as claimed.

It follows that $t = 33$ in what follows.

Suppose P_θ for the remainder of the section.

Note that $\operatorname{Fav}(E(r))$ and the result follows.

Here $f_\sharp \mu$ is used in the estimate below.

In this case $\mathbb{R}^n \setminus \{a\}$ holds throughout.

The value of $|\mathcal{K}_n + \Gamma| \sim \mathcal{P}_\Gamma(\mathcal{K}_n)$ by the previous lemma.

We then obtain $\pi_\theta(x)$ as claimed.

Recall that $c > 0$ in what follows.

It follows that $a \in A$ for the remainder of the section.

Suppose $\gamma \in \Gamma$ and the result follows.

Note that $\Omega = \mathbb{R}^n$ is used in the estimate below.

Here $P_a : E \to S^{n-1}$ holds throughout.

In this case $a, x, y \in B(x, R)$ by the previous lemma.

The value of Γ_i as claimed.

We then obtain s, t in what follows.

Recall that $m = 1$ for the remainder of the section.

It follows that $|b_1 - a_1|$ and the result follows.

Suppose $b_1 - a_1 < d_\lambda$ is used in the estimate below.

Note that $d_\lambda = \lambda - b_1 - t_0 = \lambda - a_1 - s_0 \geq 0$ holds throughout.

Here $a_1 \neq b_1$ by the previous lemma.

In this case $|b_2 - a_2|$ as claimed.

The value of $x = 0$ in what follows.

We then obtain $\mathbf{x} = (x_1, x_2)$ for the remainder of the section.

Recall that $(\Gamma + \mathbf{b})$ and the result follows.

It follows that $s = m$ is used in the estimate below.

Suppose $\widetilde{\pi_{\alpha\sharp}\mu} \ll h$ holds throughout.

Note that $r \in (0, 1)$ by the previous lemma.

Here $\psi = \mathcal{H}^{n-1}$ as claimed.

In this case $\nu(\mathcal{K}_n) = 1$ in what follows.

The value of $(0, 1/4)$ for the remainder of the section.

We then obtain $n - 2$ and the result follows.

Recall that A/I is used in the estimate below.

It follows that (A, J) holds throughout.

Suppose $\Gamma_{\mathbb{Z}[X]}(Y)$ by the previous lemma.

Note that $f(C) \in \mathcal{D}'$ as claimed.

Here $(K_0 \to K)^*$ in what follows.

In this case $A = \mathbb{Z}[x_1, \ldots, x_n]$ for the remainder of the section.

The value of $\mathcal{D} \hookrightarrow \mathcal{C}$ and the result follows.

We then obtain $\mathcal{C}' = \mathcal{C}$ is used in the estimate below.

Recall that $(A, I), (B, J)$ holds throughout.

It follows that $x \in I$ by the previous lemma.

Suppose $A \to A''$ as claimed.

Note that $M_{(A,I)} \to N_{(A,I)}$ in what follows.

Here $N \otimes^{\mathbb{L}}_A M' \to N \otimes^{\mathbb{L}}_A M \to N \otimes^{\mathbb{L}}_A M''$ for the remainder of the section.

In this case $\Gamma^*_{A/I}(I/I^2) \to J^{[*]}/J^{[*+1]}$ and the result follows.

The value of $\Gamma^*_{A/I}(I/I^2) \to \bigoplus J^{[*]}/J^{[*+1]}$ is used in the estimate below.

We then obtain $A \to S$ holds throughout.

Recall that $(B, J, \delta) = (A[X], IA[X], \gamma)$ by the previous lemma.

It follows that $B/J \cong A''/I$ as claimed.

Suppose $\Delta^{1} \times \Delta^{n}$ in what follows.

Note that $[a, b]$ for the remainder of the section.

Here $A \to C$ and the result follows.

In this case $A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{Q} \to D \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{Q}$ is used in the estimate below.

The value of $R/p \to R/p$ holds throughout.

We then obtain $(\mathbb{F}_{p}\{x\}, (x))$ by the previous lemma.

Recall that $M\{n\}$ as claimed.

It follows that $Q_{n} \in A[X_{0}, \ldots, X_{n-1}]$ in what follows.

Suppose $\prod_{(r,y) \in E} \delta^{r}(y/d)$ for the remainder of the section.

Note that $(A\{y\}/^{\mathbb{L}}(d, y))[p^{-1}]$ and the result follows.

Here $\deg(\delta^{j}(z_{2})) = p^{j}$ is used in the estimate below.

In this case (C, d) holds throughout.

The value of $S(R)$ by the previous lemma.

We then obtain $\partial \gamma_{R} = \partial R$ as claimed.

Recall that $\mathbb{R}^{2^{n}-1}$ in what follows.

It follows that $n' \gg n$ for the remainder of the section.

Suppose γ_{AB} and the result follows.

Note that I_{B}' is used in the estimate below.

Here $\partial I_{B}'$ holds throughout.

In this case 2^{L} by the previous lemma.

The value of I_{l} as claimed.

We then obtain $x_{l} = 1$ in what follows.

Recall that $x_{A} = 1$ for the remainder of the section.

It follows that $R_{J_{r}'} \neq R_{J_{r}}$ and the result follows.

Suppose $2 \leftrightarrow B$ is used in the estimate below.

Note that $x^{A} = 110$ holds throughout.

Here J_{r} by the previous lemma.

In this case σ_{ABC} as claimed.

The value of $f(111)_{4} = 0$ in what follows.

We then obtain $\gamma_{I_{l}}$ for the remainder of the section.

Recall that $w \in \mathbb{R}^{m}$ and the result follows.

It follows that $J_{1} = A$ is used in the estimate below.

Suppose y_{r} holds throughout.

Note that $\tilde{f}(x) = f(x)$ by the previous lemma.

Here A, BC, G as claimed.

In this case ABG, C, D in what follows.

The value of $n_k \geq k$ for the remainder of the section.

We then obtain $b^T x \geq 0$ and the result follows.

Recall that $\sum_{i=1}^{m} q_i$ is used in the estimate below.

It follows that $I \ni n + 1$ holds throughout.

Suppose $\S A + \S AB \geq \S B$ by the previous lemma.

Note that $I \cup J$ as claimed.

Here $n = 5$ in what follows.

In this case $I \smallsetminus J$ for the remainder of the section.

The value of $E \subset V \times V$ and the result follows.

We then obtain $\partial V \subseteq V$ is used in the estimate below.

Recall that $C(W) \subseteq E$ holds throughout.

It follows that $[n]$ by the previous lemma.

Suppose Sym_{n+1} as claimed.

Note that $S_{I^{\complement}} = S_I$ in what follows.

Here G^{σ} for the remainder of the section.

In this case $\sigma(I) = \{\sigma(i) : i \in I\}$ and the result follows.

The value of $qS \geq 0$ is used in the estimate below.

We then obtain q'_I holds throughout.

Recall that $\tilde{q} = Nq$ by the previous lemma.

It follows that $w = 2, 4, \ldots, n$ as claimed.

Suppose $w = n, n-2, \ldots$ in what follows.

Note that $l = 2$ for the remainder of the section.

Here $l = 3$ and the result follows.

In this case 5.5 is used in the estimate below.

The value of $n - i + 1$ holds throughout.

We then obtain $I, J \subseteq [n]$ by the previous lemma.

Recall that $\Pi(M)$ as claimed.

It follows that $\Pi(M_n^{\mathrm{SQEC}})$ in what follows.

Suppose 3rd for the remainder of the section.

Note that $1/(n/2)!$ and the result follows.

Here $l = 4$ is used in the estimate below.

In this case $-\tilde{S}_1 - 5\tilde{S}_2 + 5\tilde{S}_3 \geq 0$ holds throughout.

The value of $\S A + \S BC - \S ABC \geq 0$ by the previous lemma.

We then obtain $\tilde{S}_1 + \tilde{S}_2 - \tilde{S}_3 \geq 0$ as claimed.

Recall that $-\tilde{S}_1 + 2\tilde{S}_2 - \tilde{S}_3 \geq 0$ in what follows.

It follows that $-\S CD + \S BCD + \S ACD - \S ABCD \geq 0$ for the remainder of the section.

Suppose $\epsilon_{\Delta_V \Delta_W}$ and the result follows.

Note that Λ_L is used in the estimate below.

Here v_B holds throughout.

In this case $\lambda_L \leq \frac{2\pi}{\beta}$ by the previous lemma.

The value of T_+ as claimed.

We then obtain T_- in what follows.

Recall that λ_L^- for the remainder of the section.

It follows that $\beta_\pm = \beta(1 \mp \ell\Omega)$ and the result follows.

Suppose $x^- \to \lambda x^-$ is used in the estimate below.

Note that $x^+ \to \bar{\lambda} x^+$ holds throughout.

Here $SL(2,R) \times U(1)$ by the previous lemma.

In this case $SL(2,R) \times SL(2,R)$ as claimed.

The value of H^2 in what follows.

We then obtain $SL(2,R)$ for the remainder of the section.

Recall that AdS_3 and the result follows.

It follows that $\tilde{h}_\psi = h_\psi - q^2/k$ is used in the estimate below.

Suppose h_ψ holds throughout.

Note that q_ψ by the previous lemma.

Here $\tilde{h}_i$ as claimed.

In this case $m = 1.05$ in what follows.

The value of $m = 10$ for the remainder of the section.

We then obtain $c = l = 1$ and the result follows.

Recall that $j^i = \frac{1}{2}(\delta x^- \nabla^i \delta x^+ - \delta x^+ \nabla^i \delta x^-)$ is used in the estimate below.

It follows that 4122 holds throughout.

Suppose $U^*_{\mathscr{J}}$ by the previous lemma.

Note that $\beta^\uparrow$ as claimed.

Here $\partial \mathcal{M}_m$ in what follows.

In this case $G = (V, E)$ for the remainder of the section.

The value of $U \subset V$ and the result follows.

We then obtain $U^*_{\mathscr{J}\complement}$ is used in the estimate below.

Recall that $\mathcal{S} = \Gamma(\mathcal{W})$ holds throughout.

It follows that $\mathbf{w} \in \mathbb{W}$ by the previous lemma.

Suppose $(A\sigma, B\sigma, C\sigma, O\sigma)$ as claimed.

Note that $\mathcal{W}_{\mathfrak{m}'} \neq \mathbf{x}_{\mathcal{W}}$ in what follows.

Here $\mathbf{w}' = \mathbf{x}_{\mathcal{W}_{\mathfrak{m}}}$ for the remainder of the section.

In this case $\{A, \sigma\}$ and the result follows.

The value of G_2^1 is used in the estimate below.

We then obtain $\mathbb{P} = \pi(\rho)$ holds throughout.

Recall that $\beta(\partial V \cap U) = \mathscr{I}$ by the previous lemma.

It follows that $\mathscr{I}(f)$ as claimed.

Suppose $Y = \mathscr{C}^*_{\mathscr{J}} \setminus X$ in what follows.

Note that $\mathscr{I}(e) \triangle \mathscr{I}(f) = \mathscr{I}(e) \cup \mathscr{I}(f)$ for the remainder of the section.

Here $\mathscr{L}_2$ and the result follows.

In this case $\operatorname{Ker} \Gamma^{\intercal} = \{\mathbf{0}\}$ is used in the estimate below.

The value of $\mathscr{I}'$ holds throughout.

We then obtain B' by the previous lemma.

Recall that $\mathcal{W}_{\mathfrak{m}'} \subseteq \mathcal{W}_{\mathfrak{m}}$ as claimed.

It follows that A_1 in what follows.

Suppose $(1, 1, 1)$ for the remainder of the section.

Note that $\Phi_{6 \to 5}$ and the result follows.

Here $\mathscr{R}_{\text{SSA}} = \mathscr{R}$ is used in the estimate below.

In this case $a + b$ holds throughout.

The value of $\{k\}, \{i, j\}$ by the previous lemma.

We then obtain $|\Psi_{AF}\rangle$ as claimed.

Recall that $g \mapsto U_g \left|\Psi_{AF}\right\rangle$ in what follows.

It follows that $M - N \in \mathscr{P}(\mathcal{H})$ for the remainder of the section.

Suppose $\{|x\rangle\}_x$ and the result follows.

Note that $\mathscr{C}(A \to B)$ is used in the estimate below.

Here $\mathrm{H}(A)_\rho$ holds throughout.

In this case U_1, U_2 by the previous lemma.

The value of $\sigma \in \mathscr{D}(\mathcal{H})$ as claimed.

We then obtain $n_{\lambda} > 1$ in what follows.

Recall that $\mathcal{G}^{\otimes n}$ for the remainder of the section.

It follows that $q_{\widehat{MM}}$ and the result follows.

Suppose $\varepsilon < 1$ is used in the estimate below.

Note that $x \leq y$ holds throughout.

Here $\mathcal{C} \in (\mathfrak{E}, \mathfrak{D})$ by the previous lemma.

In this case d_A as claimed.

The value of $\pi \in S(N)$ in what follows.

We then obtain $k = 0, \ldots, \lfloor N/2 \rfloor$ for the remainder of the section.

Recall that ρ'_A and the result follows.

It follows that $\Psi_A^{\otimes n}$ is used in the estimate below.

Suppose $x^{\alpha} \leq x$ holds throughout.

Note that $\{|\psi_{A,n}\rangle\}$ by the previous lemma.

Here $\mathcal{S}^{f,t}$ as claimed.

In this case $s' \in [0,1]$ in what follows.

The value of $\overline{\sigma}_A$ for the remainder of the section.

We then obtain $(1 - \delta^{F,T})^{-s} < 2^s$ and the result follows.

Recall that $\{|\boldsymbol{b}_F^{l'}\rangle\}_{l'=1}^{L}$ is used in the estimate below.

It follows that $-sH(AF)_{\xi} + \frac{s^2}{2}V(AF)_{\xi}$ holds throughout.

Suppose P_A by the previous lemma.

Note that ρ_{AF} as claimed.

Here ${\rho'_{AF}}^{\otimes n}$ in what follows.

In this case $a = (a^i)_{i=1}^{k}$ for the remainder of the section.

The value of $F \circ L[V^{-1}]$ and the result follows.

We then obtain $\hat{\eta} = (\bar{\eta}_1, \ldots, \bar{\eta}_{d-l}) \in \mathbb{R}^{d-l}$ is used in the estimate below.

Recall that $\theta \in \mathcal{E}$ holds throughout.

It follows that $D^F(\theta_0 \| \hat{\theta}) - D^F(\theta_0 \| \theta^*) = Tx + O(x^2) < 0$ by the previous lemma.

Suppose $\Xi(v_1, \ldots, v_l)$ as claimed.

Note that $\phi_{\mathcal{M}}^{(e)}(\bar{\theta}) := U(\bar{\theta}, \kappa(\bar{\theta}))$ in what follows.

Here $\theta^{(n_m)}$ for the remainder of the section.

In this case $D^F(\theta_t \| \theta_0) > D^F(\theta_1 \| \theta_0)$ and the result follows.

The value of $\mathcal{X}_1 \times \mathcal{X}_2 \times \mathcal{X}_3$ is used in the estimate below.

We then obtain $\xi(x_i)_* \in \mathbb{R}^l$ holds throughout.

Recall that $(f_{n-1-j}(x_0))_{j=1}^{l}$ by the previous lemma.

It follows that $f_1, \ldots, f_{n_1 n_2 - 1}$ as claimed.

Suppose ρ_i in what follows.

Note that $\{\theta_n^*(\mathcal{M}, \mathcal{E})\}$ for the remainder of the section.

Here $\theta_{(1)}$ and the result follows.

In this case $\theta_{(t_2(\lambda)-1),\lambda}$ is used in the estimate below.

The value of Q_Y holds throughout.

We then obtain $\frac{d}{d\tau}\hat{F}[P_Y](0) < 0$ by the previous lemma.

Recall that $P_Y^{(t)}$ as claimed.

It follows that $g_i(x, y) := \delta_{i,x}$ in what follows.

Suppose $A \subset \{1, \ldots, m\}$ for the remainder of the section.

Note that $\mathbb{R}^{n_1 n_3 (n_2 - 1)}$ and the result follows.

Here $(\mathbb{R}^{n_1(n_2-1)}, \mu, D^{\mu})$ is used in the estimate below.

In this case $\rho_R \otimes \rho_B^{(t+1)}$ holds throughout.

The value of $\mathbb{R}^{d_A^2 d_B^2 d_C - 1}$ by the previous lemma.

We then obtain d_A^2 as claimed.

Recall that $b_{k+1} := b_k$ in what follows.

It follows that $\theta_0 \in \hat{\mathcal{M}}_\lambda \setminus \mathcal{M}_\lambda$ for the remainder of the section.

Suppose $n_1 - 1$ and the result follows.

Note that $\mathcal{X} := \{1, \ldots, n_1\}$ is used in the estimate below.

Here $(W \cdot Q_X)(x, y) := \sum_{x \in \mathcal{X}} W(y|x) Q_X(x)$ holds throughout.

In this case $\widehat{Q}_{X,*}$ by the previous lemma.

The value of $P_{\theta,Y}$ as claimed.

We then obtain $W \cdot \widehat{Q}_{X,*} = Q_{Y,*}$ in what follows.

Recall that $\theta^i = -H(W_i) + H(W_{n_2})$ for the remainder of the section.

It follows that $f_{n_1}, \ldots, f_{n_2 - 1}$ and the result follows.

Suppose x_1, x_2 is used in the estimate below.

Note that $\mathcal{X} = \{1, 2, 3, 4\}$ holds throughout.

Here $P_{j,Y}$ by the previous lemma.

In this case C_4 as claimed.

The value of $\{1, 2, 4\}$ in what follows.

We then obtain $\epsilon \to 0$ for the remainder of the section.

Recall that $\epsilon \geq 0.3972$ and the result follows.

It follows that C_{**} is used in the estimate below.

Suppose $\{W_j\}_{j=1}^{n_1}$ holds throughout.

Note that $n_2^2 - 1$ by the previous lemma.

Here $\sigma = \rho_\theta$ as claimed.

In this case $n_2^2 \times n_2^2$ in what follows.

The value of $O(n_2^4)$ for the remainder of the section.

We then obtain $(W(y|x))_{x,y}$ and the result follows.

Recall that $y = n_1, \ldots, n_2$ is used in the estimate below.

It follows that $Q_{X,\epsilon}$ holds throughout.

Suppose $x = 1, \ldots, n_1$ by the previous lemma.

Note that $\{(h_{x,j})_{j=1}^{n_1-1}\}_{x=1}^{n_1-1}$ as claimed.

Here $D(W_x \| Q_{Y,*}) = D(W_x \| \overline{Q}_{Y,*}) + D(\overline{Q}_{Y,*} \| Q_{Y,*})$ in what follows.

In this case f_{x_0} for the remainder of the section.

The value of $Q_{Y,*} = Q_{Y,t_*}, Q_{Y,0}$ and the result follows.

We then obtain $x \in \mathcal{X} \setminus \{x_0\}$ is used in the estimate below.

Recall that $\widehat{Q}_{X,2}$ holds throughout.

It follows that $m_\phi = 10^{-5} m_\chi$ by the previous lemma.

Suppose $|\tilde{\Psi}\rangle$ as claimed.

Note that $\mu = m_\chi m_N / (m_\chi + m_N)$ in what follows.

Here $f(\mathbf{v})$ for the remainder of the section.

In this case $q > 1/\Delta x$ and the result follows.

The value of N_0 is used in the estimate below.

We then obtain σ_V holds throughout.

Recall that 10^8 by the previous lemma.

It follows that σ_ϕ as claimed.

Suppose 2.9×10^{-7} in what follows.

Note that 1.0×10^{-4} for the remainder of the section.

Here 0.9 and the result follows.

In this case $N = AN_A$ is used in the estimate below.

The value of $R \propto A^2 N_A$ holds throughout.

We then obtain $j_1(x)$ by the previous lemma.

Recall that $\mathbf{v}_e = 0$ as claimed.

It follows that $m_\chi \to \infty$ in what follows.

Suppose $(y_\chi y_\phi)^2$ for the remainder of the section.

Note that $\sin(\mathbf{q} \cdot \mathbf{\Delta x}) \simeq q \, \Delta x \cos\theta$ and the result follows.

Here $\tilde{v}_e$ is used in the estimate below.

In this case $F'_{\mu\nu}$ holds throughout.

The value of g_B by the previous lemma.

We then obtain 100 GeV as claimed.

Recall that $T^{\rm dec}_{\nu_L}$ in what follows.

It follows that T_γ for the remainder of the section.

Suppose $\theta_{\rm e} \approx 42^\circ$ and the result follows.

Note that $\theta_{\rm x} = \pi/2$ is used in the estimate below.

Here $\phi(t)/\phi_0$ holds throughout.

In this case $F(\mathbf{q})$ by the previous lemma.

The value of $\varepsilon^{(r)}$ as claimed.

We then obtain $10 - 100$ in what follows.

Recall that $\phi(x) = \tanh(gx)$ for the remainder of the section.

It follows that $\{\sigma\}$ and the result follows.

Suppose $g > 1$ is used in the estimate below.

Note that $\mu_\sigma = N_a/N$ holds throughout.

Here σ_β^{DMFT} by the previous lemma.

In this case μ_σ as claimed.

The value of $g < \sqrt{2}$ in what follows.

We then obtain $G' = QG$ for the remainder of the section.

Recall that $\langle (J_{ij})^2 \rangle = N^{-1}$ and the result follows.

It follows that $\eta = 0.8$ is used in the estimate below.

Suppose $g \to g(1 + \delta^2)$ holds throughout.

Note that $\sigma_\mu = 0$ by the previous lemma.

Here $[-1, 1]^3$ as claimed.

In this case $\phi(x) = \max(0, x)$ in what follows.

The value of $\phi''(x)$ for the remainder of the section.

We then obtain τ_h and the result follows.

Recall that $\tau_{switch} \sim \alpha^2 \langle \phi(h)^2 \rangle$ is used in the estimate below.

It follows that $\partial_t \mathbf{h}(t) = F[G(\mathbf{h})]$ holds throughout.

Suppose $\Delta\tau = \tau_E - \tau_I > 0$ by the previous lemma.

Note that $\Delta_h = \int Dx \phi(\sqrt{\Delta_h} x)^2$ as claimed.

Here $\Delta = C_\phi(\Delta)$ in what follows.

In this case $Nd = N(1 - \mu_\sigma)$ for the remainder of the section.

The value of $e_\nu^\mu = \delta_\nu^\mu$ and the result follows.

We then obtain $N \to \infty$ is used in the estimate below.

Recall that N_a holds throughout.

It follows that $\hat{l}_\mu = 0$ by the previous lemma.

Suppose $\phi_\mu^{'} \equiv \phi'(f_\mu)$ as claimed.

Note that $N = 3, g = 2.0$ in what follows.

Here $s_{E/I}$ for the remainder of the section.

In this case $X = (X_\ell)_{\ell=1}^n = (X_1, X_2, \ldots, X_n)$ and the result follows.

The value of $X_\ell = i$ is used in the estimate below.

We then obtain $j \in \partial i$ holds throughout.

Recall that c^* by the previous lemma.

It follows that $\lambda_{s,N}[\nu^*]$ as claimed.

Suppose ν^* in what follows.

Note that $\lambda_{s,N}$ for the remainder of the section.

Here $f(X_l) = k_{X_l}$ and the result follows.

In this case $N - 2$ is used in the estimate below.

The value of $C_n \sim 0$ holds throughout.

We then obtain $N = 11$ by the previous lemma.

Recall that $k_3 = N - 2 = 9$ as claimed.

It follows that $k_4 = N - 3 = 8$ in what follows.

Suppose $\nu^* = (\nu_{12}^*, \nu_{23}^*, \nu_{34}^*, \nu_{44}^*)$ for the remainder of the section.

Note that $N = 15$ and the result follows.

Here $\lambda_{s,N}'$ is used in the estimate below.

In this case $\lambda_{s,\infty}$ holds throughout.

The value of $s > s_c$ by the previous lemma.

We then obtain $\tilde{\epsilon}_{\tilde{s}}$ as claimed.

Recall that $(-\log 2/2, -\log 2/2$ in what follows.

It follows that $\tilde{s} \gg 0$ for the remainder of the section.

Suppose $b \to \infty$ and the result follows.

Note that $p_{b1} = 1/b$ is used in the estimate below.

Here $\sum_{i,j \in \{1,b\}} \nu_{ij} = 1$ holds throughout.

In this case ν_{11} by the previous lemma.

The value of $s_c = -\log 2$ as claimed.

We then obtain $\lambda_{s,b}$ in what follows.

Recall that $b = 250$ for the remainder of the section.

It follows that $s^- \equiv s < s_c$ and the result follows.

Suppose $b^{-\gamma}$ is used in the estimate below.

Note that $\sim 160meV$ holds throughout.

Here $C_{2z}T$ by the previous lemma.

In this case N_M as claimed.

The value of $e_{Y_1} = e_{Y_2}$ in what follows.

We then obtain $|\Delta_{e_Y=+}| = |\Delta_{e_Y=-}| = \Delta_0$ for the remainder of the section.

Recall that $V_{pp\sigma}$ and the result follows.

It follows that 5.229 is used in the estimate below.

Suppose $\mathbf{d}_{\alpha s}^{r(i)}$ holds throughout.

Note that $i, j \in x, y, z$ by the previous lemma.

Here $E_{\pm} = \pm v_f k$ as claimed.

In this case $\mathbf{p} = \mathbf{K}_1 + \delta\mathbf{p}$ in what follows.

The value of τ'_{α} for the remainder of the section.

We then obtain $\eta\mathbf{K}_D^l - \mathbf{Q}$ and the result follows.

Recall that $\mathbf{k}_1 = -\mathbf{k}, \mathbf{k}'_1 = -\mathbf{k}'$ is used in the estimate below.

It follows that ζ^a holds throughout.

Suppose D_i by the previous lemma.

Note that $\eta = -$ as claimed.

Here $\mathbf{k} = \Gamma_M$ in what follows.

In this case $\mathbf{p} = \mathbf{k} + \eta\mathbf{K}_D - \mathbf{Q}'_{l\eta}$ for the remainder of the section.

The value of z^2 and the result follows.

We then obtain S_{rel}^- is used in the estimate below.

Recall that $|\lambda_{C_{2z}}| = 1$ holds throughout.

It follows that U_{0} by the previous lemma.

Suppose $\mathbf{k} = 0$ as claimed.

Note that Φ_{+-} in what follows.

Here $\mu = \nu = y$ for the remainder of the section.

In this case $\beta_{\mathbf{k},n}^{\dagger}$ and the result follows.

The value of $\mathcal{S}^{10} = s_{x}$ is used in the estimate below.

We then obtain $|\Delta_{\mathbf{k},+}| = |\Delta_{\mathbf{k},-}| = \tilde{\Delta}_{0,k}$ holds throughout.

Recall that $\xi \approx 10nm$ by the previous lemma.

It follows that $\beta(\varphi_{0})$ as claimed.

Suppose $\beta(\varphi_{0}) = \beta_{0}$ in what follows.

Note that -4.5 for the remainder of the section.

Here $\beta_{0} \gg 1$ and the result follows.

In this case $\mathcal{O}(\Omega^{2})$ is used in the estimate below.

The value of D/r holds throughout.

We then obtain 10^{16} by the previous lemma.

Recall that $\sqrt{M/R^{3}}$ as claimed.

It follows that $\varphi_{0} = 10^{-5}$ in what follows.

Suppose $\bar{I} - \bar{Q}$ for the remainder of the section.

Note that 10% and the result follows.

Here $f(\varphi)$ is used in the estimate below.

In this case φ^{a} holds throughout.

The value of $F_{\mu\nu} = \nabla_{\mu} X_{\nu} - \nabla_{\nu} X_{\mu}$ by the previous lemma.

We then obtain $\mu_{X} = 1.6 \times 10^{-11}$ as claimed.

Recall that $g^{tt} < 0$ in what follows.

It follows that $g^{tt} > 0$ for the remainder of the section.

Suppose $F^{\rho\mu}$ and the result follows.

Note that $2\mu_{X}^{2} x \to 4V(x)$ is used in the estimate below.

Here $f_{\mu\nu}$ holds throughout.

In this case $\lambda\varphi^{3}$ by the previous lemma.

The value of $M = 0$ as claimed.

We then obtain $\nabla^{(0)}$ in what follows.

Recall that $X = -\nabla_{\mu}\varphi\nabla^{\mu}\varphi/2$ for the remainder of the section.

It follows that μ_{φ}^{2} and the result follows.

Suppose $\alpha = \lambda$ is used in the estimate below.

Note that $T < 0$ holds throughout.

Here $f(\varphi) = \lambda^2(1 - e^{-3/2\varphi^2})/12$ by the previous lemma.

In this case $\varphi \to -\varphi$ as claimed.

The value of $R = 0$ in what follows.

We then obtain ≥ 300 for the remainder of the section.

Recall that $\hat{n}_a$ and the result follows.

It follows that $t_e = t$ is used in the estimate below.

Suppose $e^A_{ij}(\hat{\Omega})$ holds throughout.

Note that $z(f, \hat{n})$ by the previous lemma.

Here $Q(f, \hat{\Omega})$ as claimed.

In this case ${}_{\pm s}Y_{LM}(\hat{\Omega})$ in what follows.

The value of $c^I_{00} = 1$ for the remainder of the section.

We then obtain $P_- = (Q - iU)$ and the result follows.

Recall that $E(f, \hat{\Omega})$ is used in the estimate below.

It follows that $|Q|^2 + |U|^2 \leq I^2$ holds throughout.

Suppose $X \in \{E,\, B\}$ by the previous lemma.

Note that $A^{LM}_{\ell\ell'}$ as claimed.

Here $A^L_{\ell\ell'}$ in what follows.

In this case $|\ell - \ell'| \leq L \leq \ell + \ell'$ for the remainder of the section.

The value of $\left\langle A^{LM}_{\ell\ell'}(A^{LM}_{\ell\ell'})^* \right\rangle$ and the result follows.

We then obtain $\langle z^*_{\ell m} z_{\ell' m'} \rangle$ is used in the estimate below.

Recall that $fL \gg 1$) holds throughout.

It follows that $\hat{\Omega} = (\theta, \phi)$ by the previous lemma.

Suppose $I(\hat{\Omega})$ as claimed.

Note that $\mathcal{Z}^{P_\pm}_{\ell\ell' m m'}$ in what follows.

Here $X = E$ for the remainder of the section.

In this case $A^{LM}_{\ell,\ell'}$ and the result follows.

The value of I, Q, U is used in the estimate below.

We then obtain $\{\ell', \ell\}$ holds throughout.

Recall that ${}^{(ab)}\Gamma^E_{LM}$ by the previous lemma.

It follows that $LM\ell\ell'$ as claimed.

Suppose $c^{E,B}_{4M}$ in what follows.

Note that $F^{\times}(\hat{\Omega})$ for the remainder of the section.

Here $X \in \{I, P_{+}, P_{-}\}$ and the result follows.

In this case $O(\log(k + \Delta))$ is used in the estimate below.

The value of $(\Delta + 2)$ holds throughout.

We then obtain $L(G)$ by the previous lemma.

Recall that $\mathtt{out}_{v}$ as claimed.

It follows that $x \in D(C)$ in what follows.

Suppose $t_0 \geq t^*$ for the remainder of the section.

Note that C_0 and the result follows.

Here $\pi(G, C) = 0$ is used in the estimate below.

In this case $e \in D(C_t) \cap D(C_{t+1})$ holds throughout.

The value of $\mathtt{out}_{u,t-1}(v) = \perp$ by the previous lemma.

We then obtain $H = G_t(B)$ as claimed.

Recall that $e \in \overline{D}(C_{t-1}) \cap D(\hat{C}_{t-1})$ in what follows.

It follows that $\lambda_{t'}(e)$ for the remainder of the section.

Suppose $V(G) - D$ and the result follows.

Note that $C_t(v) = o$ is used in the estimate below.

Here $t - (t^* + \phi + 1)$ holds throughout.

In this case $V(H_{t'})$ by the previous lemma.

The value of $r_u(v)$ as claimed.

We then obtain $\mathrm{d}_H(u)$ in what follows.

Recall that $F_v^g = \{\{u, v\} \in F_v \mid \mathrm{d}_H(u) \leq \mathrm{d}_H(v)\}$ for the remainder of the section.

It follows that $b_{v,u}(i) = 1$ and the result follows.

Suppose c_e is used in the estimate below.

Note that $\rho_0 = c$ holds throughout.

Here $((1 + \epsilon)\Delta)$ by the previous lemma.

In this case $m_v = \perp$ as claimed.

The value of $isincidentonatleastonevertexin$ in what follows.

We then obtain $inducedbythenodesthatstartaphaseinround$ for the remainder of the section.

Recall that $asitmayhavebeenremovedduringtheroundinterval$ and the result follows.

It follows that $.Ifnode$ is used in the estimate below.

Suppose $)where$ holds throughout.

Note that $A \in \mathbb{R}^{n \times n}$ by the previous lemma.

Here $\boldsymbol{d} \in \mathbb{R}^m$ as claimed.

In this case $(0, 2)$ in what follows.

The value of $k = 0, 1, 2, \ldots$ for the remainder of the section.

We then obtain $\boldsymbol{x}^{(k)} \to \boldsymbol{x}^*$ and the result follows.

Recall that $\boldsymbol{x}^{(2)} = \mathbf{0}$ is used in the estimate below.

It follows that $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^m$ holds throughout.

Suppose $P = D^{-1}$ by the previous lemma.

Note that $x_i^{(k)}$ as claimed.

Here $\omega^{(0)} \in (0, 2)$ in what follows.

In this case $\omega^{(k)} \to 0$ for the remainder of the section.

The value of $\rho \in (0, 1)$ and the result follows.

We then obtain ε_ω is used in the estimate below.

Recall that $c_1 \in (0, 1)$ holds throughout.

It follows that $\omega^{(0)} = 1$ by the previous lemma.

Suppose $h^{(k+1)} = 2$ as claimed.

Note that k' in what follows.

Here $\overline{d}^{(l')} < -2$ for the remainder of the section.

In this case $\overline{s}_m^{(k-1)} > \overline{s}_m^{(k-2)}$ and the result follows.

The value of $\boldsymbol{x}_0$ is used in the estimate below.

We then obtain $\mathsf{sprandsym}$ holds throughout.

Recall that $\boldsymbol{b} = A\hat{\boldsymbol{x}} - \boldsymbol{y}$ by the previous lemma.

It follows that $\kappa = 10^4$ as claimed.

Suppose $\omega = 1.77...$ in what follows.

Note that $A \in \mathbb{R}^{10000 \times 10000}$ for the remainder of the section.

Here $\sigma = \min(\mathrm{diag}(A)) = 3030.3...$ and the result follows.

In this case $\omega = 0.1, 0.2, \ldots, 1.9$ is used in the estimate below.

The value of $\boldsymbol{x} \geq \mathbf{0}$ holds throughout.

We then obtain $\{\boldsymbol{x}^{(k)}\}_{k=0}^{\infty}$ by the previous lemma.

Recall that $\omega = 1.7$ as claimed.

It follows that (k, M) in what follows.

Suppose $\mathcal{Y} = \{0, 1\}$ for the remainder of the section.

Note that $k_N / N \to 0$ and the result follows.

Here $k = k_N$ is used in the estimate below.

In this case L_2 holds throughout.

The value of $\alpha > 1$ by the previous lemma.

We then obtain $d \leq 10$ as claimed.

Recall that $\mathcal{Y} \subseteq \mathbb{R}$ in what follows.

It follows that $k \leq N$ for the remainder of the section.

Suppose $x \in \operatorname{supp}(\mu)$ and the result follows.

Note that l_Y is used in the estimate below.

Here $1 - \delta$ holds throughout.

In this case $N \geq 1$ by the previous lemma.

The value of (β, B) as claimed.

We then obtain $M = M_o N^{\frac{2\alpha}{2\alpha+1}} (\log \frac{1}{\delta})^{\frac{1}{2\alpha+1}} \leq N$ in what follows.

Recall that $M = N^{\gamma}$ for the remainder of the section.

It follows that $m_1, \ldots, m_L$ and the result follows.

Suppose $\Theta(N^{\frac{2\alpha}{2\alpha+1} - \gamma})$ is used in the estimate below.

Note that $\mathbf{X} = X_{1:N} = \{X_i\}_{i=1}^{N}$ holds throughout.

Here (σ, S) by the previous lemma.

In this case $\zeta := \frac{\sigma}{d} \wedge 1$ as claimed.

The value of $\zeta = \frac{\sigma}{d}$ in what follows.

We then obtain $|\mathbf{X}_m| = n \to \infty$ for the remainder of the section.

Recall that $\mathbb{E}[\log U_{kn}^{\infty}(x)] = \Psi(k) - \log p(x)$ and the result follows.

It follows that $\phi_k^{(f)}(u)$ is used in the estimate below.

Suppose $\mathcal{L}^{-1}(F(p))(u)$ holds throughout.

Note that $\alpha \in \mathbb{R}$ by the previous lemma.

Here $(k, M, \frac{M}{2})$ as claimed.

In this case $(1, M, \frac{M}{2}, M)$ in what follows.

The value of $k_{\mathsf{base}} \in \{1, 2, 3, 4, 5\}$ for the remainder of the section.

We then obtain $\boldsymbol{F}(\boldsymbol{S}, \boldsymbol{T})$ and the result follows.

Recall that $\boldsymbol{B}_{id}$ is used in the estimate below.

It follows that $\boldsymbol{B}_{exp}$ holds throughout.

Suppose $\boldsymbol{\alpha} \in \mathbb{R}^{80}$ by the previous lemma.

Note that $\boldsymbol{\beta} \in \mathbb{R}^{64}$ as claimed.

Here $\boldsymbol{n}_i$ in what follows.

In this case $\boldsymbol{t}_{i}$ for the remainder of the section.

The value of Φ_{b} and the result follows.

We then obtain $\boldsymbol{p} \in \mathbb{R}^{6}$ is used in the estimate below.

Recall that $\boldsymbol{R} \in SO(3)$ holds throughout.

It follows that $\Pi : \mathbb{R}^{3} \to \mathbb{R}^{2}$ by the previous lemma.

Suppose $\boldsymbol{T}'$ as claimed.

Note that M_{k} in what follows.

Here $\boldsymbol{C}_{k} = \{(\boldsymbol{I}_{i}, \{l_{n}\}_{i})\}_{i=1}^{M_{k}}$ for the remainder of the section.

In this case $\boldsymbol{I}'$ and the result follows.

The value of ω_{n} is used in the estimate below.

We then obtain $\omega_{\boldsymbol{\alpha}} = 0.5$ holds throughout.

Recall that $\omega_{\boldsymbol{\beta}} = 2.0$ by the previous lemma.

It follows that $\lambda_{pho} = 1.92$ as claimed.

Suppose $\lambda_{lan} = 1.6e{-}3$ in what follows.

Note that $\boldsymbol{I}'(\hat{\boldsymbol{x}})$ for the remainder of the section.

Here $\boldsymbol{D}'$ and the result follows.

In this case $\boldsymbol{Z}_{\boldsymbol{S}}'$ is used in the estimate below.

The value of $\boldsymbol{Z}_{max} = \max(\boldsymbol{Z}_{\boldsymbol{S}_{i}})$ holds throughout.

We then obtain $K = 50k$ by the previous lemma.

Recall that $1/7$ as claimed.

It follows that $\boldsymbol{\alpha}_{gt}$ in what follows.

Suppose $\boldsymbol{\beta}_{gt}$ for the remainder of the section.

Note that $1.0e{-}4$ and the result follows.

Here $50k$ is used in the estimate below.

In this case $\mathcal{D}_{1}$ holds throughout.

The value of p^{2} by the previous lemma.

We then obtain $g^{1}, \ldots, g^{\mathsf{y}}$ as claimed.

Recall that $2\mathsf{d}$ in what follows.

It follows that $\iota_{\mathbb{C}^{d}}$ for the remainder of the section.

Suppose $\mathcal{B} \in \mathfrak{B}$ and the result follows.

Note that $\mathcal{A} \subset \mathcal{C} \subset [\mathsf{n}]$ is used in the estimate below.

Here $\mathcal{DEC}[\mathcal{A}]$ holds throughout.

In this case U_{D} by the previous lemma.

The value of $\in \mathbb{F}_q^{\times}$ as claimed.

We then obtain $R = GU_S \in \mathbb{F}_q^{\bar{\mathsf{n}}}$ in what follows.

Recall that $\bar{\mathsf{n}} \times \mathsf{n}$ for the remainder of the section.

It follows that $\mathsf{n} \geq \mathsf{r} > \mathsf{t} > 0$ and the result follows.

Suppose $2(\mathsf{r} - \mathsf{t})/\mathsf{n}$ is used in the estimate below.

Note that $, and$ holds throughout.

Here $, and a$ by the previous lemma.

In this case $can be considered as a stabilizer code with the stabilizer$ as claimed.

The value of $, the user makes the measurement given by the POVM$ in what follows.

We then obtain $, and sends the$ for the remainder of the section.

Recall that $, the state reduced density with respect to$ and the result follows.

It follows that $, the protocol does not have the matrix$ is used in the estimate below.

Suppose $-EAMDS code when it is$ holds throughout.

Note that $on the system$ by the previous lemma.

Here $is column orthogonal to$ as claimed.

In this case $-secure EASPIR protocol with rate$ in what follows.

The value of $, we have$ for the remainder of the section.

We then obtain $, and the second relation shows that the MMSP$ and the result follows.

Recall that $components by$ is used in the estimate below.

It follows that $.The case with$ holds throughout.

Suppose $, the relation holds for$ by the previous lemma.

Note that $\rho \to \infty$ as claimed.

Here $\rho = \rho_0$ in what follows.

In this case $\{\boldsymbol{\theta}_k\}$ for the remainder of the section.

The value of $|\mathcal{I}_k| = b$ and the result follows.

We then obtain $\mathcal{I} \leftarrow$ is used in the estimate below.

Recall that $g = 0$ holds throughout.

It follows that $0.5 < \gamma \leq 1$ by the previous lemma.

Suppose $m = 5c^2 + 2L^2r^2 + 2LGr$ as claimed.

Note that δ_0 in what follows.

Here $\mathbb{E}\|P_C(\boldsymbol{\theta}_k) - \boldsymbol{\theta}_*\|^2$ for the remainder of the section.

In this case ζ_k and the result follows.

The value of $P_{C_s}(\boldsymbol{\theta})$ is used in the estimate below.

We then obtain 20% holds throughout.

Recall that $\gamma = \frac{4}{3}$ by the previous lemma.

It follows that C_u as claimed.

Suppose $p = q = 64$ in what follows.

Note that $\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_*\|^2$ for the remainder of the section.

Here $\mathcal{O}(pb)$ and the result follows.

In this case $\rho_1 = 1 \times 10^{-2}$ is used in the estimate below.

The value of $\boldsymbol{\theta}_* \in C'$ holds throughout.

We then obtain $\|\nabla g(\boldsymbol{x}_k)\| \leq \|\nabla g(\boldsymbol{x}_{k-1})\|$ by the previous lemma.

Recall that $\sum_{i=1}^{+\infty} b_i < +\infty$ as claimed.

It follows that $k = 0, 1, ...$ in what follows.

Suppose $F[P_C(\boldsymbol{\theta}_{i_k - 1})] - F(\boldsymbol{\theta}_*) \to 0$ for the remainder of the section.

Note that $b_n = b_1 n^{-\beta}$ and the result follows.

Here $\beta = 1$ is used in the estimate below.

In this case $\sqrt{a + b} \leq \sqrt{a} + \sqrt{b}$ holds throughout.

The value of $L_\delta(\tilde{y}_i - \tilde{\boldsymbol{x}}_i^T \boldsymbol{\theta})$ by the previous lemma.

We then obtain $\|\boldsymbol{\theta}_{\text{true}}\|_2 = 2$ as claimed.

Recall that $\boldsymbol{X}_i \in \mathbb{R}^{p \times p}$ in what follows.

It follows that $Bernoulli(p_i)$ for the remainder of the section.

Suppose μ, Σ and the result follows.

Note that $\theta \in \Theta$ is used in the estimate below.

Here $g_\theta(x) := g(\theta, x)$ holds throughout.

In this case $\theta_0 \in \Theta$ by the previous lemma.

The value of $q = 1$ as claimed.

We then obtain $v \in S$ in what follows.

Recall that $W(q)$ for the remainder of the section.

It follows that $\Delta(v, X)$ and the result follows.

Suppose $g_\theta(y) := y^\top X \theta$ is used in the estimate below.

Note that $X > z_{1-\alpha}$ holds throughout.

Here $\alpha = 0.025$ by the previous lemma.

In this case $f''(\theta) \leq 1$ as claimed.

The value of $2^p - 1$ in what follows.

We then obtain $\{H_i\}_{i=1}^I$ for the remainder of the section.

Recall that $b_j = 1$ and the result follows.

It follows that $\hat{\lambda}_i(\alpha)$ is used in the estimate below.

Suppose $\mathcal{F}_\tau$ holds throughout.

Note that S_τ by the previous lemma.

Here θ_1 as claimed.

In this case (θ_3, θ_4) in what follows.

The value of $\text{Bern}(p_i)$ for the remainder of the section.

We then obtain λ^* and the result follows.

Recall that $\theta_3 = -1$ is used in the estimate below.

It follows that $(p_3 = 0)$ holds throughout.

Suppose $(\theta, \theta, \theta, -1)$ by the previous lemma.

Note that (μ, σ^2) as claimed.

Here L_X^2 in what follows.

In this case $\text{Beta}(\alpha, \beta)$ for the remainder of the section.

The value of $\hat{u}(\theta)$ and the result follows.

We then obtain $c_1, c_2, \ldots, c_n$. is used in the estimate below.

Recall that $h_k(x)$ holds throughout.

It follows that $\mu(x)$ by the previous lemma.

Suppose $Y = \min(T_1, T_2, \ldots, T_n)$ as claimed.

Note that c_j in what follows.

Here $\mathbf{T} = (T_1, \ldots, T_n)'$ for the remainder of the section.

In this case $S_{\mathbf{T}}(0, \cdots, 0) = 1$ and the result follows.

The value of $t_j \in (0, \infty)$ is used in the estimate below.

We then obtain $S_{\mathbf{T}}(t_1, \cdots, t_n)$ holds throughout.

Recall that T_i by the previous lemma.

It follows that $T_j < T_i$ as claimed.

Suppose $h(t \,|\, j) = -\frac{d}{dt} \log S(t \,|\, j) = f(t \,|\, j)/S(t \,|\, j)$ in what follows.

Note that $S_j(t_j)$ for the remainder of the section.

Here $i \neq j$, $j = 1, \ldots, n$ and the result follows.

In this case $\prod_{j=1}^{n} S_j(t_j)$ is used in the estimate below.

The value of c_1 holds throughout.

We then obtain $Y = \min(T_1, T_2)$ by the previous lemma.

Recall that g_2 as claimed.

It follows that $\frac{1}{b} \ln \frac{c}{a}$ in what follows.

Suppose $\frac{1}{b} \ln \frac{c}{a(1-kc)}$ for the remainder of the section.

Note that $\frac{c}{a_1 e^{-b_1 x} + a_2 e^{b_2 x} + c}$ and the result follows.

Here e_x is used in the estimate below.

In this case $g_2(x)$ holds throughout.

The value of M_{GM} by the previous lemma.

We then obtain e_0 as claimed.

Recall that $x > 0$ in what follows.

It follows that $b = 0.1$ for the remainder of the section.

Suppose $b = 0.85$ and the result follows.

Note that $a = 0.4$ is used in the estimate below.

Here $p(x) = c/\left(ae^{bx} + c\right)$ holds throughout.