The value of B_n/C_n, is used in the estimate below.

We then obtain $\tau = \theta/2\pi + 4\pi i/g^2$. holds throughout.

Recall that A_n; by the previous lemma.

It follows that A_μ: as claimed.

Suppose $x_2 + ix_3 = re^{i\theta}$, in what follows.

Note that $\mathcal{A} = A + i\phi$. for the remainder of the section.

Here n_k; and the result follows.

In this case $\lambda_m^{n_m}\lambda_{m-1}^{n_{m-1}}\cdots\lambda_1^{n_1}$: is used in the estimate below.

The value of (D_n), holds throughout.

We then obtain B_k. by the previous lemma.

Recall that $\lambda'' = 0$; as claimed.

It follows that r'_k: in what follows.

Suppose $l - k + \lambda_k$, for the remainder of the section.

Note that g_*. and the result follows.

Here $\frac{1}{2}(\sum_{k=i}^{m} n_k)$; is used in the estimate below.

In this case $l + 1$: holds throughout.

The value of S_{e1221}, by the previous lemma.

We then obtain ρ_{odd}. as claimed.

Recall that C_{16}; in what follows.

It follows that $(\rho_{\text{even}} + X_S^{-1}\rho_{\text{even}}\,;\rho_{\text{odd}} + Y_S\rho_{\text{odd}})_B$: for the remainder of the section.

Suppose $(\lambda_{even}\,;\rho_{odd})$, and the result follows.

Note that $L(C1)$. is used in the estimate below.

Here C_1; holds throughout.

In this case $L(D2)$: by the previous lemma.

The value of (b), as claimed.

We then obtain $c22$. in what follows.

Recall that IIB; for the remainder of the section.

It follows that $(\lambda, \rho)_C = (\lambda_{odd}, \rho_{even})_C$: and the result follows.

Suppose CB_{eo}, is used in the estimate below.

Note that $\mathcal{I} \in \mathbb{Z}$. holds throughout.

Here $(\psi(0), \psi'(0))$; by the previous lemma.

In this case $(h_{R,t})$: as claimed.

The value of $\mathcal{H}_1$, in what follows.

We then obtain $\mathbf{y} = (y, y')$. for the remainder of the section.

Recall that $\ell_2 \subset \ell_2^{\circ}$; and the result follows.

It follows that $J^* : \mathcal{H}_b \to \mathcal{H}_b$: is used in the estimate below.

Suppose $\ell_2(t)$, holds throughout.

Note that $C_0^{\infty}(\mathbb{R}^+, \mathbb{C}^n)$. by the previous lemma.

Here $(0, u') \in \ell_D$; as claimed.

In this case $\ell^+(E) \cap \ell^-(E) \neq \{0\}$: in what follows.

The value of $(h - E)\psi$, for the remainder of the section.

We then obtain $\ell(t)$. and the result follows.

Recall that $U_1(t)$; is used in the estimate below.

It follows that $P_1 x_j = x_j$: holds throughout.

Suppose $\sigma\left(h_t, \mathbb{T}^1\right)$, by the previous lemma.

Note that $(u_1, \cdots, u_j)$. as claimed.

Here $\ell_t^{\sharp}\{(\Theta x, \Pi x),\ x \in \mathbb{C}\}$; in what follows.

In this case $\mathbf{k} \in \mathbb{Z}^{d-1}$: for the remainder of the section.

The value of $L^2(\Gamma) = H^{s=0}(\gamma)$, and the result follows.

We then obtain $\chi(x)$. is used in the estimate below.

Recall that $(H - E)\psi = 0$; holds throughout.

It follows that $H^2(\Omega^-)$: by the previous lemma.

Suppose $R_{t,D} := (H_{t,D} - \Sigma)^{-1}$, as claimed.

Note that $\ell_{R,t}^{\pm}(E)$. in what follows.

Here $\widetilde{H}_t$; for the remainder of the section.

In this case $L^2(\mathbb{R}^+ \times \mathbb{R})$: and the result follows.

The value of $\gamma^D \psi_D = 0$, is used in the estimate below.

We then obtain $H^{-1/2}(\Gamma) \times H^{1/2}(\Gamma)$. holds throughout.

Recall that $G_{\Sigma} f$; by the previous lemma.

It follows that LSwHom(H): as claimed.

Suppose G^R, in what follows.

Note that $\phi : G' \to H$. for the remainder of the section.

Here (u, v); and the result follows.

In this case $\phi(v) \in L(v)$: is used in the estimate below.

The value of H_c, holds throughout.

We then obtain $\pi_S^{-1}(i) = \{(v, i) \mid v \in v(H)\}$. by the previous lemma.

Recall that $(u, 0)$; as claimed.

It follows that L_S: in what follows.

Suppose $u, v \in V$, for the remainder of the section.

Note that $x_i \sim y_i$. and the result follows.

Here $u_{i-1} \not\sim v_i$; is used in the estimate below.

In this case B_a: holds throughout.

The value of $V(H)^3$, by the previous lemma.

We then obtain x_3. as claimed.

Recall that $i \neq i'$; in what follows.

It follows that $i = \pi_B((i, v))$: for the remainder of the section.

Suppose $\phi : H^k \to H$, and the result follows.

Note that $(i, h, j)(i', h', j')$. is used in the estimate below.

Here $a', b' \in V$; holds throughout.

In this case (y, x): by the previous lemma.

The value of $x_a x_b$, as claimed.

We then obtain r_a. in what follows.

Recall that $x_a b$; for the remainder of the section.

It follows that r_b: and the result follows.

Suppose $X = \mathrm{maj}(x_1, x_2, x_3)$, is used in the estimate below.

Note that $(a_i, x_i) \sim (b_i, y_i)$. holds throughout.

Here $A \leq a_i$; by the previous lemma.

In this case (a_2, b_1, c_1): as claimed.

The value of $\phi(a_2, d, d) = a_2$, in what follows.

We then obtain $G \cong F$. for the remainder of the section.

Recall that $u \in U \setminus \{1\}$; and the result follows.

It follows that $v \in V(\Gamma)$: is used in the estimate below.

Suppose $\gamma_{n+1}(G) = [\gamma_n(G), G]$, holds throughout.

Note that kI_H. by the previous lemma.

Here $p(b, c) = b + c$; as claimed.

In this case $p(b, c) = b + c(t - 1)$: in what follows.

The value of $\widetilde{G} \to G$, for the remainder of the section.

We then obtain M_3. and the result follows.

Recall that $\mathcal{D}_{\mathbb{F}_p H}$; is used in the estimate below.

It follows that $\mathbb{F}_p I_G$: holds throughout.

Suppose $A = H_1 \cap H_2$, by the previous lemma.

Note that u_i. as claimed.

Here $\mathcal{D}_{\mathbb{F}_p[H]}$; in what follows.

In this case $c_k \in \mathbb{F}_p N$: for the remainder of the section.

The value of $\widetilde{H_1}$, and the result follows.

We then obtain $i \in \{1, 2\}$. is used in the estimate below.

Recall that u_2; holds throughout.

It follows that $u_2 = v_2^{n_2}$: by the previous lemma.

Suppose $r_{\mathrm{ab}}(\widetilde{H_1}) + r_{\mathrm{ab}}(\widetilde{H_2}) - 1$, as claimed.

Note that ϕ_i. in what follows.

Here $\widetilde{H_i} \cong H_i$; for the remainder of the section.

In this case $\widetilde{H}_{\widehat{p}}$: and the result follows.

The value of w_1, is used in the estimate below.

We then obtain $\widetilde{H}_{\widehat{p}} \coprod \mathbb{Z}_p$. holds throughout.

Recall that $d(\widetilde{H}_{\widehat{p}}) = d(\overline{H})$; by the previous lemma.

It follows that $q|_G$: as claimed.

Suppose $u \equiv (uv^{-1})^c \mod [U, U]$, in what follows.

Note that $\gamma_k W_{\widehat{p}}$. for the remainder of the section.

Here $\Gamma \setminus \{e\} = \Delta_1 \cup \Delta_2$; and the result follows.

In this case $\mathbb{Q}(\sqrt{1 - 2m^k})$: is used in the estimate below.

The value of $m \geq 3$,, holds throughout.

We then obtain J_K. by the previous lemma.

Recall that $n > 3$; as claimed.

It follows that $\mu = \sqrt{2}$: in what follows.

Suppose $k = p^r$, for the remainder of the section.

Note that $(n = 1)$. and the result follows.

Here $p_i \mid h_{(m,p_i)}$; is used in the estimate below.

In this case $i = 1$: holds throughout.

The value of $, where$, by the previous lemma.

We then obtain $, then.$ as claimed.

Recall that , $whichisacontradictionto$; in what follows.

It follows that $doesnotdivide$: for the remainder of the section.

Suppose $Then$, and the result follows.

Note that $where.$ is used in the estimate below.

Here , for; holds throughout.

In this case $.Henceweget$: by the previous lemma.

The value of $andhencewehave$, as claimed.

We then obtain $.Takingmoduloover4onbothsides, weget.$ in what follows.

Recall that $isco-primeto$; for the remainder of the section.

It follows that $isoddpositiveinteger, weget$: and the result follows.

Suppose $musthaveprimeidealswhosenormsare$, is used in the estimate below.

Note that $.Hencetheprimedecompositionof.$ holds throughout.

Here $.Considertheideal$; by the previous lemma.

In this case $forsomeoddprime$: as claimed.

The value of $dividesclassnumbersof$, in what follows.

We then obtain $betheclassnumbersof.$ for the remainder of the section.

Recall that $suchthat$; and the result follows.

It follows that $isaprimewhichdividesanoddinteger$: is used in the estimate below.

Suppose $H^3(G, \mathbb{T})$, holds throughout.

Note that $\omega_1 \otimes_{\text{stack}} \omega_2$. by the previous lemma.

Here $\text{Index}(\omega \circ \mu)$; as claimed.

In this case $\Gamma = \mathbb{Z}$: in what follows.

The value of $\mathcal{A}_\Gamma$, for the remainder of the section.

We then obtain $\tau(\Phi(I)) = \Phi(\tau(I))$. and the result follows.

Recall that $\text{Index}_{1d}^{\mathcal{A}_{L_0},U}(\phi) = \text{Index}^{\mathcal{A},U}(\omega)$; is used in the estimate below.

It follows that $\gamma_{0;1}^{\Psi}$: holds throughout.

Suppose $\alpha_\lambda(g) = \exp(2\pi ik/n)$, by the previous lemma.

Note that $\text{Index}_{1d\ \text{trans}}(\omega) = \text{Index}_{1d\ \text{trans}}(\omega \circ \gamma_{0;1}^{\Phi'})$. as claimed.

Here $x \in \text{Cone}_\sigma^W(\alpha_0, \theta)$; in what follows.

In this case $[\pi_0 \circ \alpha_0 \circ \Theta(A_L), v_R]$: for the remainder of the section.

The value of $\tau_{\mathcal{B}}$, and the result follows.

We then obtain $b_g^R \in \text{Cone}_R^W(\alpha_0, \theta)$. is used in the estimate below.

Recall that $h_{g} \in \mathrm{IAC}_{L}(\theta, \alpha_{0}, \Theta)$; holds throughout.

It follows that $W_{g,1}, u_{R,1}(g,h), v$: by the previous lemma.

Suppose $W_{g,1} \delta_{g}^{\sigma} W_{g,1}^{\dagger} \in \mathrm{IAC}_{\sigma}(\alpha_{0}, \theta, \Theta)$, as claimed.

Note that $n \in \mathbb{N}$. in what follows.

Here n_{0}; for the remainder of the section.

In this case $\mathrm{Index}(\omega \circ \gamma_{0;1}^{H}) = \mathrm{Index}(\omega)$: and the result follows.

The value of $a \in \mathrm{IAC}_{L}(\alpha_{0}, \theta, \Theta)$, is used in the estimate below.

We then obtain $x \in \mathrm{IAC}_{L \times R}^{W}(\alpha_{0}, \theta, \Theta)$. holds throughout.

Recall that $B_{\sigma} \in \mathcal{U}(\mathcal{A})$; by the previous lemma.

It follows that $\omega \circ \gamma_{0;1}^{H} \circ \tilde{\beta}_{g,2} = \omega \circ \gamma_{0;1}^{H}$: as claimed.

Suppose $\{1,2,3,4\}$, in what follows.

Note that $\pi_{0} = \bigotimes_{i=1}^{4} \pi_{0,i}$. for the remainder of the section.

Here $(\mathcal{H}_{0}, \pi_{0} \circ i_{V}, \Omega_{0})$; and the result follows.

In this case $B_{n,L}$: is used in the estimate below.

The value of $\Xi^{(s)}(Z,t)$, holds throughout.

We then obtain $X \subset U, X \subset L, X \subset R$. by the previous lemma.

Recall that $\mathcal{K} := \mathrm{age}(\mathcal{A})$; as claimed.

It follows that $\mathrm{age}(\mathcal{I})$: in what follows.

Suppose $\overline{i}' := (i_{k} : k < s) \in X^{s}$, for the remainder of the section.

Note that $\mathrm{Diag}_{\mathcal{A}}(\overline{x})$. and the result follows.

Here $\mathcal{A} \cong_{L} \mathcal{A}'$; is used in the estimate below.

In this case $\mathrm{qftp}^{\mathcal{A}}(\overline{a})$: holds throughout.

The value of $h : B \to C$, by the previous lemma.

We then obtain $L = \{E, <\}$. as claimed.

Recall that $A, B \subseteq \mathcal{M}$; in what follows.

It follows that $\alpha(e,x) = x$: for the remainder of the section.

Suppose $h : \mathcal{A} \to \mathcal{B}$, and the result follows.

Note that $Y \vDash \mathrm{EMtp}(X)$. is used in the estimate below.

Here $||\mathcal{I}||^{+}$; holds throughout.

In this case $\mathcal{I} = (\mathbb{Z}, p, s, <)$: by the previous lemma.

The value of $a_{i} \prec a_{j}$, as claimed.

We then obtain $s \neq t$. in what follows.

Recall that $a_{i} \neq a_{j}$; for the remainder of the section.

It follows that $(g(i) \mid i \in \mathcal{A})$: and the result follows.

Suppose $\eta, \nu \in \omega^{<\omega}$, is used in the estimate below.

Note that $\operatorname{age}(\mathcal{A}) = \operatorname{age}(\mathcal{I}_{\mathrm{eq}})$. holds throughout.

Here $B = \{b_i : i < \omega\}$; by the previous lemma.

In this case $(\mathcal{B}_{ba}, \prec)$.: as claimed.

The value of $,itfollowsimmediatelythat$, in what follows.

We then obtain $bestructuresandsuppose$. for the remainder of the section.

Recall that $andfixgenerators$,; and the result follows.

It follows that $.Ifthereexistsan$: is used in the estimate below.

Suppose $homogeneousforthecoloring$, holds throughout.

Note that $isthedesiredhomogeneouscopyof$. by the previous lemma.

Here $hasapreimagethatissomewherein$; as claimed.

In this case $with$: in what follows.

The value of $(X, \Delta) \to Z$, for the remainder of the section.

We then obtain (X, Δ). and the result follows.

Recall that $f \colon X' \to X$; is used in the estimate below.

It follows that $e \in \{1, -1\}$: holds throughout.

Suppose $h \colon W \to T'$, by the previous lemma.

Note that $D = \sum_i d_i D_i$. as claimed.

Here $Y_1 \to X$; in what follows.

In this case $g \colon V \to Y$: for the remainder of the section.

The value of B_Y, and the result follows.

We then obtain $K_{X'} + \Delta' = \psi^*(K_X + \Delta)$. is used in the estimate below.

Recall that $S^\nu \to S$; holds throughout.

It follows that $a(E, X, \Delta) < a(E, X', \Delta')$: by the previous lemma.

Suppose E', as claimed.

Note that $m\Delta = f_*(mf^*(K_X + \Delta)) - mK_X$. in what follows.

Here n''; for the remainder of the section.

In this case Σ': and the result follows.

The value of $V' = f^{-1}(V)$, is used in the estimate below.

We then obtain π'. holds throughout.

Recall that $\left(V', (\Delta' + t_0 A')|_{V'}\right)$; by the previous lemma.

It follows that $(K_{X'} + \Delta' + t_0 A')$: as claimed.

Suppose π_Y, in what follows.

Note that $\lfloor \Gamma \rfloor$. for the remainder of the section.

Here $K_{X'} + \Theta' = f^*(K_X + \Delta)$; and the result follows.

In this case $\mathrm{mult}_{Q_j}(\pi_Y^* D_j) \leq \beta$: is used in the estimate below.

The value of $(X, \Delta + tB)$, holds throughout.

We then obtain $m(K_{X'} + \Delta')$. by the previous lemma.

Recall that $G \sim IA$; as claimed.

It follows that $2n''eD$: in what follows.

Suppose $B' \geq 0$, for the remainder of the section.

Note that $\pi' \colon X' \dashrightarrow W$. and the result follows.

Here $\mathcal{F}(R, \mathcal{G})$; is used in the estimate below.

In this case $\mathcal{F}(R)$: holds throughout.

The value of S_0, by the previous lemma.

We then obtain $\mathcal{F}(\mathcal{R}, \mathcal{G})$. as claimed.

Recall that $P_{\mu\nu}$; in what follows.

It follows that $\tilde{b} = \frac{4}{3} q^2 \Lambda$: for the remainder of the section.

Suppose $E(r) = -A'(r)$, and the result follows.

Note that $r \to \infty$. is used in the estimate below.

Here $\mathcal{H}(P)$; holds throughout.

In this case $P <<$: by the previous lemma.

The value of $r \to 0$, as claimed.

We then obtain $r_h = (8Mq^2)^{1/3}$. in what follows.

Recall that (i, j, k); for the remainder of the section.

It follows that $T_H = \frac{\kappa}{2\pi}$: and the result follows.

Suppose $\beta = 1/T_H$, is used in the estimate below.

Note that $r_H <<$. holds throughout.

Here (C); by the previous lemma.

In this case ξ^μ: as claimed.

The value of A_E, in what follows.

We then obtain $n_+ = n_- = n \geq 1$. for the remainder of the section.

Recall that $\beta = \frac{4M\tilde{\Lambda}}{3}$; and the result follows.

It follows that $r^{1+p}/\sqrt{\eta - \beta r}$: is used in the estimate below.

Suppose $\delta = \tilde{\Lambda}/3$, holds throughout.

Note that $||R||^{2} \to \infty$. by the previous lemma.

Here $Z_{\pm} = R_{1} \pm R_{2}$; as claimed.

In this case $L^{2}\left(0, \infty \right)$: in what follows.

The value of c_{5}, for the remainder of the section.

We then obtain $\sigma = \sqrt{1 + 4\bar{\lambda}}/2$. and the result follows.

Recall that $6 \leq \bar{\lambda}^{2}$; is used in the estimate below.

It follows that Q_{1}, Q_{5}: holds throughout.

Suppose $a = 0, 1, ..., n - 1$, by the previous lemma.

Note that $\star_{3} \, dA = dH$. as claimed.

Here T^{4}; in what follows.

In this case $(n - 1)$: for the remainder of the section.

The value of k_{a}^{2}, and the result follows.

We then obtain Π_{ab}^{2}. is used in the estimate below.

Recall that q_{a}; holds throughout.

It follows that $\lambda = 10^{-5}$: by the previous lemma.

Suppose $[x_{1}, \cdots, x_{\mathbf{n_{0}}}]$, as claimed.

Note that $z = 0$. in what follows.

Here $\{k_{\bar{a}}^{0,1}, k_{0}^{2}\}$; for the remainder of the section.

In this case $Q_{\max} \leftarrow$: and the result follows.

The value of $\bar{r}_{a}^{i}$, is used in the estimate below.

We then obtain $\{r_{A}^{(p)}, \sigma_{A}^{(p)}\}$. holds throughout.

Recall that $\delta r_{A}^{(p)}$; by the previous lemma.

It follows that f': as claimed.

Suppose $2/d$, in what follows.

Note that $\mathcal{N}(0, 1)$. for the remainder of the section.

Here `generation_update`; and the result follows.

In this case $\sigma_{A} \to (\texttt{factor_random_update}) \, \sigma_{A}$: is used in the estimate below.

The value of 200, holds throughout.

We then obtain 1st. by the previous lemma.

Recall that -0.6923; as claimed.

It follows that -61: in what follows.

Suppose -48.5265, for the remainder of the section.

Note that $Q_{5} \simeq 410.$ and the result follows.

Here 1.1×10^{4}; is used in the estimate below.

In this case $(1 + 1)$: holds throughout.

The value of $\gamma \neq 0$, by the previous lemma.

We then obtain $Z_{n}.$ as claimed.

Recall that Z_{3}; in what follows.

It follows that $\mathbb{C}^{3}$: for the remainder of the section.

Suppose $2L^{2}$, and the result follows.

Note that $\hat{\Delta}_{v}.$ is used in the estimate below.

Here $\varepsilon_{v} = 0$; holds throughout.

In this case h_{v}: by the previous lemma.

The value of $\hat{N}_{f}$, as claimed.

We then obtain $h_{f}.$ in what follows.

Recall that B_{f}; for the remainder of the section.

It follows that $C_{<f,f^{\prime}>}$: and the result follows.

Suppose $\hat{N}_{<f,f^{\prime}>}$, is used in the estimate below.

Note that $-10L^{2}.$ holds throughout.

Here $a = 1, 2, 3$; by the previous lemma.

In this case $A^{(a)}$: as claimed.

The value of H_{int}, in what follows.

We then obtain $H_{v}.$ for the remainder of the section.

Recall that N_{i}; and the result follows.

It follows that $p_{\sigma} = \frac{1}{N_{\text{i}}}$: is used in the estimate below.

Suppose $\mathcal{D} = 4$, holds throughout.

Note that $A^{\prime (a)}_{v}.$ by the previous lemma.

Here $\hat{\Delta}^{\prime}_{v}$; as claimed.

In this case $h^{\prime}_{v}$: in what follows.

The value of $h^{\prime}_{f}$, for the remainder of the section.

We then obtain $\{\sqrt{3} + 1, 2, 1, 0, 1 - \sqrt{3}\}.$ and the result follows.

Recall that $N_{\text{FPL}} = \binom{2l}{l}$; is used in the estimate below.

It follows that $n \approx \sigma l$: holds throughout.

Suppose $\mathcal{C}_{x,y}$, by the previous lemma.

Note that T_c. as claimed.

Here $\theta/z = -1$; in what follows.

In this case $\rho_{xx} \sim \tau_{tr}^{-1} \sim T$: for the remainder of the section.

The value of $\cot \Theta_H \sim \tau_h^{-1} \sim T^2$, and the result follows.

We then obtain $\tilde{\sigma} \propto T^{-1}$. is used in the estimate below.

Recall that $\rho_{xx} \propto \alpha T$; holds throughout.

It follows that $3 + 1$: by the previous lemma.

Suppose S_{GR}, as claimed.

Note that $F = \mathrm{d}A$. in what follows.

Here ψ_I; for the remainder of the section.

In this case $z = 1$: and the result follows.

The value of A_t, is used in the estimate below.

We then obtain $T = U(1)/(4\pi)$. holds throughout.

Recall that $B = 0$; by the previous lemma.

It follows that k/μ: as claimed.

Suppose $B/\mu^2 = 1/10$, in what follows.

Note that x, y. for the remainder of the section.

Here ω_0; and the result follows.

In this case $1 < \beta \leq 2$: is used in the estimate below.

The value of $\sigma_0, \tilde{\sigma}, n$, holds throughout.

We then obtain $x_3 = 0$. by the previous lemma.

Recall that $x_1 \leq 1$; as claimed.

It follows that $x_1 = 1$: in what follows.

Suppose $\beta = x_3 + x_2 \leq 1$, for the remainder of the section.

Note that $T \gg W$. and the result follows.

Here $x_3 \neq 0$; is used in the estimate below.

In this case $\sqrt{a_1 T^2 + a_2 B^2}$: holds throughout.

The value of $|\psi_0\rangle$, by the previous lemma.

We then obtain $N = 10$. as claimed.

Recall that $N = 4, L = 11$; in what follows.

It follows that $S(\theta_1, \ldots, \theta_N)$: for the remainder of the section.

Suppose $\tau_{\mathrm{cr}} \approx 0.33137171$, and the result follows.

Note that $\ell \geq 2$. is used in the estimate below.

Here $\tau^{\mathrm{QSL}} > 0$; holds throughout.

In this case $\sigma^{+}_{k_1 \cdots k_N}$: by the previous lemma.

The value of $\kappa \pm \square_a$, as claimed.

We then obtain $U(1)$. in what follows.

Recall that $\left\{ e^{i 2\pi (s-1)/L},\ s = 1 \ldots, L \right\}$; for the remainder of the section.

It follows that S_L: and the result follows.

Suppose $\forall\ N \in \mathbb{N}$, is used in the estimate below.

Note that $\beta \to \infty$. holds throughout.

Here $S_{\mathrm{eff}}[\rho]$; by the previous lemma.

In this case $\gamma > \frac{1}{2}$: as claimed.

The value of $\gamma = \frac{1}{2}$, in what follows.

We then obtain $\tilde{x}_a = -x_a$. for the remainder of the section.

Recall that $\tau \to 0$; and the result follows.

It follows that $(\tau_{\mathrm{cr}} - \tau)$: is used in the estimate below.

Suppose $N = 8$, holds throughout.

Note that A_p. by the previous lemma.

Here $[0, N-1]$; as claimed.

In this case $\rho^{\mathscr{D}}_{\mathrm{unif.}}$: in what follows.

The value of $\varepsilon > 0$, for the remainder of the section.

We then obtain $\delta > 0$. and the result follows.

Recall that $\{P_j\}_{j \in \mathbb{N}}$; is used in the estimate below.

It follows that $|x| < 1$: holds throughout.

Suppose $e^{itH_{\mathrm{XY}}} |\psi_0\rangle$, by the previous lemma.

Note that h_{2n+1}. as claimed.

Here $T = 0$; in what follows.

In this case (a): for the remainder of the section.

The value of $G^{s_1}_{ij}, G^{s_2}_{ij}$, and the result follows.

We then obtain (s_1, s_2). is used in the estimate below.

Recall that Z_C; holds throughout.

It follows that $\det(g_C^{s_1, s_2})$: by the previous lemma.

Suppose $G^{s}_{ij}(\theta)$, as claimed.

Note that $\pi/3$. in what follows.

Here S_2; for the remainder of the section.

In this case N_{seed}: and the result follows.

The value of $\theta = 10$, is used in the estimate below.

We then obtain $\phi = 0.1$. holds throughout.

Recall that $\ln(L)$; by the previous lemma.

It follows that $\Delta_\tau = 0.5$: as claimed.

Suppose 0.95(5), in what follows.

Note that f + b. for the remainder of the section.

Here $B^s(\tau_2, \tau_1)$; and the result follows.

In this case $\gamma_i = (1/(1 - e^{-2i\alpha\tilde{\sigma}_i}) - G^s_{i,i}(\tau))^{-1}$: is used in the estimate below.

The value of $\boldsymbol{v}_i$, holds throughout.

We then obtain $\mathcal{O}(N_C^2)$. by the previous lemma.

Recall that g^{s_1,s_2}_{C+i}; as claimed.

It follows that $\tau = \theta$: in what follows.

Suppose $G^s(\theta)$, for the remainder of the section.

Note that $\lambda = 0.0006$. and the result follows.

Here $\approx 10^{-6}$; is used in the estimate below.

In this case $L_{\max} = 60$: holds throughout.

The value of $L = 6$, by the previous lemma.

We then obtain $L = 3, 6, 9, 12, 15, 18$. as claimed.

Recall that $N_\lambda/(N_\lambda + 1)$; in what follows.

It follows that $L = 3, 6, 9, ..., 60$: for the remainder of the section.

Suppose *, and the result follows.

Note that $\mathcal{U}_h$. is used in the estimate below.

Here $T_1 G$; holds throughout.

In this case $U_g U_h = U_{gh}$: by the previous lemma.

The value of $\Delta = H^2$, as claimed.

We then obtain $\psi \in \mathcal{H}$. in what follows.

Recall that $t \in (-\tau, \tau)$; for the remainder of the section.

It follows that U, V: and the result follows.

Suppose $s : G \to \tilde{G}$, is used in the estimate below.

Note that $\mathcal{H}_j = \mathbb{C}^{2j+1}$. holds throughout.

Here $\mathrm{SO}(2) \cong \mathrm{U}(1)$; by the previous lemma.

In this case $-\mathrm{i}A(\mathfrak{so}(2m)) \subset \mathfrak{u}(\mathcal{H})$: as claimed.

The value of B_{jk}, in what follows.

We then obtain $\mathbb{H}_m = \mathbb{R}^{2m+1} = \mathbb{R}^{2m} \times \mathbb{R}$. for the remainder of the section.

Recall that $X \in \mathfrak{sp}(2m, \mathbb{R})$; and the result follows.

It follows that $n \in \mathbb{N}_0^m$: is used in the estimate below.

Suppose $\mathcal{O}(1/L)$, holds throughout.

Note that $\sigma_z = \operatorname{diag}(1, -1)$. by the previous lemma.

Here $\Delta^{(n)}$; as claimed.

In this case $\mathrm{O}(1,3)$: in what follows.

The value of $p \in \mathcal{M}_m$, for the remainder of the section.

We then obtain $M_{0i} = -M_{i0} = K_i$. and the result follows.

Recall that $d(g_n, 1) \to 0$; is used in the estimate below.

It follows that $\psi = \chi_W$: holds throughout.

Suppose $[0, 1]$, by the previous lemma.

Note that $PX_k = p_k X_k$. as claimed.

Here $\dim(\mathcal{H}) \geq 3$; in what follows.

In this case $\mathcal{P} : \omega(\rho, \psi, p) \mapsto \omega(\rho, 0, p)$: for the remainder of the section.

The value of $\tilde{d}(\tilde{g}, 1_{\tilde{G}}) \geq d(g, 1_G)$, and the result follows.

We then obtain E_{jk}. is used in the estimate below.

Recall that $t_g \sim 10^{-8} s$; holds throughout.

It follows that U_g: by the previous lemma.

Suppose $\mathrm{dW}_{k,s}$, as claimed.

Note that ϵ^2. in what follows.

Here $(-)$; for the remainder of the section.

In this case $2N^2 - 1$: and the result follows.

The value of $\Lambda(\theta, \phi)$, is used in the estimate below.

We then obtain $s = t/t_g$. holds throughout.

Recall that T_1; by the previous lemma.

It follows that $\gamma_z = t_g(2T_1 - T_2)/4T_1T_2$: as claimed.

Suppose $\epsilon = \sqrt{\lambda}$, in what follows.

Note that (k). for the remainder of the section.

Here $\Delta\mathcal{H}_{\sigma}^{\mathrm{ng}}$; and the result follows.

In this case ibm_oslo: is used in the estimate below.

The value of σ_{kk}, holds throughout.

We then obtain $\mathcal{H}^{\mathrm{ng}}$. by the previous lemma.

Recall that $\theta = \pi$; as claimed.

It follows that $\mathcal{N} = 15000$: in what follows.

Suppose $\sim 10^{-3} - 10^{-2}$, for the remainder of the section.

Note that 0.25. and the result follows.

Here $\sim 10^{-2}$; is used in the estimate below.

In this case $\mathcal{H}_{\chi}^{\mathrm{ng}} = \mathcal{H}(\rho^{\mathrm{ng}}, \chi)$: holds throughout.

The value of $\Delta\mathcal{H}_{\chi}^{\mathrm{ng}} < \Delta\mathcal{H}_{\chi}^{\mathrm{ibm}}$, by the previous lemma.

We then obtain $n = 9$. as claimed.

Recall that $\mathrm{R}_{xy}(\phi) = \cos(\phi)\mathrm{X} + \sin(\phi)\mathrm{Y}$; in what follows.

It follows that $\mathrm{N}(\theta, \phi)$: for the remainder of the section.

Suppose $\mathrm{U}(\theta, \phi)$, and the result follows.

Note that $\leq N_s$. is used in the estimate below.

Here $50\% - 60\%$; holds throughout.

In this case $\Delta V/V$: by the previous lemma.

The value of $\Delta H = H_f - H_i$, as claimed.

We then obtain H_i. in what follows.

Recall that μ_0; for the remainder of the section.

It follows that J_{ij}: and the result follows.

Suppose ΔS_L, is used in the estimate below.

Note that $\mathbf{e}_i$. holds throughout.

Here $\mathbf{e}_j$; by the previous lemma.

In this case R_{ij}: as claimed.

The value of R_{ij}/a, in what follows.

We then obtain μ_B. for the remainder of the section.

Recall that T_C; and the result follows.

It follows that $M(T) = (1 - \frac{T}{T_C})^{\beta}$: is used in the estimate below.

Suppose $J(r_{ij})$, holds throughout.

Note that r_{ij}. by the previous lemma.

Here $\boldsymbol{s}_i$; as claimed.

In this case $\mathbf{p}_i$: in what follows.

The value of $\boldsymbol{r}_i$, for the remainder of the section.

We then obtain ω_i. and the result follows.

Recall that $n_{\text{qubits}} = \log_2 \mathcal{C}$; is used in the estimate below.

It follows that S_P: holds throughout.

Suppose $a_{\{n_{B_i}\}_k} \neq 0$, by the previous lemma.

Note that $i, j \in 1 \dots N_o$. as claimed.

Here $\sum_i \tilde{\lambda}_i = N_o - N_p$; in what follows.

In this case $S_c \leq \max\{-\ln \nu, -\ln(1-\nu)\}$: for the remainder of the section.

The value of $\alpha_g = 2$, and the result follows.

We then obtain $N_o = 16$. is used in the estimate below.

Recall that α_g; holds throughout.

It follows that S_{opt}: by the previous lemma.

Suppose $U/t, V/t \approx 4$, as claimed.

Note that $k \in \{1, 2, \dots Q\}$. in what follows.

Here $\epsilon, \delta > 0$; for the remainder of the section.

In this case $n_{\text{bit}} = \log_2 2^{nH(P_k)}$: and the result follows.

The value of $n_1 < n_2$, is used in the estimate below.

We then obtain $n > n'$. holds throughout.

Recall that $\sigma = C^{\beta}$; by the previous lemma.

It follows that $\sigma(n)$: as claimed.

Suppose $\bar{n}_{B_i}$, in what follows.

Note that $S_c^h \leq S_P$. for the remainder of the section.

Here S_{CUE}; and the result follows.

In this case $P_j^2 = U_{jk} U_{jl}^* U_{jm} U_{jn}^* a_k a_l^* a_m a_n^*$: is used in the estimate below.

The value of $D_{2,B} = S_{P_B} / \log(Q)$, holds throughout.

We then obtain $S_P = \alpha N_p S_c$. by the previous lemma.

Recall that $\alpha \approx 0.8$; as claimed.

It follows that $\langle x \rangle = 2(1-\nu) N_p$: in what follows.

Suppose $k_1 + q$, for the remainder of the section.

Note that x_{fs}. and the result follows.

Here $N_p \gamma$; is used in the estimate below.

In this case $p \to -p$: holds throughout.

The value of $1 + 1$, by the previous lemma.

We then obtain $F_{\mu\nu}(x, t)$. as claimed.

Recall that $\beta = \sqrt{4\pi}$; in what follows.

It follows that $E_\theta = \frac{e}{2\pi}\theta$: for the remainder of the section.

Suppose $m_q \equiv \mathcal{E}(p = 0)$, and the result follows.

Note that $\theta = \pi - \varepsilon.$ is used in the estimate below.

Here $\pi_1 \pi_2$; holds throughout.

In this case 0.5: by the previous lemma.

The value of π_1, π_2, as claimed.

We then obtain $\pi_1.$ in what follows.

Recall that $\beta \neq \sqrt{4\pi}$; for the remainder of the section.

It follows that $A_1(x) = 0$: and the result follows.

Suppose $a, b = 1, 2$, is used in the estimate below.

Note that $\frac{e\beta}{2\pi}\phi = E + \frac{e\theta}{2\pi} \equiv E_T.$ holds throughout.

Here $\{b(k), b^\dagger(k')\} = \{c(k), c^\dagger(k')\} = \delta(k - k')$; by the previous lemma.

In this case $\hat{p}|p\rangle = p|p\rangle$: as claimed.

The value of $p/m \ll 1$, in what follows.

We then obtain $m_q.$ for the remainder of the section.

Recall that $m_\pi / m_q > 2$; and the result follows.

It follows that $I^i_C = C\ddot{\phi}_i$: is used in the estimate below.

Suppose $\phi^i_J + \phi^i_C = \Phi_{\text{ext}}$, holds throughout.

Note that $\omega_{k,j}.$ by the previous lemma.

Here $\chi = 1$; as claimed.

In this case $\lambda(\Phi_{\text{ext}})$: in what follows.

The value of A_L, for the remainder of the section.

We then obtain $\varepsilon = 0.$ and the result follows.

Recall that c_p; is used in the estimate below.

It follows that $n \in [i + 1, i + N]$: holds throughout.

Suppose $D' = 100$, by the previous lemma.

Note that $n = 2.$ as claimed.

Here $SU(2)$; in what follows.

In this case $\mathbf{P}(\omega, \vec{v})$: for the remainder of the section.

The value of $\hat{\mathbf{e}}_1$, and the result follows.

We then obtain $(\sin 2\theta_V,\ 0,\ -\cos 2\theta_V).$ is used in the estimate below.

Recall that $v^\rho(\vec{v}) = (1, \vec{v})$; holds throughout.

It follows that $\text{diag}(+1,\ -1,\ -1,\ -1)$: by the previous lemma.

Suppose $\omega > 0$, as claimed.

Note that μ^{-1}. in what follows.

Here $\Omega_i > 0$; for the remainder of the section.

In this case v^z: and the result follows.

The value of $u = \omega$, is used in the estimate below.

We then obtain $\mathbf{Y} = -\mu \mathbf{J}^z$. holds throughout.

Recall that $(0, 0, 1)$; by the previous lemma.

It follows that Ω_i: as claimed.

Suppose $\hat{\mathbf{e}}_3^{\mathrm{II}} = \hat{\mathbf{e}}_3$, in what follows.

Note that $\hat{\mathbf{e}}_3^{\mathrm{II}} = \cos\Theta \hat{\mathbf{e}}_3^{\mathrm{III}} - \sin\Theta \hat{\mathbf{e}}_1^{\mathrm{III}}$. for the remainder of the section.

Here $\Theta = \pi$; and the result follows.

In this case $\partial_t \Theta = \Omega_i \sin\Theta$: is used in the estimate below.

The value of $\mathsf{O}^{\mathrm{II}} \ll 1$, holds throughout.

We then obtain $P_3^{\mathrm{III}} = \cos\Theta'$. by the previous lemma.

Recall that $\lambda^\rho = (\mu,\, 0,\, 0,\, 0.5\mu)$; as claimed.

It follows that $v^x = v^y = 0$: in what follows.

Suppose $5\pi/6$,, for the remainder of the section.

Note that $\times (1 + 0.5 v^z)$. and the result follows.

Here $\Delta \vec{v}$; is used in the estimate below.

In this case $|\mathsf{Q}_{v^x, v^y}|$: holds throughout.

The value of v_x, by the previous lemma.

We then obtain J_3^z. as claimed.

Recall that $\mathbf{H}_{\mathrm{vac}}$; in what follows.

It follows that $\mathrm{Im}(\mathsf{C}\,\mathsf{D}^*) = 0$: for the remainder of the section.

Suppose σ_8, and the result follows.

Note that ρ_0. is used in the estimate below.

Here R_{200}; holds throughout.

In this case $c_{200} = R_{200}/r_s$: by the previous lemma.

The value of $X(\hat{\mathbf{n}})$, as claimed.

We then obtain $\boldsymbol{\alpha}(\hat{\mathbf{n}}) = \boldsymbol{\nabla}_{\hat{\mathbf{n}}} \phi(\hat{\mathbf{n}})$. in what follows.

Recall that $\Sigma_{\mathrm{cl}}(r)$; for the remainder of the section.

It follows that $d_{A,\mathrm{cl}}$: and the result follows.

Suppose $x = r/r_s = \theta/\theta_s$, is used in the estimate below.

Note that κ_t. holds throughout.

Here $\boldsymbol{\ell} = (\ell_x, \ell_y)$; by the previous lemma.

In this case $z = 0.7$: as claimed.

The value of θ_s, in what follows.

We then obtain $N_L^{(1)}$. for the remainder of the section.

Recall that $L = 5000$; and the result follows.

It follows that $\ell_{\mathrm{max}}^{\mathrm{CMB}} = 4000$: is used in the estimate below.

Suppose $L \sim 1100$, holds throughout.

Note that 1000. by the previous lemma.

Here $100 \leq L \leq 5000$; as claimed.

In this case $M = 4, 7$: in what follows.

The value of $M_{200} = 4 \times 10^{14} M_{\odot}, z = 0.7$, for the remainder of the section.

We then obtain $\hat{\kappa}_0$. and the result follows.

Recall that $\hat{\kappa}_L$; is used in the estimate below.

It follows that $\langle \kappa_0 \rangle$: holds throughout.

Suppose 13%, by the previous lemma.

Note that x_{max}. as claimed.

Here $\chi(z) = \int_0^z dz' \frac{1}{H(z')} = (1+z) d_A(z)$; in what follows.

In this case $\mathrm{Ci}(x)$: for the remainder of the section.

The value of $s = 0, t = 1$, and the result follows.

We then obtain $\langle g(\hat{n}) \rangle_{\text{fixed deflection}}$. is used in the estimate below.

Recall that m_c; holds throughout.

It follows that $\psi_C(x) = \gamma_2 \gamma_4 \, \bar{\psi}^T(x)$: by the previous lemma.

Suppose $\bar{\Delta}_5$, as claimed.

Note that $\mu_u + 2\mu_d$. in what follows.

Here $r - g$; for the remainder of the section.

In this case P_{lep}: and the result follows.

The value of μ_{ν_e}, is used in the estimate below.

We then obtain $m_c = 2.3$. holds throughout.

Recall that $\eta_V = G_V / G_S$; by the previous lemma.

It follows that $\eta_V = 0.5$: as claimed.

Suppose μ_B^c, in what follows.

Note that $\mu_B = 850$. for the remainder of the section.

Here $\eta_V = \eta_D = 0$; and the result follows.

In this case $\eta_V = 0.9$: is used in the estimate below.

The value of $n_B = \partial P / \partial \mu_B$, holds throughout.

We then obtain $n_f = \sum_c n_{fc}$. by the previous lemma.

Recall that $g_{\omega B}(n)$; as claimed.

It follows that b_τ: in what follows.

Suppose I_{3B}, for the remainder of the section.

Note that E_B^*. and the result follows.

Here $B = N, P$; is used in the estimate below.

In this case m_e: holds throughout.

The value of $P(r = R) = 0$, by the previous lemma.

We then obtain $\mu_{\nu_e}/T = 2$. as claimed.

Recall that $\mu_{\nu_e} = \alpha T$; in what follows.

It follows that $2.3 M_\odot$: for the remainder of the section.

Suppose $T = 30$, and the result follows.

Note that $\bar{\Delta}_0$. is used in the estimate below.

Here n_e; holds throughout.

In this case $q \in Q$: by the previous lemma.

The value of $r \in R$, as claimed.

We then obtain $\mathrm{NN}_2(1; R) = \{0, 1, 2\} = \mathrm{NN}_3(1; R)$. in what follows.

Recall that $t \in \mathbb{R}_+$; for the remainder of the section.

It follows that $c(R) = n + 1$: and the result follows.

Suppose $R \subset \mathbb{R}^n$, is used in the estimate below.

Note that $O\big(c_m(R)^{O(1)} \cdot |R| \log(\Delta(R))\big)$. holds throughout.

Here $\mathcal{S}_i(p, \mathcal{T}(R))$; by the previous lemma.

In this case $p \in \mathcal{T}(R)$: as claimed.

The value of $t > \delta$, in what follows.

We then obtain $r \in \mathbb{R}_+$. for the remainder of the section.

Recall that $H(\mathcal{T}(R)) = \{i \mid C_{i-1} \neq C_i\} \cup \{l_{\max}, l_{\min}\}$; and the result follows.

It follows that $l(a_{m-1}) \leq i - 1$: is used in the estimate below.

Suppose $j = l_{\min}(\mathcal{T}(R)) - 1$, holds throughout.

Note that $W_1 = \{r\}$. by the previous lemma.

Here $i \leftarrow -\infty$; as claimed.

In this case $\eta(i) \leftarrow i$: in what follows.

The value of $q \in X \setminus W$, for the remainder of the section.

We then obtain $i \in S$. and the result follows.

Recall that $O(c(R)^2 \log(|R|))$; is used in the estimate below.

It follows that $\varrho \in L(\mathcal{T}(R), q)$: holds throughout.

Suppose $R_{l_{\min}}$, by the previous lemma.

Note that $\psi_k(t)$. as claimed.

Here q_2; in what follows.

In this case p_{m-2}: for the remainder of the section.

The value of $m - 2$, and the result follows.

We then obtain $p_{i \cdot m-1}$. is used in the estimate below.

Recall that $i = m, j = m - 1$; holds throughout.

It follows that $i = 3, j = 2$: by the previous lemma.

Suppose $\langle N, S, A, T, \{R_i\}, \Omega, \{O_i\} \rangle$, as claimed.

Note that $s^t = \langle s^{1,t}, ..., s^{k,t} \rangle$. in what follows.

Here $T(s^{t+1}|s^t, a^t)$; for the remainder of the section.

In this case $O_i(o_i^t|s^t)$: and the result follows.

The value of $\pi_i(a_i^t|h_i^t)$, is used in the estimate below.

We then obtain $X_i = \{X_i^1, ..., X_i^m\} \subseteq S$. holds throughout.

Recall that $O_i(o_i^t|s^t) = \dot{O}_i(o_i^t|x_i^t)$; by the previous lemma.

It follows that $\langle s^t, h_{-i}^t \rangle$: as claimed.

Suppose x_i, in what follows.

Note that $x_i \in X_i = \{X_i^1, X_i^2\}$. for the remainder of the section.

Here Y_i; and the result follows.

In this case a_j^t: is used in the estimate below.

The value of $x_i \in \times_{j \in |X_i|} X_i^j$, holds throughout.

We then obtain $I_i(u_i^t|l_i^t)$. by the previous lemma.

Recall that y_t; as claimed.

It follows that θ_i: in what follows.

Suppose $I_i(u_i^t|l_i^t) \in \Psi_i$, for the remainder of the section.

Note that $\Psi = \times_{i \in N} \Psi_i$. and the result follows.

Here a_{-i}; is used in the estimate below.

In this case M_i^1: holds throughout.

The value of $Q^{\pi_i}_{M_i^2}$, by the previous lemma.

We then obtain $\bar{R} = ||R||_\infty$. as claimed.

Recall that $\{I_{\theta_i}\}_{i \in N}$; in what follows.

It follows that 5×5: for the remainder of the section.

Suppose $F = 4$, and the result follows.

Note that $F = 1$. is used in the estimate below.

Here $P^2(u_i^{t-1} y_i^{t-1}, h_{-i}^{t-1} | l^t)$; holds throughout.

In this case $P^2(h_i^{t+1} | h_i^t, a_i^t)$: by the previous lemma.

The value of $\hat{a}_i^t \neq \bar{a}_i^t$, as claimed.

We then obtain $\hat{I}_{\theta_m}$. in what follows.

Recall that $S(\omega)$; for the remainder of the section.

It follows that $b, 3$: and the result follows.

Suppose $\hat{Y} : \Omega \to \{0, 1\}$, is used in the estimate below.

Note that $.Soeither$. holds throughout.

Here $isdependenton$; by the previous lemma.

In this case $overhardlabelstoinferthevalueof$: as claimed.

The value of $.Itisusedtoevaluatehowwell$, in what follows.

We then obtain $.Wedefinerandomvariables$. for the remainder of the section.

Recall that $isarandomvariable$:; and the result follows.

It follows that $), whether$: is used in the estimate below.

Suppose $.Forthisattacktowork$,, holds throughout.

Note that $isincludedintheinputdatarecordsand$. by the previous lemma.

Here $.Itmakesiteasiertoextract$; as claimed.

In this case $, classifierof$: in what follows.

The value of $issplitintotwodisjointdatasets$:, for the remainder of the section.

We then obtain $.Theseareusedtocreatethestatistic$. and the result follows.

Recall that $using$; is used in the estimate below.

It follows that $)curveasafunctionwhichmapseach$: holds throughout.

Suppose $shouldnotbeabletoinfer$, by the previous lemma.

Note that $thenforallattackmodel$. as claimed.

Here $.Let$; in what follows.

In this case $fortraining$: for the remainder of the section.

The value of $and evaluate iton 634 unseen data records in$, and the result follows.

We then obtain $and evaluate iton 1,200 unseen data records in.$ is used in the estimate below.

Recall that $and target model's$; holds throughout.

It follows that $using hard labels against a$: by the previous lemma.

Suppose $issuch that$, as claimed.

Note that $.We then say that.$ in what follows.

Here $we call$; for the remainder of the section.

In this case the: and the result follows.

The value of $e_k \doteq (\mathbf{x}_k, t_k, p_k)$, is used in the estimate below.

We then obtain $\mathbf{x}_k, t_k.$ holds throughout.

Recall that $\mathbf{x}'_k = \mathbf{W}(\mathbf{x}_k, t_k; \boldsymbol{\theta})$; by the previous lemma.

It follows that e'_k: as claimed.

Suppose $\epsilon = 1$, in what follows.

Note that $\mathcal{D}(\mathcal{E}, \boldsymbol{\theta})$. for the remainder of the section.

Here $\mathbf{f} \doteq \partial \mathbf{W}(\mathbf{x}, t; \boldsymbol{\theta})/\partial t$; and the result follows.

In this case $t_{\text{ref}} = t_1$: is used in the estimate below.

The value of $t \in [t_1, t_{N_e}]$, holds throughout.

We then obtain $\Delta t = 0.$ by the previous lemma.

Recall that $\mathcal{R} = 0$; as claimed.

It follows that $\mathcal{R} < 0$: in what follows.

Suppose $\boldsymbol{\theta} \equiv (v_x, v_y)^\top$, for the remainder of the section.

Note that $\mathcal{R} \equiv \int_0^1 \frac{d|\mathtt{J}|}{d\Delta t} = 0.$ and the result follows.

Here $t \in [0, 1]$; is used in the estimate below.

In this case $\mathtt{R}(\boldsymbol{\omega} \Delta t)$: holds throughout.

The value of $\mathcal{R}_{\mathbf{x}(0)}$, by the previous lemma.

We then obtain $\boldsymbol{\omega} = (0, 0, \omega_z)^\top.$ as claimed.

Recall that 346×260; in what follows.

It follows that $N = \{3, 10, 20\}$: for the remainder of the section.

Suppose $\lambda = 1.0$, and the result follows.

Note that $\mathbf{v}(\mathbf{x}).$ is used in the estimate below.

Here 2×6; holds throughout.

In this case $\mathbf{V} \equiv (0, 0, v_z)^\top$: by the previous lemma.

The value of $\mathbf{x}'_k = (1 - v_z/Z(\mathbf{x}))t_k.$, as claimed.

We then obtain $\mathbf{x}^{h}(t) = (x(t), y(t), 1)^{\top}$. in what follows.

Recall that 3×3; for the remainder of the section.

It follows that $\mathbf{e}_3^{\top} \mathtt{H}_{t,t+\Delta t} = (0, 0, 1)$: and the result follows.

Suppose $\omega_z = 0, \mathbf{v} = \mathbf{0}$, is used in the estimate below.

Note that $|V|$. holds throughout.

Here $V = \{have, a, great, day\}$; by the previous lemma.

In this case $\{1, 0, 0, 0\}$: as claimed.

The value of V, V^T, in what follows.

We then obtain V^T. for the remainder of the section.

Recall that $n - 1$; and the result follows.

It follows that w_i: is used in the estimate below.

Suppose w_{i+j}, holds throughout.

Note that $P(w_i)$. by the previous lemma.

Here $f(w_i)$; as claimed.

In this case b_p: in what follows.

The value of X_{pq}, for the remainder of the section.

We then obtain w_q. and the result follows.

Recall that x_{max}; is used in the estimate below.

It follows that 0.75: holds throughout.

Suppose $f(0) = 0$, by the previous lemma.

Note that $f(0)$. as claimed.

Here v_w; in what follows.

In this case $n = 3$: for the remainder of the section.

The value of $\mathcal{M} \subset \{1, ..., M\}$, and the result follows.

We then obtain $(t_1, ..., t_{i-1})$. is used in the estimate below.

Recall that t_{i+1}; holds throughout.

It follows that $(t_{i+1}, t_{i+2}, ..., t_N)$: by the previous lemma.

Suppose Θ_x, as claimed.

Note that Θ_s. in what follows.

Here $2L + 1$; for the remainder of the section.

In this case $h_{i,j}^{L} M$: and the result follows.

The value of $x_i^{LM} = h_{i,0}^{LM}$, is used in the estimate below.

We then obtain $\left(SentenceA, SentenceB \right)$. holds throughout.

Recall that $[SEP]$; by the previous lemma.

It follows that $(X_t)_{t\geq 0}$: as claimed.

Suppose $\Delta_n := t_{i+1} - t_i$, in what follows.

Note that X_{Δ_n}. for the remainder of the section.

Here $(b_l)_{0\leq l\leq 3}$; and the result follows.

In this case $(\tilde{C}, \tilde{\rho})$: is used in the estimate below.

The value of λ_0, holds throughout.

We then obtain $\pi_{|t-s|}$. by the previous lemma.

Recall that $n \geq 1$; as claimed.

It follows that $d = 1$: in what follows.

Suppose $P_t(\psi_\xi)(x) = \int_{\mathbb{R}} p_t(x,y)\psi_\xi(y)dy$, for the remainder of the section.

Note that $\|\psi\|_\infty \leq 1$. and the result follows.

Here $|b(0)| = 0 \leq b_0$; is used in the estimate below.

In this case $M_n h_n^3 \geq 1/\alpha_0$: holds throughout.

The value of $k \geq 1$, by the previous lemma.

We then obtain $(x, y, \epsilon) \mapsto p^\epsilon_{\Delta_n}(x, y)$. as claimed.

Recall that $h_n = n^{-1/(1+2\beta)}$; in what follows.

It follows that $(B'_n)_n$: for the remainder of the section.

Suppose $b \equiv 0$, and the result follows.

Note that $\dot{X}^\epsilon$. is used in the estimate below.

Here $y > 2h_n$; holds throughout.

In this case $r' > 0$: by the previous lemma.

The value of $|\partial_x A_\epsilon(x, v)| + |\partial_x B(x, v)| \leq c(1 + |v|)$, as claimed.

We then obtain $L(\int_0^s u_r dB_r) = \int_0^s L(u_r)dB_r - \int_0^s u_r dB_r$. in what follows.

Recall that $\|n_{h_n}\|_\infty \leq \frac{c}{M_n}$; for the remainder of the section.

It follows that $\limsup_n \frac{1}{h_n^2 M_n} < \infty$: and the result follows.

Suppose ℓ_{x,y,Δ_n,h_n}, is used in the estimate below.

Note that $(M_t)_t$. holds throughout.

Here $2\sqrt{s'}\sqrt{\tau \vee h^2} \leq s' + \tau \vee h^2$; by the previous lemma.

In this case $\varphi : \mathbb{R} \to \mathbb{R}$: as claimed.

The value of $F \in \mathbb{D}^{1,2}(\mathbb{R})$, in what follows.

We then obtain $n > 1$. for the remainder of the section.

Recall that $n^{-}(\mathcal{I})$; and the result follows.

It follows that $r \leq R$: is used in the estimate below.

Suppose $x_1, \ldots, x_n$, holds throughout.

Note that $f(x_1, \ldots, x_d) \leq f((x'_1, \ldots, x'_d))$. by the previous lemma.

Here $\{H_i^- : p_i \leq 1/2\} \cup \{H_i^+ : p_i > 1/2\}$.; as claimed.

In this case $\ell_\alpha^-(\mathcal{I}) \geq |\mathcal{D}_\alpha^-|$: in what follows.

The value of $J_v(\mathcal{I}) : n^+(\mathcal{I}) = v$, for the remainder of the section.

We then obtain $|\mathcal{S}^- \cap \mathcal{I}^c| \times |\mathcal{S}^+ \cap \mathcal{I}^c|$. and the result follows.

Recall that $\mathcal{S} = (1, 1)$; is used in the estimate below.

It follows that $1/2 < p_1 < 1 - 2\alpha/3$: holds throughout.

Suppose $\text{SE}(\hat{\pi}_i^j)$, by the previous lemma.

Note that $0 \leq n^+$. as claimed.

Here $H_2^+ \cap H_3^+$; in what follows.

In this case $n^+ = n^- = 15$: for the remainder of the section.

The value of $O(|\mathcal{I}|^2 \cdot \max(1, |\mathcal{I}^c|^2))$, and the result follows.

We then obtain $(p_1, p_2) = (0.19, 0.19)$. is used in the estimate below.

Recall that $c_\alpha(\mathcal{K})$; holds throughout.

It follows that $K_1 \cap K_2 \cap K_3$: by the previous lemma.

Suppose $\tilde{\ell}_\alpha^+$, as claimed.

Note that $\mathcal{S}^- = \{i \in [n] : p_i \leq 1/2\}$. in what follows.

Here $f_v(\mathcal{I}) = \max\{f_{v,u}(\mathcal{I}), u = 0, \ldots, |\mathcal{I}^c|\}$; for the remainder of the section.

In this case $i \in \mathcal{S}^- \cap \mathcal{I}$: and the result follows.

The value of $p^{r/n}$, is used in the estimate below.

We then obtain $\left\{p^{r/n} : r \in [n]\right\}$. holds throughout.

Recall that $q^{n-k+1/n} = f(\{2q_{(n-k+1)}, \ldots, 2q_{(n-s)}\})$; by the previous lemma.

It follows that $p^{0/n} \equiv 0$: as claimed.

Suppose $K^{2/2}$, in what follows.

Note that 0.0271. for the remainder of the section.

Here (0.0403); and the result follows.

In this case ϵ_ℓ: is used in the estimate below.

The value of ϕ_ℓ, holds throughout.

We then obtain $f_\ell(h_\ell; \epsilon_\ell; \phi_\ell)$. by the previous lemma.

Recall that C_L, C'_L; as claimed.

It follows that $\mathbf{X} \in \mathbb{R}^{N \times D}$: in what follows.

Suppose $\mathbf{y} \in \mathbb{R}^N$, for the remainder of the section.

Note that $w \in \mathbb{R}^D$. and the result follows.

Here $\sigma(x) = 1/(1 + e^{-x})$; is used in the estimate below.

In this case μ_{MAP}: holds throughout.

The value of $\Lambda = \beta \sum_i x_i x_i^T + \alpha I$, by the previous lemma.

We then obtain $\sigma(w^T x_i)(1 - \sigma(w^T x_i))$. as claimed.

Recall that $\sigma(w^T x_i)$; in what follows.