The value of B_n/C_n is used in the estimate below.

We then obtain $\tau = \theta/2\pi + 4\pi i/g^2$ holds throughout.

Recall that A_n by the previous lemma.

It follows that A_μ as claimed.

Suppose $x_2 + ix_3 = re^{i\theta}$ in what follows.

Note that $\mathcal{A} = A + i\phi$ for the remainder of the section.

Here n_k and the result follows.

In this case $\lambda_m^{n_m}\lambda_{m-1}^{n_{m-1}}\cdots\lambda_1^{n_1}$ is used in the estimate below.

The value of (D_n) holds throughout.

We then obtain B_k by the previous lemma.

Recall that $\lambda'' = 0$ as claimed.

It follows that r_k' in what follows.

Suppose $l - k + \lambda_k$ for the remainder of the section.

Note that g_* and the result follows.

Here $\frac{1}{2}(\sum_{k=i}^{m} n_k)$ is used in the estimate below.

In this case $l + 1$ holds throughout.

The value of S_{e1221} by the previous lemma.

We then obtain ρ_{odd} as claimed.

Recall that C_{16} in what follows.

It follows that $(\rho_{\text{even}} + X_S^{-1}\rho_{\text{even}}\,;\rho_{\text{odd}} + Y_S\rho_{\text{odd}})_B$ for the remainder of the section.

Suppose $(\lambda_{even}\,;\rho_{odd})$ and the result follows.

Note that $L(C1)$ is used in the estimate below.

Here C_1 holds throughout.

In this case $L(D2)$ by the previous lemma.

The value of (b) as claimed.

We then obtain $c22$ in what follows.

Recall that IIB for the remainder of the section.

It follows that $(\lambda, \rho)_C = (\lambda_{odd}, \rho_{even})_C$ and the result follows.

Suppose CB_{eo} is used in the estimate below.

Note that $\mathcal{I} \in \mathbb{Z}$ holds throughout.

Here $(\psi(0), \psi'(0))$ by the previous lemma.

In this case $(h_{R,t})$ as claimed.

The value of $\mathcal{H}_1$ in what follows.

We then obtain $\mathbf{y} = (y, y')$ for the remainder of the section.

Recall that $\ell_2 \subset \ell_2^{\circ}$ and the result follows.

It follows that $J^* : \mathcal{H}_b \to \mathcal{H}_b$ is used in the estimate below.

Suppose $\ell_2(t)$ holds throughout.

Note that $C_0^{\infty}(\mathbb{R}^+, \mathbb{C}^n)$ by the previous lemma.

Here $(0, u') \in \ell_D$ as claimed.

In this case $\ell^+(E) \cap \ell^-(E) \neq \{0\}$ in what follows.

The value of $(h - E)\psi$ for the remainder of the section.

We then obtain $\ell(t)$ and the result follows.

Recall that $U_1(t)$ is used in the estimate below.

It follows that $P_1 x_j = x_j$ holds throughout.

Suppose $\sigma\left(h_t, \mathbb{T}^1\right)$ by the previous lemma.

Note that $(u_1, \cdots, u_j)$ as claimed.

Here $\ell_t^{\sharp}\{(\Theta x, \Pi x),\ x \in \mathbb{C}\}$ in what follows.

In this case $\mathbf{k} \in \mathbb{Z}^{d-1}$ for the remainder of the section.

The value of $L^2(\Gamma) = H^{s=0}(\gamma)$ and the result follows.

We then obtain $\chi(x)$ is used in the estimate below.

Recall that $(H - E)\psi = 0$ holds throughout.

It follows that $H^2(\Omega^-)$ by the previous lemma.

Suppose $R_{t,D} := (H_{t,D} - \Sigma)^{-1}$ as claimed.

Note that $\ell_{R,t}^{\pm}(E)$ in what follows.

Here $\widetilde{H}_t$ for the remainder of the section.

In this case $L^2(\mathbb{R}^+ \times \mathbb{R})$ and the result follows.

The value of $\gamma^D \psi_D = 0$ is used in the estimate below.

We then obtain $H^{-1/2}(\Gamma) \times H^{1/2}(\Gamma)$ holds throughout.

Recall that $G_{\Sigma} f$ by the previous lemma.

It follows that $\mathrm{LSwHom}(H)$ as claimed.

Suppose G^R in what follows.

Note that $\phi : G' \to H$ for the remainder of the section.

Here (u, v) and the result follows.

In this case $\phi(v) \in L(v)$ is used in the estimate below.

The value of H_c holds throughout.

We then obtain $\pi_S^{-1}(i) = \{(v, i) \mid v \in v(H)\}$ by the previous lemma.

Recall that $(u, 0)$ as claimed.

It follows that L_S in what follows.

Suppose $u, v \in V$ for the remainder of the section.

Note that $x_i \sim y_i$ and the result follows.

Here $u_{i-1} \not\sim v_i$ is used in the estimate below.

In this case B_a holds throughout.

The value of $V(H)^3$ by the previous lemma.

We then obtain x_3 as claimed.

Recall that $i \neq i'$ in what follows.

It follows that $i = \pi_B((i, v))$ for the remainder of the section.

Suppose $\phi : H^k \to H$ and the result follows.

Note that $(i, h, j)(i', h', j')$ is used in the estimate below.

Here $a', b' \in V$ holds throughout.

In this case (y, x) by the previous lemma.

The value of $x_a x_b$ as claimed.

We then obtain r_a in what follows.

Recall that $x_a b$ for the remainder of the section.

It follows that r_b and the result follows.

Suppose $X = \mathrm{maj}(x_1, x_2, x_3)$ is used in the estimate below.

Note that $(a_i, x_i) \sim (b_i, y_i)$ holds throughout.

Here $A \leq a_i$ by the previous lemma.

In this case (a_2, b_1, c_1) as claimed.

The value of $\phi(a_2, d, d) = a_2$ in what follows.

We then obtain $G \cong F$ for the remainder of the section.

Recall that $u \in U \setminus \{1\}$ and the result follows.

It follows that $v \in V(\Gamma)$ is used in the estimate below.

Suppose $\gamma_{n+1}(G) = [\gamma_n(G), G]$ holds throughout.

Note that kI_H by the previous lemma.

Here $p(b, c) = b + c$ as claimed.

In this case $p(b, c) = b + c(t - 1)$ in what follows.

The value of $\widetilde{G} \to G$ for the remainder of the section.

We then obtain M_3 and the result follows.

Recall that $\mathcal{D}_{\mathbb{F}_p H}$ is used in the estimate below.

It follows that $\mathbb{F}_p I_G$ holds throughout.

Suppose $A = H_1 \cap H_2$ by the previous lemma.

Note that u_i as claimed.

Here $\mathcal{D}_{\mathbb{F}_p[H]}$ in what follows.

In this case $c_k \in \mathbb{F}_p N$ for the remainder of the section.

The value of $\widetilde{H_1}$ and the result follows.

We then obtain $i \in \{1, 2\}$ is used in the estimate below.

Recall that u_2 holds throughout.

It follows that $u_2 = v_2^{n_2}$ by the previous lemma.

Suppose $r_{\mathrm{ab}}(\widetilde{H_1}) + r_{\mathrm{ab}}(\widetilde{H_2}) - 1$ as claimed.

Note that ϕ_i in what follows.

Here $\widetilde{H_i} \cong H_i$ for the remainder of the section.

In this case $\widetilde{H}_{\widehat{p}}$ and the result follows.

The value of w_1 is used in the estimate below.

We then obtain $\widetilde{H}_{\widehat{p}} \coprod \mathbb{Z}_p$ holds throughout.

Recall that $d(\widetilde{H}_{\widehat{p}}) = d(\overline{H})$ by the previous lemma.

It follows that $q|_G$ as claimed.

Suppose $u \equiv (uv^{-1})^c \mod [U, U]$ in what follows.

Note that $\gamma_k W_{\widehat{p}}$ for the remainder of the section.

Here $\Gamma \setminus \{e\} = \Delta_1 \cup \Delta_2$ and the result follows.

In this case $\mathbb{Q}(\sqrt{1 - 2m^k})$ is used in the estimate below.

The value of $m \geq 3$, holds throughout.

We then obtain J_K by the previous lemma.

Recall that $n > 3$ as claimed.

It follows that $\mu = \sqrt{2}$ in what follows.

Suppose $k = p^r$ for the remainder of the section.

Note that $(n = 1)$ and the result follows.

Here $p_i \mid h_{(m,p_i)}$ is used in the estimate below.

In this case $i = 1$ holds throughout.

The value of $, where$ by the previous lemma.

We then obtain $, then$ as claimed.

Recall that , $whichisacontradictionto$ in what follows.

It follows that $doesnotdivide$ for the remainder of the section.

Suppose $Then$ and the result follows.

Note that $where$ is used in the estimate below.

Here , for holds throughout.

In this case $.Henceweget$ by the previous lemma.

The value of $andhencewehave$ as claimed.

We then obtain $.Takingmoduloover4onbothsides, weget$ in what follows.

Recall that $isco-primeto$ for the remainder of the section.

It follows that $isoddpositiveinteger, weget$ and the result follows.

Suppose $musthaveprimeidealswhosenormsare$ is used in the estimate below.

Note that $.Hencetheprimedecompositionof$ holds throughout.

Here $.Considertheideal$ by the previous lemma.

In this case $forsomeoddprime$ as claimed.

The value of $dividesclassnumbersof$ in what follows.

We then obtain $betheclassnumbersof$ for the remainder of the section.

Recall that $suchthat$ and the result follows.

It follows that $isaprimewhichdividesanoddinteger$ is used in the estimate below.

Suppose $H^3(G, \mathbb{T})$ holds throughout.

Note that $\omega_1 \otimes_{\text{stack}} \omega_2$ by the previous lemma.

Here $\text{Index}(\omega \circ \mu)$ as claimed.

In this case $\Gamma = \mathbb{Z}$ in what follows.

The value of $\mathcal{A}_\Gamma$ for the remainder of the section.

We then obtain $\tau(\Phi(I)) = \Phi(\tau(I))$ and the result follows.

Recall that $\text{Index}_{1d}^{\mathcal{A}_{L_0},U}(\phi) = \text{Index}^{\mathcal{A},U}(\omega)$ is used in the estimate below.

It follows that $\gamma_{0;1}^{\Psi}$ holds throughout.

Suppose $\alpha_\lambda(g) = \exp(2\pi ik/n)$ by the previous lemma.

Note that $\text{Index}_{1d\ \text{trans}}(\omega) = \text{Index}_{1d\ \text{trans}}(\omega \circ \gamma_{0;1}^{\Phi'})$ as claimed.

Here $x \in \text{Cone}_\sigma^W(\alpha_0, \theta)$ in what follows.

In this case $[\pi_0 \circ \alpha_0 \circ \Theta(A_L), v_R]$ for the remainder of the section.

The value of $\tau_{\mathcal{B}}$ and the result follows.

We then obtain $b_g^R \in \text{Cone}_R^W(\alpha_0, \theta)$ is used in the estimate below.

Recall that $h_{g} \in \mathrm{IAC}_{L}(\theta, \alpha_{0}, \Theta)$ holds throughout.

It follows that $W_{g,1}, u_{R,1}(g,h), v$ by the previous lemma.

Suppose $W_{g,1} \delta_{g}^{\sigma} W_{g,1}^{\dagger} \in \mathrm{IAC}_{\sigma}(\alpha_{0}, \theta, \Theta)$ as claimed.

Note that $n \in \mathbb{N}$ in what follows.

Here n_{0} for the remainder of the section.

In this case $\mathrm{Index}(\omega \circ \gamma_{0;1}^{H}) = \mathrm{Index}(\omega)$ and the result follows.

The value of $a \in \mathrm{IAC}_{L}(\alpha_{0}, \theta, \Theta)$ is used in the estimate below.

We then obtain $x \in \mathrm{IAC}_{L \times R}^{W}(\alpha_{0}, \theta, \Theta)$ holds throughout.

Recall that $B_{\sigma} \in \mathcal{U}(\mathcal{A})$ by the previous lemma.

It follows that $\omega \circ \gamma_{0;1}^{H} \circ \tilde{\beta}_{g,2} = \omega \circ \gamma_{0;1}^{H}$ as claimed.

Suppose $\{1,2,3,4\}$ in what follows.

Note that $\pi_{0} = \bigotimes_{i=1}^{4} \pi_{0,i}$ for the remainder of the section.

Here $(\mathcal{H}_{0}, \pi_{0} \circ i_{V}, \Omega_{0})$ and the result follows.

In this case $B_{n,L}$ is used in the estimate below.

The value of $\Xi^{(s)}(Z,t)$ holds throughout.

We then obtain $X \subset U, X \subset L, X \subset R$ by the previous lemma.

Recall that $\mathcal{K} := \mathrm{age}(\mathcal{A})$ as claimed.

It follows that $\mathrm{age}(\mathcal{I})$ in what follows.

Suppose $\overline{i}' := (i_{k} : k < s) \in X^{s}$ for the remainder of the section.

Note that $\mathrm{Diag}_{\mathcal{A}}(\overline{x})$ and the result follows.

Here $\mathcal{A} \cong_{L} \mathcal{A}'$ is used in the estimate below.

In this case $\mathrm{qftp}^{\mathcal{A}}(\overline{a})$ holds throughout.

The value of $h : B \to C$ by the previous lemma.

We then obtain $L = \{E, <\}$ as claimed.

Recall that $A, B \subseteq \mathcal{M}$ in what follows.

It follows that $\alpha(e,x) = x$ for the remainder of the section.

Suppose $h : \mathcal{A} \to \mathcal{B}$ and the result follows.

Note that $Y \vDash \mathrm{EMtp}(X)$ is used in the estimate below.

Here $||\mathcal{I}||^{+}$ holds throughout.

In this case $\mathcal{I} = (\mathbb{Z}, p, s, <)$ by the previous lemma.

The value of $a_{i} \prec a_{j}$ as claimed.

We then obtain $s \neq t$ in what follows.

Recall that $a_{i} \neq a_{j}$ for the remainder of the section.

It follows that $(g(i) \mid i \in \mathcal{A})$ and the result follows.

Suppose $\eta, \nu \in \omega^{<\omega}$ is used in the estimate below.

Note that $\operatorname{age}(\mathcal{A}) = \operatorname{age}(\mathcal{I}_{\mathrm{eq}})$ holds throughout.

Here $B = \{b_i : i < \omega\}$ by the previous lemma.

In this case $(\mathcal{B}_{ba}, \prec)$. as claimed.

The value of $,itfollowsimmediatelythat$ in what follows.

We then obtain $bestructuresandsuppose$ for the remainder of the section.

Recall that $andfixgenerators$, and the result follows.

It follows that $.Ifthereexistsan$ is used in the estimate below.

Suppose $homogeneousforthecoloring$ holds throughout.

Note that $isthedesiredhomogeneouscopyof$ by the previous lemma.

Here $hasapreimagethatissomewherein$ as claimed.

In this case $with$ in what follows.

The value of $(X, \Delta) \to Z$ for the remainder of the section.

We then obtain (X, Δ) and the result follows.

Recall that $f \colon X' \to X$ is used in the estimate below.

It follows that $e \in \{1, -1\}$ holds throughout.

Suppose $h \colon W \to T'$ by the previous lemma.

Note that $D = \sum_i d_i D_i$ as claimed.

Here $Y_1 \to X$ in what follows.

In this case $g \colon V \to Y$ for the remainder of the section.

The value of B_Y and the result follows.

We then obtain $K_{X'} + \Delta' = \psi^*(K_X + \Delta)$ is used in the estimate below.

Recall that $S^\nu \to S$ holds throughout.

It follows that $a(E, X, \Delta) < a(E, X', \Delta')$ by the previous lemma.

Suppose E' as claimed.

Note that $m\Delta = f_*(mf^*(K_X + \Delta)) - mK_X$ in what follows.

Here n'' for the remainder of the section.

In this case Σ' and the result follows.

The value of $V' = f^{-1}(V)$ is used in the estimate below.

We then obtain π' holds throughout.

Recall that $\left(V', (\Delta' + t_0 A')|_{V'}\right)$ by the previous lemma.

It follows that $(K_{X'} + \Delta' + t_0 A')$ as claimed.

Suppose π_Y in what follows.

Note that $\lfloor \Gamma \rfloor$ for the remainder of the section.

Here $K_{X'} + \Theta' = f^*(K_X + \Delta)$ and the result follows.

In this case $\operatorname{mult}_{Q_j}(\pi_Y^* D_j) \leq \beta$ is used in the estimate below.

The value of $(X, \Delta + tB)$ holds throughout.

We then obtain $m(K_{X'} + \Delta')$ by the previous lemma.

Recall that $G \sim IA$ as claimed.

It follows that $2n''eD$ in what follows.

Suppose $B' \geq 0$ for the remainder of the section.

Note that $\pi' \colon X' \dashrightarrow W$ and the result follows.

Here $\mathcal{F}(R, \mathcal{G})$ is used in the estimate below.

In this case $\mathcal{F}(R)$ holds throughout.

The value of S_0 by the previous lemma.

We then obtain $\mathcal{F}(\mathcal{R}, \mathcal{G})$ as claimed.

Recall that $P_{\mu\nu}$ in what follows.

It follows that $\tilde{b} = \frac{4}{3} q^2 \Lambda$ for the remainder of the section.

Suppose $E(r) = -A'(r)$ and the result follows.

Note that $r \to \infty$ is used in the estimate below.

Here $\mathcal{H}(P)$ holds throughout.

In this case $P <<$ by the previous lemma.

The value of $r \to 0$ as claimed.

We then obtain $r_h = (8Mq^2)^{1/3}$ in what follows.

Recall that (i, j, k) for the remainder of the section.

It follows that $T_H = \frac{\kappa}{2\pi}$ and the result follows.

Suppose $\beta = 1/T_H$ is used in the estimate below.

Note that $r_H <<$ holds throughout.

Here (C) by the previous lemma.

In this case ξ^μ as claimed.

The value of A_E in what follows.

We then obtain $n_+ = n_- = n \geq 1$ for the remainder of the section.

Recall that $\beta = \frac{4M\tilde{\Lambda}}{3}$ and the result follows.

It follows that $r^{1+p}/\sqrt{\eta - \beta r}$ is used in the estimate below.

Suppose $\delta = \tilde{\Lambda}/3$ holds throughout.

Note that $||R||^{2} \to \infty$ by the previous lemma.

Here $Z_{\pm} = R_{1} \pm R_{2}$ as claimed.

In this case $L^{2}\left(0, \infty \right)$ in what follows.

The value of c_{5} for the remainder of the section.

We then obtain $\sigma = \sqrt{1 + 4\bar{\lambda}}/2$ and the result follows.

Recall that $6 \leq \bar{\lambda}^{2}$ is used in the estimate below.

It follows that Q_{1}, Q_{5} holds throughout.

Suppose $a = 0, 1, ..., n - 1$ by the previous lemma.

Note that $\star_{3}\, dA = dH$ as claimed.

Here T^{4} in what follows.

In this case $(n - 1)$ for the remainder of the section.

The value of k_{a}^{2} and the result follows.

We then obtain Π_{ab}^{2} is used in the estimate below.

Recall that q_{a} holds throughout.

It follows that $\lambda = 10^{-5}$ by the previous lemma.

Suppose $[x_{1}, \cdots, x_{\mathbf{n_{0}}}]$ as claimed.

Note that $z = 0$ in what follows.

Here $\{k_{\bar{a}}^{0,1}, k_{0}^{2}\}$ for the remainder of the section.

In this case $Q_{\max} \leftarrow$ and the result follows.

The value of $\bar{r}_{a}^{i}$ is used in the estimate below.

We then obtain $\{r_{A}^{(p)}, \sigma_{A}^{(p)}\}$ holds throughout.

Recall that $\delta r_{A}^{(p)}$ by the previous lemma.

It follows that f' as claimed.

Suppose $2/d$ in what follows.

Note that $\mathcal{N}(0, 1)$ for the remainder of the section.

Here `generation_update` and the result follows.

In this case $\sigma_{A} \to (\texttt{factor_random_update})\, \sigma_{A}$ is used in the estimate below.

The value of 200 holds throughout.

We then obtain 1st by the previous lemma.

Recall that -0.6923 as claimed.

It follows that -61 in what follows.

Suppose -48.5265 for the remainder of the section.

Note that $Q_{5} \simeq 410$ and the result follows.

Here 1.1×10^{4} is used in the estimate below.

In this case $(1 + 1)$ holds throughout.

The value of $\gamma \neq 0$ by the previous lemma.

We then obtain Z_{n} as claimed.

Recall that Z_{3} in what follows.

It follows that $\mathbb{C}^{3}$ for the remainder of the section.

Suppose $2L^{2}$ and the result follows.

Note that $\hat{\Delta}_{v}$ is used in the estimate below.

Here $\varepsilon_{v} = 0$ holds throughout.

In this case h_{v} by the previous lemma.

The value of $\hat{N}_{f}$ as claimed.

We then obtain h_{f} in what follows.

Recall that B_{f} for the remainder of the section.

It follows that $C_{<f,f^{\prime}>}$ and the result follows.

Suppose $\hat{N}_{<f,f^{\prime}>}$ is used in the estimate below.

Note that $-10L^{2}$ holds throughout.

Here $a = 1, 2, 3$ by the previous lemma.

In this case $A^{(a)}$ as claimed.

The value of H_{int} in what follows.

We then obtain H_{v} for the remainder of the section.

Recall that N_{i} and the result follows.

It follows that $p_{\sigma} = \frac{1}{N_{\text{i}}}$ is used in the estimate below.

Suppose $\mathcal{D} = 4$ holds throughout.

Note that $A^{\prime (a)}_{v}$ by the previous lemma.

Here $\hat{\Delta}^{\prime}_{v}$ as claimed.

In this case $h^{\prime}_{v}$ in what follows.

The value of $h^{\prime}_{f}$ for the remainder of the section.

We then obtain $\{\sqrt{3} + 1, 2, 1, 0, 1 - \sqrt{3}\}$ and the result follows.

Recall that $N_{\text{FPL}} = \binom{2l}{l}$ is used in the estimate below.

It follows that $n \approx \sigma l$ holds throughout.

Suppose $\mathcal{C}_{x,y}$ by the previous lemma.

Note that T_c as claimed.

Here $\theta/z = -1$ in what follows.

In this case $\rho_{xx} \sim \tau_{tr}^{-1} \sim T$ for the remainder of the section.

The value of $\cot \Theta_H \sim \tau_h^{-1} \sim T^2$ and the result follows.

We then obtain $\tilde{\sigma} \propto T^{-1}$ is used in the estimate below.

Recall that $\rho_{xx} \propto \alpha T$ holds throughout.

It follows that $3 + 1$ by the previous lemma.

Suppose S_{GR} as claimed.

Note that $F = \mathrm{d}A$ in what follows.

Here ψ_I for the remainder of the section.

In this case $z = 1$ and the result follows.

The value of A_t is used in the estimate below.

We then obtain $T = U(1)/(4\pi)$ holds throughout.

Recall that $B = 0$ by the previous lemma.

It follows that k/μ as claimed.

Suppose $B/\mu^2 = 1/10$ in what follows.

Note that x, y for the remainder of the section.

Here ω_0 and the result follows.

In this case $1 < \beta \leq 2$ is used in the estimate below.

The value of $\sigma_0, \tilde{\sigma}, n$ holds throughout.

We then obtain $x_3 = 0$ by the previous lemma.

Recall that $x_1 \leq 1$ as claimed.

It follows that $x_1 = 1$ in what follows.

Suppose $\beta = x_3 + x_2 \leq 1$ for the remainder of the section.

Note that $T \gg W$ and the result follows.

Here $x_3 \neq 0$ is used in the estimate below.

In this case $\sqrt{a_1 T^2 + a_2 B^2}$ holds throughout.

The value of $|\psi_0\rangle$ by the previous lemma.

We then obtain $N = 10$ as claimed.

Recall that $N = 4, L = 11$ in what follows.

It follows that $S(\theta_1, \ldots, \theta_N)$ for the remainder of the section.

Suppose $\tau_{\mathrm{cr}} \approx 0.33137171$ and the result follows.

Note that $\ell \geq 2$ is used in the estimate below.

Here $\tau^{\mathrm{QSL}} > 0$ holds throughout.

In this case $\sigma^{+}_{k_1 \cdots k_N}$ by the previous lemma.

The value of $\kappa \pm \square_a$ as claimed.

We then obtain $U(1)$ in what follows.

Recall that $\left\{ e^{i2\pi(s-1)/L},\ s = 1 \ldots, L \right\}$ for the remainder of the section.

It follows that S_L and the result follows.

Suppose $\forall\ N \in \mathbb{N}$ is used in the estimate below.

Note that $\beta \to \infty$ holds throughout.

Here $S_{\mathrm{eff}}[\rho]$ by the previous lemma.

In this case $\gamma > \frac{1}{2}$ as claimed.

The value of $\gamma = \frac{1}{2}$ in what follows.

We then obtain $\tilde{x}_a = -x_a$ for the remainder of the section.

Recall that $\tau \to 0$ and the result follows.

It follows that $(\tau_{\mathrm{cr}} - \tau)$ is used in the estimate below.

Suppose $N = 8$ holds throughout.

Note that A_p by the previous lemma.

Here $[0, N-1]$ as claimed.

In this case $\rho^{\mathscr{D}}_{\mathrm{unif.}}$ in what follows.

The value of $\varepsilon > 0$ for the remainder of the section.

We then obtain $\delta > 0$ and the result follows.

Recall that $\{P_j\}_{j \in \mathbb{N}}$ is used in the estimate below.

It follows that $|x| < 1$ holds throughout.

Suppose $e^{itH_{\mathrm{XY}}} |\psi_0\rangle$ by the previous lemma.

Note that h_{2n+1} as claimed.

Here $T = 0$ in what follows.

In this case (a) for the remainder of the section.

The value of $G^{s_1}_{ij}, G^{s_2}_{ij}$ and the result follows.

We then obtain (s_1, s_2) is used in the estimate below.

Recall that Z_C holds throughout.

It follows that $\det(g_C^{s_1, s_2})$ by the previous lemma.

Suppose $G^{s}_{ij}(\theta)$ as claimed.

Note that $\pi/3$ in what follows.

Here S_2 for the remainder of the section.

In this case N_{seed} and the result follows.

The value of $\theta = 10$ is used in the estimate below.

We then obtain $\phi = 0.1$ holds throughout.

Recall that $\ln(L)$ by the previous lemma.

It follows that $\Delta_\tau = 0.5$ as claimed.

Suppose 0.95(5) in what follows.

Note that $\mathrm{f} + \mathrm{b}$ for the remainder of the section.

Here $B^s(\tau_2, \tau_1)$ and the result follows.

In this case $\gamma_i = (1/(1 - e^{-2i\alpha\tilde{\sigma}_i}) - G^s_{i,i}(\tau))^{-1}$ is used in the estimate below.

The value of $\boldsymbol{v}_i$ holds throughout.

We then obtain $\mathcal{O}(N_C^2)$ by the previous lemma.

Recall that $g_{C+i}^{s_1,s_2}$ as claimed.

It follows that $\tau = \theta$ in what follows.

Suppose $G^s(\theta)$ for the remainder of the section.

Note that $\lambda = 0.0006$ and the result follows.

Here $\approx 10^{-6}$ is used in the estimate below.

In this case $L_{\max} = 60$ holds throughout.

The value of $L = 6$ by the previous lemma.

We then obtain $L = 3, 6, 9, 12, 15, 18$ as claimed.

Recall that $N_\lambda/(N_\lambda + 1)$ in what follows.

It follows that $L = 3, 6, 9, ..., 60$ for the remainder of the section.

Suppose $\star$ and the result follows.

Note that $\mathcal{U}_h$ is used in the estimate below.

Here $T_1 G$ holds throughout.

In this case $U_g U_h = U_{gh}$ by the previous lemma.

The value of $\Delta = H^2$ as claimed.

We then obtain $\psi \in \mathcal{H}$ in what follows.

Recall that $t \in (-\tau, \tau)$ for the remainder of the section.

It follows that U, V and the result follows.

Suppose $s : G \to \tilde{G}$ is used in the estimate below.

Note that $\mathcal{H}_j = \mathbb{C}^{2j+1}$ holds throughout.

Here $\mathrm{SO}(2) \cong \mathrm{U}(1)$ by the previous lemma.

In this case $-\mathrm{i}A(\mathfrak{so}(2m)) \subset \mathfrak{u}(\mathcal{H})$ as claimed.

The value of B_{jk} in what follows.

We then obtain $\mathbb{H}_m = \mathbb{R}^{2m+1} = \mathbb{R}^{2m} \times \mathbb{R}$ for the remainder of the section.

Recall that $X \in \mathfrak{sp}(2m, \mathbb{R})$ and the result follows.

It follows that $n \in \mathbb{N}_0^m$ is used in the estimate below.

Suppose $\mathcal{O}(1/L)$ holds throughout.

Note that $\sigma_z = \operatorname{diag}(1, -1)$ by the previous lemma.

Here $\Delta^{(n)}$ as claimed.

In this case $\mathrm{O}(1, 3)$ in what follows.

The value of $p \in \mathcal{M}_m$ for the remainder of the section.

We then obtain $M_{0i} = -M_{i0} = K_i$ and the result follows.

Recall that $d(g_n, 1) \to 0$ is used in the estimate below.

It follows that $\psi = \chi_W$ holds throughout.

Suppose $[0, 1]$ by the previous lemma.

Note that $PX_k = p_k X_k$ as claimed.

Here $\dim(\mathcal{H}) \geq 3$ in what follows.

In this case $\mathcal{P} : \omega(\rho, \psi, p) \mapsto \omega(\rho, 0, p)$ for the remainder of the section.

The value of $\tilde{d}(\tilde{g}, 1_{\tilde{G}}) \geq d(g, 1_G)$ and the result follows.

We then obtain E_{jk} is used in the estimate below.

Recall that $t_g \sim 10^{-8} s$ holds throughout.

It follows that U_g by the previous lemma.

Suppose $\mathrm{dW}_{k,s}$ as claimed.

Note that ϵ^2 in what follows.

Here $(-)$ for the remainder of the section.

In this case $2N^2 - 1$ and the result follows.

The value of $\Lambda(\theta, \phi)$ is used in the estimate below.

We then obtain $s = t / t_g$ holds throughout.

Recall that T_1 by the previous lemma.

It follows that $\gamma_z = t_g (2T_1 - T_2) / 4 T_1 T_2$ as claimed.

Suppose $\epsilon = \sqrt{\lambda}$ in what follows.

Note that (k) for the remainder of the section.

Here $\Delta \mathcal{H}_\sigma^{\mathrm{ng}}$ and the result follows.

In this case ibm_oslo is used in the estimate below.

The value of σ_{kk} holds throughout.

We then obtain $\mathcal{H}^{\mathrm{ng}}$ by the previous lemma.

Recall that $\theta = \pi$ as claimed.

It follows that $\mathcal{N} = 15000$ in what follows.

Suppose $\sim 10^{-3} - 10^{-2}$ for the remainder of the section.

Note that 0.25 and the result follows.

Here $\sim 10^{-2}$ is used in the estimate below.

In this case $\mathcal{H}_{\chi}^{\mathrm{ng}} = \mathcal{H}(\rho^{\mathrm{ng}}, \chi)$ holds throughout.

The value of $\Delta\mathcal{H}_{\chi}^{\mathrm{ng}} < \Delta\mathcal{H}_{\chi}^{\mathrm{ibm}}$ by the previous lemma.

We then obtain $n = 9$ as claimed.

Recall that $\mathrm{R}_{xy}(\phi) = \cos(\phi)\mathrm{X} + \sin(\phi)\mathrm{Y}$ in what follows.

It follows that $\mathrm{N}(\theta, \phi)$ for the remainder of the section.

Suppose $\mathrm{U}(\theta, \phi)$ and the result follows.

Note that $\leq N_s$ is used in the estimate below.

Here $50\% - 60\%$ holds throughout.

In this case $\Delta V / V$ by the previous lemma.

The value of $\Delta H = H_f - H_i$ as claimed.

We then obtain H_i in what follows.

Recall that μ_0 for the remainder of the section.

It follows that J_{ij} and the result follows.

Suppose ΔS_L is used in the estimate below.

Note that $\mathbf{e}_i$ holds throughout.

Here $\mathbf{e}_j$ by the previous lemma.

In this case R_{ij} as claimed.

The value of R_{ij}/a in what follows.

We then obtain μ_B for the remainder of the section.

Recall that T_C and the result follows.

It follows that $M(T) = (1 - \frac{T}{T_C})^{\beta}$ is used in the estimate below.

Suppose $J(r_{ij})$ holds throughout.

Note that r_{ij} by the previous lemma.

Here $\boldsymbol{s}_i$ as claimed.

In this case $\mathbf{p}_i$ in what follows.

The value of $\boldsymbol{r}_i$ for the remainder of the section.

We then obtain ω_i and the result follows.

Recall that $n_{\text{qubits}} = \log_2 \mathcal{C}$ is used in the estimate below.

It follows that S_P holds throughout.

Suppose $a_{\{n_{B_i}\}_k} \neq 0$ by the previous lemma.

Note that $i, j \in 1 \dots N_o$ as claimed.

Here $\sum_i \tilde{\lambda}_i = N_o - N_p$ in what follows.

In this case $S_c \leq \max\{-\ln \nu, -\ln(1-\nu)\}$ for the remainder of the section.

The value of $\alpha_g = 2$ and the result follows.

We then obtain $N_o = 16$ is used in the estimate below.

Recall that α_g holds throughout.

It follows that S_{opt} by the previous lemma.

Suppose $U/t, V/t \approx 4$ as claimed.

Note that $k \in \{1, 2, \dots Q\}$ in what follows.

Here $\epsilon, \delta > 0$ for the remainder of the section.

In this case $n_{\text{bit}} = \log_2 2^{nH(P_k)}$ and the result follows.

The value of $n_1 < n_2$ is used in the estimate below.

We then obtain $n > n'$ holds throughout.

Recall that $\sigma = C^{\beta}$ by the previous lemma.

It follows that $\sigma(n)$ as claimed.

Suppose $\bar{n}_{B_i}$ in what follows.

Note that $S_c^h \leq S_P$ for the remainder of the section.

Here S_{CUE} and the result follows.

In this case $P_j^2 = U_{jk} U_{jl}^* U_{jm} U_{jn}^* a_k a_l^* a_m a_n^*$ is used in the estimate below.

The value of $D_{2,B} = S_{P_B} / \log(Q)$ holds throughout.

We then obtain $S_P = \alpha N_p S_c$ by the previous lemma.

Recall that $\alpha \approx 0.8$ as claimed.

It follows that $\langle x \rangle = 2(1-\nu) N_p$ in what follows.

Suppose $k_1 + q$ for the remainder of the section.

Note that x_{fs} and the result follows.

Here $N_p \gamma$ is used in the estimate below.

In this case $p \to -p$ holds throughout.

The value of $1 + 1$ by the previous lemma.

We then obtain $F_{\mu\nu}(x, t)$ as claimed.

Recall that $\beta = \sqrt{4\pi}$ in what follows.

It follows that $E_\theta = \frac{e}{2\pi}\theta$ for the remainder of the section.

Suppose $m_q \equiv \mathcal{E}(p=0)$ and the result follows.

Note that $\theta = \pi - \varepsilon$ is used in the estimate below.

Here $\pi_1 \pi_2$ holds throughout.

In this case 0.5 by the previous lemma.

The value of π_1, π_2 as claimed.

We then obtain π_1 in what follows.

Recall that $\beta \neq \sqrt{4\pi}$ for the remainder of the section.

It follows that $A_1(x) = 0$ and the result follows.

Suppose $a, b = 1, 2$ is used in the estimate below.

Note that $\frac{e\beta}{2\pi}\phi = E + \frac{e\theta}{2\pi} \equiv E_T$ holds throughout.

Here $\{b(k), b^\dagger(k')\} = \{c(k), c^\dagger(k')\} = \delta(k - k')$ by the previous lemma.

In this case $\hat{p}|p\rangle = p|p\rangle$ as claimed.

The value of $p/m \ll 1$ in what follows.

We then obtain m_q for the remainder of the section.

Recall that $m_\pi / m_q > 2$ and the result follows.

It follows that $I_C^i = C\ddot{\phi}_i$ is used in the estimate below.

Suppose $\phi_J^i + \phi_C^i = \Phi_{\text{ext}}$ holds throughout.

Note that $\omega_{k,j}$ by the previous lemma.

Here $\chi = 1$ as claimed.

In this case $\lambda(\Phi_{\text{ext}})$ in what follows.

The value of A_L for the remainder of the section.

We then obtain $\varepsilon = 0$ and the result follows.

Recall that c_p is used in the estimate below.

It follows that $n \in [i+1, i+N]$ holds throughout.

Suppose $D' = 100$ by the previous lemma.

Note that $n = 2$ as claimed.

Here $SU(2)$ in what follows.

In this case $\mathbf{P}(\omega, \vec{v})$ for the remainder of the section.

The value of $\hat{\mathbf{e}}_1$ and the result follows.

We then obtain $(\sin 2\theta_V,\, 0,\, -\cos 2\theta_V)$ is used in the estimate below.

Recall that $v^\rho(\vec{v}) = (1, \vec{v})$ holds throughout.

It follows that $\text{diag}(+1,\, -1,\, -1,\, -1)$ by the previous lemma.

Suppose $\omega > 0$ as claimed.

Note that μ^{-1} in what follows.

Here $\Omega_{i} > 0$ for the remainder of the section.

In this case v^{z} and the result follows.

The value of $u = \omega$ is used in the estimate below.

We then obtain $\mathbf{Y} = -\mu \mathbf{J}^{z}$ holds throughout.

Recall that $(0, 0, 1)$ by the previous lemma.

It follows that Ω_{i} as claimed.

Suppose $\hat{\mathbf{e}}_{3}^{\mathrm{II}} = \hat{\mathbf{e}}_{3}$ in what follows.

Note that $\hat{\mathbf{e}}_{3}^{\mathrm{II}} = \cos\Theta \hat{\mathbf{e}}_{3}^{\mathrm{III}} - \sin\Theta \hat{\mathbf{e}}_{1}^{\mathrm{III}}$ for the remainder of the section.

Here $\Theta = \pi$ and the result follows.

In this case $\partial_{t}\Theta = \Omega_{i} \sin\Theta$ is used in the estimate below.

The value of $\mathsf{O}^{\mathrm{II}} \ll 1$ holds throughout.

We then obtain $P_{3}^{\mathrm{III}} = \cos\Theta'$ by the previous lemma.

Recall that $\lambda^{\rho} = (\mu,\, 0,\, 0,\, 0.5\mu)$ as claimed.

It follows that $v^{x} = v^{y} = 0$ in what follows.

Suppose $5\pi/6$, for the remainder of the section.

Note that $\times(1 + 0.5v^{z})$ and the result follows.

Here $\Delta\vec{v}$ is used in the estimate below.

In this case $|\mathsf{Q}_{v^{x}, v^{y}}|$ holds throughout.

The value of v_{x} by the previous lemma.

We then obtain J_{3}^{z} as claimed.

Recall that $\mathbf{H}_{\mathrm{vac}}$ in what follows.

It follows that $\operatorname{Im}(\mathsf{C}\,\mathsf{D}^{*}) = 0$ for the remainder of the section.

Suppose σ_{8} and the result follows.

Note that ρ_{0} is used in the estimate below.

Here R_{200} holds throughout.

In this case $c_{200} = R_{200}/r_{s}$ by the previous lemma.

The value of $X(\hat{\mathbf{n}})$ as claimed.

We then obtain $\boldsymbol{\alpha}(\hat{\mathbf{n}}) = \boldsymbol{\nabla}_{\hat{\mathbf{n}}}\phi(\hat{\mathbf{n}})$ in what follows.

Recall that $\Sigma_{\mathrm{cl}}(r)$ for the remainder of the section.

It follows that $d_{A,\mathrm{cl}}$ and the result follows.

Suppose $x = r/r_{s} = \theta/\theta_{s}$ is used in the estimate below.

Note that κ_t holds throughout.

Here $\boldsymbol{\ell} = (\ell_x, \ell_y)$ by the previous lemma.

In this case $z = 0.7$ as claimed.

The value of θ_s in what follows.

We then obtain $N_L^{(1)}$ for the remainder of the section.

Recall that $L = 5000$ and the result follows.

It follows that $\ell_{\mathrm{max}}^{\mathrm{CMB}} = 4000$ is used in the estimate below.

Suppose $L \sim 1100$ holds throughout.

Note that 1000 by the previous lemma.

Here $100 \leq L \leq 5000$ as claimed.

In this case $M = 4, 7$ in what follows.

The value of $M_{200} = 4 \times 10^{14} M_{\odot}, z = 0.7$ for the remainder of the section.

We then obtain $\hat{\kappa}_0$ and the result follows.

Recall that $\hat{\kappa}_L$ is used in the estimate below.

It follows that $\langle \kappa_0 \rangle$ holds throughout.

Suppose 13% by the previous lemma.

Note that x_{max} as claimed.

Here $\chi(z) = \int_0^z dz' \frac{1}{H(z')} = (1+z) d_A(z)$ in what follows.

In this case $\mathrm{Ci}(x)$ for the remainder of the section.

The value of $s = 0, t = 1$ and the result follows.

We then obtain $\langle g(\hat{n}) \rangle_{\text{fixed deflection}}$ is used in the estimate below.

Recall that m_c holds throughout.

It follows that $\psi_C(x) = \gamma_2 \gamma_4 \, \bar{\psi}^T(x)$ by the previous lemma.

Suppose $\bar{\Delta}_5$ as claimed.

Note that $\mu_u + 2\mu_d$ in what follows.

Here $r - g$ for the remainder of the section.

In this case P_{lep} and the result follows.

The value of μ_{ν_e} is used in the estimate below.

We then obtain $m_c = 2.3$ holds throughout.

Recall that $\eta_V = G_V / G_S$ by the previous lemma.

It follows that $\eta_V = 0.5$ as claimed.

Suppose μ_B^c in what follows.

Note that $\mu_B = 850$ for the remainder of the section.

Here $\eta_V = \eta_D = 0$ and the result follows.

In this case $\eta_V = 0.9$ is used in the estimate below.

The value of $n_B = \partial P / \partial \mu_B$ holds throughout.

We then obtain $n_f = \sum_c n_{fc}$ by the previous lemma.

Recall that $g_{\omega B}(n)$ as claimed.

It follows that b_τ in what follows.

Suppose I_{3B} for the remainder of the section.

Note that E_B^* and the result follows.

Here $B = N, P$ is used in the estimate below.

In this case m_e holds throughout.

The value of $P(r = R) = 0$ by the previous lemma.

We then obtain $\mu_{\nu_e}/T = 2$ as claimed.

Recall that $\mu_{\nu_e} = \alpha T$ in what follows.

It follows that $2.3 M_\odot$ for the remainder of the section.

Suppose $T = 30$ and the result follows.

Note that $\bar{\Delta}_0$ is used in the estimate below.

Here n_e holds throughout.

In this case $q \in Q$ by the previous lemma.

The value of $r \in R$ as claimed.

We then obtain $\mathrm{NN}_2(1; R) = \{0, 1, 2\} = \mathrm{NN}_3(1; R)$ in what follows.

Recall that $t \in \mathbb{R}_+$ for the remainder of the section.

It follows that $c(R) = n + 1$ and the result follows.

Suppose $R \subset \mathbb{R}^n$ is used in the estimate below.

Note that $O\big(c_m(R)^{O(1)} \cdot |R| \log(\Delta(R))\big)$ holds throughout.

Here $\mathcal{S}_i(p, \mathcal{T}(R))$ by the previous lemma.

In this case $p \in \mathcal{T}(R)$ as claimed.

The value of $t > \delta$ in what follows.

We then obtain $r \in \mathbb{R}_+$ for the remainder of the section.

Recall that $H(\mathcal{T}(R)) = \{i \mid C_{i-1} \neq C_i\} \cup \{l_{\max}, l_{\min}\}$ and the result follows.

It follows that $l(a_{m-1}) \leq i - 1$ is used in the estimate below.

Suppose $j = l_{\min}(\mathcal{T}(R)) - 1$ holds throughout.

Note that $W_1 = \{r\}$ by the previous lemma.

Here $i \leftarrow -\infty$ as claimed.

In this case $\eta(i) \leftarrow i$ in what follows.

The value of $q \in X \setminus W$ for the remainder of the section.

We then obtain $i \in S$ and the result follows.

Recall that $O(c(R)^2 \log(|R|))$ is used in the estimate below.

It follows that $\varrho \in L(\mathcal{T}(R), q)$ holds throughout.

Suppose $R_{l_{\min}}$ by the previous lemma.

Note that $\psi_k(t)$ as claimed.

Here q_2 in what follows.

In this case p_{m-2} for the remainder of the section.

The value of $m-2$ and the result follows.

We then obtain $p_{i \cdot m-1}$ is used in the estimate below.

Recall that $i = m, j = m-1$ holds throughout.

It follows that $i = 3, j = 2$ by the previous lemma.

Suppose $\langle N, S, A, T, \{R_i\}, \Omega, \{O_i\} \rangle$ as claimed.

Note that $s^t = \langle s^{1,t}, ..., s^{k,t} \rangle$ in what follows.

Here $T(s^{t+1}|s^t, a^t)$ for the remainder of the section.

In this case $O_i(o_i^t|s^t)$ and the result follows.

The value of $\pi_i(a_i^t|h_i^t)$ is used in the estimate below.

We then obtain $X_i = \{X_i^1, ..., X_i^m\} \subseteq S$ holds throughout.

Recall that $O_i(o_i^t|s^t) = \dot{O}_i(o_i^t|x_i^t)$ by the previous lemma.

It follows that $\langle s^t, h_{-i}^t \rangle$ as claimed.

Suppose x_i in what follows.

Note that $x_i \in X_i = \{X_i^1, X_i^2\}$ for the remainder of the section.

Here Y_i and the result follows.

In this case a_j^t is used in the estimate below.

The value of $x_i \in \times_{j \in |X_i|} X_i^j$ holds throughout.

We then obtain $I_i(u_i^t|l_i^t)$ by the previous lemma.

Recall that y_t as claimed.

It follows that θ_i in what follows.

Suppose $I_i(u_i^t|l_i^t) \in \Psi_i$ for the remainder of the section.

Note that $\Psi = \times_{i \in N} \Psi_i$ and the result follows.

Here a_{-i} is used in the estimate below.

In this case M_i^1 holds throughout.

The value of $Q_{M_i^2}^{\pi_i}$ by the previous lemma.

We then obtain $\bar{R} = ||R||_\infty$ as claimed.

Recall that $\{I_{\theta_i}\}_{i \in N}$ in what follows.

It follows that 5×5 for the remainder of the section.

Suppose $F = 4$ and the result follows.

Note that $F = 1$ is used in the estimate below.

Here $P^2(u_i^{t-1} y_i^{t-1}, h_{-i}^{t-1} | l^t)$ holds throughout.

In this case $P^2(h_i^{t+1} | h_i^t, a_i^t)$ by the previous lemma.

The value of $\hat{a}_i^t \neq \bar{a}_i^t$ as claimed.

We then obtain $\hat{I}_{\theta_m}$ in what follows.

Recall that $S(\omega)$ for the remainder of the section.

It follows that $b, 3$ and the result follows.

Suppose $\hat{Y} : \Omega \to \{0, 1\}$ is used in the estimate below.

Note that $.Soeither$ holds throughout.

Here $isdependenton$ by the previous lemma.

In this case $overhardlabelstoinferthevalueof$ as claimed.

The value of $.Itisusedtoevaluatehowwell$ in what follows.

We then obtain $.Wedefinerandomvariables$ for the remainder of the section.

Recall that $isarandomvariable$: and the result follows.

It follows that $), whether$ is used in the estimate below.

Suppose $.Forthisattacktowork$, holds throughout.

Note that $isincludedintheinputdatarecordsand$ by the previous lemma.

Here $.Itmakesiteasiertoextract$ as claimed.

In this case $, classifierof$ in what follows.

The value of $issplitintotwodisjointdatasets$: for the remainder of the section.

We then obtain $.Theseareusedtocreatethestatistic$ and the result follows.

Recall that $using$ is used in the estimate below.

It follows that $)curveasafunctionwhichmapseach$ holds throughout.

Suppose $shouldnotbeabletoinfer$ by the previous lemma.

Note that $thenforallattackmodel$ as claimed.

Here $.Let$ in what follows.

In this case $fortraining$ for the remainder of the section.

The value of $and evaluate iton 634 unseen data records in$ and the result follows.

We then obtain $and evaluate iton 1,200 unseen data records in$ is used in the estimate below.

Recall that $and target model's$ holds throughout.

It follows that $using hard labels against a$ by the previous lemma.

Suppose $issuch that$ as claimed.

Note that $.We then say that$ in what follows.

Here $we call$ for the remainder of the section.

In this case the and the result follows.

The value of $e_k \doteq (\mathbf{x}_k, t_k, p_k)$ is used in the estimate below.

We then obtain $\mathbf{x}_k, t_k$ holds throughout.

Recall that $\mathbf{x}'_k = \mathbf{W}(\mathbf{x}_k, t_k; \boldsymbol{\theta})$ by the previous lemma.

It follows that e'_k as claimed.

Suppose $\epsilon = 1$ in what follows.

Note that $\mathcal{D}(\mathcal{E}, \boldsymbol{\theta})$ for the remainder of the section.

Here $\mathbf{f} \doteq \partial \mathbf{W}(\mathbf{x}, t; \boldsymbol{\theta}) / \partial t$ and the result follows.

In this case $t_{\text{ref}} = t_1$ is used in the estimate below.

The value of $t \in [t_1, t_{N_e}]$ holds throughout.

We then obtain $\Delta t = 0$ by the previous lemma.

Recall that $\mathcal{R} = 0$ as claimed.

It follows that $\mathcal{R} < 0$ in what follows.

Suppose $\boldsymbol{\theta} \equiv (v_x, v_y)^\top$ for the remainder of the section.

Note that $\mathcal{R} \equiv \int_0^1 \frac{d|\mathtt{J}|}{d\Delta t} = 0$ and the result follows.

Here $t \in [0, 1]$ is used in the estimate below.

In this case $\mathtt{R}(\boldsymbol{\omega} \Delta t)$ holds throughout.

The value of $\mathcal{R}_{\mathbf{x}(0)}$ by the previous lemma.

We then obtain $\boldsymbol{\omega} = (0, 0, \omega_z)^\top$ as claimed.

Recall that 346×260 in what follows.

It follows that $N = \{3, 10, 20\}$ for the remainder of the section.

Suppose $\lambda = 1.0$ and the result follows.

Note that $\mathbf{v}(\mathbf{x})$ is used in the estimate below.

Here 2×6 holds throughout.

In this case $\mathbf{V} \equiv (0, 0, v_z)^\top$ by the previous lemma.

The value of $\mathbf{x}'_k = (1 - v_z / Z(\mathbf{x})) t_k.$ as claimed.

We then obtain $\mathbf{x}^{h}(t) = (x(t), y(t), 1)^{\top}$ in what follows.

Recall that 3×3 for the remainder of the section.

It follows that $\mathbf{e}_3^{\top} \mathtt{H}_{t,t+\Delta t} = (0, 0, 1)$ and the result follows.

Suppose $\omega_z = 0, \mathbf{v} = \mathbf{0}$ is used in the estimate below.

Note that $|V|$ holds throughout.

Here $V = \{have, a, great, day\}$ by the previous lemma.

In this case $\{1, 0, 0, 0\}$ as claimed.

The value of V, V^T in what follows.

We then obtain V^T for the remainder of the section.

Recall that $n - 1$ and the result follows.

It follows that w_i is used in the estimate below.

Suppose w_{i+j} holds throughout.

Note that $P(w_i)$ by the previous lemma.

Here $f(w_i)$ as claimed.

In this case b_p in what follows.

The value of X_{pq} for the remainder of the section.

We then obtain w_q and the result follows.

Recall that x_{max} is used in the estimate below.

It follows that 0.75 holds throughout.

Suppose $f(0) = 0$ by the previous lemma.

Note that $f(0)$ as claimed.

Here v_w in what follows.

In this case $n = 3$ for the remainder of the section.

The value of $\mathcal{M} \subset \{1, ..., M\}$ and the result follows.

We then obtain $(t_1, ..., t_{i-1})$ is used in the estimate below.

Recall that t_{i+1} holds throughout.

It follows that $(t_{i+1}, t_{i+2}, ..., t_N)$ by the previous lemma.

Suppose Θ_x as claimed.

Note that Θ_s in what follows.

Here $2L + 1$ for the remainder of the section.

In this case $h_{i,j}^{L} M$ and the result follows.

The value of $x_i^{LM} = h_{i,0}^{LM}$ is used in the estimate below.

We then obtain $\left(SentenceA, SentenceB \right)$ holds throughout.

Recall that $[SEP]$ by the previous lemma.

It follows that $(X_t)_{t\geq 0}$ as claimed.

Suppose $\Delta_n := t_{i+1} - t_i$ in what follows.

Note that X_{Δ_n} for the remainder of the section.

Here $(b_l)_{0\leq l\leq 3}$ and the result follows.

In this case $(\tilde{C}, \tilde{\rho})$ is used in the estimate below.

The value of λ_0 holds throughout.

We then obtain $\pi_{|t-s|}$ by the previous lemma.

Recall that $n \geq 1$ as claimed.

It follows that $d = 1$ in what follows.

Suppose $P_t(\psi_\xi)(x) = \int_{\mathbb{R}} p_t(x,y)\psi_\xi(y)dy$ for the remainder of the section.

Note that $\|\psi\|_\infty \leq 1$ and the result follows.

Here $|b(0)| = 0 \leq b_0$ is used in the estimate below.

In this case $M_n h_n^3 \geq 1/\alpha_0$ holds throughout.

The value of $k \geq 1$ by the previous lemma.

We then obtain $(x, y, \epsilon) \mapsto p^\epsilon_{\Delta_n}(x, y)$ as claimed.

Recall that $h_n = n^{-1/(1+2\beta)}$ in what follows.

It follows that $(B'_n)_n$ for the remainder of the section.

Suppose $b \equiv 0$ and the result follows.

Note that $\dot{X}^\epsilon$ is used in the estimate below.

Here $y > 2h_n$ holds throughout.

In this case $r' > 0$ by the previous lemma.

The value of $|\partial_x A_\epsilon(x, v)| + |\partial_x B(x, v)| \leq c(1 + |v|)$ as claimed.

We then obtain $L(\int_0^s u_r dB_r) = \int_0^s L(u_r)dB_r - \int_0^s u_r dB_r$ in what follows.

Recall that $\|n_{h_n}\|_\infty \leq \frac{c}{M_n}$ for the remainder of the section.

It follows that $\limsup_n \frac{1}{h_n^2 M_n} < \infty$ and the result follows.

Suppose ℓ_{x,y,Δ_n,h_n} is used in the estimate below.

Note that $(M_t)_t$ holds throughout.

Here $2\sqrt{s'}\sqrt{\tau \vee h^2} \leq s' + \tau \vee h^2$ by the previous lemma.

In this case $\varphi : \mathbb{R} \to \mathbb{R}$ as claimed.

The value of $F \in \mathbb{D}^{1,2}(\mathbb{R})$ in what follows.

We then obtain $n > 1$ for the remainder of the section.

Recall that $n^{-}(\mathcal{I})$ and the result follows.

It follows that $r \leq R$ is used in the estimate below.

Suppose $x_1, \ldots, x_n$ holds throughout.

Note that $f(x_1, \ldots, x_d) \leq f((x'_1, \ldots, x'_d))$ by the previous lemma.

Here $\{H_i^- : p_i \leq 1/2\} \cup \{H_i^+ : p_i > 1/2\}$. as claimed.

In this case $\ell_\alpha^-(\mathcal{I}) \geq |\mathcal{D}_\alpha^-|$ in what follows.

The value of $J_v(\mathcal{I}) : n^+(\mathcal{I}) = v$ for the remainder of the section.

We then obtain $|\mathcal{S}^- \cap \mathcal{I}^c| \times |\mathcal{S}^+ \cap \mathcal{I}^c|$ and the result follows.

Recall that $\mathcal{S} = (1, 1)$ is used in the estimate below.

It follows that $1/2 < p_1 < 1 - 2\alpha/3$ holds throughout.

Suppose $\mathrm{SE}(\hat{\pi}_i^j)$ by the previous lemma.

Note that $0 \leq n^+$ as claimed.

Here $H_2^+ \cap H_3^+$ in what follows.

In this case $n^+ = n^- = 15$ for the remainder of the section.

The value of $O(|\mathcal{I}|^2 \cdot \max(1, |\mathcal{I}^c|^2))$ and the result follows.

We then obtain $(p_1, p_2) = (0.19, 0.19)$ is used in the estimate below.

Recall that $c_\alpha(\mathcal{K})$ holds throughout.

It follows that $K_1 \cap K_2 \cap K_3$ by the previous lemma.

Suppose $\tilde{\ell}_\alpha^+$ as claimed.

Note that $\mathcal{S}^- = \{i \in [n] : p_i \leq 1/2\}$ in what follows.

Here $f_v(\mathcal{I}) = \max\{f_{v,u}(\mathcal{I}), u = 0, \ldots, |\mathcal{I}^c|\}$ for the remainder of the section.

In this case $i \in \mathcal{S}^- \cap \mathcal{I}$ and the result follows.

The value of $p^{r/n}$ is used in the estimate below.

We then obtain $\left\{p^{r/n} : r \in [n]\right\}$ holds throughout.

Recall that $q^{n-k+1/n} = f(\{2q_{(n-k+1)}, \ldots, 2q_{(n-s)}\})$ by the previous lemma.

It follows that $p^{0/n} \equiv 0$ as claimed.

Suppose $K^{2/2}$ in what follows.

Note that 0.0271 for the remainder of the section.

Here (0.0403) and the result follows.

In this case ϵ_ℓ is used in the estimate below.

The value of ϕ_ℓ holds throughout.

We then obtain $f_\ell(h_\ell; \epsilon_\ell; \phi_\ell)$ by the previous lemma.

Recall that C_L, C'_L as claimed.

It follows that $\mathbf{X} \in \mathbb{R}^{N \times D}$ in what follows.

Suppose $\mathbf{y} \in \mathbb{R}^N$ for the remainder of the section.

Note that $w \in \mathbb{R}^D$ and the result follows.

Here $\sigma(x) = 1/(1 + e^{-x})$ is used in the estimate below.

In this case μ_{MAP} holds throughout.

The value of $\Lambda = \beta \sum_i x_i x_i^T + \alpha I$ by the previous lemma.

We then obtain $\sigma(w^T x_i)(1 - \sigma(w^T x_i))$ as claimed.

Recall that $\sigma(w^T x_i)$ in what follows.