The value of $\Omega \subset \mathbf{R}^d$, is used in the estimate below.

We then obtain $\epsilon > 0$. holds throughout.

Recall that $A, B \subset \mathbf{R}^d$; by the previous lemma.

It follows that $W(0,T)$: as claimed.

Suppose $(\omega, f, u) \in \mathfrak{Y}(\Omega) \times H_0^1(\Omega) \times \mathrm{U}$, in what follows.

Note that $2\gamma \bar{u} = \chi_\omega \bar{p}$. for the remainder of the section.

Here $y = y^{u,f,w}$.; and the result follows.

In this case $J(\omega, \cdot, y)$: is used in the estimate below.

The value of $\|f\|_{H_0^1(\Omega)} \leq \liminf_{k\to\infty} \|f_{n_k}\|_{H_0^1(\Omega)} \leq 1$, holds throughout.

We then obtain $t \to 0^+$. by the previous lemma.

Recall that $\tilde{\varphi}_n \rightharpoonup \varphi$; as claimed.

It follows that $s \in [0,1]$: in what follows.

Suppose a_{ij}, b_{ij}, for the remainder of the section.

Note that $D\mathcal{J}_1(\omega, f)$. and the result follows.

Here $\bar{p}^{f,\omega} \in L_2(0,T; H^3(\Omega))$; is used in the estimate below.

In this case $\partial_{x_j}(\bar{y}^{f,\omega}(t)\partial_t \bar{p}^{f,\omega}(t)) \in L_1(\Omega)$: holds throughout.

The value of $D\mathcal{J}_2(\omega)(X) = 0$, by the previous lemma.

We then obtain $\mathbf{S}_0(-f) = \mathbf{S}_0(f)$. as claimed.

Recall that $\tau \in [0, \tau_X]$; in what follows.

It follows that $f_h \in V_h$: for the remainder of the section.

Suppose $(\mathrm{id} + t\nabla \mathcal{J}_{1,h}(\omega, f_h))(\omega)$, and the result follows.

Note that β_n. is used in the estimate below.

Here $\psi_h^0 \in V_h(\Omega)$; holds throughout.

In this case Π_h: by the previous lemma.

The value of $y_0(x) = \max(\sin(3\pi x), 0)^2$, as claimed.

We then obtain 10^3. in what follows.

Recall that $\sigma(x) = (1 - \max(\sin(9\pi x), 0))\chi_{\{x<0.5\}}(x) + 10^{-3}$; for the remainder of the section.

It follows that $\mathcal{G} : [0,\tau] \times \mathfrak{V}_1 \to \mathbf{R}$: and the result follows.

Suppose $(0, \tau]$, is used in the estimate below.

Note that $g(t) := \sup_{y \in \mathfrak{V}} \mathcal{G}(t, y)$. holds throughout.

Here $N \in \mathbb{N}$; by the previous lemma.

In this case $\mathbb{R}^d$: as claimed.

The value of $f : \mathbb{R}^d \to \mathbb{R}$, in what follows.

We then obtain $f(y) \geq f(x) + \langle g, y - x \rangle$. for the remainder of the section.

Recall that $\mathcal{N}_i^+ = \{v_j \in \mathcal{V} : \ (v_i, v_j) \in \mathcal{E}\}$; and the result follows.

It follows that $v_j \in \mathcal{V} \setminus \{v_i\}$: is used in the estimate below.

Suppose $\mathcal{R} = \mathcal{V} \setminus \mathcal{A}$, holds throughout.

Note that $x_i^* \leftarrow$. by the previous lemma.

Here $\mathcal{N}_i^-$; as claimed.

In this case x_i^*: in what follows.

The value of $x_j[k] \in \mathcal{S}_i^{\mathrm{dist}}[k] \setminus \{x_i[k]\}$, for the remainder of the section.

We then obtain $\|g_i(x)\|_2 \leq L$. and the result follows.

Recall that $v_i, v_j \in \mathcal{R}$; is used in the estimate below.

It follows that $\mathcal{S}_i^{mm}[k]$: holds throughout.

Suppose $v_j \in (\mathcal{N}_i^- \cap \mathcal{R}) \cup \{v_i\}$, by the previous lemma.

Note that $v_j \in \mathcal{N}_i^{\mathrm{mm}}[k]$. as claimed.

Here $x \notin \mathcal{C}(\epsilon)$; in what follows.

In this case $t \in (0, \infty)$: for the remainder of the section.

The value of $\mathbb{R}^d \setminus \mathcal{B}(\hat{x},\ s^*(\xi))$, and the result follows.

We then obtain $\|z_i[k] - \hat{x}\|_2$. is used in the estimate below.

Recall that $x_0 \notin \mathcal{C}_i(\epsilon)$; holds throughout.

It follows that $x \in [-1, 1]$: by the previous lemma.

Suppose $f(x) = \frac{1}{|\mathcal{R}|} \sum_{v_i \in \mathcal{R}} f_i(x)$, as claimed.

Note that b_i. in what follows.

Here $x - \hat{x}$; for the remainder of the section.

In this case $k \geq k_1^*$: and the result follows.

The value of k_2^*, is used in the estimate below.

We then obtain $b_i > 0$. holds throughout.

Recall that $\mathcal{I}_z[k]$; by the previous lemma.

It follows that $k^* = \max\{k_1^*, k_2^*, k_3^*\}$: as claimed.

Suppose $\rho = \sum_{\alpha} A_\alpha \otimes B_\alpha$, in what follows.

Note that $\sum_{\alpha} A_\alpha \otimes B_\alpha$. for the remainder of the section.

Here $d = d_1 \cdots d_n$; and the result follows.

In this case $\operatorname{tr}(\chi C_T) \geqslant 0$: is used in the estimate below.

The value of $s \times s$, holds throughout.

We then obtain $t = 2$. by the previous lemma.

Recall that $i = 1, 2$; as claimed.

It follows that z_i: in what follows.

Suppose ω^*, for the remainder of the section.

Note that $\tau_1, \ldots, \tau_d$. and the result follows.

Here $(\tau_1, \ldots, \tau_d) \in C_s(p)$; is used in the estimate below.

In this case p_i: holds throughout.

The value of $d, s \in \mathbb{N}$, by the previous lemma.

We then obtain σ_1, σ_2. as claimed.

Recall that $T_1 = C_1(\ell)$; in what follows.

It follows that D_1: for the remainder of the section.

Suppose $\mathcal{D} \subseteq \mathcal{C}(\ell)$, and the result follows.

Note that $n \times n$. is used in the estimate below.

Here $\mathcal{Q}_A$; holds throughout.

In this case $r_A(a)(x)$: by the previous lemma.

The value of a, b, as claimed.

We then obtain $\mathcal{Q}_A, \mathcal{Q}_B, \mathcal{A}_A, \mathcal{A}_B$. in what follows.

Recall that $\sum_r v_r^* v_r = 1$; for the remainder of the section.

It follows that i, j, k, l: and the result follows.

Suppose $z_{(a,x)}$, is used in the estimate below.

Note that $I_d \otimes q_B(b)(y)$. holds throughout.

Here $d_i = 2$; by the previous lemma.

In this case $10^{10^{100}}$: as claimed.

The value of $\tau_n(\rho) \succcurlyeq 0$, in what follows.

We then obtain $\operatorname{tr}(\rho^k)$. for the remainder of the section.

Recall that $t \gg n^{(d-1)/d}$; and the result follows.

It follows that $t \geq 1$: is used in the estimate below.

Suppose $t \in \mathbb{N}$, holds throughout.

Note that $t := t(n) = o(\tau_{\mathcal{P}}(n))$. by the previous lemma.

Here $\tau_{\mathcal{P}_H} \leq n^{(d-1)/d}$; as claimed.

In this case $(3\Delta/2 + o(\Delta))n$: in what follows.

The value of $U_t \cup V_t$, for the remainder of the section.

We then obtain $H \subseteq G_t^{(s)}$. and the result follows.

Recall that $\ell \in [k]$; is used in the estimate below.

It follows that $\mathbb{E}X_t^{(x)} = (1 - o(1))\frac{t^x}{x!n^{x-1}}$: holds throughout.

Suppose $I_t(i)^{(x)} = 1$, by the previous lemma.

Note that $\{v_1, v_2, \ldots, v_{\ell-1}\}$. as claimed.

Here $w_1, \ldots, w_h$; in what follows.

In this case $H = C_4$: for the remainder of the section.

The value of H_{i-1}, and the result follows.

We then obtain $w_H(v) \geq \delta$. is used in the estimate below.

Recall that $\delta - 1$; holds throughout.

It follows that $T = V(H_m) \setminus S$: by the previous lemma.

Suppose $x_i \in T, y_i \notin T$, as claimed.

Note that $i = 0$. in what follows.

Here $\phi(e)$; for the remainder of the section.

In this case $d(u, v)$: and the result follows.

The value of $i \in \{0, 1, \ldots, m\}$, is used in the estimate below.

We then obtain $w_i(v) = w_{i-1}(v)$. holds throughout.

Recall that $u \in S(x, H_i; t)$; by the previous lemma.

It follows that $w_i(u) = w$: as claimed.

Suppose $v \in V(H')$, in what follows.

Note that G_{m-1}. for the remainder of the section.

Here $G_t = G_t^{(1,s)}$; and the result follows.

In this case $f(c)$: is used in the estimate below.

The value of $\mathrm{Recol}(H)$, holds throughout.

We then obtain K_k. by the previous lemma.

Recall that $\phi(v)$; as claimed.

It follows that $\phi(y)$: in what follows.

Suppose C', for the remainder of the section.

Note that W_r. and the result follows.

Here $W \cdot C \cdot W^{-1}$; is used in the estimate below.

In this case $[X]$: holds throughout.

The value of $Y_1 \in [X_1]$, by the previous lemma.

We then obtain $[C] = [C']$. as claimed.

Recall that $[\psi(C)] = [\beta_{W_r}(\phi(C))]$; in what follows.

It follows that $x_0 = x_\ell$: for the remainder of the section.

Suppose $raev$, and the result follows.

Note that $\Pi(\mathcal{U}; (r), (r, x_1, x_2, \ldots, x_\ell))$. is used in the estimate below.

Here ϕ^{-1}; holds throughout.

In this case $W \cdot \phi_{v_i}^{-1}(v_i) \cdot U$: by the previous lemma.

The value of $C_2 = U_2 \cdot X^{-1} \cdot V_2$, as claimed.

We then obtain $C_2' = X^{-1} \cdot V_2 \cdot U_1$. in what follows.

Recall that $[\psi(C)] = [\beta_{W_v}(\phi(C)]$; for the remainder of the section.

It follows that $C = T_u \cdot e \cdot T_v^{-1}$: and the result follows.

Suppose $[\![C]\!]$, is used in the estimate below.

Note that $X = I \cdot J$. holds throughout.

Here $[A^d W] = [W B^d]$; by the previous lemma.

In this case I_W: as claimed.

The value of $T \cdot A_f^{-d}$, in what follows.

We then obtain $B = \beta_S(B_f)$. for the remainder of the section.

Recall that $T \neq T'$; and the result follows.

It follows that W_0: is used in the estimate below.

Suppose $(W_v)_{v \in G}$, holds throughout.

Note that $|W_v| \leq 1$. by the previous lemma.

Here $Re(s) > 1 - \frac{1}{(\log q)^R}$; as claimed.

In this case me^{2m}: in what follows.

The value of $q < x^\theta$, for the remainder of the section.

We then obtain $t = 0$. and the result follows.

Recall that $\left(\frac{M}{n}\right)$; is used in the estimate below.

It follows that $A > 2$: holds throughout.

Suppose $m \geq 1$, by the previous lemma.

Note that $q < x^{\frac{233}{462}}$. as claimed.

Here $L'(s, \chi_D) \ll \log^2 D$; in what follows.

In this case θ': for the remainder of the section.

The value of $q \leq x^\theta$, and the result follows.

We then obtain $q < x^{\theta'}$. is used in the estimate below.

Recall that χ_C; holds throughout.

It follows that $\chi_D(n) = \chi_g(n)\chi_{D'}(n)$,: by the previous lemma.

Suppose $l \leq D$, as claimed.

Note that $x = D'y$. in what follows.

Here $H > k2^{k-1}$; for the remainder of the section.

In this case $Q_j(y)$: and the result follows.

The value of $y' \leq g$, is used in the estimate below.

We then obtain $\theta < \frac{58}{115}(1 - \frac{1}{r})$. holds throughout.

Recall that S_k; by the previous lemma.

It follows that $o(1)$: as claimed.

Suppose e_i, in what follows.

Note that $\lambda \ll 1$. for the remainder of the section.

Here $m = 3, 4, 5$; and the result follows.

In this case Z, Z_3, W, X, V, U: is used in the estimate below.

The value of X_i, holds throughout.

We then obtain $\beta = .973$. by the previous lemma.

Recall that $M_{38,801} \geq 7.9310494$; as claimed.

It follows that H_3: in what follows.

Suppose $f \colon \mathcal{M} \to \mathbb{R}, h \colon \mathcal{M} \to \mathbb{R}^l$, for the remainder of the section.

Note that $\dot{c}(t) \in T_{c(t)}\mathcal{M}$. and the result follows.

Here $\operatorname{Exp}_p \colon T_p\mathcal{M} \to \mathcal{M}$; is used in the estimate below.

In this case $\mathbb{R}^m \ni s := -g(x)$: holds throughout.

The value of $\mathbb{A}(x) := \{i \mid g_i(x) = 0\}$, by the previous lemma.

We then obtain $\mathbb{I} := \{1, \ldots, m\}$. as claimed.

Recall that $\nabla \hat{e}(w)[\Delta w] = (0_x, 0, 0, 0)$; in what follows.

It follows that $(z_{k+1}, s_{k+1}) > 0$: for the remainder of the section.

Suppose $[\hat{\mathcal{A}}]_{ij} := \left\langle \mathcal{A}u_j, u_i \right\rangle_x$, and the result follows.

Note that $\alpha_n := \langle r_n, \mathcal{A}r_n \rangle_x / \langle \mathcal{A}p_n, \mathcal{A}p_n \rangle_x$. is used in the estimate below.

Here $\mathrm{T}_\eta\left(\xi\right) \in T_{\mathrm{R}_x(\eta)}\mathcal{M}$; holds throughout.

In this case $t \mapsto \gamma(t) := \mathrm{R}_x\left(t\eta\right)$: by the previous lemma.

The value of $a_0, a_1, \delta_{a_0,a_1} > 0$, as claimed.

We then obtain $\Xi > 0$. in what follows.

Recall that $v_k > 0$; for the remainder of the section.

It follows that $\gamma(t) = \mathrm{R}_{p_k}(t\eta)$: and the result follows.

Suppose $w_k \to w^*$, is used in the estimate below.

Note that $y(\alpha) = y + \alpha \Delta y$. holds throughout.

Here $\gamma_k \in (0.5, \gamma_{k-1})$; by the previous lemma.

In this case $\rho \leq \left\| F(w) \right\| / \sqrt{m}$: as claimed.

The value of $\mathcal{A}_x$, in what follows.

We then obtain $\left\| \mathcal{A}_x y \right\|_x = \| \hat{\mathcal{A}}_x \hat{y} \|_2$. for the remainder of the section.

Recall that $\{ \dot{e}_i \}_{i=1}^m$; and the result follows.

It follows that $\{ \Delta w_k \}$: is used in the estimate below.

Suppose $[\hat{\nabla} F(w_k)]^{-1}$, holds throughout.

Note that $\alpha \Delta w$. by the previous lemma.

Here $\omega(\cdot)$; as claimed.

In this case $\sigma = 0$: in what follows.

The value of $\times 10^{-7}$, for the remainder of the section.

We then obtain $\Omega \subset \mathbb{R}^N$. and the result follows.

Recall that u_0; is used in the estimate below.

It follows that $\nu > 0$: holds throughout.

Suppose $0 < \chi < \frac{N}{N-2}$, by the previous lemma.

Note that $(t, x) \in (s, s+T) \times \Omega$. as claimed.

Here $\sup_{t \geq s} \| u(t, \cdot; s, u_0) \|_\infty < \infty$; in what follows.

In this case $a_{\chi, \mu}$: for the remainder of the section.

The value of $\int_\Omega u^{-p}(t, x; s, u_0) dx$, and the result follows.

We then obtain $T(u_0) > 0$. is used in the estimate below.

Recall that $(u, v) = (u^*(t, x), v^*(t, x))$; holds throughout.

It follows that $\chi_0 > 1$: by the previous lemma.

Suppose $N \leq 2$, as claimed.

Note that $p > 1$. in what follows.

Here $X_p^\alpha = D(A_p^\alpha)$; for the remainder of the section.

In this case $C_{p, \beta, \epsilon} > 0$: and the result follows.

The value of $\inf_{x \in \bar{\Omega}} u_0(x) > 0$, is used in the estimate below.

We then obtain $a_0(t, x), b_0(t, x)$. holds throughout.

Recall that $0 < T \ll 1$; by the previous lemma.

It follows that $\mathcal{M}(\tilde{u}_0, a, b)$: as claimed.

Suppose $v := \nu A^{-1} u,$, in what follows.

Note that $t \in [s, s+T],$. for the remainder of the section.

Here $R > 0$; and the result follows.

In this case $\tilde{M}_1^* > 0$: is used in the estimate below.

The value of $\beta \in (0, \frac{1}{2})$, holds throughout.

We then obtain $t_n - s_n \geq T_n + 1$. by the previous lemma.

Recall that $\int_\Omega u_1^{-p}(x)dx \leq M_1^*$; as claimed.

It follows that $u \in \mathcal{E}$: in what follows.

Suppose $(-\infty, 0)$, for the remainder of the section.

Note that $\mathcal{T}(T)$. and the result follows.

Here $T > 0$,; is used in the estimate below.

In this case $\mathcal{L}_{CS} =:$ holds throughout.

The value of k_μ, by the previous lemma.

We then obtain $[1, 2]$. as claimed.

Recall that b_μ; in what follows.

It follows that $[6, 7, 8]$: for the remainder of the section.

Suppose [9], and the result follows.

Note that $S_{eff}[b, A]$. is used in the estimate below.

Here $S_{eff}[b, A] = S_{eff}[b] + S'_{eff}[b, A]$; holds throughout.

In this case $S_{\text{eff}}\,[b] = -i\,\mathrm{Tr}\ln(\not{p} - m - \not{\gamma}_5)$: by the previous lemma.

The value of $S_{\text{eff}}^{(2)}[b, A]$, as claimed.

We then obtain $\Pi^{\alpha\mu\nu}$. in what follows.

Recall that $\Pi_{\mathbf{r}}^{\mu\alpha\nu}$; for the remainder of the section.

It follows that $x_0 \to -ix_0, p_0 \to ip_0, b_0 \to ib_0, d^4x \to -id^4x$: and the result follows.

Suppose $d^4p \to id^4p$, is used in the estimate below.

Note that I_1. holds throughout.

Here I_2; by the previous lemma.

In this case $\epsilon = 3 - D$: as claimed.

The value of $D = 3$, in what follows.

We then obtain $[5, 17]$. for the remainder of the section.

Recall that $n = 1$; and the result follows.

It follows that $B = eA$: is used in the estimate below.

Suppose $\left(e^2/16\pi^2\right)$, holds throughout.

Note that $\Delta k_\beta = \left(3e^2/16\pi^2\right) b_\beta$. by the previous lemma.

Here k_β; as claimed.

In this case $K^0 \bar{K}^0$: in what follows.

The value of $B^0 \bar{B}^0$, for the remainder of the section.

We then obtain $\{\cdot, \cdot\}$. and the result follows.

Recall that $N = 2$; is used in the estimate below.

It follows that a^0: holds throughout.

Suppose E^0, by the previous lemma.

Note that $\boldsymbol{\gamma} = \boldsymbol{\Gamma}/|\boldsymbol{\Gamma}|$. as claimed.

Here $\theta_{\mathbf{e}\boldsymbol{\gamma}}$; in what follows.

In this case $\pm\pi/2$: for the remainder of the section.

The value of $\Gamma_{11} = \Gamma_{22} \in \mathbb{R}$, and the result follows.

We then obtain K^0. is used in the estimate below.

Recall that 1.511 ± 0.147; holds throughout.

It follows that $(2.62 \pm 0.38) \times 10^{-3}$: by the previous lemma.

Suppose m_-, as claimed.

Note that $w_0 = \left(1 \pm \sqrt{1 - 4|w_I|^2}\right)/2$. in what follows.

Here $\zeta = 1$; for the remainder of the section.

In this case $|\mathbf{b}_\star| = 1$: and the result follows.

The value of $x = \zeta \sin^2 \theta_{\mathbf{b}_\star \mathbf{d}}/(2 \cos \theta_{\mathbf{b}_\star \boldsymbol{\gamma}})$, is used in the estimate below.

We then obtain $\mathbf{b}(0) = \mathbf{0}$. holds throughout.

Recall that $r = 0.2$; by the previous lemma.

It follows that $(\boldsymbol{\gamma}, \mathbf{e} \times \boldsymbol{\gamma})$: as claimed.

Suppose $|\mathbf{b}(0)| = 1$, in what follows.

Note that $|\mathbf{b}|$. for the remainder of the section.

Here $\mathbf{b}(0) = -\boldsymbol{\gamma}$; and the result follows.

In this case $|\mathbf{b}(\tau)|$: is used in the estimate below.

The value of $\tau \to \infty$, holds throughout.

We then obtain $\mathbf{d} = \boldsymbol{\gamma}$. by the previous lemma.

Recall that $\operatorname{Tr} \mathbb{J}$; as claimed.

It follows that $|\cos \theta| \ll 1$: in what follows.

Suppose $g - 2$, for the remainder of the section.

Note that $d = 2 + 1$. and the result follows.

Here $\mathcal{N} = 2$; is used in the estimate below.

In this case $D_\mu = \partial_\mu + ieA_\mu$: holds throughout.

The value of $m_\sigma < m_A$, by the previous lemma.

We then obtain $\rho \to v.$ as claimed.

Recall that I_ν; in what follows.

It follows that $\mathcal{O}\left(N^6\right)$: for the remainder of the section.

Suppose $\mathcal{H}(\mathbf{x})$, and the result follows.

Note that $r \sim m^{-1}.$ is used in the estimate below.

Here $\mathcal{P}_\mu{}^\nu \equiv \text{diag}\,(1,-1,1)$; holds throughout.

In this case $n \geq 2$: by the previous lemma.

The value of $Z = +m$, as claimed.

We then obtain $Z = M.$ in what follows.

Recall that $c_{\Gamma\Gamma\Gamma} = c_{\bar{\Gamma}\Gamma\Gamma}$; for the remainder of the section.

It follows that S_{AHM}: and the result follows.

Suppose $\mathcal{O}\left(N^{1/2}\right)$, is used in the estimate below.

Note that $E_{\mathbf{p}} \equiv \sqrt{\mathbf{p}^2 + \mathfrak{m}^2}.$ holds throughout.

Here $P^\mu = (M, 0)$; by the previous lemma.

In this case $-\mathbf{p}$: as claimed.

The value of $\cdots$, in what follows.

We then obtain $\mathbf{p} \cdot \mathbf{q}.$ for the remainder of the section.

Recall that $\nabla^2 \delta^{(2)}\left(\mathbf{x}\right)$; and the result follows.

It follows that $q \to -q$: is used in the estimate below.

Suppose $\frac{m}{M} \to 0$, holds throughout.

Note that $\mathbf{x}_i \equiv (x_i, y_i).$ by the previous lemma.

Here $\sigma \tilde{F}_\mu \Phi^* \partial^\mu \Phi$; as claimed.

In this case $\epsilon^{12} = +1$: in what follows.

The value of $-1/2$, for the remainder of the section.

We then obtain $l \to -l - q.$ and the result follows.

Recall that $n > 0$; is used in the estimate below.

It follows that $\gamma_3 = -0.00048(4)$: holds throughout.

Suppose $T\bar{T}$, by the previous lemma.

Note that $\gamma_5.$ as claimed.

Here $N_s a$; in what follows.

In this case $(\vec{x}, z)$: for the remainder of the section.

The value of $N_{t,c}(\beta)$, and the result follows.

We then obtain $E_0(T)$. is used in the estimate below.

Recall that $\xi^1 = X^1$; holds throughout.

It follows that $\frac{D-3}{2}$: by the previous lemma.

Suppose $T_{c,NG}$, as claimed.

Note that $K = \Delta(g)X$. in what follows.

Here $\gamma_3 K^4$; for the remainder of the section.

In this case K^4: and the result follows.

The value of $1/R$, is used in the estimate below.

We then obtain $\beta_c = 0.76141330(6)$. holds throughout.

Recall that $\beta < \beta_r$; by the previous lemma.

It follows that $\sum_{\langle i,j \rangle}$: as claimed.

Suppose Δ_ϵ, in what follows.

Note that $V(r)$. for the remainder of the section.

Here $N_{t,c} = 12$; and the result follows.

In this case $N_t = 24$: is used in the estimate below.

The value of $\beta = 0.751800$, holds throughout.

We then obtain σN_t. by the previous lemma.

Recall that $N_p = 3N_s^2 L$; as claimed.

It follows that $K_0'(z)/K_0(z)$: in what follows.

Suppose $x = \frac{\pi}{3\sigma_0 N_t^2}$, for the remainder of the section.

Note that $e^{-\sigma_0 N_s^2}$. and the result follows.

Here $\beta = 0.758266$; is used in the estimate below.

In this case $SU(6)$: holds throughout.

The value of $\sigma_c = 10.083(8)$, by the previous lemma.

We then obtain t_c. as claimed.

Recall that $\tau_b \neq \tau_d$; in what follows.

It follows that $\phi(t, x)$: for the remainder of the section.

Suppose $\partial_t \phi(t, x)$, and the result follows.

Note that $\rho \propto (\partial_t \phi(t, x))^2 + ...$ is used in the estimate below.

Here $\vec{x} = \vec{x}_P$; holds throughout.

In this case $\partial_t \psi(t, \vec{x}_P)$: by the previous lemma.

The value of $\psi(x, t_c)$, as claimed.

We then obtain $\psi(t_0)$. in what follows.

Recall that $\psi(t) = e^{i\hat{H}(t-t_c)}\psi(t_c)$; for the remainder of the section.

It follows that $\dot{q} = \infty$: and the result follows.

Suppose t_i, is used in the estimate below.

Note that $dt < 0$. holds throughout.

Here (t_a, t_f); by the previous lemma.

In this case τ_d: as claimed.

The value of (τ_c, τ_d), in what follows.

We then obtain $q(\tau)$. for the remainder of the section.

Recall that K_Z; and the result follows.

It follows that $dt = -d\tau$: is used in the estimate below.

Suppose $d\tau > 0$, holds throughout.

Note that $\tau = \tau_c$. by the previous lemma.

Here $\tau_d - \tau_c$; as claimed.

In this case $\tau_c + n\epsilon$: in what follows.

The value of K_0, for the remainder of the section.

We then obtain $(\tau - \epsilon, \tau + \epsilon)$. and the result follows.

Recall that $\tau_c + \epsilon, \tau_d$; is used in the estimate below.

It follows that $\tau_c + N\epsilon = \tau_d$: holds throughout.

Suppose t_{in}, by the previous lemma.

Note that $\phi_p(t)$. as claimed.

Here $T = \Delta T = \tau_c - \tau_b = \tau_d - \tau_c > 0$; in what follows.

In this case $n = 10$: for the remainder of the section.

The value of $q_{\mathrm{F}} = 1 - \alpha$, and the result follows.

We then obtain 75%. is used in the estimate below.

Recall that $\mathrm{P}(\mathrm{ent}|Q) \geq q_{\mathrm{B}}$; holds throughout.

It follows that $\mathrm{P}(Q \in \mathcal{O}_{\mathrm{acc}}|\mathrm{sep}) \approx 0$: by the previous lemma.

Suppose σ_0, as claimed.

Note that $R_{\mu_1} \cdots R_{\mu_N} = \pm 1$. in what follows.

Here $\sigma_j^2 = 1 - T_j^2$; for the remainder of the section.

In this case $n_j = n$: and the result follows.

The value of 2.5%, is used in the estimate below.

We then obtain 4.2%. holds throughout.

Recall that $q \geq q_{\min}$; by the previous lemma.

It follows that $E_{\mathrm{W}}^{\infty} = T_{1} - T_{2} - \cdots - T_{M} + 1$: as claimed.

Suppose $n = 4$, in what follows.

Note that $\mathrm{P}(\mathrm{ent} | E_{\mathrm{W}}) \geq 97.5\%$. for the remainder of the section.

Here $\mathrm{P}(S < \mathcal{B} | \mathrm{ent})$; and the result follows.

In this case $N = 4$: is used in the estimate below.

The value of $\mathrm{P}(\mathrm{sep}) = 1/17$, holds throughout.

We then obtain $\bar{p} = 0.8$. by the previous lemma.

Recall that $n_{2} = 4$; as claimed.

It follows that 0.137: in what follows.

Suppose $q_{\mathrm{B}} \geq 70\%$, for the remainder of the section.

Note that $\mathrm{G}_{\mathrm{acc}}(Q)$. and the result follows.

Here $\mathcal{O}_{\mathrm{acc}} = \{2.25, 3\}$; is used in the estimate below.

In this case $E_{\mathrm{W}} < 0$: holds throughout.

The value of $(1 \pm T_{i})/2$, by the previous lemma.

We then obtain $S = 1$. as claimed.

Recall that ρ_{j}; in what follows.

It follows that $\bar{\rho}_{\mathrm{sep}} = \sum_{j} \rho_{j} \mathrm{P}(\rho_{j} | \mathrm{sep})$: for the remainder of the section.

Suppose $S^{\infty} \leq 1$, and the result follows.

Note that $10^{-8} \cdot \Phi_{0}$. is used in the estimate below.

Here T_{2}; holds throughout.

In this case $\{0, 1\}$: by the previous lemma.

The value of τ_{i}, as claimed.

We then obtain $n = 0$. in what follows.

Recall that $\Phi / \Phi_{0} \in [0, 0.5]$; for the remainder of the section.

It follows that $\Delta = |\omega_{q} - \omega_{r}|$: and the result follows.

Suppose $\sim \omega_{q}^{-1/2}$, is used in the estimate below.

Note that M'. holds throughout.

Here $S_{\Phi}(\omega) = 2\pi A_{\Phi}^{2} / \omega$; by the previous lemma.

In this case $\delta \omega_{q} \simeq |\partial \omega_{q} / \partial I_{c}| \delta I_{c}$: as claimed.

The value of $\cdot 10^{-3}$, in what follows.

We then obtain $\delta P_{m} \leq 1/(2\sqrt{K})$. for the remainder of the section.

Recall that $\Phi^{\ast}$; and the result follows.

It follows that $0 - 1$: is used in the estimate below.

Suppose $\Phi = \Phi_0/2$, holds throughout.

Note that f_q. by the previous lemma.

Here k_B; as claimed.

In this case $\Phi^*/\Phi_0 = 0.442$: in what follows.

The value of $|ij\rangle$, for the remainder of the section.

We then obtain $\Gamma_N = N\Gamma$. and the result follows.

Recall that $P_{|10\ldots0\rangle}$; is used in the estimate below.

It follows that $M = 24$: holds throughout.

Suppose $\overline{\tau_l}$, by the previous lemma.

Note that $\sigma_0 = \sigma_1 = 1.0$. as claimed.

Here n_i; in what follows.

In this case $\hat{\Phi}_{jkl}$: for the remainder of the section.

The value of $10^{-8}\Phi_0$, and the result follows.

We then obtain $d\vec{l}$. is used in the estimate below.

Recall that ω_m; holds throughout.

It follows that $F_{\mathrm{FB}} = -gM\gamma\dot{x}$: by the previous lemma.

Suppose $1/\gamma$, as claimed.

Note that n_c. in what follows.

Here $y_c(t)$; for the remainder of the section.

In this case f_{fb}: and the result follows.

The value of $A[\omega]$, is used in the estimate below.

We then obtain $\omega_m/\gamma \gg 1$. holds throughout.

Recall that n_m; by the previous lemma.

It follows that $n_{\mathrm{add}} < 4$: as claimed.

Suppose $A_0/2\pi = 6.7$, in what follows.

Note that ω_f. for the remainder of the section.

Here $\omega_m/2\pi = 8.14$; and the result follows.

In this case $n_m^T \simeq 205$: is used in the estimate below.

The value of $\Delta_f/2\pi = -20$, holds throughout.

We then obtain $\delta f_2/2\pi \simeq 400$. by the previous lemma.

Recall that $G/2\pi \simeq 104$; as claimed.

It follows that $n_m \simeq 2.9 \pm 0.3$: in what follows.

Suppose X_{bax}, for the remainder of the section.

Note that $x_c = \chi_c \sqrt{\kappa} x_{\mathrm{c,in}}$. and the result follows.

Here $a = \alpha(t) + \alpha_f(t) + \tilde{a}(t)$; is used in the estimate below.

In this case $-\sqrt{2} g_0 a^\dagger a$: holds throughout.

The value of $\Delta_f + \omega_m$, by the previous lemma.

We then obtain $+\omega_m$. as claimed.

Recall that $s(t)$; in what follows.

It follows that $\cos s(t) \simeq 1$: for the remainder of the section.

Suppose ϕ_+, and the result follows.

Note that $D \equiv \sqrt{A^2 + B^2 + 2AB\cos(\varphi - \varphi')}$. is used in the estimate below.

Here $\omega_c + \omega_m + \Delta_f$; holds throughout.

In this case $arg(\chi_c[-\omega_f]) = -\phi_1$: by the previous lemma.

The value of $\frac{1}{\sqrt{N}}$, as claimed.

We then obtain $\frac{\hbar^2}{mL^2}$. in what follows.

Recall that $\hat{T}(x)$; for the remainder of the section.

It follows that $\hat{W}(x_i - x_j)$: and the result follows.

Suppose $n_1 + n_2 + \cdots + n_M = N$, is used in the estimate below.

Note that $|n_1, n_2, \ldots, n_M\rangle$. holds throughout.

Here $\{\phi_i(x,t)\}_{i=1}^{M}$; by the previous lemma.

In this case $|\psi(t)\rangle$: as claimed.

The value of $C_{\bar{n}}(t)$, in what follows.

We then obtain $E^N_{\lambda\to\infty} = \frac{N^2}{2}$. for the remainder of the section.

Recall that 8.0; and the result follows.

It follows that $x_{\max} = +32$: is used in the estimate below.

Suppose $\rho^{(1)}(x_1'|x_1;t)$, holds throughout.

Note that $p = 2$. by the previous lemma.

Here $\rho^{(3)}(x_1, x_2, x_3 = x_{\mathrm{ref}}; t)$; as claimed.

In this case $|g^{(p)}(x_1, \ldots, x_p; t)| = 1$: in what follows.

The value of $x_1, \ldots, x_p$, for the remainder of the section.

We then obtain $g^{(2)}(0,0)$. and the result follows.

Recall that $M = 1$; is used in the estimate below.

It follows that $\sim 10^{-6}$: holds throughout.

Suppose $R_{ic}(t)$, by the previous lemma.

Note that $R_c(t)$. as claimed.

Here V_{ex}^{oc}; in what follows.

In this case $V_{ex} = \frac{dR(t)}{dt}$: for the remainder of the section.

The value of $N = 5$, and the result follows.

We then obtain $\rho^{(2)}(x_1, x_2)$. is used in the estimate below.

Recall that $t = 0.0$; holds throughout.

It follows that 5.0: by the previous lemma.

Suppose $\rho^{(3)}(x_1, x_2, x_3 = 0.0)$, as claimed.

Note that $x_1 = -x_2$. in what follows.

Here $\mathbb{Z}_2$; for the remainder of the section.

In this case $W^{\pm}, Z$: and the result follows.

The value of $O(3)$, is used in the estimate below.

We then obtain $n = c^{\dagger}c$. holds throughout.

Recall that s_{xy}^{k}; by the previous lemma.

It follows that $V = 2t$: as claimed.

Suppose $[Q_x, H^{\mathbb{Z}_2}] = 0$, in what follows.

Note that $H^{U(1)}$. for the remainder of the section.

Here $H = H_e + H_o$; and the result follows.

In this case $1, \cdots, 2N_t$: is used in the estimate below.

The value of s_{xy}^{-}, holds throughout.

We then obtain w_A. by the previous lemma.

Recall that $[C_i]$; as claimed.

It follows that $(2+1)-$: in what follows.

Suppose $G_x = 0$, for the remainder of the section.

Note that $\bar{\psi}\psi = \sum_x (-1)^x n_x$. and the result follows.

Here χ_{EE}; is used in the estimate below.

In this case 2×2: holds throughout.

The value of $m = 2^{N_f}$, by the previous lemma.

We then obtain $A^{\otimes N_f}$. as claimed.

Recall that $n = 2^{3N_f} - 2^{2N_f}$; in what follows.

It follows that $T \cdot N_t$: for the remainder of the section.

Suppose s_i, and the result follows.

Note that n_{w}.

Here 10×10; holds throughout.

In this case 0.1: by the previous lemma.

The value of $L = 8$, as claimed.

We then obtain $\beta t = 0.1$. in what follows.

Recall that $L^{1.94}$; for the remainder of the section.

It follows that $t_2 = 0$: and the result follows.

Suppose c_k, is used in the estimate below.

Note that ϵ_k. holds throughout.

Here $\mathcal{H}_{N_X}$; by the previous lemma.

In this case $\mathbf{k}_3$: as claimed.

The value of $\mathbf{k}_s$, in what follows.

We then obtain $R_3(t_1, t_2, t_3)$. for the remainder of the section.

Recall that $\mathcal{H}_{N_X \pm 1}$; and the result follows.

It follows that $\mathcal{E}_n^{(P)}$: is used in the estimate below.

Suppose $|\eta\rangle_B$, holds throughout.

Note that $\Phi_0^{(\eta)} \equiv \langle \mathrm{FS}|X_0 X_0|\eta\rangle/\sqrt{2}$. by the previous lemma.

Here X_0; as claimed.

In this case $\mathcal{S}_{\mathrm{ESA}}$: in what follows.

The value of $\phi_0^{(n=0)} = 1$, for the remainder of the section.

We then obtain $\mathcal{E}_n$. and the result follows.

Recall that $n = N/La$; is used in the estimate below.

It follows that $a = 1$: holds throughout.

Suppose t_c/L, by the previous lemma.

Note that $\mathcal{E}_n \to \mathcal{E}_n - nU$. as claimed.

Here $t_d = 4\epsilon_c$; in what follows.

In this case $1/u$: for the remainder of the section.

The value of $|\Phi_0^{(\eta)}|^2$, and the result follows.

We then obtain $\mathcal{E}_B$. is used in the estimate below.

Recall that $-\omega_1$; holds throughout.

It follows that $(-\omega_1, \omega_3) \approx (\mathcal{E}_A, \mathcal{E}_R)$: by the previous lemma.

Suppose $(-\omega_1, \omega_3)$, as claimed.

Note that $k - k_p + k_h \geq -k.$ in what follows.

Here $|k_p k_h\rangle_1$; for the remainder of the section.

In this case $z = 10$: and the result follows.

The value of V_L, is used in the estimate below.

We then obtain $a^2 \dot{D}_+.$ holds throughout.

Recall that t_m; by the previous lemma.

It follows that M_2: as claimed.

Suppose $v^2 \gg G(M_1 + M_2)/b$, in what follows.

Note that $\langle L_i \rangle = 0.$ for the remainder of the section.

Here $\mu = \hat{\boldsymbol{u}}.\hat{\boldsymbol{v}}$; and the result follows.

In this case $|\mu|$: is used in the estimate below.

The value of $\xi = P(|\mu|) - 1$, holds throughout.

We then obtain $\hat{L}_i(n).$ by the previous lemma.

Recall that $\mathrm{Tr}[A] = 1$; as claimed.

It follows that θ_a: in what follows.

Suppose $\cos \alpha_a = \hat{\boldsymbol{v}}.\hat{\boldsymbol{J}}^{\boldsymbol{a}}$, for the remainder of the section.

Note that $\lambda_i.$ and the result follows.

Here $f(M, \boldsymbol{r}, \boldsymbol{v})$; is used in the estimate below.

In this case $\boldsymbol{v}_0$: holds throughout.

The value of $M + dM$, by the previous lemma.

We then obtain $[\delta_{ij} + 3\hat{v_0}_i \hat{v_0}_j].$ as claimed.

Recall that m_{DM}; in what follows.

It follows that 2160^3: for the remainder of the section.

Suppose Ω_Λ, and the result follows.

Note that $\mu = \hat{\boldsymbol{L}}.\hat{\boldsymbol{v}}.$ is used in the estimate below.

Here $a \gtrsim 1$; holds throughout.

In this case 0.04: by the previous lemma.

The value of $\hat{\boldsymbol{e}}_1, \hat{\boldsymbol{e}}_2, \hat{\boldsymbol{e}}_3$, as claimed.

We then obtain $n^3.$ in what follows.

Recall that $D_i = \sinh 2\pi i k_i / n$; for the remainder of the section.

It follows that $|\cos \alpha| \sim 1$: and the result follows.

Suppose $\hat{\boldsymbol{J}}.\hat{\boldsymbol{e}}_2$, is used in the estimate below.

Note that $\mu_\nu = 0.$ holds throughout.

Here $\nu_R = (\nu_{eR}, \nu_{\mu R}, \nu_{\tau R})$; by the previous lemma.

In this case H_L: as claimed.

The value of $V_e = \sqrt{2} G_F (n_e - 0.5 n_n)$, in what follows.

We then obtain ν_R. for the remainder of the section.

Recall that $\nu_{\alpha L} \leftrightarrow \nu_{\alpha R}$; and the result follows.

It follows that $V_e + \dot{\phi} (> 0)$: is used in the estimate below.

Suppose r_0, holds throughout.

Note that $B(r) = B_0 (r_0 / r)^3$. by the previous lemma.

Here Y_e; as claimed.

In this case $\ddot{\phi}(r)$: in what follows.

The value of $r \lesssim 100$, for the remainder of the section.

We then obtain $r = r_0$. and the result follows.

Recall that $x = \mu, \tau$; is used in the estimate below.

It follows that $\mathrm{m}^{-1} \gtrsim \dot{\phi} \gtrsim -2$: holds throughout.

Suppose $B_0 = 10^{12}$, by the previous lemma.

Note that ν_τ. as claimed.

Here 3×10^4; in what follows.

In this case $|U_{e3}| \approx 0.02$: for the remainder of the section.

The value of $|U_{e1}|^2 \Phi_{ix}$, and the result follows.

We then obtain $\dot{\phi} > 0$. is used in the estimate below.

Recall that 8.2×10^{-15}; holds throughout.

It follows that $\nu_{\alpha L} \leftrightarrow \nu'_R$: by the previous lemma.

Suppose $15 M_\odot$, as claimed.

Note that $\Phi_{ie} \to 0$. in what follows.

Here Φ_e^{IO}; for the remainder of the section.

In this case 30%: and the result follows.

The value of 100%, is used in the estimate below.

We then obtain N_t. holds throughout.

Recall that F_i; by the previous lemma.

It follows that $f(R)$: as claimed.

Suppose $\tilde{g}_{\mu\nu}$, in what follows.

Note that $\{\psi\}$. for the remainder of the section.

Here $(-, +, +, +)$; and the result follows.

In this case $-3\mu^4/2\lambda$: is used in the estimate below.

The value of $\chi \ll M$, holds throughout.

We then obtain $\tilde{\chi}^2/M^2$. by the previous lemma.

Recall that $(\alpha_{hs}\tilde{\chi} + \lambda_{hs}\tilde{\chi}^2)\tilde{\phi}^2$; as claimed.

It follows that $\rho_\psi = y\tilde{\phi}_m\langle\bar{\tilde{\psi}}\tilde{\psi}\rangle$: in what follows.

Suppose $\tilde{\phi}_m$, for the remainder of the section.

Note that $v^2 = 6\mu^2/\lambda$. and the result follows.

Here $1/M^2$; is used in the estimate below.

In this case $\mu_\chi \ll \mu$: holds throughout.

The value of $\rho_\psi/(\mu^2 v^2)$, by the previous lemma.

We then obtain $\mu^2 v^2$. as claimed.

Recall that $A^2(\chi)$; in what follows.

It follows that $b = 1$: for the remainder of the section.

Suppose μ_χ, and the result follows.

Note that $\tilde{\chi} = \tilde{\chi}_m + \delta\tilde{\chi}$. is used in the estimate below.

Here $\delta\tilde{\phi}$; holds throughout.

In this case μ_χ^2/μ^2: by the previous lemma.

The value of v^2/M^2, as claimed.

We then obtain μ^2. in what follows.

Recall that $\sim \sqrt{\rho_\psi}/M$; for the remainder of the section.

It follows that $\beta \sim 1$: and the result follows.

Suppose $\sim 10^{-26}$ g/cm^3, is used in the estimate below.

Note that ρ_ψ. holds throughout.

Here $\delta\chi = s\cos\theta$; by the previous lemma.

In this case $r > R$: as claimed.

The value of x', in what follows.

We then obtain x_{right}. for the remainder of the section.

Recall that $[n] = \{1, \ldots, n\}$; and the result follows.

It follows that x_j: is used in the estimate below.

Suppose $(c_1, \ldots, c_k) \in \mathbb{R}^k$, holds throughout.

Note that $O(kn \log n)$. by the previous lemma.

Here $C_i = Y_i$; as claimed.

In this case $x_{i,r-1}$: in what follows.

The value of $\vec{C}'$, for the remainder of the section.

We then obtain $M(\vec{x}) = (y_1, ..., y_k)$. and the result follows.

Recall that $\min_{j \in \{1,2,...,k\}}\{d(x, y_j)\} = d(x, y_i)$; is used in the estimate below.

It follows that $\mathbb{P}\mathrm{r}[E]$: holds throughout.

Suppose y_j, by the previous lemma.

Note that $O(1)$. as claimed.

Here $C'_{i'-1}$; in what follows.

In this case $d_{i,j}$: for the remainder of the section.

The value of $x_{i-1,\hat{1}} \geq x_{i-1,\hat{2}} \geq \ldots \geq x_{i-1,\hat{n'}}$, and the result follows.

We then obtain X'_{i-1}. is used in the estimate below.

Recall that $> 0.4 \cdot D(C_i)$; holds throughout.

It follows that $d(x_{i,l}, c_{i-1}) > 5d(x_{i,l}, c_i)$: by the previous lemma.

Suppose $D(C_i)$, as claimed.

Note that $\gamma \geq 1$. in what follows.

Here (γ/γ'); for the remainder of the section.

In this case $d(x_i, x') > d(x_i, c_i)$: and the result follows.

The value of $1 \leq i < j \leq n$, is used in the estimate below.

We then obtain $\{c_i, ..., x_{i,r}, ..., c_j\}$. holds throughout.

Recall that $d(x_{i,r}, x_{i+1,l}) > (\gamma - 1)d(x_{i,r}, c_i)$; by the previous lemma.

It follows that $k - 2$: as claimed.

Suppose $d(x_{i,l}, c_i) = y$, in what follows.

Note that $[0, 2 - \sqrt{3}]$. for the remainder of the section.

Here $\mathbf{x}_i$; and the result follows.

In this case $u(p)$: is used in the estimate below.

The value of $\Lambda = \mathrm{diag}([\sigma_1^2, ...\sigma_d^2])$, holds throughout.

We then obtain $\mathbb{E}(I^k)$. by the previous lemma.

Recall that $K = 3$; as claimed.

It follows that $Q \in \mathbb{R}^{N \times K}$: in what follows.

Suppose $\bar{P} = \frac{1}{M} \sum_{m=1}^{M} P_{m,*,*} \in \mathbb{R}^{N \times K}$, for the remainder of the section.

Note that $j = 1, ..., J$. and the result follows.

Here $2(N_l - 1)\Delta^2_{\mathcal{D}_l} \in \mathcal{F}$; is used in the estimate below.

In this case 95.80 ± 1: holds throughout.

The value of 90.95 ± 2, by the previous lemma.

We then obtain $50.80 \pm 2.$ as claimed.

Recall that 88.09 ± 2; in what follows.

It follows that 92.89 ± 2: for the remainder of the section.

Suppose 0.8, and the result follows.

Note that $87.06 \pm 1.9.$ is used in the estimate below.

Here 91.44 ± 1.2; holds throughout.

In this case $\gamma \in \{0.5, 0.7, 0.9\}$: by the previous lemma.

The value of $\{\mathbf{x}_i, y_i\}_{i=1}^{N_l}$, as claimed.

We then obtain $\mu_-.$ in what follows.

Recall that $\tilde{X}_i^- \sim \mathcal{N}(\mu_-, \sigma^2)$; for the remainder of the section.

It follows that $\texttt{Var}(I_i^+)$: and the result follows.

Suppose $2(N_l - 1)\Delta_{\mathcal{D}}^2 \in \mathcal{F}$, is used in the estimate below.

Note that $1, ...K.$ holds throughout.

Here 95.74 ± 1; by the previous lemma.

In this case 57.50 ± 2: as claimed.

The value of 81.03 ± 3, in what follows.

We then obtain $T \leq 5.$ for the remainder of the section.

Recall that $\textit{89.14} \pm 2$; and the result follows.

It follows that $\textit{82.91} \pm 3$: is used in the estimate below.

Suppose h_z, holds throughout.

Note that $N_p.$ by the previous lemma.

Here $p_k \in \{+1, -1\}$; as claimed.

In this case $\mathcal{E}' = \{e'_k\}_{k=1}^{N_e}$: in what follows.

The value of $b_k = p_k$, for the remainder of the section.

We then obtain $G(\boldsymbol{\theta})$. and the result follows.

Recall that s_k; is used in the estimate below.

It follows that $h_z < 0$: holds throughout.

Suppose $\nabla \cdot \mathbf{v} = \partial_x \mathbf{v}_x + \partial_y \mathbf{v}_y$, by the previous lemma.

Note that $N_e(\mathbf{x}) \doteq \sum_k \delta(\mathbf{x} - \mathbf{x}'_k).$ as claimed.

Here $|\det(\mathtt{J})| < 1$; in what follows.

In this case $|\det(\mathtt{J})| \in [0, \infty)$: for the remainder of the section.

The value of $\det(\mathtt{J}(e_k)) = s_k^2$, and the result follows.

We then obtain $\mathbf{x}_p.$ is used in the estimate below.

Recall that $\mathbf{x}, t$; holds throughout.

It follows that $\boldsymbol{\theta} \equiv \boldsymbol{\omega} = (\omega_1, \omega_2, \omega_3)^\top$: by the previous lemma.

Suppose $\det(\mathtt{J}) = 1$, as claimed.

Note that $R(\boldsymbol{\theta})$. in what follows.

Here $dt = 4$; for the remainder of the section.

In this case $\lambda_{\text{def}} = 5$: and the result follows.

The value of > 80, is used in the estimate below.

We then obtain 10–20. holds throughout.

Recall that $h_z \approx 0$; by the previous lemma.

It follows that $\mathbf{x}(t)$: as claimed.

Suppose $\boldsymbol{\theta} = \boldsymbol{\omega} = (\omega_1, \omega_2, \omega_3)^\top$, in what follows.

Note that $t_k \|\boldsymbol{\omega}\|$. for the remainder of the section.

Here $\det(\mathtt{J}_k) \approx 1 + 3(x_k\omega_2 - y_k\omega_1)t_k$; and the result follows.

In this case $\mathbf{x}_k' = (1 - t_k h_z)\mathbf{x}_k$: is used in the estimate below.

The value of $SO(2)$, holds throughout.

We then obtain $\beta_k = 1 + t_k s$. by the previous lemma.

Recall that T_K; as claimed.

It follows that w_0: in what follows.

Suppose $T = 20$, for the remainder of the section.

Note that 0.432 ± 0.018. and the result follows.

Here 1.918 ± 0.006; is used in the estimate below.

In this case 1.912 ± 0.021: holds throughout.

The value of 0.485 ± 0.020, by the previous lemma.

We then obtain 1.927 ± 0.006. as claimed.

Recall that 5.596 ± 0.030; in what follows.

It follows that 4.597 ± 0.070: for the remainder of the section.

Suppose $\mathcal{P}, \mathcal{Q}$, and the result follows.

Note that $\mathcal{F}^{-1}$. is used in the estimate below.

Here $K^d \times \mathrm{C} \times \mathrm{C}$; holds throughout.

In this case $K = 90$: by the previous lemma.

The value of s_t, as claimed.

We then obtain $K_t \leftarrow K_t + 1$. in what follows.

Recall that (K_t, R_{curr}); for the remainder of the section.

It follows that $g(K, s_p) > \alpha$: and the result follows.

Suppose K_{t+1}, is used in the estimate below.

Note that 918 ± 10. holds throughout.

Here 0.155 ± 0.003; by the previous lemma.

In this case 13.53 ± 13.08: as claimed.

The value of 2.844 ± 0.070, in what follows.

We then obtain 0.276 ± 0.083. for the remainder of the section.

Recall that 0.316 ± 0.015; and the result follows.

It follows that 1.130 ± 0.023: is used in the estimate below.

Suppose 1.290 ± 0.022, holds throughout.

Note that 5.627 ± 0.073. by the previous lemma.

Here x_1; as claimed.

In this case 128×128: in what follows.

The value of $f : X \to \mathbb{R}$, for the remainder of the section.

We then obtain $\mathsf{Hom}(-, \mathbb{R}_+)$. and the result follows.

Recall that (F, μ, η); is used in the estimate below.

It follows that $a \in X$: holds throughout.

Suppose $[\phi] = 1$, by the previous lemma.

Note that (D, μ, η). as claimed.

Here $\mu^D : D^2 \to D$; in what follows.

In this case $\mathbb{R}_+$: for the remainder of the section.

The value of $\otimes^L : LF \to FL$, and the result follows.

We then obtain $[\phi]$. is used in the estimate below.

Recall that $\otimes^L_X$; holds throughout.

It follows that $\otimes^L : LD \to DL$: by the previous lemma.

Suppose $(\cdot)$, as claimed.

Note that $f : X \to Y$. in what follows.

Here $\{1, \ldots, n\}$; for the remainder of the section.

In this case $\otimes^M_X$: and the result follows.

The value of $d_1, \ldots, d_n$, is used in the estimate below.

We then obtain $\phi' : F' \to F''$. holds throughout.

Recall that $\phi : F \to G$; by the previous lemma.

It follows that $\phi : \Phi \to \Psi$: as claimed.

Suppose ϕ_S, in what follows.

Note that $I_O : O \to O$. for the remainder of the section.

Here $\tau : T \to U$; and the result follows.

In this case $\sigma, \tau : S \to T$: is used in the estimate below.

The value of $\Psi\sigma = \Psi\tau$, holds throughout.

We then obtain $\mu^M : MM \to M$. by the previous lemma.

Recall that $\otimes_X^L(\bar{f})(\bar{a}) = 0$; as claimed.

It follows that $\mathsf{supp} \otimes_X^L (\bar{f}) = \prod_{i=1}^{n} \mathsf{supp}\, f_i$: in what follows.

Suppose $Fh(f) \in FY$, for the remainder of the section.

Note that $f_i \in FFX = \mathbb{R}^{\mathbb{R}^X}$. and the result follows.

Here $p(\mathbf{X})$; is used in the estimate below.

In this case $\{X_i, Z_i, Y_i, R_i\}_{i=1}^{N_s}$: holds throughout.

The value of R_i, by the previous lemma.

We then obtain $N_{pop} >> N_s$. as claimed.

Recall that $D(\cdot, \cdot)$; in what follows.

It follows that $\mu = \frac{1}{N_{pop}} \sum_i Y_i$: for the remainder of the section.

Suppose $g(\cdot) : \mathcal{R} \mapsto [0, 1]$, and the result follows.

Note that $\phi(X)^\top \beta + (Z + \nu)$. is used in the estimate below.

Here N_s; holds throughout.

In this case $k(X_i, X_j)$: by the previous lemma.

The value of $\phi(X_i)^\top \theta = \sum_j c_j k(X_i, X_j)$, as claimed.

We then obtain i^{th}. in what follows.

Recall that $X_i = X_j$; for the remainder of the section.

It follows that l^{th}: and the result follows.

Suppose $\mathbf{K} = \mathbf{VAU}^\top$, is used in the estimate below.

Note that $(w^{(pop)\top} \mathbf{V}_{pop} - w_s^\top \mathbf{V}_s)$. holds throughout.

Here $exp(-||X_i - X_j||^2/b)$; by the previous lemma.

In this case $k(X_i, X_j) = e^{-\left((1-0)^2 + (0-0)^2\right)} = e^{-1} \approx 0.37$: as claimed.

The value of $k(, 2)$, in what follows.

We then obtain $N_s = 2,052$. for the remainder of the section.

Recall that $kpop + mf$; and the result follows.

It follows that $\{X_i, Z_i, Y_i, R_i\}$: is used in the estimate below.

Suppose $Y - \phi(X)^\top \beta$, holds throughout.

Note that $\mu_s(w)$. by the previous lemma.

Here $p(X = x)$; as claimed.

In this case $\phi(x)$: in what follows.

The value of $g(\cdot) : \mathbb{R} \mapsto [0, 1]$, for the remainder of the section.

We then obtain $\phi(X) = X = [\text{female, college}]$. and the result follows.

Recall that $b = 2D$; is used in the estimate below.

It follows that R^2: holds throughout.

Suppose $X_{ij}, j = 1, \ldots, p$, by the previous lemma.

Note that c_j^2. as claimed.

Here $\delta_0(.)$; in what follows.

In this case $\mathbf{B}_g$: for the remainder of the section.

The value of $\omega_{gj,k}$, and the result follows.

We then obtain $IG(.)$. is used in the estimate below.

Recall that s^2; holds throughout.

It follows that $Y_{i,k}$: by the previous lemma.

Suppose $\Phi^{-1}(.)$, as claimed.

Note that τ^2. in what follows.

Here $\text{sgn}(\cdot)$; for the remainder of the section.

In this case $n \to \infty$: and the result follows.

The value of $d_{gj}(t) = I\{P_{gj}(H_0|\mathbf{X}, \mathbf{Y}) < t\}$, is used in the estimate below.

We then obtain $PIP_{gj}^{\mathcal{M}_S}$. holds throughout.

Recall that $\mathcal{M}_{\mathcal{S}}$; by the previous lemma.

It follows that $r \leftarrow 1$: as claimed.

Suppose σ_Y^2, in what follows.

Note that $n = 500$. for the remainder of the section.

Here $\beta_{gj,k} = 0$; and the result follows.

In this case n_{test}: is used in the estimate below.

The value of $\vec{M}_{gj}$, holds throughout.

We then obtain $k \neq k'$. by the previous lemma.

Recall that Σ_{kk}^{-1}; as claimed.

It follows that $\xi_{gj}|\gamma_{gj}, d_0, d_1 \sim N(d_0 + d_1 A_{gj}, 1)_-$: in what follows.

Suppose (σ^2, τ^2), for the remainder of the section.

Note that $\mathbf{X}^T \mathbf{X} = n\mathbf{I}$. and the result follows.

Here $M = (1 - D_n)\hat{\beta}^{LS}_{gj,k}$; is used in the estimate below.

In this case $\hat{\beta}^{LS}_{gj,k} < 0$: holds throughout.

The value of $\sqrt{n}|\hat{\beta}^{LS}_{gj,k}| = O_p(1)$, by the previous lemma.

We then obtain $l_{gj,k} \to 1$. as claimed.

Recall that τ_b; in what follows.

It follows that $\mathcal{Z} \subseteq \mathcal{V}$: for the remainder of the section.

Suppose $\mathbf{Y}_i$, and the result follows.

Note that $\mathcal{D} = (\mathcal{T}, h)$. is used in the estimate below.

Here $Z_5 = Z_6 = d$; holds throughout.

In this case $|\mathcal{Z}|$: by the previous lemma.

The value of $u, v \in P_{m-1}$, as claimed.

We then obtain $b = \min\{h(v) - h(\mathrm{Pa}_v)\}$. in what follows.

Recall that $\max_{v \in \mathcal{Z}} \|\mathbf{S}(v)\|_{\mathrm{op}} \in O(1)$; for the remainder of the section.

It follows that $O_{\mathbb{P}}(\cdot)$: and the result follows.

Suppose $p^{-1}\mathbb{E}[\|\mathbf{Y}_i\|^2 | Z_1, \ldots, Z_n]$, is used in the estimate below.

Note that $n = 5079$. holds throughout.

Here $O(npr)$; by the previous lemma.

In this case $\mathbf{Y}_1, \ldots, \mathbf{Y}_n \in \mathbb{R}^p$: as claimed.

The value of $u \neq v$, in what follows.

We then obtain $w' \in P_m$. for the remainder of the section.

Recall that P_0; and the result follows.

It follows that $\hat{h}(w_m) = \hat{\alpha}(u_m, v_m) \geq \hat{h}(w_{m+1})$: is used in the estimate below.

Suppose $l \in B$, holds throughout.

Note that z_l. by the previous lemma.

Here $S_v \neq [n]$; as claimed.

In this case $A' \subseteq S_v$: in what follows.

The value of $m^* + 1$, for the remainder of the section.

We then obtain $(\mathcal{Z}, d_{\mathcal{Z}})$. and the result follows.

Recall that Δ_1; is used in the estimate below.

It follows that $\mathbf{E}_j \in \mathbb{R}^p$: holds throughout.

Suppose $|\mathcal{Z}| - 1$, by the previous lemma.

Note that $p = 1000.$ as claimed.

Here $d_{\cos}(\cdot, \cdot)$; in what follows.

In this case $\hat{\alpha}_{\cos}(i, j)$: for the remainder of the section.