

EVOLUTION AND EVOLVABILITY IN CHANGING ENVIRONMENTS

By

Rosangela Canino-Koning

A DISSERTATION

Submitted to
Michigan State University
in partial fulfillment of the requirements
for the degree of

Computer Science – Doctor of Philosophy

2017

ABSTRACT

EVOLUTION AND EVOLVABILITY IN CHANGING ENVIRONMENTS

By

Rosangela Canino-Koning

prep a layman's abstract if I have time

rework abstract

Copyright by
ROSANGELA CANINO-KONING
2017

This thesis is dedicated to Kendall Koning.

ACKNOWLEDGMENTS

Your acknowledgments here

acknowledgments

TABLE OF CONTENTS

LIST OF TABLES	ix
LIST OF FIGURES	x
KEY TO ABBREVIATIONS	xv
TODO LIST	xvi
CHAPTER 1 INTRODUCTION - CHANGE, ADAPTATION AND THE EVOLUTION OF EVOLVABILITY	1
1.1 Evolvability and Evolutionary Potential - Why Study It	1
1.2 What is Evolvability (and why is it so hard to pin down)	2
1.3 Changing Environments and Evolvability	3
1.4 Historical Conceptions of Evolvability	4
1.4.1 Modern Synthesis	4
1.4.2 Evolvability as a Distinct Concept	6
1.4.3 Theoretical Frameworks for the Evolution of Evolvability	7
1.5 So, what do I mean by Evolvability?	8
1.5.1 What is Modularity?	9
1.5.1.1 Measuring Modularity	10
1.5.2 What is Robustness?	11
1.5.2.1 Measuring Robustness to Mutation	12
1.5.3 Predicting Short-Term Evolvability with Landscape Metrics	14
1.5.4 Expected Value of Fitness Landscapes	16
1.6 Digital Evolution	16
1.6.1 Avida	17
CHAPTER 2 CHANGING ENVIRONMENTS PROMOTE RAPID ADAPTATION IN DIGITAL ORGANISMS	20
2.1 Background	20
2.1.1 Evolvability and Genetic Architecture	21
2.1.1.1 Mutational Landscapes	22
2.2 Methods	23
2.2.1 Experimental Design	23
2.2.1.1 Cyclic and Stochastic Changing Environments	23
2.3 Results and Discussion	25
2.3.1 Cyclic Changing Environments	26
2.3.1.1 Evolutionary History and Population Structure	26
2.3.1.2 Genetic Architecture	28
2.3.1.3 Nearby mutational landscape	29
2.3.2 Stochastic Changing Environments	32
2.4 Conclusion	35

CHAPTER 3 CHANGING ENVIRONMENTS AND THE EVOLUTION OF HORIZONTAL GENE TRANSFER	39
3.1 Background	39
3.1.1 Origins of Horizontal Gene Transfer in nature	39
3.2 Methods	40
3.2.1 HGT in Avida	40
3.2.1.1 Environmental Conditions	41
3.2.2 Experimental Design	42
3.3 Results and Discussion	44
3.3.1 Changing environments elevate HGT use	44
3.3.2 HGT derives most benefit from on-cycle fragments, but not all	45
3.3.2.1 Expected Fitness Effects	46
3.4 Conclusion	49
CHAPTER 4 HORIZONTAL GENE TRANSFER VS OTHER TYPES OF MUTATION	51
4.1 Background	51
4.2 Methods	52
4.2.1 Experimental Design	52
4.2.1.1 Alternative Mutagenic Instructions	52
4.2.1.2 Comparing HGT to Other Mutation Types	54
4.3 Results and Discussion	55
4.3.1 Other mutation types are not elevated in response to HGT	55
4.3.2 Mutation fitness effect correlate with mutagen use	57
4.3.3 HGT mutations increase evolved probability of beneficial phenotype switching	57
4.4 Conclusion	58
CHAPTER 5 CHANGING ENVIRONMENTS AND LONG TERM EVOLVABILITY	59
5.1 Background	59
5.2 Methods	60
5.2.1 Experimental Design	60
5.2.1.1 Simple Cyclic Changing Environments	60
5.2.1.2 Complex Cyclic Changing Environment	61
5.2.1.3 Complex Cyclic Changing Environment with HGT mutagen	62
5.2.2 Measuring Task Discovery and Task Performance	63
5.2.2.1 Task Performance	63
5.2.2.2 Task Discovery	63
5.3 Results and Discussion	63
5.3.1 Long-Term Evolvability in Simple Cyclic Changing Environments . .	65
5.3.1.1 Benign changing environments outperform harsh environments in task discovery	65
5.3.1.2 Benign changing environments outperform harsh environments in task performance	66

5.3.1.3	Harsh changing environments drive populations across the mutational landscape	67
5.3.2	Long Term Evolvability and Complex Changing Environments	68
5.3.2.1	Benign changing environments outperform harsh environments in task discovery and task performance	68
5.3.3	Long Term Evolvability and Horizontal Gene Transfer	69
5.3.3.1	New environments do not promote the use of HGT	70
5.3.3.2	Fragment grazing bonus reduces task discovery and task performance	70
5.3.3.3	Benign changing environments outperform harsh environments in task discovery and task performance in populations with HGT	73
5.4	Conclusion	73
CHAPTER 6 CONCLUSION		74
6.1	Limitations of Cyclic Changing Environments	74
APPENDICES		76
APPENDIX A	Experimentally Deriving Parameters for Changing Environment Cycle Lengths	77
APPENDIX B	Experimentally Deriving Parameters for HGT Recombination Probability and Bonus Levels	80
APPENDIX C	Changing Environments Elevate All Instruction Use Due to Repeated Bottlenecks	83
BIBLIOGRAPHY		84

LIST OF TABLES

Table 2.1: Experimental Treatments	24
Table 3.1: Experimental Treatments - Evolution of HGT	43
Table 3.2: Experimental Treatments - Effects of HGT	44
Table 4.1: Experimental Treatments - Mutation Types	55
Table 5.1: Experimental Treatments - Simple Cyclic Changing Environment	61
Table 5.2: Experimental Treatments - Complex Cyclic Changing Environment	62
Table 5.3: Experimental Treatments - Complex Cyclic Changing Environments with HGT	64

LIST OF FIGURES

Figure 2.9: A survey of the single-step mutational neighborhood in the stochastic changing environment around organisms that performed the fluctuating task. Again, in the static and benign treatments, values are comparable to the cyclic changing environment [TODO stats]. However, in the harsh treatment, the means for both loss of the fluctuating task (EQU) and loss of both task were slight reduced [TODO stats]. This result indicates that in the context of a harsh treatment, stochastic environmental change is less effective at moving organisms to areas of the fitness landscape where they can more easily switch task expression.	36
Figure 2.10: A survey of the two-step mutational neighborhood in the stochastic changing environment of the organisms that lost EQU function in the one-step survey. Similarly to the result in Fig 2.9, we found that the fraction of organisms regaining the fluctuating task from a single additional mutation in the harsh treatment were reduced compared to the cyclic harsh treatment[TODO stats]. This result confirms that the harsh stochastic environment is less effective than the cyclic harsh at promoting evolvability.	37
Figure 2.11: Number of functional and vestigial sites by treatment in a stochastic changing environment. The vestigial site counts remain comparable to the cyclic environment [TODO stats], however, there was a reduction in functional site counts for XOR+EQU overlapping sites in the stochastic harsh environment as compared to the cyclic harsh environment [TODO stats], as well as an overall reduction in the number of vestigial sites.	38
Figure 3.1: The HGT process. Organisms can execute instructions that trigger uptake from the environment. When uptake occurs, there is an experimenter-defined chance that either it will yield a boost to speed of execution or, alternatively, that the fragment will be integrated into the genome.	42
Figure 3.2: bleh	46
Figure 3.3: bleh	47
Figure 3.4: bleh	48
Figure 3.5: bleh	48
Figure 3.6: bleh	49
Figure 4.1: bleh	56

Figure 4.2: bleh	57
Figure 4.3: bleh	58
Figure 5.1: Number of new logic-77 tasks discovered beginning in phase 2 . .	66
Figure 5.2: Number of new logic-77 tasks performed beginning in phase 2 . .	67
Figure 5.3: Number of new logic-77 tasks discovered over time. [TODO DESCRIPTION AND STATS]	68
Figure 5.4: Number of logic-77 tasks performed over time. [TODO DESCRIPTION AND STATS]	69
Figure 5.5: Number of new logic-77 tasks discovered post-reward. [TODO DESCRIPTION AND STATS]	69
Figure 5.6: HGT Use, pre and post introduction of new logic-77 tasks. [TODO DESCRIPTION AND STATS]	70
Figure 5.7: Task Performance, with and without HGT Grazing Bonus. [TODO DESCRIPTION AND STATS]	71
Figure 5.8: Task Discovery, with and without HGT Grazing Bonus. [TODO DESCRIPTION AND STATS]	71
Figure 5.9: HGT-Bonus vs Task Performance and Discovery. [TODO DESCRIPTION AND STATS]	72
Figure 5.10: HGT-noBonus vs Task Performance and Discovery. [TODO DESCRIPTION AND STATS]	72
Figure A.1: Survey of task execution for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]	78
Figure A.2: Survey of task execution for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]	79
Figure B.1: Survey of hgt use for a series of recombination probabilities. [TODO DESCRIPTION AND STATS]	81
Figure B.2: Survey of task execution for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]	82

Figure B.3: Survey of comparable bonus values for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]	82
Figure C.1: bleh	83

KEY TO ABBREVIATIONS

CE	Changing Environment
CV_A	Additive Genetic Coefficient of Genetic Variation
D_g	Genomic Diffusion Rate
D_p	Phenotypic Diffusion Rate
h^2	Narrow-sense Heritability
H^2	Broad-sense Heritability
HGT	Horizontal Gene Transfer
m_S	Spatial Modularity
ME	Mutational Event
MRI	Mutation Rate Increase
r_P	Phenotypic Robustness
r_G	Genotypic Robustness
MWW	Mann Whitney Wilcoxon

TODO LIST

■ prep a layman's abstract if I have time	ii
■ rework abstract	ii
■ acknowledgments	v
■ revise this	3
■ MORE HERE about other things CE can or can't do	4
■ remove "proposal"	8
■ revise this paragraph for my thesis statement.	8
■ what goals? (see following sentence)	17
■ clarify	22
Figure: A diagram of how the environments are placed. stochastic as jumbled, cyclic as periodic. Harsh is red/blue, Benign is white/blue.	23
■ look for un-anchored "this"s	26
■ fix figure below so that it's two separate figures. Separate the mean entropy to its own figure.	26
■ clarify wording	28
■ polish this figure, task layout, make three separate figures	29
■ stats for figure	29
■ add figure of the nearby landscape showing how mutations relate 1 and 2 steps out., a couple of nice figures showing both changes in detrimental mutations and phenotypic distribution.	29
■ fix wording to clarify	29
■ Todo, clarify per Charles's comment above	31
■ stats for figure	31
■ stats for figure	31
■ Add to caption per Charles's comment above	31
■ add stats for this paragraph	31
■ TODO - add paragraph describing/speculating about why Benign loses EQU almost as well as Harsh	31
■ stats	32
■ STATS for drop in SCE-Harsh D_p	32
■ reword below	32
■ stats	33
■ stats	33
■ stats	33
■ stats	33
■ stats	33
■ stats	34
■ stats	34
■ add a sentence emphasizing that it's an indirect effect.	36
■ More about how this question is still open	39
■ look for and hunt down passive voice	41
■ Citation TODOs	41

orange	rename table	44
green	stats	45
green	stats	46
green	stats	47
green	stats	48
green	stats	49
orange	review the below. too strong? too informal?	50
orange	[TODO CITE]	51
orange	stats	56
orange	stats	56
green	stats	56
orange	caption	56
green	stats	57
orange	caption	57
green	stats	58
orange	caption	58
magenta	ch5 - background - fill this in about the prior research done in this area	59
magenta	CH5 methods - more details about each treatment	60
orange	update table to match HGT	62
magenta	TODO fill this in with more detail of each treatment	62
magenta	TODO fill this in with more detail of each treatment	63
red	correct second half of the table with info about the bonus structures B0 and B0.8	64
orange	stats	65
green	stats	65
orange	caption	65
orange	stats	65
orange	make a figure to highlight this?	65
orange	rename the populations A and B	65
orange	stats	66
orange	stats	66
orange	stats	66
orange	stats	66
green	stats	66
orange	caption	66
orange	stats	67
orange	stats	67
orange	stats	67
orange	REPLACE THIS WITH THE ONE THAT HAS THE VIOLIN PLOT?	67
green	stats	67
orange	caption	67
green	stats	68
orange	caption	68
green	stats	68
orange	caption	68
orange	stats	69

■ move below to the introduction about why we care about HGT	69
■ stats	70
■ stats	70
■ caption	70
Figure: HGT Use Bonus 0, split by pre and post introduction of new logic-77 tasks	70
■ stats	70
■ hypothesis here?	70
■ stats	70
■ caption	70
■ stats	70
■ caption	70
■ stats	71
■ caption	71
■ stats	71
■ caption	71
■ stats	73
■ fill this in, recapitulating the argument and framing about balancing of fitness effects.	73
■ FILL THIS IN - Synthesis of all results and how it all fits together to paint a picture of how CE works.	74
■ tweak below?	75
■ stats	77
■ caption	77
■ stats	77
■ caption	77
■ stats	80
■ caption	80
■ stats	81
■ caption	81
■ fix up figure to include CE uptake value, plus less terrible	81
■ stats	81
■ caption	81
■ rework figure	83
■ stats	83
■ caption	83

CHAPTER 1

INTRODUCTION - CHANGE, ADAPTATION AND THE EVOLUTION OF EVOLVABILITY

1.1 Evolvability and Evolutionary Potential - Why Study It

The evolutionary potential of a genome is a controversial and nuanced topic, ultimately measurable only in retrospect once the evolutionary success of its descendants is known. Questions relating to evolutionary potential, however, are some of the biggest in Evolutionary Biology: What selective pressures drive organisms to become more evolvable? What aspects of genetic architecture influence evolutionary potential? How do we go about predicting longer-term evolutionary success? And how do features of the environment, such as complexity, change, and periodicity drive and constrain movement across mutational landscapes?

The evolution of sex, multi-cellularity, and other major transitions are characterized by significant changes in genetic architecture that appear to have facilitated the transitions[1]. The adaptive radiations that accompanied Metazoan evolution were also accompanied by changes in genetic architecture that were carried along as species diversified and colonized new ecological niches[2]. The vast diversity of species and their complex ecological interplay depends fundamentally on the ability of populations to not only adapt to their environment, but also create new niches and rapidly explore and exploit their environment as it changes around them. Evolvability has many subtle forms that are produced in different types of changing environment.

Within evolutionary computation, evolvability is also fundamental. The “representation problem”, which influences every aspect of evolutionary search, can be characterized as a problem of how to design the underlying genetic encoding such that genomes can not only express complex solutions, but can also be mutated in meaningful ways[3]. In particular,

good designs for genetic representations often involve increasing the probability that a recombination between potential solutions can produce a result that is not only viable, but more fit than either parent. The entire goal of the representation problem is to improve evolvability so that better solutions can be found. By definition, systems that exhibit good characteristics in evolvability produce good solutions more quickly, while avoiding premature convergence[4] so adaptive evolution continues to as high a level as possible. Beyond the representation problem, many of the barriers to complexity are actually barriers to evolvability.

1.2 What is Evolvability (and why is it so hard to pin down)

In its most abstract sense, evolvability appears to be a simple concept: the ability of genetic systems to produce adaptive variation. However, the devil is in the details. How exactly do genetic systems generate adaptive variation? How do we measure this potential? Should all forms of variation count as evolvability? At what time-scales does evolvability act? And finally, how did it evolve in the first place? That is, are evolvable features under some form of direct selection, or are they by-products of other processes?

Evolvability, in its details, must mean different things at different evolutionary scopes and timescales. Depending on your perspective, evolvability can describe the response to selection at the population level[5, 6], the ability of populations to adapt to changing conditions[7], larger phenomena such as variability generation[8], exploration of neutral spaces and robustness[9, 10], generation of novel features[11, 12], or even the potential to generate the larger clade-level innovations[2] and major transitions[1]. Beyond that, there is a lot of confusion and controversy about the definitions and components of evolvability even within any one of these scopes[13].

Finally, it is unclear whether evolvability is acted upon by direct selection, or whether it is a byproduct of other traits that are selected upon, or some combination of the two. At the individual level, its possible that some traits that support evolvability, such as robustness of

developmental or cell processes[2] could have been selected for directly in response to adverse environmental conditions. However, at the population level, traits like neutral variation generation are more likely to have hitchhiked on the genomes of the adaptive variants that they produced. Finally, at the clade level, genetic structures that produced populations of adaptive variants with robust and flexible genetic architectures would have been more successful at adaptive radiations[3], and thus go on to found whole branches of life with those traits[2].

Of course, we must be careful when invoking selection at higher levels than the individual. While there is some evidence to support clade-level selection in the evolution of evolvability [14], caution should be applied when attributing evolutionary outcomes to higher levels of selection when random chance or lower levels of selection are adequately explanatory. Specifically, we need to be careful to avoid falling into the trap of adaptationism[15] by assuming that evolvability is an end in itself. Selection can only act on organisms and populations as they exist, and it is an error to assume that patterns identified in hind-sight are predictive of future evolution.

1.3 Changing Environments and Evolvability

Sustained directional selection adjusts the composition of phenotypes and genotypes in a population [16], typically moving that population across the mutational landscape to local regions of higher fitness. When populations find a fitness peak, they tend to cluster there, and exploration of that region of the landscape slows dramatically.

revise
this

In changing environments, however, the direction of selection is not fixed and peaks are not stable. Instead, as the environment changes, populations are driven to explore new regions of the mutational landscape [17, 18]. As they proceed, populations accumulate and carry with the genetic material acquired in prior explorations and adaptations, and use this history as raw material for new adaptation [19]. Indeed, earlier work has shown that changing environments promote evolvability in many contexts, without compromising

robustness [20, 21]. Strength of selection is also an important component of this exploration, since the harshness of the environment drives the speed with which organisms adapt to new conditions [22].

MORE HERE about other things CE can or can't do

1.4 Historical Conceptions of Evolvability

Evolvability is described at many different scopes and levels in the literature, each with varying amounts of detail and predictive power. As such, it may be best to avoid attempting to unify the concept, and rather acknowledge that evolvability is not a singular idea, but rather an overlapping and interrelated set of concepts relating to adaptation and evolutionary potential. In order to synthesize the large field of evolvability and understand how the distinct scopes and ideas connect together, a historical narrative is clearly useful.

1.4.1 Modern Synthesis

The evolution of evolvability as a formalized theory originated with Dawkins[3] and Albrech[11], though the underlying concept (as the response to selection, measured by heritability) existed much earlier, in the work of Fisher[5] and Wright[16]. Fishers fundamental theorem of the response of a population to selection identified narrow-sense heritability (h^2) as a measure for how evolvable populations were. Evolvability as heritability (h^2) is a measure of the portion of the phenotypic variation in a population that can be accounted for by additive genetic effects. h^2 therefore is the component that directly relates to a populations response to selection[6].

$$h^2 = \frac{Var_A}{Var_P} \quad (1.1)$$

In contrast to narrow-sense heritability (h^2), broad-sense heritability (H^2) refers to the entire genetic contribution to a populations variance, including dominance and epistasis. Because of these other contributors, it is unsuitable for isolating the response to selection.

As a measure of evolvability, narrow-sense heritability (h^2) was also used as a term in the breeders equation, in order to estimate the response of a population to artificial selection.

$$R = h^2 S \quad (1.2)$$

Heritability, however, is not an ideal predictor for the response to selection because it fails to integrate factors such as the population distribution of variability in a trait[6]. Heritability, being scaled by total population variation in a trait, would predict the same response to selection regardless of whether the standard deviation of variance of that trait was large or small, or where the mean of that trait lay.

Houle advocated for an alternative genetic variability measure that suffered from fewer of these problems: the Additive Genetic Coefficient of Genetic Variation (CV_A).

$$CV_A = 100 \sqrt{\frac{V_A}{\bar{X}}} \quad (1.3)$$

Using CV_A as the measure of genetic variability is superior to narrow-sense heritability because it scales additive genetic variance by the trait mean, rather than by total population variation. Thus, the additive variation component isn't overwhelmed by large population trait variance[23]. Since life-history (fitness-related) traits tend to have large population variances, h^2 predicts that life-history traits have low heritability and thus low response to selection[24]. CV_A , however, being scaled by trait mean, predicts much higher response to selection for life-history traits[25, 6].

CV_A still suffers from significant drawbacks as predictors for adaptation and evolvability in a larger sense [25]. Both h^2 and CV_A measures predict the response to selection based on the expressed trait variation in a population, under the current environmental conditions. They say nothing of the potential for cryptic variation that may be revealed in different genetic background, nor do they address differences in genetic architecture that may promote faster adaptation. Ultimately, CV_A is best when examining the short-term response to

selection in artificially-selected populations, in static environments, with low mutational load[6].

Clearly, such short-term, population-based measures are unsuitable for measuring larger patterns of the evolution of evolvability, especially over the long term.

1.4.2 Evolvability as a Distinct Concept

Dawkins, in his foundational paper on evolvability and evolutionary constraint [3], re-framed the problem of evolvability in the context of computational evolution and development. Dawkins described a generative genetic system based on a few alleles, and rules that governed development based on the traits encoded in the alleles. Each allele would govern the execution of a generative rule, and the rules would interact with each other as they produced the phenotype. As he added new kinds of rules (constraints) into the generative process, he showed that the system produced more and more complexity.

Dawkins used this example to draw parallels to biological generative developmental systems and how evolutionary constraints in development allow for more complex and robust phenotypes. Dawkins identified a few key themes that underlay the more powerful features of developmental systems. These systems would be organized in such a way as to facilitate cumulative effects. That is, innovations in constraints can build upon each other and are cumulative in evolutionarily interesting ways[3].

Dawkins hypothesized that these kinds of generative developmental systems, or embryologies were the basis for evolvability, and that they must have evolved as a result of their intrinsic power to produce adaptive variation. Dawkins further suggested that the genetic systems that persisted were those that facilitated adaptive radiations into new or otherwise empty ecological niches.

Alberch followed up Dawkins ideas with a more thorough accounting of how, exactly, these kinds of evolvable traits translate into an analyzable phenotype space[11]. Alberch dismantled the concept of a simplistic, hierarchical genotype-to-phenotype mapping function

and emphasized that developmental and cell metabolic systems are strongly dynamical, nonlinear systems, for which genes are just one part of the regulatory cycle. Because of the dynamic nature of cell processes, it was clear that the gene-centric, population genetics view was inadequate to fully describe the complexity of the processes involved, and how they translated complex parameters into phenotypes. To that end, a new framework for analysis was required.

Alberch introduced the concept of parameter spaces to describe the variation in genotypic parameters that results in distinct phenotypes, while addressing the lack of one-to-one correlation between alleles (parameters) and phenotype. Parameter spaces are multidimensional spaces, divided by parameter thresholds (bifurcation boundaries) that form borders between phenotypes. The domains bounded by these thresholds include all of the parameter combinations that produce a given phenotype. Larger domains can be described as more stable than smaller domains, because there are larger ranges of neutral variation available before organisms tip into a different phenotype. Populations with distinct phenotypes and varying parameters can thus be visualized as blobs occupying areas in parameter space.

Alberch contended that the evolvability potential of a dynamical system is encapsulated by the properties of the parameter space. Specifically, the topology of the bifurcation boundaries govern the ease with which the systems can produce both neutral and adaptive variation. Alberch asserted that the generative systems must have undergone selection that favors those systems that provide a good balance between exploration and stability, but provided no mechanism for that selection.

Dawkins and Alberch laid out a compelling case for the role generative developmental systems in facilitating evolvability, but their theoretical frameworks were far from complete.

1.4.3 Theoretical Frameworks for the Evolution of Evolvability

The Wagner and Altenberg paper on the evolution of evolvability significantly expanded the theoretical framework behind the evolution of the genotype-phenotype map[8]. The authors

draw on knowledge from computational evolution to inform their perspective on evolvability, since the problem of evolvability is central to the representation problem in evolutionary computer science.

Initially, Wagner and Altenberg emphasized a distinction between variation and variability. *Variation* is the realized diversity in a population, which is a concept that lies firmly within population genetics and the gene-centric modern synthesis. *Variability*, on the other hand, is a concept that they introduced to describe the ability to generate new phenotypes in response to mutation or environmental change. Variability is a metric associated with a local neighborhood in a genotype to phenotype map, and depends on features of that map, including pleiotropy and modularity, and robustness and flexibility of biological processes.

Wagner and Altenbergs paper led to a vast proliferation of new work exploring the evolution of evolvability. Of particular note is the Kirschner and Gerhart 1998 paper[2], which explored metazoan evolution for examples of traits that, in combination, acted to increase evolvability. The authors found numerous examples of new, evolvable features coinciding with adaptive radiations. The authors also develop a case for a combination of direct selection upon the individual for evolvability-enhancing features, and those traits persisting as by-products as a result of adaptive radiations, setting the stage for the evolution of more and more complex evolvable features.

1.5 So, what do I mean by Evolvability?

As I described above, evolvability is a series of distinct, but overlapping concepts that are generally concerned with adaptation, variation, and/or novelty generation. For the purposes of my research, I am using the Wagner/Altenberg conception of evolvability, which focuses on variability (i.e., the generation of adaptive variation in response to mutation). Variability depends primarily on the organization and interrelation of the components of the genome; that is, the genetic architecture, and the resulting genotype-to-phenotype map.

remove "proposal"

revise this paragraph for my thesis statement.

The major features that influence this metric for evolvability appear to be modularity of functional components and phenotypic robustness to mutation and environmental perturbation. While there are other architectural features that are also likely to contribute to evolvability, they will not be the focus of this proposal.

1.5.1 What is Modularity?

Modularity is the degree to which traits are both self-contained and decoupled from each other. Modular organization can appear at different scales, from the reduction of overlap between unrelated gene regions (spatial modularity[26]), to the decoupling the mutational effects on distinct traits (functional modularity[8]), to the composition of groups of related trait complexes (variational modularity [27, 28]).

Features such as evolvability and robustness are thought to rely heavily on modularity[8]. For example, traits with high functional modularity will have low pleiotropy and therefore should be able to evolve independently—a critical feature if individual traits need to quickly respond to changes in selection. Additionally, modular traits may be more easily re-purposed or co-opted by other traits to add new function[28]. Conversely, spatially modular genomic regions, because they are more self-contained, tend to better resist disruption from recombination, thus increasing robustness[26].

The relationship between modularity and pleiotropy is complex. At small scales, spatial modularity acts to directly reduce pleiotropy by reducing the number of traits affected by a single locus[26]. However, at higher scales, modularity may rely on pleiotropic links within groups of related trait complexes to enable those groups to evolve and optimize in concert[27].

Despite the benefits described above, modularity, like many other variational trait complexes, may not be an unmitigated boon for evolvability. High levels of functional modularity may reduce the overall evolvability of a genotype by reducing the incidence of mutations of large effect and reducing the size of mutational targets[29]. Reducing the incidence of large

changes reduces the likelihood of the development of entirely new traits as a result of relatively few mutations. Thus, the evolvability benefit of modularity may be mediated by the scale and degree to which it occurs.

1.5.1.1 Measuring Modularity

At the phenotypic level, modularity is assessed based on the functional independence of traits and trait complexes. Spatial modularity is correlated with functional modularity, though it is possible to have spatially modular genomes that are not functionally modular and vice-versa[30, 31].

For the purposes of this research, I will focus on spatial modularity. Spatial modularity may be measured by calculating the proportion of traits that are affected by a given site in the genome, normalized by the number of sites that code for a trait [26]. Trivially, this can be measured by performing knock-out experiments to identify the sites that contribute to particular function.

To measure Spatial Modularity, m_S :

1. count the total number of traits expressed in a genome: T
2. identify the number of sites that code for any trait: set K
3. count the number of items in set K : k
4. count the number of traits coded for by each site within set K : t_k ;
5. calculate the inverse of the average number of traits coded for per site to reflect the level of spatial modularity (m_S) of coding regions of a genome

$$m_S = \frac{1}{\frac{1}{k} \sum_{i=1}^k \frac{t_k}{T}} \quad (1.4)$$

1.5.2 What is Robustness?

Much like evolvability, robustness is a set of overlapping concepts concerned with the ability of a genotype to maintain a given phenotype despite an unexpected disruption[10, 32]. Most commonly, robustness is studied in regard to either perturbations in the environment or else mutational disruptions. In the first case, the evolution of robustness to environmental disturbances depends heavily on the flexibility and decoupling of gene regulatory or signaling pathways[2]. For example, a gene-regulatory or signaling pathway that is loosely coupled may make use of signaling from multiple incoming paths, rather than depending on a single, rigid precursor. This type of arrangement is more likely to continue to function even if some part of the signaling path is disrupted. An example of this kind of robust arrangement is nerve conduction in vertebrates where axons connect several cells, thus routing signals in parallel, and avoiding single points of failure[2].

For the purposes of my research, I will focus on the second case: mutational robustness. Distinct from robustness to environmental perturbation, robustness against mutation depends largely on degeneracy, redundancy, and regulatory decoupling[10]. Degeneracy refers to a many-to-one relationship between an encoding and a product, such that several codes can produce a single output. Thus, there is a chance that mutations in the code will not alter the product. One example of this feature is codon degeneracy in biological organisms, where, depending on the hydropathy of the amino-acid, single, or even double mutations in some positions of the encoding do not affect the binding of the encoded amino-acid[33].

Similarly, redundancy refers to the duplication of function in multiple places in the genome, such that mutations altering function in one copy of a gene do not alter function in the other copy. Redundancy may also refer to redundancy of function within genes, such that if a mutation occurs in one portion of a gene, other neighboring portions of the protein will compensate, and the protein will retain its structure and function.[34]

Finally, regulatory decoupling allows for more than one kind regulatory precursor to provide inputs for a process[34]. Thus, if mutation were to damage one set of precursors,

others can take their place and preserve function. An example of this kind of architecture is in the production of the acetate precursor for the Krebs cycle, which produces ATP in all aerobic organisms[35]. Acetate can be derived from either carbohydrates, lipids, or proteins, thus if any of those pathways are damaged by mutation, acetate can still be produced from other sources, and ATP production can continue.

It is worth noting that many of the architectural features that confer robustness to processes and genomes are based on arrangements of modular structures[36, 37]. In this way, much of robustness is facilitated by the evolution of modularity.

1.5.2.1 Measuring Robustness to Mutation

Robustness to mutation can be assessed in multiple ways, either from the perspective of a specific phenotype, a specific genotype, or combinations of the two. From the perspective of an individual genotype, you can assess its robustness by calculating the proportion of mutations that produce a phenotype that is different from the one expressed by the target genotype[38]. In most cases it is easiest to focus on single-step mutations (to cover the 1-neighborhood in the fitness landscape), but sampling from the full distribution of mutation combinations that occur naturally will produce a more exact results.

To measure Genotypic Robustness, r_G , of a genotype G :

1. count the number of loci in the genome: n
2. count the number of possible alleles at a given site: D
3. enumerate all possible single-step mutants that may arise from the given genotype (or sample from a more realistic distribution): $n(D - 1)$;
4. count those mutants that prove to be neutral phenotypic variants: R_G ;
5. calculate the proportion of neutral phenotypic variants to reflect the probability of a neutral variant being produced by this genotype in response to mutation.

$$r_G = \frac{R_G}{n(D-1)} \quad (1.5)$$

Genotypic robustness is trivially negatively correlated with genotypic evolvability, because each neutral variant in the 1-neighborhood of a genotype is, by definition, not of a different phenotype. However, the inverse is not necessarily the case, because each non-neutral neighbor phenotype may not be unique. Therefore, a non-robust genotype may not necessarily have high evolvability if its neighborhood is dominated by a single or few distinct phenotypes [38].

From the perspective of the phenotype, robustness may be assessed by taking the average genotypic robustness across the phenotype.

To measure Phenotypic robustness, r_P :

1. count the number of distinct neutral genetic variants that produce a given phenotypic trait in a population, ($SetK : k$);
2. calculate the proportion of neutral variants produced by single-step mutations r_G , averaged over all of the neutral genetic variants to reflect the probability of a neutral genotype currently in the population producing another neutral genotype in response to mutation.

$$r_P = \frac{1}{k} \sum_{i=1}^k r_G \quad (1.6)$$

Unlike genotypic robustness, higher phenotypic robustness has been shown to correlate with phenotypic evolvability in cases where the possible number of neutral variants in a phenotype (the frequency of the phenotype) is high[38]. With increasing numbers of neutral variants, the number of potential unique phenotypes in the 1-neighborhood of the phenotype increases.

These measures of robustness are each limited in that they do not address realized population composition, the shape of the mutational landscape, nor the expected frequency of

the target phenotype. In particular, the correlation of phenotypic robustness with evolvability depends on the expected phenotypic frequency [38]. Thus, if the frequency is unknown, phenotypic robustness may not predict evolvability.

Further, different populations may have vastly different numbers of realized neutral variants for a given phenotype. Factors such as gene-flow, bottle-necking, linkage dis-equilibrium, founder effects, and sexual selection may strongly affect overall diversity in populations, including the neutral diversity for a particular phenotypic trait that we are concerned with[11].

For this reason, while population level metrics may cause a phenotype to appear to be non-robust, this apparent value may be the result of the amount and type of realized diversity present in a given population, rather than the robustness of that phenotype as predicted by its potential neutral network[11].

1.5.3 Predicting Short-Term Evolvability with Landscape Metrics

As indicated above, the features that confer robustness may also promote evolvability by allowing for greater neutral genetic diversity within a given phenotype. The larger the number of distinct genotypes with the same phenotype in a connected region of the fitness landscape, the more exploration of the genotype space that can be done without decreasing organismal fitness. As a population diffuses through such a neutral region, more potential phenotypes become available in few mutational steps[9].

Historically, predicting this robust-yet-evolvable quality has been challenging. Previously-used measures for robustness that focus on counting the proportion of unique genotypes that compose a phenotype (Phenotypic Robustness [38]) are limited in their ability to predict the evolvability of a population, especially where phenotypic frequency is unknown.

In contrast, we will use Genomic Diffusion Rate, which is the probability that an offspring will be different from its parent, while expressing a neutral or positive fitness effect. This metric may be used to characterize overall population evolvability as it approximates the overall rate in which entirely new genotypes are encountered[39].

To calculate the **Genomic Diffusion Rate** (D_g) in the local neighborhood of a genotype, first calculate its *Fidelity* (F), or the probability of an offspring sharing this genotype with its parent, by measuring the probability that a single locus is not mutated ($1 - \mu$) and raising it to the power of the genome length (l). Next, measure the proportion of 1-step mutants that are neutral or beneficial when compared to the parent (p_ν) as well as those that are detrimental or lethal (p_d), which must sum to one ($p_\nu + p_d = 1$). The *Neutral Fidelity* (F_ν) of a genotype is thus the probability that no harmful mutations occur, assuming no epistasis. Finally, subtracting Fidelity from Neutral Fidelity will yield the overall probability of producing an neutral offspring with a different genotype, yet neutral or better fitness (D_g).

$$F = (1 - \mu)^l \quad (1.7)$$

$$F_\nu = (1 - \mu p_d)^l \quad (1.8)$$

$$D_g = F_\nu - F \quad (1.9)$$

Measures of neutral exploration, however, only show part of the picture. While some form of neutrality is necessary for exploring a fitness landscape, new phenotypes must be discovered to achieve higher local evolvability. In order to assess evolvability more specifically, we introduce a related measure, the **Phenotypic Diffusion Rate** (D_p), which represents the probability that an offspring will be fitness-neutral, but also express a different phenotype than its parent. To do so, we must first measure the proportion of one-step mutants that are *phenotypically* neutral as compared to their parent ($p_{p\nu}$) and follow a similar procedure as above, first calculating the probability that a phenotype-changing mutation will occur (μ_{pheno}), then the phenotypic-level fidelity ($F_{p\nu}$).

$$\mu_{pheno} = \mu(1 - p_{p\nu}) \quad (1.10)$$

$$F_{p\nu} = (1 - \mu_{pheno})^l \quad (1.11)$$

$$D_p = F_\nu - F_{p\nu} \quad (1.12)$$

The difference between the overall neutral fidelity and the phenotype-preserving neutral fidelity ($F_\nu - F_{p\nu}$) yields the phenotypic diffusion rate.

1.5.4 Expected Value of Fitness Landscapes

In the context of changing environments, the expected fitness value ($E(w)$), and thus the neutrality, of a mutant in the mutational landscape will vary depending on the environmental context. So, in one environment, a mutant may be highly fit, but the same allele may be highly deleterious in a different environment. In order to address this variation, all metrics must be normalized by the probability that a particular environment will occur (P_i). That is, the nearby mutational landscape must be evaluated in each possible environment, yielding a traditional fitness landscape. Then, the set of fitnesses of each mutant (w_i) in each environment must be aggregated according to the probability of that environment occurring.

$$E(w) = \sum_{i=1}^e w_i P_i \quad (1.13)$$

1.6 Digital Evolution

Digital Evolution uses self-replicating computer programs as model organisms to study evolutionary dynamics [40]. Unlike theoretical simulations, digital organisms have a fully functional genome that direct them to self-replicate, mutate, and compete with their peers for resources and space in which to reproduce. Because digital organisms undergo genetic mutations (i.e., variation) that are passed on to their offspring (inheritance), and their survival is based on the actions they take (differential selection), they undergo evolution by natural selection.

Digital organisms do not suffer from many of the drawbacks of experimentation on natural organisms. Three of the advantages of digital organisms are particularly relevant for our study. First, the rates of reproduction in digital systems are much faster than in even the

most rapidly-reproducing physical organisms; we can process generations of organisms in seconds, rather than the hours required for the fastest biological organisms under sustained conditions [41, 42], or the weeks to years needed for more complex multicellular organisms [43, 44].

Second, using digital organisms allows us to tightly control and verify experimental conditions. For example, in physical organisms, factors such as mutation rate can generally be measured only after the fact, or coarsely altered through mutagens. In digital organisms, however, we can not only control mutation rates with fine-grained precision, but also types and probabilities of different types mutations (e.g., substitutions vs. insertions vs. deletions). Furthermore, we are also able to track and replay the evolutionary history of every organism at any point in time to verify that unusual or unexpected results do not represent measurement error. This ability to exactly replicate evolutionary results at an individual organism level is firmly out of reach for experiments with physical organisms.

Finally, we can precisely and perfectly map the mutational landscape around the genome of a digital organism, and identify the role of every site in its genome[39]; such exhaustive techniques are not feasible in even the simplest physical organisms. All of these factors make digital organisms ideal for studying the effects of changing environments on the mutational landscape.

1.6.1 Avida

Throughout the rest of this thesis, I use the Avida digital evolution platform to explore the effects of changing environments on the evolvability of populations of digital organisms.

Avida is a software platform for performing evolution experiments with digital organisms in a virtual world.

An Avida organism is composed of a circular genome of assembly-like computer instructions that are executed in a virtual CPU (Fig 1.1). Populations of these organisms are placed in a toroidal world in individual cells where they are allowed to execute, reproduce, compete

what
goals?
(see fol-
lowing
sentence)




Figure 1.1: **An example virtual CPU from Avida**, with a circular genome (blue), three registers (purple), input and output handlers (tan), and an instruction pointer (yellow) indicating the next instruction to be executed.

for space, mutate, and evolve.

Organisms in Avida are self-replicating, and experience mutation. The genomes of the initial default organisms contain all of the instructions necessary for reproduction. However, the instructions are not copied into an offspring with perfect fidelity. By default, the reproductive copy instruction is faulty, meaning that it will probabilistically introduce errors (mutations) into the offspring genomes. These offspring organisms execute their own genomes even when different from their parent, and in turn pass on their inherited mutations, along with new mutations, to their own offspring (i.e., variation in the systems is heritable).

Avida worlds can be space- or resource-constrained. Avida allows the experimenter to configure many aspects of the environment, thus subjecting the organisms to various kinds of selective pressures. In many cases, these environments will include resources that can be metabolized by performing specific functions or activities, resulting in a boost to execution speed that gives the organisms a competitive advantage. However, even without explicit

external pressures, organisms still experience an implicit pressure to execute more quickly and efficiently. The organisms that run fastest are typically able to also reproduce fastest, and thus out-compete their peers for space.

Avida is available for download without cost from <http://avida.devosoft.org/>, and specific versions along with data-files to reproduce the experiments described in this thesis may be found at <https://github.com/voidptr/avida> and <https://github.com/voidptr/dissertation>.

CHAPTER 2

CHANGING ENVIRONMENTS PROMOTE RAPID ADAPTATION IN DIGITAL ORGANISMS

2.1 Background

The interaction between an environment and possible genomes can be mathematically expressed by a fitness landscape. Fitness landscapes are a mathematical tool to map genetic sequences to reproductive fitness. Many studies have examined the important role that different types of fitness landscapes play on evolutionary dynamics and outcomes, both in biological populations [45, 46, 47, 48] and in evolutionary computation settings [49, 50, 51]. However, real-world fitness landscapes are far more complex and varied than the limited or idealized models that are used in most of these studies. Neighboring regions of real landscapes can have starkly different properties from each other based on the effects of and interactions among mutations; as such, a local region of a fitness landscape around a genotype is commonly referred to as its mutational landscape.

Examples of the type of properties that we are interested in include robustness, epistasis, and modularity, all of which are measurements of how information is organized inside of a genome and commonly categorized as components of an organism’s “genetic architecture”. Isolated pockets in a landscape can often be characteristically different from the landscape as a whole due to the amount and organization of genetic information. In fact, in most natural fitness landscapes, the vast majority of neighborhoods consist entirely of non-replicating genomes with zero fitness (and thus no genetic information), making life itself appear to be a rare exception [52].

Evolution on these convoluted landscapes is clearly limited to those regions that have non-zero fitness, with a selective pressure for fitness to increase. Beyond that, however, populations can evolve toward neighborhoods with specific local properties based on the

evolutionary forces acting upon the populations. For example, high mutation rates drive populations toward neighborhoods with a higher fraction of neutral mutations in an effect dubbed survival of the flattest [21]. Similarly, sexual populations tend toward regions of the fitness landscape with more modularity [26] and more negative epistasis [53] than otherwise equivalent asexual populations.

Understanding these dynamics is of broad interest. It is important to evolutionary computation, given the strong influence of local landscape properties on the quality of the final solutions that an evolving population is able to obtain. Its relevance to evolutionary biology is equally obvious – the local landscape that a population occupies will influence the selective forces at play in the population, creating a feedback cycle between these two important evolutionary factors [54, 55]. Disentangling such interactions is likely to provide further insights into fundamental evolutionary dynamics. Computational artificial life systems have the advantage of being able to bridge these two realms: they have unconstrained evolutionary dynamics similar to natural systems, while maintaining the ability to rapidly perform experiments and collect any data we need about populations or their local landscapes.

2.1.1 Evolvability and Genetic Architecture

As described in Chapter 1, evolvability refers to a series of distinct but overlapping concepts that are generally concerned with adaptation, variation, and/or novelty generation [13]. Depending on your perspective, evolvability can describe the response to selection at the population level[5, 6], the ability of populations to adapt to changing conditions[7], larger phenomena such as variability generation[8], exploration of neutral spaces and robustness[9, 10], generation of novel features[11, 12], or even the potential to generate clade-level innovations[2] and major transitions[1]. For the purposes this chapter, we will focus on evolvability as the capability of genomes to generate adaptive variation in response to mutation.

In the short-term, this kind of evolvability determines a population’s response to selection. This kind of evolvability depends primarily on the organization and interrelation of

information in the genome; that is, the genetic architecture, and the resulting genotype-to-phenotype map [8]. An example of evolvable architecture can be found in some bacterial genomes that contain highly mutable genome regions, called contingency loci. Small sets of insertions or deletions to these regions create transcription frameshifts that alter the expression of nearby coding regions, thus allowing populations to easily switch phenotypes via minor mutations. Contingency loci are most often seen in the genomes of pathogens, which are subject to frequent environmental shifts caused by the host immune system [56]. Thus, these populations are able to produce large amounts of heritable variation despite the reduction in population diversity resulting from population bottlenecks.

2.1.1.1 Mutational Landscapes

Properties of genetic architectures such as evolvability and robustness are determined by the shape of the resulting mutational landscape (local fitness landscape around a genotype, accessible in a single mutation) [38]. Robust genetic architectures that can tolerate more mutations without altering their phenotype reside in mutational landscapes that connect to more neutral mutants. Similarly, architectures that more easily switch phenotypes in response to mutation without substantial reduction in fitness, reside in more evolvable regions of genotype-space.

It is worth noting that not all regions of the mutational landscape are equally accessible. Some genome regions may be more resistant to mutation than others [57], thereby altering the probabilities of mutations occurring that lead into certain regions of the mutational landscape. This kind of differential probability may therefore moderate a population's diffusion through the mutational landscape.

Further, response to selection is likely to be weaker in regions of the landscape where there are fewer available mutations that provide potentially adaptive traits, whereas response to selection will be stronger in regions where there are many adaptive variants available within a few mutational steps [11, 58]. This differential response to selection therefore constrains

the ability of populations to diffuse across a fitness landscape.

In order to assess the potential of different regions of the fitness landscape to promote or hinder evolvability, we will use both the **Genomic Diffusion Rate** (D_g)^{1.9} and the **Phenotypic Diffusion Rate** (D_p)^{1.12}, as normalized across changing environments^{1.13}.

2.2 Methods

2.2.1 Experimental Design

In order to examine the dynamics and mechanisms of evolving populations in changing environments, we performed two sets of experiments. We subjected populations of evolving digital organisms to a set of cyclic changing environments, and a set stochastic changing environments. The cyclic environments were designed to simulate predictable cycles of change, such as day/night or seasonal cycles, whereas the stochastic environments represent less predictable oscillations in environmental states, such as random weather patterns, or climactic changes. These experiments allow organisms to adapt to a predictable set of environments, and explores short-term evolvability dynamics. See Table 2.1

Missing figure

A diagram of how the environments are placed. stochastic as jumbled, cyclic as periodic. Harsh is red/blue, Benign is white/blue.

2.2.1.1 Cyclic and Stochastic Changing Environments

For the cyclic environment, we subjected a total of 150 replicate populations of digital organisms to two different treatments of two-phase cyclically changing environments, plus a static control. The environment cycles between 500 updates of reward, and 500 updates of punishment; as such each full cycle is 1,000 updates, or roughly 30 generations. We chose this cycle length after surveying a series of possible values in order to determine an optimal

Table 2.1: **Experimental Treatments**

Treatment	Changing Environment	Rewarded Tasks	
		XOR	EQU
Control	None (static)	constant 2^3	constant 2^5
CCE Benign	Cyclic	constant 2^3	benign fluctuating 0 or 2^5
CCE Harsh	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5
SCE Benign	Stochastic	constant 2^3	benign fluctuating 0 or 2^5
SCE Harsh	Stochastic	constant 2^3	harsh fluctuating -2^5 or 2^5

Experimental treatments. Four types of changing environment, plus a static control. In the first two treatments, the environment switches in a predictable cycle, whereas in the second two, the environment switches at random intervals.

length of time. That is, long enough to allow adaptation to occur and spread through the population, but short enough to reduce the effects of drift destroying vestigial genetic information. For more details about this survey, please refer to Appendix A.

In the static control, there is no cycle. Rather, the rewards remain constant. The first phase of the experiment extends for 200 cycles, or 200,000 updates, approximately 6,000 generations.

The stochastic changing environment experiment is similar to the cyclic environment, except that rather than the environment toggling every 500 updates, the environmental switch happens randomly, with a 0.002 probability of changing on every update. This averages, in the long term, to approximately one switch every 500 updates, but in the short term, the environmental switches are unpredictable.

We set up the system to detect organisms that performed XOR or EQU, two challenging

bitwise logical tasks. In the static control, XOR is rewarded with a CPU speed (and thus fitness) multiple of 8, while EQU is rewarded with a CPU speed multiple of 32. In the harsh treatment, as the cycle progresses, the XOR reward remains constant, while the EQU reward cycles between a 32-fold bonus and a correspondingly harsh 32-fold penalty (i.e., CPU speed is divided by 32 when EQU is performed in the off cycle). The benign treatment is nearly identical to the harsh treatment, except that the reward merely goes away in the off-cycle as opposed to incurring a severe penalty.

In both environments, we identify EQU as the *Fluctuating Task*. XOR, because it is rewarded continuously, is the *Backbone Task*, and is used as a background for comparing the separation or intertwining of functional genetic components in the evolution of EQU. Further, the 4-fold difference in reward level between XOR and EQU encourages the evolution and maintenance of EQU when possible.

For all of the experiments described in this section, we held the individual genomes at a fixed length of 121¹ instructions, but tested the new genomes for mutations after each successful replication event at a substitution probability of 0.00075 per site. We configured the Avida world to have local interactions on a toroidal grid that is 60-by-60 cells (3600 cells in total), and we seeded the initial populations with an ancestor that was previously evolved to perform XOR and EQU under a static reward. The genetic architecture for performing XOR and EQU is tightly intertwined in this ancestral organism, as it was evolved with no selective pressure for modularity.

2.3 Results and Discussion

Our experiments demonstrate that digital organisms that were evolved in changing environments differ substantially from those that evolved in static environments in a number of ways. These differences include the number of mutations that fix in the lineage from

¹As part of our initial controls, we hand-wrote an organism with separated sections that performed XOR and EQU. This hand-written organism had 121 instructions and as such we used this genome length as a constraint for the evolve organisms as well.

the ancestor (the “phylogenetic depth”), key metrics of their genetic architecture, and the presence of reservoirs of pseudogenes that change the nearby mutational landscape. These features represent adaptation to the larger regime of repeated environmental switching. We also show that while populations evolved in cyclic environments are slightly better adapted to change than those that evolved in stochastic environments, in most measures of adaptation and short-term evolvability, these differences are generally not significant. This result indicates that while regular periodicity may offer a slight advantage for adaptation, stochastic environments perform similarly in most respects.

look
for un-
anchored
”this”s

2.3.1 Cyclic Changing Environments

We will begin by examining the characteristics of populations evolved in cyclic changing environments.

2.3.1.1 Evolutionary History and Population Structure

Evolution in the harsh cyclic changing environment resulted in many more mutations fixing, and thus populations with substantially higher phylogenetic depth as compared to those evolved in static or benign environments. At each environmental shift, adaptive mutations rapidly swept and fixed in the populations. (Fig 2.1)

The populations that evolved in the control and benign environments displayed more genetic diversity as compared to those evolved in the harsh cyclic environment, which underwent a bottleneck at each cycle shift. Because a selective sweep reduces current diversity within a population, the smaller number of sweeps in the benign and control treatments led populations in them to have higher standing diversity for most of their evolutionary history than those populations from the harsh changing environment. Despite this higher standing diversity in the benign and control treatments, regions of low diversity are still evident in the genomes of these populations, implying purifying selection on the traits encoded at these sites (see Fig 2.2).

Phylogenetic Depth and Last Coalescence




Figure 2.1: **Phylogenetic depth over time** of a sample population evolved in each of the three treatments of the cyclic changing environments. White horizontal lines mark the depth of the most recent common ancestor, and discontinuities in this line indicate that the most recent common ancestor has changed, and thus that a sweep occurred, or that a competing clade went extinct. The control treatments had a mean of 18 sweeps (STD=9.05), the benign treatments had a mean of 21 (STD=19.05), and the harsh treatments had a mean of 88 sweeps (STD=23.37). Note the difference in scales between y-axes: the control-evolved population has a maximum depth of 400 mutational steps from ancestor, while the harsh-evolved has upward of 1100.

fix figure below so that it's two separate figures. Separate the mean entropy to its own figure.




Figure 2.2: **Population Per-site Entropy over time** of a representative sample population. Each vertical slice represents the per-site entropy of the population at each update, both by genetic locus (upper), and overall population mean (lower). Hotter colors (red/orange/yellow) indicate greater diversity at this locus, while cooler colors (blues) indicate the a locus is more consistent across the population. Mean population entropy indicates the relative diversity of the population at any given time, while the per-site entropy shows where in the genomes the population diversity is located.

2.3.1.2 Genetic Architecture

The selective shifts in both benign and harsh changing environments result in qualitatively

clarify
wording

different architectural styles from the static control environment. The task arrangements evolved under both experimental treatments are much more scattered throughout the genome than in the control. Specifically, the bulk of the sites responsible for performing the fluctuating task (EQU) did not overlap with the backbone task (XOR), except for a core region, which represents portions of the tasks that are shared between XOR and EQU. (Fig 2.3)

In contrast, the architecture of XOR and EQU remain tightly intertwined in the control, and site positions do not change substantially over the course of the experiment. In the benign treatment, many more regions that perform the fluctuating task (XOR) are scattered throughout the genome, but site positions remain relatively fixed throughout the run after an initial adaptive phase. In the harsh treatment, not only are the active sites scattered, but the positions of active sites change and proliferate wildly over time.

Interestingly, populations evolved in both the benign and harsh treatments also show development of a large reservoir of formerly functional, now vestigial, sites; that is, sites that remain unchanged from when they were previously active in performing a task, but were disabled by a mutation elsewhere and are thus now neutral. These vestigial pseudogene-like sites appear to be important for allowing the organisms to quickly re-adapt as the fluctuations in the environment restore the previously-rewarded functions. (Fig 2.4)

2.3.1.3 Nearby mutational landscape

add figure of the nearby landscape showing how mutations relate 1 and 2 steps out., a couple of nice figures showing both changes in detrimental mutations and phenotypic distribution.

In order to identify the role that these pseudogene-like structures play, we performed a survey of the single-step mutational landscape surrounding the most abundant genotype at the end of the experiment for each replicate population. This landscape contained 3,025

polish
this figure, task layout, make three separate figures

stats for figure

fix word-ing to clarify




Figure 2.3: **Varying genetic architecture of XOR and EQU over time** for the final dominant genotype in a randomly selected replicate. Proceeding from the left of each figure, each vertical slice represents an organism along the line-of-descent to the final dominant. Positions along the Y-axis represent each genome locus; loci in an organism are colored based on the tasks that they code for. Sites in **red** are active sites that code for the XOR task only, sites in **blue** are active sites for the EQU task only, and **purple** sites code for both XOR and EQU. Knockouts to the sites in black are lethal to the organism. Sites in the lighter colors (tan, light blue, lavender) represent vestigial sites for XOR only, EQU only, or both tasks, respectively. As we proceed from left to right, we can see the evolutionary history of the final dominant genotype.

distinct mutants (121 loci with 25 possible mutations per locus) in each of the 50 replicates per treatment, for a total of nearly 450,000 mutants surveyed. We found that the availability




Figure 2.4: **Number of functional and vestigial sites by treatment.** The harsh environment has a significantly larger number of vestigial sites for the fluctuating (EQU) task compared to the benign treatment or control, while having a comparable number of functional sites (One-Way ANOVA $F(X, YYY) = ZZ.ZZ$, $p << 0.000QQ$).

of reservoirs of vestigial sites shifted the change-evolved organisms' position in the mutational neighborhood, such that a task that was lost due to mutation remains more accessible via one or two additional mutational steps. (Fig 2.5, 2.6)

Todo, clarify per Charles's comment above

We also measured the proportion of non-deleterious mutants in the nearby fitness landscape. We found that between all treatments, this proportion remained approximately the same. However, we found that the proportion of these mutants with different (potentially adaptive) phenotypes increased in the changing environments. In this way, the organisms from the changing environment treatments have an advantage over organisms from the control runs in terms of the short-term evolvability of the fluctuating task. This result indicates real adaptation, not only to resources in their local environment, but a direct adaptation to the environmental change. (Fig 2.7)

stats for figure
stats for figure
Add to caption per Charles's comment above
add stats for this




Figure 2.5: **A survey of the single-step mutational neighborhood** around organisms that performed the fluctuating task. Note that in both the benign and harsh treatments, there were significantly more mutants that lost the EQU task as compared to the control (Wilcoxon Rank Sum Test: Z = X.XX and Y.YY respectively, $p << 0.000ZZ$). This result indicates that it was easier for the organisms in both treatments to turn off the EQU task in response to one mutation.

TODO - add paragraph describing/speculating about why Benign loses EQU almost as well as Harsh

stats

2.3.2 Stochastic Changing Environments

Contrary to our expectations, stochastic changing environments were no more effective at promoting evolvability than cyclically changing environments. In all measures of evolvability, the stochastic treatments performed similarly to the cyclically changing environments. There was a significant reduction [TODO STATS] in the Phenotypic Diffusion Rate (D_P) between the cyclic and stochastic harsh changing environments. Overall, D_P showed much larger variances, but settled on a lower mean in the stochastic harsh treatment as compared to the cyclic harsh, indicating a much lower probability of the population producing offspring that

STATS
for drop
in SCE-
Harsh
 D_p
reward
below




Figure 2.6: **A survey of the two-step mutational neighborhood** of the organisms that lost EQU function in the one-step survey. We found that in both the harsh and benign treatments, there were significantly more organisms that regained function in response to mutation than the control. (Wilcoxon Rank Sum Test: $Z = X.XX$ and $Y.YY$ respectively, $p << 0.000Z$). This result indicates that it was easier for the organisms in both fluctuating environments to regain the task in response to one additional mutation.

would switch phenotypes neutrally. (Fig 2.8)

stats

Similarly, both the overall fraction of 1-step mutants that lost EQU, and the fraction of 2nd-step regaining of EQU, were slightly reduced in comparison to the cyclic treatments. This result indicates that stochastic harsh environment was slightly less effective at promoting evolution toward areas of the mutational landscape where such mutations were common. (Fig 2.9, 2.10)

stats

The greatest differences between the cyclic and stochastic treatments appeared in the number of functional and vestigial sites. While the functional site counts in the stochastic environment were, overall, relatively similar to those in the cyclic environment, the pattern was different for the vestigial sites. In the stochastic harsh treatment in particular, there was

stats

stats




Figure 2.7: **Genomic and Phenotypic Diffusion Rates**, showing the probabilities of producing offspring that are genotypically (D_g) or phenotypically (D_p) distinct from the parent, while not reducing fitness. Note that while overall neutral exploration capacity remains relatively stable between treatments, phenotypic exploration capacity is increased in both treatments, but especially in the Harsh treatment. (Wilcoxon Rank Sum Test: $Z = XX$ and XX respectively, $p << 0.0001$). This result indicates that changing environments promote the phenotypic evolvability of populations in particular.

a small, but significant reduction in the number of XOR+EQU overlapping functional sites [stats] as compared to the cyclic treatment. Further, there was an overall reduction in the number of EQU-only vestigial sites. These features may provide a clue, indicating that architectural features that would promote the retention of EQU were less prevalent in the harsh stochastic environmental treatment. (Fig 2.11)

Together, from these measures, we conclude that stochastic environments exert slightly less evolutionary pressure to move toward regions of the mutational landscape that are more congenial to neutral phenotypic exploration and evolvability. We hypothesize that this dynamic may be due to the randomly-occurring environmental changes may either occur too rapidly for a response to selection, or too slowly, such that drift may cause the information




Figure 2.8: **Genomic and Phenotypic Diffusion Rates** in stochastic changing environments, showing the probabilities of producing offspring that are genotypically and phenotypically different from the parent, while remaining fitness neutral or better. As in the cyclic environment D_g remains stable, at comparable levels [todo stats], however, the mean is significantly lower[TODO stats]. This result shows that stochastic environments are not as effective as cyclic environments at increasing the probability that organisms will produce phenotypically different, yet neutral offspring.

contained in vestigial sites to mutate away. While the environment, on average, experiences as many changes as in the cyclic experiment, the distribution of the length of those environment periods may be very different. Thus, we can conclude that our stochastic changing environment is not more effective than a cyclic changing environment, and under harsh conditions, may actually be slightly worse for promoting the evolution of evolvability.

2.4 Conclusion

In cyclic changing environments, the direction of selection shifts frequently, and periodically drives populations to not only explore new regions of the genetic landscape, but also to carry with them vestigial genetic information about previous environmental conditions. Thus, the resulting populations are not only adapted to the current environment, but also to the meta-environment of cyclic change. Because of their evolutionary history, the genomes




Figure 2.9: **A survey of the single-step mutational neighborhood** in the stochastic changing environment around organisms that performed the fluctuating task. Again, in the static and benign treatments, values are comparable to the cyclic changing environment [TODO stats]. However, in the harsh treatment, the means for both loss of the fluctuating task (EQU) and loss of both task were slight reduced [TODO stats]. This result indicates that in the context of a harsh treatment, stochastic environmental change is less effective at moving organisms to areas of the fitness landscape where they can more easily switch task expression.

contain vestigial fragments of genetic material that were adapted to prior environments. As this exploration proceeds, mutations accumulate in the population, each creating a link to a new region of the mutational landscape. As these links accumulate, they form a reservoir of mobility for the population to quickly shift to new phenotypes as dictated by current selective conditions. In this way, the accumulation of vestigial or pseudogene-like regions acts as an indirect adaptation to the larger pattern of changing selective forces.

add a sentence emphasizing that it's an indirect effect.

By contrast, in static (non-changing) environments, the majority of neutral mutations do not connect to as many phenotypically-interesting regions of genotype-space. There are far fewer pseudogene-like regions available that could regain functionality should conditions




Figure 2.10: **A survey of the two-step mutational neighborhood** in the stochastic changing environment of the organisms that lost EQU function in the one-step survey. Similarly to the result in Fig 2.9, we found that the fraction of organisms regaining the fluctuating task from a single additional mutation in the harsh treatment were reduced compared to the cyclic harsh treatment[TODO stats]. This result confirms that the harsh stochastic environment is less effective than the cyclic harsh at promoting evolvability.

change. Thus, populations evolved in static environments are less evolvable in the short-term.

Surprisingly, stochastically changing environments are slightly less effective at exploration than cyclic changing environments, even if, on average, the amount of time spent in each environment was equal. We hypothesize that this result is because of more opportunity for drift to destroy the information contained in vestigial regions, as well as potentially fewer opportunities for populations to respond to selection.

Stochastic Changing Environment Only EQU-Performing




Figure 2.11: **Number of functional and vestigial sites by treatment** in a stochastic changing environment. The vestigial site counts remain comparable to the cyclic environment [TODO stats], however, there was a reduction in functional site counts for XOR+EQU overlapping sites in the stochastic harsh environment as compared to the cyclic harsh environment [TODO stats], as well as an overall reduction in the number of vestigial sites.

CHAPTER 3

CHANGING ENVIRONMENTS AND THE EVOLUTION OF HORIZONTAL GENE TRANSFER

3.1 Background

Horizontal Gene Transfer (HGT) is a broad term for the non-reproductive transfer of genetic material between organisms. Organisms may uptake genes directly from the environment (transformation via natural competence [59]), or else receive them via bacterial conjugation[60] or viral infection (transduction [61, 62]). In the case of transformation, the fragments may either be decomposed inside the recipient cell for their nutrients, or recombined into their genomes.

HGT has had a profound impact on the evolutionary history of both prokaryotes and eukaryotes, with one study showing approximately 81% of genes in the sample "being involved in HGT at some point in their history" [63]. For example, HGT appears to be the primary mechanism by which antibiotic resistance is conferred [64, 65] since most antibiotics are sourced from the environment, and the organisms that develop them are themselves resistant to the compounds. However, the origins and evolution of HGT mechanisms remain unclear.

3.1.1 Origins of Horizontal Gene Transfer in nature

More about how this question is still open

In prokaryotes, "transformation" is an HGT mechanism by which organisms spontaneously uptake the DNA of dead organisms in the environment. Competent organisms benefit in several ways. 1) DNA is composed of a 5-carbon sugar, a phosphate, and nitrogenous bases, materials that are useful for DNA synthesis and repair. 2) The organisms may also benefit from taking up gene fragments that confer new adaptive functionality into the

genome [66]. However, it is unclear whether the origins of transformation functions were developed solely in order to obtain nutrients (the Grazing Hypothesis), or if the acquisition of new functionality was selected for as well. While grazing for gene fragments as nutrients certainly conveys an advantage, the possibility of integrating these gene fragments is likely to be deleterious to organisms more often than it is beneficial [67]. For example, since the DNA would be originating from dead organisms, DNA fragments may be of low quality, and recombine deleterious mutations, or even remove competence altogether [67]. Alternately, errors in homologous recombination may apply fragments to random locations in the genome.

In order to test the Grazing Hypothesis, , and address the question of whether there are circumstances where gene fragment integration may be beneficial, we subjected populations of evolving digital organisms to a harsh changing environment, where there is a strong selective pressure to quickly switch phenotype. We supplied organisms with an instruction that performs Horizontal Gene Transfer (`HGT-Uptake`). That is, the instruction triggers uptake of a genetic fragment from the environment, and there is a probability that, rather than metabolizing the fragment for a bonus to execution speed, the fragment will instead be homologously recombined into the organism’s genome. We show that in harsh changing environments, without any kind of bonus, organisms increase use of HGT as compared to execution in a static environment.

3.2 Methods

In this chapter, we use Avida to test hypotheses about the origins of Horizontal Gene Transfer.

3.2.1 HGT in Avida

In Avida, HGT is triggered by the `HGT-Uptake` instruction that, when executed, attempts to uptake a genome fragment from the individual cell reservoirs in the environment. As organisms die in Avida runs with HGT enabled, fragments of their genomes will accumulate

in reservoirs. The fragments in the reservoirs will deteriorate over time, with older genomes disappearing from the reservoir as new ones enter. Fragments for uptake will be randomly selected from the reservoir.

look for and hunt down passive voice

We have implemented a configurable probability that, when a fragment is taken up, it is not metabolized; instead, homologous recombination may occur (Figure 3.1). We performed experiments to derive bonus levels, fragment sizes, and recombination probabilities to arrive at a maximum use level for the HGT instruction. See Appendix B for more details.

For all experiments described in this chapter, we used a 10% recombination probability. Please refer to Appendix B for more information about the selection of this probability value. We also required three instructions as the minimal homologous match length on either side of the fragment. Three instructions on either side of the fragment (26^3 unique sequences) approximates the number of unique values keyed by 7 nucleotides (4^7 unique sequences). For the purposes of homologous recombination in plasmid cloning in *E. coli*, 20bp is enough to assume identity[68].

Citation
TODOs

3.2.1.1 Environmental Conditions

All experiments in this chapter compare outcomes between static environments, and harsh changing environments. In a similar manner to the experiments listed in the previous chapter (Chapter 2, Table 2.1), task rewards in the harsh changing environment switch from a positive to a negative reward. For the HGT experiments, we did not establish a backbone task that was always rewarded. Rather, we divided the Logic-9 environment into two halves, and alternated positive and negative rewards for each task in each set. Thus, for the first phase of the cycle, the NOT, AND, OR, NOR, and EQU instructions were punished at -2^1 , -2^2 , -2^3 , -2^4 , and -2^5 respectively, while NAND, ORN, ANDN, and XOR were rewarded at 2^1 , 2^2 , 2^3 , and 2^4 . In the second phase of the cycle, these rewards flip, such that NOT, AND, OR, NOR, and EQU are rewarded, whereas NAND, ORN, ANDN, and XOR are punished.




Figure 3.1: The HGT process. Organisms can execute instructions that trigger uptake from the environment. When uptake occurs, there is an experimenter-defined chance that either it will yield a boost to speed of execution or, alternatively, that the fragment will be integrated into the genome.

Each complete cycle lasts 1,000 updates, and the whole experimental run continues for 200,000 updates.

The static environment rewards executions of all the Logic-9 tasks at their default levels, with no reward switching.

3.2.2 Experimental Design

For the treatments corresponding to the first set of hypotheses on the origins of horizontal gene transfer, we subjected four populations of evolving digital organisms with HGT to

Table 3.1: **Experimental Treatments - Evolution of HGT**

Treatment	Changing Env. Type	HGT Action	Bonus
Control	Static	None	n/a
HGT B0.0 (Natural Competence No Bonus)	Static	10% Recombination Probability	n/a
HGT B0.0 CE (Natural Competence No Bonus)	Harsh Cyclic	10% Recombination Probability	n/a
HGT B0.8 (Natural Competence with Bonus)	Static	10% Recombination Probability otherwise Bonus Allocation	$2^{0.8}$ per Uptake
HGT B0.8 CE (Natural Competence with Bonus)	Harsh Cyclic	10% Recombination Probability otherwise Bonus Allocation	$2^{0.8}$ per Uptake

Experimental treatments. Four treatments corresponding to the combination of two factors: Static vs Changing Environment, and Grazing Bonus vs No Grazing Bonus, plus a non-HGT control, where the HGT instruction is inert.

harsh changing environments (Table 3.1), plus a fifth non-HGT control. The treatments correspond to the combination of two factors: static vs changing environment, and grazing bonus vs no bonus.

For the second set of hypotheses, where we identify the mechanisms that promote the use of HGT, we manipulate the content of the reservoirs to contain fragments drawn from specific phases in the cyclically changing environment, such that fragments either match or do not match the environment (Table 3.2. We then measure HGT use, as well as average

Table 3.2: Experimental Treatments - Effects of HGT

Treatment	Fragment Source
HGT	organism death
HGT-Both	sampled from organisms from both phases
HGT-OnPhase	sampled from organisms from the matching phase
HGT-OffPhase	sampled from organisms from the non-matching phase

Experimental treatments. Four treatments corresponding to the sources of fragments. The first treatment uses the default fragment source (dead organisms from the environment). The second treatment samples a population for organisms corresponding to both phases, and injects those into the reservoirs. The third and fourth treatments sample the population, but only inject fragments from the matching and non-matching phases, respectively.

fitness effects of the HGT mutations, and the fraction of mutations that lead to beneficial phenotype switches.

3.3 Results and Discussion

Our results show that both an uptake bonus and changing environment promote the use of HGT, and that the increases in uptake are largely additive. Further, we found that while on average, HGT mutations are neutral, that the majority of the benefits conveyed by HGT in changing environments come from fragments originating in periods of the cycle where the environment matched the current environment. This result indicates that it is the information content of the fragment that is valuable. Finally, we find that providing only fragments from the matching cycle elevates uptake rates significantly.

3.3.1 Changing environments elevate HGT use

We measured HGT fragment uptake in four conditions (see Table 3.1), plus of pair of non-HGT controls. Without a bonus, in a static environment, fragment uptake is depressed to a low level as compared to the rate of the non-HGT control, where the HGT-Uptake instruc-

rename
table

tion does nothing. This indicates that HGT in a static environment is largely deleterious (Figure 3.2). However, in a harsh changing environment, fragment uptake is elevated. This shows that in the context of a harsh changing environment, the integration of new genetic material is more beneficial than in the static environment. We also found that, regardless of whether the environment is static or changing, when a bonus to fragment uptake is provided (analogous to the nutritive benefit granted by natural competence in biological organisms), fragment uptake also increases.

In order to investigate the relationship between the effects granted by a nutritive bonus and the benefit of fragment recombination in a changing environment, we selected a bonus level that increased fragment uptake in a static environment to a level comparable to the increase seen in changing environments without a bonus. We then combined these factors, giving a bonus, plus using a changing environment. We saw that the resulting uptake level increases largely additively. This indicates that the benefits granted by a grazing bonus are largely independent of the benefits conveyed by integrating new genetic material.

Thus, not only is HGT evolution possible absent a bonus, the benefit stacks with that of a grazing bonus, proving a more likely scenario by which HGT might evolve.

stats

3.3.2 HGT derives most benefit from on-cycle fragments, but not all

In order to understand the basis of the beneficial nature of HGT in changing environments, we performed a series of treatments where we replaced the fragments in the cell reservoirs with fragments that originated at the end of the matching phase of the cycle, the off-phase of the cycle, and from both phases, plus a non-replaced control. (Figures 3.3, 3.6, 3.4, 3.5) We then measured the levels of HGT use, the fitness effect of HGT mutations, as well as the fraction of beneficial phenotype-switching mutations. In the both-phase and non-replaced control, the mean fitness effect was largely neutral, while the on-phase fragments had a significantly positive mean fitness benefit, and the off-phase fragments were largely negative.




Figure 3.2: bleh

Further, the control, both-phase and on-phase had a larger fraction of beneficial phenotype-switching mutants , whereas the off-phase treatment had few to no beneficial phenotype switches. This indicates that it is not only genetic disruption that is beneficial, but that the majority of the benefit of HGT derives from taking up and integrating useful fragments of information from time-periods where organisms were already adapted to the environment, thus bypassing the need to evolve a new phenotype from scratch.

3.3.2.1 Expected Fitness Effects

We measured the average fitness effects of fragments in a randomly selected replicate of the non-replaced HGT control treatment at the end of the last environmental cycle, and generated a distribution of fragments sorted the fragments by age and fitness-effect (Figures 3.3). We can see a clear pattern of more beneficial fitness effects from fragments of organisms originating in the matching cycle phase.




Figure 3.3: bleh

In order to quantify this result, we performed experiments where we replaced the fragments in the reservoir to contain only fragments originating in the matching phase, the off-phase, and mixture of both phases. We measured the mean expected fitness effects of fragments in these treatments (Figure 3.4). We found that HGT mutations are, on average, neutral in both the non-replaced control and the replaced-”both” treatment. The average fitness effect in the ”on-phase” treatment was mildly beneficial, while in the ”off-phase” treatment, the effect was strongly deleterious.

In our environments, the largest fitness benefits should occur when a mutation leads to the acquisition of a new rewarded task, or the loss of a punished task. Mutations that convey phenotypic change should have the largest impact. We identified the phenotypic effect of each fragment by counting the number of times that fragments produced a beneficial phenotype change, vs all HGT mutations (Figure 3.5). We see that a significantly larger proportion of




Figure 3.4: bleh

fragments in the "on-phase" treatment produce beneficial phenotype changes, as compared to fewer in the normal HGT and "both" treatments, and virtually zero in the "off-cycle" treatment.




Figure 3.5: bleh

These results suggests that any direct benefit of using HGT derives from taking up fragments that match the cycle phase of the environment of the affected organism, and thus likely contains information that relates to that environment.

Further, if fragments from the matching phase are indeed beneficial, we would expect

to see an increase in HGT use in those treatments, as compared to those where the fitness effects are mixed or deleterious. And indeed (Figure 3.6), we do observe just such an increase. Thus we can conclude that HGT in our changing environments is most beneficial when the fragments in the environment contain information that would be beneficial in that environment. However, even when no such information is available, HGT use is not significantly depressed as compared to the control treatment. This suggests that despite the lack of exclusively matched environmental information, that fragments could still provide some benefit.




Figure 3.6: bleh

3.4 Conclusion

Our experiments confirm our expectations about the evolution of horizontal gene transfer, and match what we would expect to see in nature. First, we find that grazing bonuses, such as those conveyed by the uptake of nutrient-rich DNA fragments in natural systems, cause an elevation of use of the horizontal gene transfer instruction, regardless of the potential risk of recombining uptaken gene fragments.

Further, when there is direct selection for phenotypic change - such as in a changing environment - we observe an increase in the use of the HGT instruction, even when no grazing bonus is offered. This indicates that the possibility of recombination itself is being selected for. Beyond this, we find that it is not only the fact of the recombination that is beneficial, but the information contained in the fragment. We found that most of the benefit conveyed by HGT derived from fragments that match the current environment, but also that when we filter so that only those fragments are available, that use of the HGT instruction increases even further.

In natural systems, this pattern is evident in the acquisition of antibiotic resistance in bacteria, where resistance is acquired by the spread of resistance genes via horizontal gene transfer. Therefore, our results provide a key insight into how horizontal gene transfer mechanisms could be selected for, and evolve in natural systems.

review the below. too strong? too informal?

It is worth noting that most prior studies of HGT have been observational studies that look at existing traits and phylogenies and attempt to make inferences about the evolution of those traits on that basis. Performing experimental evolution studies of the evolution of HGT in organic systems is something that, to a first approximation, we simply can't study in a wet system. Digital evolution, on the other hand, allows us to test many hypotheses about the evolution of HGT, such as the explicit benefits and consequences of different types of mutations, as well as gives us an opportunity to compare granular fragment grazing bonuses and quantify their effects on evolutionary outcomes. These are experiments that we would not be able to perform any other way.

CHAPTER 4

HORIZONTAL GENE TRANSFER VS OTHER TYPES OF MUTATION

4.1 Background

In both natural and artificial systems, populations evolve mechanisms to reduce or eliminate mutation[69], despite evidence that higher mutation rates may be optimal[70]. In artificial systems, prior research has failed to show an increase in endogenously-controlled mutation rates, when such an increase would be beneficial for long-term evolution, such as in a changing environment. However, Horizontal Gene Transfer through transformation seems to be exception to this rule.

[TODO
CITE]

In nature, transformation is a highly regulated and refined mechanism for taking up DNA fragments from the environment [71, 72, 73]. Clearly, evolution is acting to maintain and preserve these functions, despite the risk of bad recombinations. In the previous chapter, we showed that HGT is directly selected for in harsh changing environments. Despite the neutral to slightly deleterious fitness effects of recombination, HGT use is upregulated. What makes HGT different from other types of mutations? Is it the fitness effect? The information content of fragments? The probability for beneficial phenotype changes? How does HGT compare to other types of mutations in these respects? Can these differences account for the increased selection for HGT?

In order to address these question, we created instructions that create different kinds of mutational effects with similar per-instruction impact to HGT. We then performed experiments with these mutagenic instructions under the same conditions as our evaluation of HGT, in static vs. changing environments. We found that not only are these instructions triggered at significantly lowered rates as compared to HGT, but that the use rate correlates significantly with the amount of useful information conveyed by the mutation.

4.2 Methods

In this chapter, we use Avida to compare mutational effects of HGT to other types of mutations. As in the previous chapter, HGT is triggered by the HGT-Uptake instruction, and fragments are uptaken from reservoirs in the environment.

All experiments in this chapter compare outcomes between static environments and harsh changing environments. Similarly to the experiments in the previous chapter, task rewards in the harsh changing environment switch from a positive to a negative reward between the divided set of tasks in the Logic-9 environment. Please refer to Table 3.1 for more details. Each complete cycle lasts 1,000 updates, and the whole experimental run continues for 200,000 updates. Also as in the previous chapter, the static environment rewards executions of all the Logic-9 tasks at their default levels, with no reward switching.

4.2.1 Experimental Design

4.2.1.1 Alternative Mutagenic Instructions

In order to perform experiments comparing HGT to other types of mutagens, we modified the functioning of the HGT instruction to perform a new set of mutagenic functions which should have the same raw, per-instruction effect as the normal HGT instruction. All mutagens have the same 10% execution probability and viable recombination site requirements as the default HGT instruction.

- **HGT (Intact Fragment):** The default HGT operation. We uptake a fragment from the environment. We apply a 10% probability that the fragment is recombined into the organism’s executing genome. The recombination occurs homologously - that is, three instructions at the beginning and end of the fragment must match exactly with locations on the genome. We use the first valid match. We find valid matches by selecting a random location on the genome, then search from there. We loop around the whole genome, past the end and into the beginning, until we find a for the front

three instructions of the fragment. Then, we begin searching for a match for the back three instructions of the fragment, starting at the edge of the front match. If a valid back-end match site is not found, we scan forward, looking for the next front-end match and repeat the process until all possible match sites on the genome are exhausted. If no match is found, the recombination fails. If recombination succeeds, it replaces the content between the selected beginning and end match sites. This may result in the genome growing or shrinking, depending on the distance between the matches, and the length of the fragment.

- **HGT Shuffle:** We uptake a fragment from the environment, and a valid recombination site is found, as above. Prior to recombination, we shuffle the fragment, with the order of each instruction randomized. We then insert the fragment at the selected site.
- **HGT Random:** We uptake a fragment from the environment, and find a valid recombination site, as above. Prior to recombination, we entirely replace the fragment with an equal number of randomly selected instructions, and insert it at the selected site.
- **Mutation Event - Sampled:** We uptake a fragment from the environment, and find a valid recombination site, as above. Rather than recombine, we apply a number of point mutations in random locations throughout the genome. The number of applied mutations matches the number of instructions in the fragment. The mutations applied are sampled, with replacement, from the selected fragment. This mutagenic instruction is analogous to the the **HGT Shuffle** mutagen, except that instead of having a shuffled fragment be applied in a single location, the instructions from the fragment are sampled and applied randomly throughout the genome.
- **Mutation Event - Random:** We uptake a fragment from the environment, and find a valid recombination site, as above. Rather than recombine, we apply a number of random point mutations, matching the number of instructions in the fragment. This

mutagenic instruction is analogous to the **HGT Random** mutagen, except that, again, its effects are applied randomly throughout the genome instead of at a single location.

- **Mutation Rate Increase:** We uptake a fragment from the environment, and find a valid recombination site, as above. Rather than recombine, we increase the point-mutation rate for the organism, such that it will experience an expected additional number of point mutations matching the number of instructions in the uptaken fragment.
- **Die:** We uptake a fragment from the environment, and find a valid recombination site, as above. Rather than recombine, the organism dies.

4.2.1.2 Comparing HGT to Other Mutation Types

For the experiments comparing the mutational effects of different types of mutations, we provided populations of evolving digital organisms with instructions that endogenously trigger different kinds of mutation events (see above), then subjected them to both static and harsh changing environments (Table 4.1).

The goal of each treatment is to compare the use rates of each of the different mutagenic instruction across several factors. We can examine the effects of localization (concentrating mutations in a single area) by comparing the HGT–Random vs the ME–Random treatments. We can measure the effect of time-concentration (all mutations happening at once vs spread out over time) by comparing the ME–Random and MRI treatments. We can examine the effect of information content by comparing the HGT, HGT–Shuffle, and ME–Sampled treatments. We can identify a basement-level for mutagenic instruction use by comparing against the use of the entirely deleterious Die instruction. Finally, we can compare all these factors with their environment by comparing the static vs changing environment treatments.

Table 4.1: **Experimental Treatments - Mutation Types**

Treatment	Environment	Mutagenic Instructions
HGT	Static	HGT
HGT CE	Changing	
HGT-Shuffle	Static	HGT-Shuffle
HGT-Shuffle CE	Changing	
HGT-Random	Static	HGT-Random
HGT-Random CE	Changing	
ME-Sampled	Static	Mutation Event - Sampled
ME-Sampled CE	Changing	
ME-Random	Static	Mutation Event - Random
ME-Random CE	Changing	
MRI	Static	Mutation Rate Increase
MRI CE	Changing	
Die	Static	Die
Die CE	Changing	

Experimental treatments. Two types of environment (static vs changing environment), for each mutagen. No bonus was given for performing any of the mutagen instructions.

4.3 Results and Discussion

Our research shows that only mutations containing useful information are elevated in response to changing environments. Intact-fragment HGT is used most, with shuffled-fragment (HGT-Shuffle) used less, and the other, non information-bearing mutagen types used at the lowest levels. Further, we find a positive correlation between expected fitness and mutagenic instruction use. That is, mutagenic instructions that have higher expected fitness effects are used more. Finally, we see a strong positive correlation between the fraction of mutations from a mutagenic instruction that provides a beneficial phenotype-altering change, and the use of that instruction.

4.3.1 Other mutation types are not elevated in response to HGT

In order to compare the use of HGT against other kinds of mutation, we compared rates of non-bonus HGT fragment uptake against a execution of series of other mutation types, in

both static and changing environments. 4.1 As expected, in the static environment, endogenously-controlled performance of all types of mutation (including HGT) is strongly suppressed. However, in harsh changing environment, HGT use dominates the other mutation types. Of particular interest is how the use of the HGT-like instructions are used most by order of information content. The intact-HGT instruction is used most, followed by the HGT-Shuffle instruction, and then followed by HGT-Random, where no information remains. The latter performs comparably to the remaining mutation types: Mutation Events: ME-Sampled, ME-Random , MRI (Mutation Rate Increase), and the Die control. This indicates that the information content of the fragment is an important predictor of the use of HGT.

Also of note is that the only difference between the ME-Sampled and HGT-Shuffle treatments is the localization of the mutation effect (HGT-Shuffle fragment insertion is applied to a single location, whereas with ME-Sampled, the instructions are scattered randomly throughout the genome), however their use rates are significantly different. This indicates that not only is information content important, but that localization (clustering) of the mutation effect also matters.




Figure 4.1: bleh

4.3.2 Mutation fitness effect correlate with mutagen use

Prior research has failed to show an increase in endogenously-controlled mutation rates, when such an increase would be beneficial for long-term evolution, such as in a changing environment. Why should HGT mutations be different? In order to address this question, we measured the average fitness effect of each mutation, and compared it to that mutation's usage rate in changing environments. We observed a correlation between the average fitness effect, and the use of the mutagen. We observed many more positive or neutral fitness effects from HGT-like instructions than the other mutagen types, which exhibited primarily neutral or negative mutation effects. This indicates that mutagen use is under direct selection^{4.2}




Figure 4.2: bleh

4.3.3 HGT mutations increase evolved probability of beneficial phenotype switching

Further, we measured the proportion of mutations that produce beneficial phenotype changes.^{4.3} In the context of a changing environment, a mutation that switches an organism's phenotype to match the environment should be strongly selected for. Indeed, we found that this proportion was significantly and substantially higher for HGT-like mutations than other mutation types, and that there was strong correlation between the proportion of beneficial phenotype-switching mutations and mutagen use. This correlation is much stronger than

the correlation with fitness effect, indicating that the magnitude of the fitness effect is less important to selection than the sign of the effect.




Figure 4.3: bleh

stats

caption

4.4 Conclusion

Our experiments expand on our understanding of the evolution of horizontal gene transfer, and how it relates to other types of mutations in the context of changing environments. As expected, we find that the fitness effects of different types of mutations vary significantly, and that they predict the levels of use of that mutagen. In particular, we find a strong correlation between the fraction of mutations that produce a beneficial phenotype change, and the use of that mutagen. Thus, in the context of a changing environment, where there are strong selective pressures to change your phenotype, levels of HGT are elevated, while levels of other mutagens remain low.

This result matches prior research that shows that populations will tend to depress the use of mutagens, even though, in the long-term, higher mutation rates are optimal [CITE]. This result is further evidence that selection tends to act in the short term. Thus, even though certain kinds of mutagens and architectural features may promote long-term evolvability, these features are hitchhiking on short-term adaptive benefits. Thus, our research shows that even though evolvability may evolve, it is not necessarily selected for in the long-term.

CHAPTER 5

CHANGING ENVIRONMENTS AND LONG TERM EVOLVABILITY

5.1 Background

For longer timescales, evolvability is concerned more with variability generation and exploration of neutral spaces. Populations that exhibit this kind of evolvability would possess genomes with genetic architectures that more easily traverse the mutational landscape along neutral roads and thereby discover new fitness peaks while avoiding crossing fitness valleys. This kind of evolvability would allow populations to more easily colonize new ecological niches and form new clades[2, 12].

Despite some common features, the relationship between short-term and long-term evolvability is not obvious. Architectural features and evolutionary pressures that might convey short-term evolvability may not be the same as those that confer longer-term evolvability[13]. For example, features that promote rapid adaptation to a harsh fluctuating environment might reduce fitness in constant or benign fluctuating environments as compared to that of wild-type invaders. Alternately, the adaptation to harsh fluctuating environments and the resulting bottlenecks would potentially reduce diversity to the point where large amounts of neutral novelty generation could not occur.

ch5 - background - fill this in about the prior research done in this area

Prior research has shown that...

In this chapter, we examine how changing environments or novel mutagens affect long-term evolutionary potential. We introduce change-evolved populations from prior experiments with changing environments, and the evolution of HGT, to an entirely new environment, where we can assess long-term variability generation in the form of new task discovery and exploration.

5.2 Methods

5.2.1 Experimental Design

In order to examine the long-term dynamics and mechanisms of evolving populations in changing environments, we performed three sets of experiments. For each set of experiments, we take populations evolved in conditions where short-term evolvability pressures dominate, and we introduce the resulting populations to a completely new environment, with a new set of rewarded bitwise tasks to perform: Logic-77. These new tasks use three bitwise inputs rather than two, and are each rewarded with a constant 1.2-fold bonus to execution. This reward structure provides a mild selective pressure to evolve these task, but the benefits to performing them do not overwhelm the existing selective pressure to continue performing the tasks rewarded by the original changing or static environment.

The first set of experiments examines the dynamics of organisms that originally evolved in simple changing environments. These are a continuation of the experiments described in Chapter 2. The second set of experiments takes populations evolved in a more complex changing environment where we evolve the organisms to switch back and forth between two halves of the Logic-9 task set. For the final set of experiments, we take populations that evolved in a complex changing environment, similar to the second set, but with the addition of the HGT mutagen. These are a continuation of experiments described in Chapter 3.

With these experiments, we will explore the characteristics of these populations, and how different kinds of environmental conditions do or do not produce long-term evolvability.

5.2.1.1 Simple Cyclic Changing Environments

We took populations evolved under the experimental conditions detailed in Chapter 2, and introduced them to an entirely new set of rewarded bitwise logical tasks: the Logic-77 environment.

CH5 methods - more details about each treatment

Table 5.1: **Experimental Treatments - Simple Cyclic Changing Environment**

Treatment	Changing Env. Type	Rewarded Tasks				
		Phase 1 (0-200,000 Updates)		Phase 2 (200,000-400,000 Updates)		
		XOR	EQU	XOR	EQU	Logic-77 (all)
Control	None (static)	constant 2^3	constant 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Benign	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	benign fluctuating 0 or 2^5	constant $2^{0.3}$
Benign Quiesce	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Harsh	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	harsh fluctuating -2^5 or 2^5	constant $2^{0.3}$
Harsh Quiesce	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$

Experimental treatments. Four types of cyclic changing environment, plus a static control. Each is split into two phases. The first phase is a normal changing environment like those found in Chapter 2, Table 2.1. The second phase introduces an additional set of tasks (Logic 77) that are rewarded at a lower rate.

As in Chapter 2, we held the individual genomes at a fixed length of 121 instructions, and applied mutations after each successful replication event at a substitution probability of 0.00075 per site. Again, we configured the Avida world to have local interactions on a toroidal grid that is 60-by-60 cells (3600 cells in total), and continued the evolution of populations where we left off in Chapter 2.

5.2.1.2 Complex Cyclic Changing Environment

The second set of experiments looks at the evolution of populations of organisms with variable-length genomes, evolved in a more complex cyclic changing environment. This

Table 5.2: Experimental Treatments - Complex Cyclic Changing Environment

Treatment	Changing Env. Type	Rewarded Tasks				
		Phase 1 (0-200,000 Updates)		Phase 2 (200,000-400,000 Updates)		
		XOR	EQU	XOR	EQU	Logic-77 (all)
Control	None (static)	constant 2^3	constant 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Benign	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	benign fluctuating 0 or 2^5	constant $2^{0.3}$
Benign Quiesce	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Harsh	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	harsh fluctuating -2^5 or 2^5	constant $2^{0.3}$
Harsh Quiesce	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$

Experimental treatments. Four types of cyclic changing environment, plus a static control. Each is split into two phases. The first phase is a normal changing environment similar to those found in Chapter 2, Table 2.1. The second phase introduces an additional set of tasks (Logic 77) that are rewarded at a lower rate.

environment and genetic structure derives from the experiments described in 3. However, these populations were evolved without a working HGT instruction.

TODO fill this in with more detail of each treatment

update
table to
match
HGT

5.2.1.3 Complex Cyclic Changing Environment with HGT mutagen

The last set of experiments looks at the evolution of populations of organisms that evolved in a more complex cyclic changing environment, while having access to the HGT mutagen. These experiments are continuations of those described in 3, and continue the evolution of

the populations evolved there.

TODO fill this in with more detail of each treatment

5.2.2 Measuring Task Discovery and Task Performance

In order to assess how populations adapt to a brand new environment, we measure Task Performance and Task Discovery.

5.2.2.1 Task Performance

Task Performance is measured by counting the number of unique non-ephemeral tasks that a population is performing. Each task may be performed only once per organism, yielding a maximum task count of 3600 at any given time. Thus, we define a non-ephemeral task as one that a population is performing at equal to or more than 0.1% level, or 3.6 times per sampling.

5.2.2.2 Task Discovery

We measure task by counting the number of unique non-ephemeral tasks that have been discovered by a population, starting at a defined point in time. Once a task is discovered, it may not become "undiscovered." Thus, task discovery counts will always increase monotonically. We measure "Overall" task discovery, counting all possible tasks - including Logic-77 which were available, but not rewarded in the first phase of experiments - and also "Post-Reward" task discovery, where we begin counting new tasks from the beginning of the second phase of the experiment, once Logic-77 rewards are applied.

5.3 Results and Discussion

Our experiments demonstrate that the harshness of fluctuation has a dramatic effect on long-term evolvability, landscape exploration and task discovery. We find that while harsh changing environments were most effective at driving short-term adaptation, these

Table 5.3: Experimental Treatments - Complex Cyclic Changing Environments with HGT

Treatment	Changing Env. Type	Rewarded Tasks				
		Phase 1 (0-200,000 Updates)		Phase 2 (200,000-400,000 Updates)		
		XOR	EQU	XOR	EQU	Logic-77 (all)
Control	None (static)	constant 2^3	constant 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Benign	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	benign fluctuating 0 or 2^5	constant $2^{0.3}$
Benign Quiesce	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Harsh	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	harsh fluctuating -2^5 or 2^5	constant $2^{0.3}$
Harsh Quiesce	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Control	None (static)	constant 2^3	constant 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Benign	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	benign fluctuating 0 or 2^5	constant $2^{0.3}$
Benign Quiesce	Cyclic	constant 2^3	benign fluctuating 0 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$
Harsh	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	harsh fluctuating -2^5 or 2^5	constant $2^{0.3}$
Harsh Quiesce	Cyclic	constant 2^3	harsh fluctuating -2^5 or 2^5	constant 2^3	constant 2^5	constant $2^{0.3}$

Experimental treatments. Four types of cyclic changing environment, plus a static control. Each is split into two phases. The first phase is a normal changing environment similar to those found in Chapter 2, Table 2.1. The second phase introduces an additional set of tasks (Logic 77) that are rewarded at a lower rate.

environments depress long-term evolvability once rewards are in place. In all experiments, we find that benign environments produce the highest levels of task discovery, far outperforming harsh changing environments. We also find that the HGT mutagen, which helps with short-term evolvability in cyclic environments, does not appear to have a beneficial effect.

5.3.1 Long-Term Evolvability in Simple Cyclic Changing Environments

5.3.1.1 Benign changing environments outperform harsh environments in task discovery

We found that once the Logic-77 reward structure is introduced, populations evolving in harsh changing environments discover many fewer tasks than those evolving in benign changing environments (Fig 5.1). We hypothesize that this effect is due to the relative differences stats in the strength of selection between the harsh changing environment and the directional selection toward the logic-77 tasks. The pressure to gain and lose the fluctuating task is much stronger than the pressure to acquire the logic-77 tasks, thereby depressing the rate at which they are found. In contrast, the benign environment, with its comparatively weaker strength of selection for EQU task gain and loss, experiences a comparatively stronger selective pressure to acquire logic-77 tasks, while still benefiting from the increased exploration rate conveyed by the benign changing environment.

Interestingly, in those populations where we removed the harsh environmental change caption as we introduce Logic-77, we see that task exploration recovers somewhat and reaches a comparable level to the control. This suggests that it is the active pressures of the harsh stats environment that depress task exploration, and not a penalty conveyed by architectural features that persist after selective pressures have changed.

Of similar interest is the comparison of task discovery rates between the benign changing environment populations BenignCE and BenignQuiescent. The BenignCE populations continue to be subject to the benign changing environment, whereas the BenignQuiescent

make a
figure to
highlight
this?
rename




Figure 5.1: Number of new logic-77 tasks discovered beginning in phase 2.

populations are only subject to directional selection toward the evolution of XOR, EQU, and the Logic-77 tasks. The task discovery rate for the BenignCE population is slightly higher than for BenignQuiescent, but the effect is not statistically significant. Both, however, still stats perform better than the control treatment. Thus, this result suggests that there may be stats architectural features conferred by evolution in the changing environment that are helpful for long-term evolvability, even after the environmental changes have stopped, and that this effect is distinct from the direct effects of the changing direction of selection.

5.3.1.2 Benign changing environments outperform harsh environments in task performance

Similar to task discovery, populations evolving in harsh changing environments perform far fewer distinct tasks than either the benign-evolved or control populations (Fig 5.2). While stats the Benign-evolving populations seem to outperform the control, this effect is not statistically significant. stats




Figure 5.2: Number of new logic-77 tasks performed beginning in phase 2.

5.3.1.3 Harsh changing environments drive populations across the mutational landscape

In the first phase of the experiments, despite the logic-77 tasks not being rewarded, both changing-environment treatments discover more new tasks than the control. The harsh environment treatments in particular discover significantly more Logic-77 tasks than either the benign treatment, or the control, despite these tasks not being rewarded (Fig 5.3. We hypothesize that this effect may be due to the large phylogenetic depth of the harsh-evolved populations, where the repeated bottlenecks drive the populations along a kind of forced march across the mutational landscape. However, as the experiment proceeds, and Logic-77 rewards are introduced, this effect disappears, and benign environments quickly dominate in task discovery.

REPLACE THIS WITH THE ONE THAT HAS THE VIOLIN PLOT?

stats

stats

caption




Figure 5.3: Number of new logic-77 tasks discovered over time. [TODO DESCRIPTION AND STATS]

5.3.2 Long Term Evolvability and Complex Changing Environments

5.3.2.1 Benign changing environments outperform harsh environments in task discovery and task performance

Similar to the results of the simple environments, Benign-evolved populations out-perform Harsh-evolving populations in complex changing environments in measures of Task Performance (Fig 5.4) and Task Discovery (Fig 5.5). Again, this result suggests that, despite the differences in genome composition and environmental complexity, Benign changing environments, with their gentler selective pressures, or possibly with more intact chunks of undisturbed functionality in their genomes, leave more room for populations to acquire new tasks without the risk of harsh punishments.

Interestingly, populations that were initially evolved in a harsh changing environment and then change ceased upon introduction of Logic-77 rewards (HarshQuiescent) seem to




Figure 5.4: Number of logic-77 tasks performed over time. [TODO DESCRIPTION AND STATS]




Figure 5.5: Number of new logic-77 tasks discovered post-reward. [TODO DESCRIPTION AND STATS]

out-perform all other environment types in task performance (Fig 5.4). This effect may be stats due to a lingering architectural effect of evolving in harsh changing environments. Further research is needed to identify a mechanism for this effect.

5.3.3 Long Term Evolvability and Horizontal Gene Transfer

The addition of mutagens are thought to promote long-term evolvability[CITE] In this section

move below to the introduction

we examine the long-term evolutionary dynamics of adding an HGT mutagen to populations introduced to a brand new, complex environment.

5.3.3.1 New environments do not promote the use of HGT

Contrary to our expectations, none of our populations previously evolved to use HGT showed a significant increase in HGT fragment uptakes as a result of being introduced to a new environment. In general, levels remain constant, and in one notable example, levels appear to decrease, though this effect is not statistically significant (Fig 5.6).




Figure 5.6: **HGT Use, pre and post introduction of new logic-77 tasks.** [TODO DESCRIPTION AND STATS]

5.3.3.2 Fragment grazing bonus reduces task discovery and task performance

In most of our environmental treatments, applying an HGT Uptake grazing bonus results in reduced levels of both task performance and task discovery as compared to non-Bonus treatments (Fig 5.7, and Fig 5.8).

Further, we see a strong negative correlation between task discovery and task performance in populations where an HGT grazing bonus is applied, regardless of environmental treatment (Fig 5.9. We see no such correlation in non-Bonus HGT populations (Fig 5.10). This result




Figure 5.7: **Task Performance, with and without HGT Grazing Bonus.** [TODO DESCRIPTION AND STATS]





Figure 5.8: **Task Discovery, with and without HGT Grazing Bonus.** [TODO DESCRIPTION AND STATS]

suggests that there is potentially an intrinsic architectural difference between populations evolved to perform HGT for a grazing bonus vs populations that use HGT without a bonus. Or it is possible that the selective pressure to perform HGT outweighs the pressure to acquire new tasks when a simple bonus is available simply by executing a single instruction. More experiments will be required to identify the mechanism behind this effect.






Figure 5.9: **HGT-Bonus** vs Task Performance and Discovery. [TODO DESCRIPTION AND STATS]




Figure 5.10: **HGT-noBonus** vs Task Performance and Discovery. [TODO DESCRIPTION AND STATS]

5.3.3.3 Benign changing environments outperform harsh environments in task discovery and task performance in populations with HGT

Again, consistent with earlier results, populations with HGT follow the same pattern as the earlier experiments. Benign-evolved populations strongly outperform Harsh-evolving in measures of Task Discovery and Task Performance (Fig. 5.7 and Fig. 5.8). These results stats show that regardless of access to an information rich mutagen, the dominant effect is one of balancing of fitness pressures.

5.4 Conclusion

Our experiments show that changing environments both directly and indirectly influence long-term evolvability by directly driving exploration across the fitness landscape, and also by creating genetic architectures that increase the rate of new task discovery in novel environments. We hypothesize that these effects are driven by the presence of the accumulated pseudogene-like structures that provide cryptic functionality. In our experiments, the increased rate of task discovery is significant, though relatively small, but we expect that future experiments focused on this specific effect might yield more conclusive results.

fill this in, recapitulating the argument and framing about balancing of fitness effects.

CHAPTER 6

CONCLUSION

FILL THIS IN - Synthesis of all results and how it all fits together to paint a picture of how CE works.

This result matches prior research that shows that populations will tend to depress the use of mutagens, even though, in the long-term, higher mutation rates are optimal [CITE].

This result is further evidence that selection tends to act in the short term. Thus, even though certain kinds of mutagens and architectural features may promote long-term evolvability, these features are hitchhiking on short-term adaptive benefits. Thus, our research shows that even though evolvability may evolve, it is not necessarily selected for in the long-term.

6.1 Limitations of Cyclic Changing Environments

Changing environments produce a set of selective pressures that speed up exploration of genotype space, while also building reservoirs of partial functionality that may be co-opted in the evolution of more complex structures. These features make changing environments useful for both their explanatory power in natural evolution, and as practical tools in the Artificial Life toolkit. Ultimately, however, cyclic changing environments only re-tread existing phenotypic ground, and though genotypic exploration is faster than under purely directional or stabilizing selection, the space explored remains constrained by the type of phenotypes that are selected. Despite this constraint, however, we see that, particularly under harsh conditions, a lot of novel genotypic ground may be explored, even without direct selection for novelty.

Even so, there must exist methods of exploring genotype space that do not suffer from these limitations at all. For example, perhaps repeated bottlenecking of populations could promote faster traversal of the fitness landscape in quasi-random directions. More ambi-

tiously, perhaps these kinds of environments could be coupled with dynamically increasing open-ended complexity goals.

Understanding the mechanisms by which select environmental conditions alter fitness landscapes is vital to understanding the forces that promote evolvability and increase complexity. In particular, understanding the role of vestigial sites may help us untangle how robustness can promote evolvability. Are these vestigial sites inactive remnants, reservoirs of function, or are they part of a complex compensatory framework supporting and buffering the expression of the phenotype? Or both? Changing environments provide one view into these dynamics, but we must explore further to find other mechanisms for exploring and exploiting genotype space.

tweak
below?

APPENDICES

APPENDIX A

EXPERIMENTALLY DERIVING PARAMETERS FOR CHANGING ENVIRONMENT CYCLE LENGTHS

Cyclic changing environments have predictable cycles and thus have consistent periods of time that allow adaptation and fixation of traits. Too short a cycle, and traits are not able to fix. Too long a cycle and traits that are no longer selected for in one phase, but are useful in the other, become increasingly vulnerable to loss through genetic drift.

In order to identify an optimal cycle time to support our experiments, we surveyed a series of cycle lengths, and measured task performance rates in both benign and harsh changing environments (Fig A.1).

We found that, of our surveyed values, a cycle time of 1,000 updates (roughly 30 generations) both provided adequate time for traits to evolve and fix (and thus be performed by the entire population), but also not enough time for drift to destroy alternate-phase traits, thus minimizing the time required to re-evolve the task. Cycle times that were too short never acquired and fixed adaptive traits in the entire population, while as cycles got longer, the fluctuating tasks would take much longer to re-evolve, indicating that the populations had drifted away from the regions of the mutational landscape that contained them (Fig A.2).

stats

caption

stats

caption




Figure A.1: **Survey of task execution** for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]




Figure A.2: **Survey of task execution** for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]

APPENDIX B

EXPERIMENTALLY DERIVING PARAMETERS FOR HGT RECOMBINATION PROBABILITY AND BONUS LEVELS

B.1 Recombination Probability Sweep

In nature, the probability of horizontal gene transfer occurring varies wildly by species, environmental conditions, and mechanism of action. In order to identify an optimal probability of HGT uptake resulting in recombination, we performed a series of experiments comparing different probabilities, coupled with a pair of basic grazing bonuses. Our initial grazing bonuses were based on the reward given for a similarly complex task - NAND - which only requires a single instruction, plus an IO instruction, to implement. NAND is typically rewarded with a bonus of 2^1 to execution speed. Thus, performing HGT, which only requires a single instruction - HGT-Uptake, was rewarded similarly.

The goal was to identify a probability of recombination that would, without a grazing bonus, result in a reduced HGT uptake level as compared to with a grazing bonus. Further, we wanted the recombination probability to not be so high as to result in mutational meltdown, thus frequently killing the populations. We found that a level of 0.1 probability met these characteristics. That is, recombination probabilities of less than 0.1 had similar HGT expression rates with and without a bonus, while probabilities above 0.1 tended to have many many fewer surviving populations (Fig B.1).

B.2 Bonus Sweep

Similarly, in nature, the amount of nutritive benefit derived from taking up DNA fragments from the environment is difficult to quantify. In Avida, when populations are presented with instructions that provide a direct boost to execution speed, use of these instructions

stats

caption




Figure B.1: **Survey of hgt use** for a series of recombination probabilities. [TODO DESCRIPTION AND STATS]

is strongly selected for. Populations rapidly fill their genomes with these instructions, to the exclusion of virtually all else. Thus, our goal in selecting potential bonus levels lay in finding a range of minimal value that would balance the presumably deleterious effects of the HGT uptake instruction, without completely overwhelming the balance of selective pressures we wished to apply to our experiments. As expected, and coupled with the recombination probability above, values around 2^1 seemed to perform the best (Fig B.2). Above a value of 2^2 , there seem to be diminishing returns.

B.3 Comparable Bonus Values

For later experiments, in order to identify the relationship between HGT uptakes prompted by grazing bonus, and uptakes prompted by evolvability pressures in changing environments, we surveyed a series of bonuses to find a value that, in a static environment, would yield an HGT fragment uptake rate similar to that found in changing environments without a grazing bonus. We found that a bonus value of $2^{0.8}$ yielded similar HGT fragment uptake levels (Fig B.3).

fix up figure to include CE uptake value, plus less terrible

stats
caption




Figure B.2: Survey of task execution for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]




Figure B.3: Survey of comparable bonus values for a series of cycle-lengths. [TODO DESCRIPTION AND STATS]

APPENDIX C

CHANGING ENVIRONMENTS ELEVATE ALL INSTRUCTION USE DUE TO REPEATED BOTTLENECKS

In Chapter 4, we observe that the Die mutagen control is also elevated in the changing environments treatment. This counterintuitive effect is due to the repeated bottlenecking of the populations subjected to the harsh changing environment.C.1. We observe that under increasingly harsh bottlenecks, rates of execution of the "die" instruction increase to levels comparable to what is seen in the changing environment treatments. Because not all instructions contained in an organism's genome are necessarily executed (due to genetic flow-control structures), it is possible for any instruction to remain dormant. Thus, not all organisms with the "die" instruction in their genome will express it. Thus, the selective pressures of the environmental change may ultimately outweigh that of the die instruction if it is initially dormant in a genome that would survive a harsh environmental change and reproduce.




Figure C.1: bleh

rework
figure
stats
caption

BIBLIOGRAPHY

BIBLIOGRAPHY

- [1] John Maynard Smith and Eors Szathmary. The major evolutionary transitions. *Nature*, 374(6519):227–32, March 1995.
- [2] Marc Kirschner and John Gerhart. Evolvability. *Proceedings of the National Academy of Sciences*, 95(15):8420–8427, 1998.
- [3] Richard Dawkins. 13 - The evolution of evolvability. In Sanjeev Kumar and Peter J. Bentley, editors, *On Growth, Form and Computers*, pages 239–255. Academic Press, London, 2003.
- [4] Lee Altenberg. The evolution of evolvability in genetic programming. *Advances in genetic programming*, 3:47–74, 1994.
- [5] Ronald Aylmer Fisher. *The genetical theory of natural selection*. Clarendon Press, Oxford, 1930.
- [6] David Houle. Comparing Evolvability and Variability of Quantitative Traits. *Genetics*, 130(1):195–204, 1992.
- [7] Terry Van Belle and David H. Ackley. Code Factoring And The Evolution Of Evolvability. In *Proceedings of the Genetic and Evolutionary Computation Conference*, GECCO '02, pages 1383–1390, San Francisco, CA, USA, 2002. Morgan Kaufmann Publishers Inc.
- [8] Gunter P. Wagner and Lee Altenberg. Perspective: complex adaptations and the evolution of evolvability. *Evolution*, pages 967–976, 1996.
- [9] Andreas Wagner. Robustness, evolvability, and neutrality. *FEBS letters*, 579(8):1772–1778, 2005.
- [10] Hiroaki Kitano. Biological robustness. *Nature Reviews Genetics*, 5(11):826–837, November 2004.
- [11] P. Alberch. From genes to phenotype: dynamical systems and evolvability. *Genetica*, 84(1):5–11, 1991.
- [12] J. F. Y Brookfield. Evolution: The evolvability enigma. *Current Biology*, 11(3):R106–R108, February 2001.
- [13] Massimo Pigliucci. Is evolvability evolvable? *Nature Reviews Genetics*, 9(1):75–82, 2008.
- [14] Samir Okasha. *Evolution and the Levels of Selection*, volume 16. Clarendon Press Oxford, 2006.

- [15] Stephen Jay Gould and Richard C. Lewontin. The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proceedings of the Royal Society of London B: Biological Sciences*, 205(1161):581–598, 1979.
- [16] Sewall Wright. Evolution in Mendelian Populations. *Genetics*, 16(2):97–159, March 1931.
- [17] Nadav Kashtan, Elad Noor, and Uri Alon. Varying environments can speed up evolution. *Proceedings of the National Academy of Sciences*, 104(34):13711–13716, August 2007.
- [18] Brian D. Connolly, Katherine J. Dickinson, Sarah P. Hammarlund, and Benjamin Kerr. Negative niche construction favors the evolution of cooperation. *Evolutionary Ecology*, pages 1–17, October 2015.
- [19] Barbara McClintock. *The significance of responses of the genome to challenge*. Singapore: World Scientific Pub. Co, 1993.
- [20] Anton Crombach and Paulien Hogeweg. Evolution of Evolvability in Gene Regulatory Networks. *PLOS Comput Biol*, 4(7):e1000112, July 2008.
- [21] Claus O. Wilke, Jia Lan Wang, Charles Ofria, Richard E. Lenski, and Christoph Adami. Evolution of digital organisms at high mutation rates leads to survival of the flattest. *Nature*, 412(6844):331–3, July 2001.
- [22] Matthew R. Goddard, H. Charles J. Godfray, and Austin Burt. Sex increases the efficacy of natural selection in experimental yeast populations. *Nature*, 434(7033):636+, March 2005. 636.
- [23] Thomas F. Hansen and David Houle. Measuring and comparing evolvability and constraint in multivariate characters. *Journal of evolutionary biology*, 21(5):1201–1219, 2008.
- [24] Trevor Price and Dolph Schluter. On the Low Heritability of Life-History Traits. *Evolution*, 45(4):853–861, 1991.
- [25] Thomas F. Hansen, Christophe Plabon, and David Houle. Heritability is not Evolvability. *Evolutionary Biology*, 38(3):258–277, June 2011.
- [26] Dusan Misevic, Charles Ofria, and Richard E Lenski. Sexual Reproduction Reshapes the Genetic Architecture of Digital Organisms. *Proceedings. Biological Sciences / The Royal Society*, 273(1585):457–464, February 2006.
- [27] Gnter P. Wagner and Jianzhi Zhang. The pleiotropic structure of the genotypephenotype map: the evolvability of complex organisms. *Nature Reviews Genetics*, 12(3):204–213, 2011.
- [28] E. Ravasz, A.L. Somera, D.A. Mongru, Z.N. Oltvai, and A.-L. Barabasi. Hierarchical organization of modularity in metabolic networks. (Reports). *Science*, 297(5586):1551+, August 2002. 1551.

- [29] Thomas F Hansen. Is modularity necessary for evolvability?: Remarks on the relationship between pleiotropy and evolvability. *Biosystems*, 69(23):83–94, May 2003.
- [30] Mihaela Pavlicev and Gnter P. Wagner. A model of developmental evolution: selection, pleiotropy and compensation. *Trends in Ecology & Evolution*, 27(6):316–322, June 2012.
- [31] Jason G. Mezey, James M. Cheverud, and Gnter P. Wagner. Is the Genotype-Phenotype Map Modular?: A Statistical Approach Using Mouse Quantitative Trait Loci Data. *Genetics*, 156(1):305–311, September 2000.
- [32] J. Arjan G. M. de Visser, Joachim Hermisson, Gnter P. Wagner, Lauren Ancel Meyers, Homayoun Bagheri-Chaichian, Jeffrey L. Blanchard, Lin Chao, James M. Cheverud, Santiago F. Elena, Walter Fontana, Greg Gibson, Thomas F. Hansen, David Krakauer, Richard C. Lewontin, Charles Ofria, Sean H. Rice, George von Dassow, Andreas Wagner, and Michael C. Whitlock. Perspective: Evolution and Detection of Genetic Robustness. *Evolution*, 57(9):1959–1972, 2003.
- [33] James M. Whitacre. Degeneracy: a link between evolvability, robustness and complexity in biological systems. *Theoretical Biology and Medical Modelling*, 7(1):6, 2010.
- [34] Andreas Wagner. Distributed robustness versus redundancy as causes of mutational robustness. *BioEssays*, 27(2):176–188, February 2005.
- [35] Jack E. Baldwin and Hans Krebs. The evolution of metabolic cycles. *Nature*, 291(5814):381–382, June 1981.
- [36] Jrg Stelling, Uwe Sauer, Zoltan Szallasi, Francis J. Doyle III, and John Doyle. Robustness of Cellular Functions. *Cell*, 118(6):675–685, September 2004.
- [37] Gunter P. Wagner, Mihaela Pavlicev, and James M. Cheverud. The road to modularity. *Nature Reviews Genetics*, 8(12):921+, December 2007. 921.
- [38] Andreas Wagner. Robustness and evolvability: a paradox resolved. *Proceedings of the Royal Society of London B: Biological Sciences*, 275(1630):91–100, 2008.
- [39] Charles Ofria and Christoph Adami. Evolution of genetic organization in digital organisms. In *Evolution as Computation*, pages 296–313. Springer, 2002.
- [40] P. McKinley, B. Cheng, C. Ofria, D. Knoester, B. Beckmann, and H. Goldsby. Harnessing Digital Evolution. *Computer*, 41(1):54–63, January 2008.
- [41] Francis J. Ryan. Evolution observed. *Scientific American*, 189:78–82, 1953.
- [42] Richard E. Lenski, Michael R. Rose, Suzanne C. Simpson, and Scott C. Tadler. Long-term experimental evolution in *Escherichia coli*. I. Adaptation and divergence during 2,000 generations. *The American Naturalist*, pages 1315–1341, 1991.
- [43] Jennifer L. Anderson, Levi T. Morran, and Patrick C. Phillips. Outcrossing and the Maintenance of Males within *C. elegans* Populations. *Journal of Heredity*, page esq003, January 2010.

- [44] S. C. Stearns, M. Ackermann, M. Doebeli, and M. Kaiser. Experimental evolution of aging, growth, and reproduction in fruitflies. *Proceedings of the National Academy of Sciences*, 97(7):3309–3313, March 2000.
- [45] Aisha I. Khan, Duy M. Dinh, Dominique Schneider, Richard E. Lenski, and Tim F. Cooper. Negative Epistasis Between Beneficial Mutations in an Evolving Bacterial Population. *Science*, 332(6034):1193–1196, June 2011.
- [46] Ivan G. Szendro, Martijn F. Schenk, Jasper Franke, Joachim Krug, and J. Arjan G. M. de Visser. Quantitative analyses of empirical fitness landscapes. *Journal of Statistical Mechanics: Theory and Experiment*, 2013(01):P01005, 2013.
- [47] Daniel M. Weinreich, Nigel F. Delaney, Mark A. DePristo, and Daniel L. Hartl. Darwinian evolution can follow only very few mutational paths to fitter proteins. *Science*, 312(5770):111–114, 2006.
- [48] Joshua R. Nahum, Peter Godfrey-Smith, Brittany N. Harding, Joseph H. Marcus, Jared Carlson-Steinermer, and Benjamin Kerr. A tortoisehare pattern seen in adapting structured and unstructured populations suggests a rugged fitness landscape in bacteria. *Proceedings of the National Academy of Sciences*, 112(24):7530–7535, June 2015.
- [49] P. Merz and B. Freisleben. Fitness landscape analysis and memetic algorithms for the quadratic assignment problem. *IEEE Transactions on Evolutionary Computation*, 4(4):337–352, November 2000.
- [50] J. Humeau, A. Liefooghe, E.-G. Talbi, and S. Verel. ParadisEO-MO: from fitness landscape analysis to efficient local search algorithms. *Journal of Heuristics*, 19(6):881–915, June 2013.
- [51] Leila Kallel, Bart Naudts, and Alex Rogers. *Theoretical aspects of evolutionary computing*. Springer Science & Business Media, 2013.
- [52] Sergey Gavrilets. *Fitness landscapes and the origin of species (MPB-41)*. Princeton University Press Princeton, NJ, 2004.
- [53] Dusan Misevic, Charles Ofria, and Richard E. Lenski. Experiments with Digital Organisms on the Origin and Maintenance of Sex in Changing Environments. *The Journal of Heredity*, 101(suppl 1):S46–S54, March 2010.
- [54] Luis Zaman, Justin R. Meyer, Suhas Devangam, David M. Bryson, Richard E. Lenski, and Charles Ofria. Coevolution drives the emergence of complex traits and promotes evolvability. *PLoS Biol*, 12(12):e1002023, 2014.
- [55] Justin R. Meyer, Devin T. Dobias, Joshua S. Weitz, Jeffrey E. Barrick, Ryan T. Quick, and Richard E. Lenski. Repeatability and Contingency in the Evolution of a Key Innovation in Phage Lambda. *Science*, 335(6067):428–432, January 2012.
- [56] Christopher D. Bayliss, Dawn Field, and E. Richard Moxon. The simple sequence contingency loci of *Haemophilus influenzae* and *Neisseria meningitidis*. *Journal of Clinical Investigation*, 107(6):657–666, March 2001.

- [57] Heewook Lee, Ellen Popodi, Haixu Tang, and Patricia L. Foster. Rate and molecular spectrum of spontaneous mutations in the bacterium *Escherichia coli* as determined by whole-genome sequencing. *Proceedings of the National Academy of Sciences*, 109(41):E2774–E2783, October 2012.
- [58] Ashley J. R. Carter, Joachim Hermisson, and Thomas F. Hansen. The role of epistatic gene interactions in the response to selection and the evolution of evolvability. *Theoretical Population Biology*, 68(3):179–196, November 2005.
- [59] Ines Chen and David Dubnau. DNA uptake during bacterial transformation. *Nature Reviews Microbiology*, 2(3):241+, March 2004. 241.
- [60] Joshua Lederberg and Edward L. Tatum. Gene recombination in *Escherichia coli*. *Nature*, 158:558, 1946.
- [61] Norton D. Zinder and Joshua Lederberg. Genetic Exchange in *Salmonella*. *Journal of Bacteriology*, 64(5):679–699, November 1952.
- [62] E. S. Lennox. Transduction of linked genetic characters of the host by bacteriophage P1. *Virology*, 1(2):190–206, July 1955.
- [63] Tal Dagan, Yael Artzy-Randrup, and William Martin. Modular Networks and Cumulative Impact of Lateral Transfer in Prokaryote Genome Evolution. *Proceedings of the National Academy of Sciences of the United States of America*, 105(29):10039–10044, 2008.
- [64] J. E. Davies. Origins, acquisition and dissemination of antibiotic resistance determinants. *Ciba Foundation Symposium*, 207:15–27; discussion 27–35, 1997.
- [65] Jos L. Martnez. Antibiotics and Antibiotic Resistance Genes in Natural Environments. *Science*, 321(5887):365–367, 2008.
- [66] Michiel Vos. Why do bacteria engage in homologous recombination? *Trends in Microbiology*, 17(6):226–232, June 2009.
- [67] R. J. Redfield, M. R. Schrag, and A. M. Dean. The Evolution of Bacterial Transformation: Sex with Poor Relations. *Genetics*, 146(1):27–38, May 1997.
- [68] Ana Paula Jacobus and Jeferson Gross. Optimal Cloning of PCR Fragments by Homologous Recombination in *Escherichia coli*. *PLOS ONE*, 10(3):e0119221, March 2015.
- [69] Sbastien Wielgoss, Jeffrey E. Barrick, Olivier Tenaillon, Michael J. Wiser, W. James Dittmar, Stphane Cruveiller, Batrice Chane-Woon-Ming, Claudine Mdigue, Richard E. Lenski, and Dominique Schneider. Mutation rate dynamics in a bacterial population reflect tension between adaptation and genetic load. *Proceedings of the National Academy of Sciences*, 110(1):222–227, January 2013.
- [70] Jeff Clune, Dusan Misevic, Charles Ofria, Richard E. Lenski, Santiago F. Elena, and Rafael Sanjun. Natural Selection Fails to Optimize Mutation Rates for Long-Term Adaptation on Rugged Fitness Landscapes. *PLoS Comput Biol*, 4(9):e1000187, 2008.

- [71] Jonathan M. Solomon and Alan D. Grossman. Who's competent and when: regulation of natural genetic competence in bacteria. *Trends in Genetics*, 12(4):150–155, April 1996.
- [72] Patrick Seitz and Melanie Blokesch. Cues and regulatory pathways involved in natural competence and transformation in pathogenic and environmental Gram-negative bacteria. *FEMS Microbiology Reviews*, 37(3):336–363, May 2013.
- [73] Laetitia Fontaine, Cline Boutry, Marie Henry de Frahan, Brigitte Delplace, Christophe Fremaux, Philippe Horvath, Patrick Boyaval, and Pascal Hols. A Novel Pheromone Quorum-Sensing System Controls the Development of Natural Competence in *Streptococcus thermophilus* and *Streptococcus salivarius*. *Journal of Bacteriology*, 192(5):1444–1454, March 2010.