

# JEE MAIN 2021

ONLINE

26<sup>th</sup> February  
1<sup>st</sup> Shift

## PHYSICS

### SECTION-A (MULTIPLE CHOICE QUESTIONS)

1. Four identical solid spheres each of mass ' $m$ ' and radius ' $a$ ' are placed with their centres on the four corners of a square of side ' $b$ '. The moment of inertia of the system about one side of square where the axis of rotation is parallel to the plane of the square is

- (a)  $\frac{8}{5}ma^2 + mb^2$  (b)  $\frac{4}{5}ma^2 + 2mb^2$   
(c)  $\frac{4}{5}ma^2$  (d)  $\frac{8}{5}ma^2 + 2mb^2$

2. If  $\lambda_1$  and  $\lambda_2$  are the wavelengths of the third member of Lyman and first member of the Paschen series respectively, then the value of  $\lambda_1 : \lambda_2$  is

- (a) 7 : 135 (b) 7 : 108 (c) 1 : 3 (d) 1 : 9

3. Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R.

**Assertion A :** An electron microscope can achieve better resolving power than an optical microscope.

**Reason R :** The de Broglie's wavelength of the electrons emitted from an electron gun is much less than wavelength of visible light.

In the light of the above statements, choose the correct answer from the options given below:

- (a) A is false but R is true.  
(b) Both A and R are true but R is not the correct explanation of A.  
(c) A is true but R is false.  
(d) Both A and R are true and R is the correct explanation of A.

4. A particle is moving with uniform speed along the circumference of a circle of radius  $R$  under the action of a central fictitious force  $F$  which is inversely proportional to  $R^3$ . Its time period of revolution will be given by

- (a)  $T \propto R^{4/3}$  (b)  $T \propto R^2$   
(c)  $T \propto R^{5/2}$  (d)  $T \propto R^{3/2}$


5. LED is constructed from Ga-As-P semiconducting material. The energy gap of this LED is 1.9 eV. Calculate the wavelength of light emitted and its colour.

$[h = 6.63 \times 10^{-34} \text{ J s and } c = 3 \times 10^8 \text{ m s}^{-1}]$

- (a) 1046 nm and red colour  
(b) 654 nm and red colour

- (c) 1046 nm and blue colour  
(d) 654 nm and orange colour

6. The temperature  $\theta$  at the junction of two insulating sheets, having thermal resistances  $R_1$  and  $R_2$  as well as top and bottom temperatures  $\theta_1$  and  $\theta_2$  (as shown in figure) is given by




- (a)  $\frac{\theta_1 R_1 + \theta_2 R_2}{R_1 + R_2}$  (b)  $\frac{\theta_2 R_2 - \theta_1 R_1}{R_2 - R_1}$   
(c)  $\frac{\theta_1 R_2 - \theta_2 R_1}{R_2 - R_1}$  (d)  $\frac{\theta_1 R_2 + \theta_2 R_1}{R_1 + R_2}$

7. The normal density of a material is  $\rho$  and its bulk modulus of elasticity is  $K$ . The magnitude of increase in density of material, when a pressure  $P$  is applied uniformly on all sides, will be

- (a)  $\frac{PK}{\rho}$  (b)  $\frac{\rho P}{K}$  (c)  $\frac{K}{\rho P}$  (d)  $\frac{\rho K}{P}$

8. Find the electric field at point  $P$  (as shown in figure) on the perpendicular bisector of a uniformly charged thin wire of length  $L$  carrying a charge  $Q$ . The distance of the point  $P$  from the centre of the rod is  $a = \frac{\sqrt{3}}{2}L$ .



- (a)  $\frac{Q}{3\pi\epsilon_0 L^2}$  (b)  $\frac{\sqrt{3}Q}{4\pi\epsilon_0 L^2}$   
(c)  $\frac{Q}{4\pi\epsilon_0 L^2}$  (d)  $\frac{Q}{2\sqrt{3}\pi\epsilon_0 L^2}$

9. A planet revolving in elliptical orbit has  
(A) a constant velocity of revolution.  
(B) has the least velocity when it is nearest to the sun.  
(C) its areal velocity is directly proportional to its velocity.  
(D) areal velocity is inversely proportional to its velocity.  
(E) to follow a trajectory such that the areal velocity is constant.

Choose the correct answer from the options given below:

- (a) C only (b) A only (c) E only (d) D only

10. Consider the combination of 2 capacitors  $C_1$  and  $C_2$ , with  $C_2 > C_1$ , when connected in parallel, the equivalent capacitance is  $\frac{15}{4}$  times the equivalent capacitance of the same connected in series. Calculate the ratio of capacitors,  $\frac{C_2}{C_1}$ .


(a)  $\frac{29}{15}$  (b)  $\frac{15}{11}$  (c)  $\frac{111}{80}$  (d)  $\frac{15}{4}$

11. In a typical combustion engine the work done by a gas

molecule is given by  $W = \alpha^2 \beta e^{\frac{-\beta x^2}{kT}}$ , where  $x$  is the displacement,  $k$  is the Boltzmann constant and  $T$  is the temperature. If  $\alpha$  and  $\beta$  are constants, dimensions of  $\alpha$  will be

(a)  $[M^2LT^{-2}]$  (b)  $[MLT^{-2}]$   
(c)  $[MLT^{-1}]$  (d)  $[M^0LT^0]$

12. Find the gravitational force of attraction between the ring and sphere as shown in the diagram, where the plane of the ring is perpendicular to the line joining the centres. If  $\sqrt{8}R$  is the distance between the centres of a ring (of mass ' $m$ ') and a sphere (mass ' $M$ ') where both have equal radius ' $R$ '.



(a)  $\frac{1}{3\sqrt{8}} \cdot \frac{GMm}{R^2}$  (b)  $\frac{2\sqrt{2}}{3} \cdot \frac{GMm}{R^2}$   
(c)  $\frac{\sqrt{8}}{27} \cdot \frac{GMm}{R^2}$  (d)  $\frac{\sqrt{8}}{9} \cdot \frac{GMm}{R^2}$

13. An alternating current is given by the equation  $i = i_1 \sin \omega t + i_2 \cos \omega t$ . The rms current will be

(a)  $\frac{1}{2}(i_1^2 + i_2^2)^{1/2}$  (b)  $\frac{1}{\sqrt{2}}(i_1^2 + i_2^2)^{1/2}$   
(c)  $\frac{1}{\sqrt{2}}(i_1 + i_2)^2$  (d)  $\frac{1}{\sqrt{2}}(i_1 + i_2)$

14. A large number of water drops, each of radius  $r$ , combine to have a drop of radius  $R$ . If the surface tension is  $T$  and mechanical equivalent of heat is  $J$ , the rise in heat energy per unit volume will be

(a)  $\frac{3T}{rJ}$  (b)  $\frac{2T}{rJ}$   
(c)  $\frac{3T}{J} \left( \frac{1}{r} - \frac{1}{R} \right)$  (d)  $\frac{2T}{J} \left( \frac{1}{r} - \frac{1}{R} \right)$

15. Assume that a tunnel is dug along a chord of the earth, at a perpendicular distance  $(R/2)$  from the earth's centre, where ' $R$ ' is the radius of the earth. The wall of

the tunnel is frictionless. If a particle is released in this tunnel, it will execute a simple harmonic motion with a time period

(a)  $\frac{1}{2\pi} \sqrt{\frac{g}{R}}$  (b)  $2\pi \sqrt{\frac{R}{g}}$  (c)  $\frac{g}{2\pi R}$  (d)  $\frac{2\pi R}{g}$

16. If two similar springs each of spring constant  $K_1$  are joined in series, the new spring constant and time period would be changed by a factor

(a)  $\frac{1}{2}, \sqrt{2}$  (b)  $\frac{1}{4}, 2\sqrt{2}$  (c)  $\frac{1}{2}, 2\sqrt{2}$  (d)  $\frac{1}{4}, \sqrt{2}$

17. Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R.

**Assertion A :** Body P having mass  $M$  moving with speed  $u$  has head-on collision elastically with another body Q having mass  $m$  initially at rest. If  $m \ll M$ , body Q will have a maximum speed equal to  $2u$  after collision.

**Reason R :** During elastic collision, the momentum and kinetic energy are both conserved.

In the light of the above statements, choose the most appropriate answer from the options given below:

(a) Both A and R are true but R is not the correct explanation of A.  
(b) A is correct but R is not correct.  
(c) Both A and R are correct and R is the correct explanation of A.  
(d) A is not correct but R is correct.


18. A short straight object of height 100 cm lies before the central axis of a spherical mirror whose focal length has absolute value  $|f| = 40$  cm. The image of object produced by the mirror is of height 25 cm and has the same orientation of the object. One may conclude from the information:

(a) Image is virtual, opposite side of convex mirror.  
(b) Image is real, same side of concave mirror.  
(c) Image is virtual, opposite side of concave mirror.  
(d) Image is real, same side of convex mirror.

19. In a Young's double slit experiment two slits are separated by 2 mm and the screen is placed one meter away. When a light of wavelength 500 nm is used, the fringe separation will be

(a) 0.75 mm (b) 0.25 mm (c) 1 mm (d) 0.50 mm

20. Five equal resistances are connected in a network as shown in figure. The net resistance between the points A and B is




(a)  $R$  (b)  $2R$  (c)  $R/2$  (d)  $3R/2$

## SECTION-B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

21. In an electrical circuit, a battery is connected to pass 20 C of charge through it in a certain given time. The potential difference between two plates of the battery is maintained at 15 V. The work done by the battery is \_\_\_\_\_ J.
22. A radiation is emitted by 1000 W bulb and it generates an electric field and magnetic field at  $P$ , placed at a distance of 2 m. The efficiency of the bulb is 1.25%. The value of peak electric field at  $P$  is  $x \times 10^{-1}$  V/m. Value of  $x$  is \_\_\_\_\_. (Rounded off to the nearest integer)  
[Take  $\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$ ,  $c = 3 \times 10^8 \text{ ms}^{-1}$ ]
23. The mass per unit length of a uniform wire is 0.135 g/cm. A transverse wave of the form  $y = -0.21 \sin(x + 30t)$  is produced in it, where  $x$  is in meter and  $t$  is in second. Then, the expected value of tension in the wire is  $x \times 10^{-2}$  N. Value of  $x$  is \_\_\_\_\_. (Round off to the nearest integer)
24. The circuit contains two diodes each with a forward resistance of  $50 \Omega$  and with infinite reverse resistance. If the battery voltage is 6 V, the current through the  $120 \Omega$  resistance is \_\_\_\_\_ mA.




25. The maximum and minimum amplitude of an amplitude modulated wave is 16 V and 8 V respectively. The modulation index for this amplitude modulated wave is  $x \times 10^{-2}$ . The value of  $x$  is \_\_\_\_\_.
26. A person standing on a spring balance inside a stationary lift measures 60 kg. The weight of that person if the lift descends with uniform downward acceleration of  $1.8 \text{ m/s}^2$  will be \_\_\_\_\_ N. [ $g = 10 \text{ m/s}^2$ ]
27. A boy pushes a box of mass 2 kg with a force  $\vec{F} = (20\hat{i} + 10\hat{j}) \text{ N}$  on a frictionless surface. If the box was initially at rest, then \_\_\_\_\_ m is displacement along the  $x$ -axis after 10 s.

28. As shown in figure, a block of mass  $\sqrt{3} \text{ kg}$  is kept on a horizontal rough surface of coefficient of friction  $\frac{1}{3\sqrt{3}}$ .

The critical force to be applied on the vertical surface as shown at an angle  $60^\circ$  with horizontal such that it does not move, will be  $3x$ . The value of  $x$  will be

$$[g = 10 \text{ m/s}^2, \sin 60^\circ = \frac{\sqrt{3}}{2}; \cos 60^\circ = \frac{1}{2}]$$







29. In a series  $LCR$  circuit, the quality factor is measured as 100. If the inductance is increased by two fold and resistance is decreased by two fold, then the quality factor after this change will be \_\_\_\_\_.
30. A container is divided into two chambers by a partition. The volume of first chamber is 4.5 litre and second chamber is 5.5 litre. The first chamber contain 3.0 moles of gas at pressure 2.0 atm and second chamber contains 4.0 moles of gas at pressure 3.0 atm. After the partition is removed and the mixture attains equilibrium, then, the common equilibrium pressure existing in the mixture is  $x \times 10^{-1}$  atm. Value of  $x$  is \_\_\_\_\_.

## CHEMISTRY







## SECTION-A (MULTIPLE CHOICE QUESTIONS)

31. The orbital having two radial as well as two angular nodes is  
(a)  $4f$  (b)  $4d$  (c)  $5d$  (d)  $3p$

32. The structure of neoprene is



33. Find A, B and C in the following reactions.



34. Compound A used as a strong oxidizing agent is amphoteric in nature. It is the part of lead storage batteries. Compound A is




35. Match List-I with List-II.

List-I Electronic configuration of elements		List-II $\Delta_f H$ in $\text{kJ mol}^{-1}$	
(A)	$1s^2 2s^2$	(i)	801
(B)	$1s^2 2s^2 2p^4$	(ii)	899
(C)	$1s^2 2s^2 2p^3$	(iii)	1314
(D)	$1s^2 2s^2 2p^1$	(iv)	1402



Choose the most appropriate answer from the options given below.

- (a) (A)  $\rightarrow$  (i), (B)  $\rightarrow$  (iii), (C)  $\rightarrow$  (iv), (D)  $\rightarrow$  (ii)  
 (b) (A)  $\rightarrow$  (i), (B)  $\rightarrow$  (iv), (C)  $\rightarrow$  (iii), (D)  $\rightarrow$  (ii)  
 (c) (A)  $\rightarrow$  (ii), (B)  $\rightarrow$  (iii), (C)  $\rightarrow$  (iv), (D)  $\rightarrow$  (i)  
 (d) (A)  $\rightarrow$  (iv), (B)  $\rightarrow$  (i), (C)  $\rightarrow$  (ii), (D)  $\rightarrow$  (iii)

36. For the given reaction



What is 'A'?

- (a)  $\text{CH}=\text{CH}-\text{NH}_2$  (b)   
 (c)  (d)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$

37. Match List-I with List-II.

List-I (Ore)		List-II (Element present)	
(A)	Kernite	(i)	Tin
(B)	Cassiterite	(ii)	Boron
(C)	Calamine	(iii)	Fluorine
(D)	Cryolite	(iv)	Zinc

Choose the most appropriate answer from the options given below.

- (a) (A)  $\rightarrow$  (i), (B)  $\rightarrow$  (iii), (C)  $\rightarrow$  (iv), (D)  $\rightarrow$  (ii)  
 (b) (A)  $\rightarrow$  (iii), (B)  $\rightarrow$  (i), (C)  $\rightarrow$  (ii), (D)  $\rightarrow$  (iv)  
 (c) (A)  $\rightarrow$  (ii), (B)  $\rightarrow$  (i), (C)  $\rightarrow$  (iv), (D)  $\rightarrow$  (iii)  
 (d) (A)  $\rightarrow$  (ii), (B)  $\rightarrow$  (iv), (C)  $\rightarrow$  (i), (D)  $\rightarrow$  (iii)

38. Which of the following vitamin is helpful in delaying the blood clotting?

- (a) Vitamin K (b) Vitamin B  
 (c) Vitamin C (d) Vitamin E

39. Given below are two statements :

**Statement I :** *o*-Nitrophenol is steam volatile due to intramolecular hydrogen bonding.

**Statement II :** *o*-Nitrophenol has high melting due to hydrogen bonding.

In the light of the above statements, choose the most appropriate answer from the options given below.

- (a) Statement I is true but statement II is false.  
 (b) Statement I is false but statement II is true.  
 (c) Both statement I and statement II are false.  
 (d) Both statement I and statement II are true.

40. On treating a compound with warm dil.  $\text{H}_2\text{SO}_4$ , gas X is evolved which turns  $\text{K}_2\text{Cr}_2\text{O}_7$  paper acidified with dil.  $\text{H}_2\text{SO}_4$  to a green compound Y. X and Y respectively are

- (a)  $\text{X} = \text{SO}_2$ ,  $\text{Y} = \text{Cr}_2(\text{SO}_4)_3$   
 (b)  $\text{X} = \text{SO}_2$ ,  $\text{Y} = \text{Cr}_2\text{O}_3$   
 (c)  $\text{X} = \text{SO}_3$ ,  $\text{Y} = \text{Cr}_2(\text{SO}_4)_3$   
 (d)  $\text{X} = \text{SO}_3$ ,  $\text{Y} = \text{Cr}_2\text{O}_3$

41. Given below are two statements : One is labelled as Assertion A and the other is labelled as Reason R.

**Assertion A :** Dipole-dipole interactions are the only non-covalent interactions, resulting in hydrogen bond formation.

**Reason R :** Fluorine is the most electronegative element and hydrogen bonds in HF are symmetrical.

In the light of the above statement choose the most appropriate answer from the options given below.

- (a) A is false but R is true.  
 (b) Both A and R are true but R is not the correct explanation of A.  
 (c) Both A and R are true and R is the correct explanation of A.  
 (d) A is true but R is false.

42. Given below are two statements :

**Statement I :** A mixture of chloroform and aniline can be separated by simple distillation.

**Statement II :** When separating aniline from a mixture of aniline and water by steam distillation, aniline boils below its boiling point.

In the light of the above statements, choose the most appropriate answer from the options given below.

- (a) Statement I is true but statement II is false.  
 (b) Both statement I and statement II are true.  
 (c) Both statement I and statement II are false.  
 (d) Statement I is false but statement II is true.

43. The presence of ozone in troposphere

- (a) protects us from greenhouse effect  
 (b) generates photochemical smog  
 (c) protects us from the X-ray radiation  
 (d) protects us from the UV radiation.


44. Statements about heavy water are given below.

- A. Heavy water is used in exchange reactions for the study of reaction mechanisms.  
 B. Heavy water is prepared by exhaustive electrolysis of water.  
 C. Heavy water has higher boiling point than ordinary water.  
 D. Viscosity of  $\text{H}_2\text{O}$  is greater than  $\text{D}_2\text{O}$ .

Choose the most appropriate answer from the options given below.



- (a) A and C only (b) A and D only  
(c) A and B only (d) A, B and C only
45. Which one of the following lanthanoids does not form  $MO_2$ ? [M is lanthanoid metal.]

(a) Dy (b) Nd (c) Pr (d) Yb



B reacts with hydroxylamine but does not give Tollen's test. Identify A and B.






- (a) 1, 1-Dichlorobutane and butanal  
(b) 2, 2-Dichlorobutane and butanal  
(c) 2, 2-Dichlorobutane and butan-2-one  
(d) 1, 1-Dichlorobutane and 2-butanone
47. An amine on reaction with benzenesulphonyl chloride produces a compound insoluble in alkaline solution. This amine can be prepared by ammonolysis of ethyl chloride. The correct structure of amine is




48. Which of the following is a false statement?

- (a) Kjeldahl's method is used for the estimation of nitrogen in an organic compound.  
(b) Phosphoric acid produced on oxidation of phosphorus present in an organic compound is precipitated as  $Mg_2P_2O_7$  by adding magnesia mixture.  
(c) Carius tube is used in the estimation of sulphur in an organic compound.  
(d) Carius method is used for the estimation of nitrogen in an organic compound.





49. Identify the major products A and B respectively in the following reactions of phenol.



50. For the given reaction



What is 'A'?



## SECTION-B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

51. For a chemical reaction,  $A + B \rightleftharpoons C + D$  ( $\Delta_r H^\circ = 80 \text{ kJ mol}^{-1}$ ) the entropy change  $\Delta_r S^\circ$  depends on the temperature  $T$  (in K) as  $\Delta_r S^\circ = 2T \text{ (J K}^{-1} \text{ mol}^{-1})$ . Minimum temperature at which it will become spontaneous is \_\_\_\_\_. K. (Integer answer)

52. 224 mL  $SO_{2(g)}$  at 298 K and 1 atm is passed through 100 mL of 0.1 M NaOH solution. The non-volatile solute produced is dissolved in 36 g of water. The lowering of vapour pressure of solution (assuming the solution is dilute) [ $P^\circ_{(H_2O)} = 24 \text{ mm of Hg}$ ] is  $x \times 10^{-2} \text{ mm of Hg}$ , the value of  $x$  is \_\_\_\_\_. (Integer answer)

53. An exothermic reaction  $X \rightarrow Y$  has an activation energy  $30 \text{ kJ mol}^{-1}$ . If energy change  $\Delta E$  during the reaction is  $-20 \text{ kJ}$ , then the activation energy for the reverse reaction in kJ is \_\_\_\_\_. (Integer answer)

54. A homogeneous ideal gaseous reaction  $AB_{2(g)} \rightleftharpoons A_{(g)} + 2B_{(g)}$  is carried out in a 25 litre flask at  $27^\circ\text{C}$ . The initial amount of  $AB_2$  was 1 mole and the equilibrium pressure was 1.9 atm. The value of  $K_p$  is  $x \times 10^{-2}$ . The value of  $x$  is \_\_\_\_\_. (Integer answer)  
[ $R = 0.08206 \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$ ]



55. The number of significant figures in  $50000.020 \times 10^{-3}$  is \_\_\_\_\_.
56. A certain gas obeys  $P(V_m - b) = RT$ . The value of  $\left(\frac{\partial Z}{\partial P}\right)_T$  is  $\frac{xb}{RT}$ . The value of  $x$  is \_\_\_\_\_. (Integer answer) ( $Z$  : compressibility factor)
57. Consider the following reaction  
 $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$ ,  $E^\circ = 1.51 \text{ V}$ .  
 The quantity of electricity required in Faraday to reduce five moles of  $\text{MnO}_4^-$  is \_\_\_\_\_. (Integer answer)
58. 3.12 g of oxygen is adsorbed on 1.2 g of platinum metal. The volume of oxygen adsorbed per gram of the adsorbent at 1 atm and 300 K in L is \_\_\_\_\_.  
 $[R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}]$

59. Dichromate ion is treated with base, the oxidation number of Cr in the product formed is \_\_\_\_\_.
60. Number of bridging CO ligands in  $[\text{Mn}_2(\text{CO})_{10}]$  is \_\_\_\_\_.

## MATHEMATICS

### SECTION-A (MULTIPLE CHOICE QUESTIONS)

61. Let  $A$  be a symmetric matrix of order 2 with integer entries. If the sum of the diagonal elements of  $A^2$  is 1, then the possible number of such matrices is  
 (a) 6 (b) 4 (c) 12 (d) 1
62. The value of  $\int_{-\pi/2}^{\pi/2} \frac{\cos^2 x}{1+3^x} dx$  is  
 (a)  $2\pi$  (b)  $\pi/2$  (c)  $\pi/4$  (d)  $4\pi$
63. The value of  $\sum_{n=1}^{100} \int_{n-1}^n e^{x-[x]} dx$ , where  $[x]$  is the greatest integer  $\leq x$ , is  
 (a)  $100(e-1)$  (b)  $100(1-e)$   
 (c)  $100(1+e)$  (d)  $100e$
64. The value of  $\lim_{h \rightarrow 0} 2 \left\{ \frac{\sqrt{3} \sin\left(\frac{\pi}{6} + h\right) - \cos\left(\frac{\pi}{6} + h\right)}{\sqrt{3}h(\sqrt{3} \cosh - \sinh)} \right\}$  is  
 (a)  $3/4$  (b)  $4/3$  (c)  $2/3$  (d)  $2/\sqrt{3}$
65. The rate of growth of bacteria in a culture is proportional to the number of bacteria present and the bacteria count is 1000 at initial time  $t = 0$ . The number of bacteria is increased by 20% in 2 hours. If the population of bacteria is 2000 after  $\frac{k}{\log_e\left(\frac{6}{5}\right)}$  hours, then  $\left(\frac{k}{\log_e 2}\right)^2$  is  
 (a) 16 (b) 8 (c) 2 (d) 4

66. If  $\frac{\sin^{-1} x}{a} = \frac{\cos^{-1} x}{b} = \frac{\tan^{-1} y}{c}$ ;  $0 < x < 1$ , then the value of  $\cos\left(\frac{\pi c}{a+b}\right)$  is  
 (a)  $\frac{1-y^2}{y\sqrt{y}}$  (b)  $\frac{1-y^2}{1+y^2}$  (c)  $1-y^2$  (d)  $\frac{1-y^2}{2y}$
67. If  $(1, 5, 35)$ ,  $(7, 5, 5)$ ,  $(1, \lambda, 7)$  and  $(2\lambda, 1, 2)$  are coplanar, then the sum of all possible values of  $\lambda$  is  
 (a)  $-\frac{39}{5}$  (b)  $-\frac{44}{5}$  (c)  $\frac{44}{5}$  (d)  $\frac{39}{5}$
68. A fair coin is tossed a fixed number of times. If the probability of getting 7 heads is equal to probability of getting 9 heads, then the probability of getting 2 heads is  
 (a)  $\frac{15}{2^8}$  (b)  $\frac{15}{2^{13}}$  (c)  $\frac{15}{2^{14}}$  (d)  $\frac{15}{2^{12}}$
69. If  $\vec{a}$  and  $\vec{b}$  are perpendicular, then  $\vec{a} \times (\vec{a} \times (\vec{a} \times (\vec{a} \times \vec{b})))$  is equal to  
 (a)  $\vec{a} \times \vec{b}$  (b)  $|\vec{a}|^4 \vec{b}$  (c)  $\frac{1}{2} |\vec{a}|^4 \vec{b}$  (d)  $\vec{0}$
70. Let  $f$  be any function defined on  $R$  and let it satisfy the condition:  $|f(x) - f(y)| \leq |x - y|^2$ ,  $\forall (x, y) \in R$   
 If  $f(0) = 1$ , then  
 (a)  $f(x) < 0$ ,  $\forall x \in R$  (b)  $f(x) = 0$ ,  $\forall x \in R$   
 (c)  $f(x)$  can take any value in  $R$   
 (d)  $f(x) > 0$ ,  $\forall x \in R$
71. In an increasing geometric series, the sum of the second and the sixth term is  $\frac{25}{2}$  and the product of the third and fifth term is 25. Then, the sum of 4<sup>th</sup>, 6<sup>th</sup> and 8<sup>th</sup> terms is equal to  
 (a) 30 (b) 26 (c) 32 (d) 35
72. The sum of the infinite series  
 $1 + \frac{2}{3} + \frac{7}{3^2} + \frac{12}{3^3} + \frac{17}{3^4} + \frac{22}{3^5} + \dots$  is equal to  
 (a)  $\frac{15}{4}$  (b)  $\frac{13}{4}$  (c)  $\frac{11}{4}$  (d)  $\frac{9}{4}$
73. The intersection of three lines  $x - y = 0$ ,  $x + 2y = 3$  and  $2x + y = 6$  is a  
 (a) None of these (b) Equilateral triangle  
 (c) Isosceles triangle (d) Right angled triangle
74. The maximum value of the term independent of  $t$  in the expansion of  $\left(tx^{1/5} + \frac{(1-x)^{1/10}}{t}\right)^{10}$ , where  $x \in (0, 1)$  is  
 (a)  $\frac{2 \cdot 10!}{3(5!)^2}$  (b)  $\frac{10!}{\sqrt{3}(5!)^2}$   
 (c)  $\frac{2 \cdot 10!}{3\sqrt{3}(5!)^2}$  (d)  $\frac{10!}{3(5!)^2}$

75. Consider the three planes

$$P_1 : 3x + 15y + 21z = 9, P_2 : x - 3y - z = 5 \text{ and}$$

$$P_3 : 2x + 10y + 14z = 5$$

Then, which one of the following is true?

- (a)  $P_2$  and  $P_3$  are parallel.  
 (b)  $P_1, P_2$  and  $P_3$  all are parallel.  
 (c)  $P_1$  and  $P_3$  are parallel.  
 (d)  $P_1$  and  $P_2$  are parallel.

76. Let  $R = \{(P, Q) \mid P \text{ and } Q \text{ are at the same distance from the origin}\}$  be a relation, then the equivalence class of  $(1, -1)$  is the set

- (a)  $S = \{(x, y) \mid x^2 + y^2 = 1\}$   
 (b)  $S = \{(x, y) \mid x^2 + y^2 = 2\}$   
 (c)  $S = \{(x, y) \mid x^2 + y^2 = 4\}$   
 (d)  $S = \{(x, y) \mid x^2 + y^2 = \sqrt{2}\}$

77. The maximum slope of the curve  $y = \frac{1}{2}x^4 - 5x^3 + 18x^2 - 19x$  occurs at the point

- (a) (2, 9) (b)  $\left(3, \frac{21}{2}\right)$  (c) (0, 0) (d) (2, 2)


78. The number of seven digit integers with sum of the digits equal to 10 and formed by using the digits 1, 2 and 3 only is

- (a) 82 (b) 42 (c) 35 (d) 77

79. The value of  $\begin{vmatrix} (a+1)(a+2) & a+2 & 1 \\ (a+2)(a+3) & a+3 & 1 \\ (a+3)(a+4) & a+4 & 1 \end{vmatrix}$  is

- (a)  $(a+2)(a+3)(a+4)$  (b) 0  
 (c)  $(a+1)(a+2)(a+3)$  (d) -2

80. In the circle given below, let  $OA = 1$  unit,  $OB = 13$  unit and  $PQ \perp OB$ . Then, the area of the triangle  $PQB$  (in square units) is



- (a)  $26\sqrt{3}$  (b)  $24\sqrt{2}$  (c)  $24\sqrt{3}$  (d)  $26\sqrt{2}$

## SECTION-B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

81. The difference between degree and order of a differential equation that represents the family of curves given by

$$y^2 = a \left( x + \frac{\sqrt{a}}{2} \right), a > 0 \text{ is } \underline{\hspace{2cm}}.$$

82. Let  $m, n \in N$  and  $\gcd(2, n) = 1$ . If

$$30 \binom{30}{0} + 29 \binom{30}{1} + \dots + 2 \binom{30}{28} + 1 \binom{30}{29} = n \cdot 2^m, \text{ then}$$

$$n + m \text{ is equal to } \underline{\hspace{2cm}}. \quad \left( \text{Here, } \binom{n}{k} = {}^nC_k \right)$$

83. If  $y = y(x)$  is the solution of the equation

$$e^{\sin y} \cos y \frac{dy}{dx} + e^{\sin y} \cos x = \cos x, y(0) = 0, \text{ then}$$

$$1 + y \left( \frac{\pi}{6} \right) + \frac{\sqrt{3}}{2} y \left( \frac{\pi}{3} \right) + \frac{1}{\sqrt{2}} y \left( \frac{\pi}{4} \right) \text{ is equal to } \underline{\hspace{2cm}}.$$

84. Let  $(\lambda, 2, 1)$  be a point on the plane which passes through the point  $(4, -2, 2)$ . If the plane is perpendicular to the line joining the points  $(-2, -21, 29)$  and  $(-1, -16, 23)$ ,

$$\text{then } \left( \frac{\lambda}{11} \right)^2 - \frac{4\lambda}{11} - 4 \text{ is equal to } \underline{\hspace{2cm}}.$$

85. The number of solutions of the equation

$$\log_4(x-1) = \log_2(x-3) \text{ is } \underline{\hspace{2cm}}.$$

86. The number of integral values of  $k$  for which the equation  $3\sin x + 4\cos x = k + 1$  has a solution,  $k \in R$  is

$$\underline{\hspace{2cm}}.$$

87. If  $\sqrt{3}(\cos^2 x) = (\sqrt{3} - 1)\cos x + 1$ , the number of solutions of the given equation when  $x \in \left[0, \frac{\pi}{2}\right]$  is

88. The sum of  $162^{\text{th}}$  power of the roots of the equation  $x^3 - 2x^2 + 2x - 1 = 0$  is

$$\underline{\hspace{2cm}}.$$

89. The area bounded by the lines  $y = ||x - 1| - 2|$  is

$$\underline{\hspace{2cm}}.$$

90. The value of the integral  $\int_0^{\pi} |\sin 2x| dx$  is

$$\underline{\hspace{2cm}}.$$

## HINTS & EXPLANATIONS

1. (d) : Moment of inertia of solid sphere about an axis

passing through centre  $= \frac{2}{5} ma^2$


$$I_1 = \frac{2}{5} ma^2 = I_2$$

By use of parallel axis theorem

$$I_3 = \frac{2}{5} ma^2 + mb^2 = I_4$$

Moment of inertia about one side

$$I = I_1 + I_2 + I_3 + I_4 = 2I_1 + 2I_3$$



$$\Rightarrow I = 2 \times \frac{2}{5} ma^2 + 2 \times \frac{2}{5} ma^2 + 2mb^2 \Rightarrow I = \frac{8ma^2}{5} + 2mb^2$$

2. (a) : For third line of Lyman series

$$\frac{1}{\lambda_1} = R \left[ \frac{1}{1^2} - \frac{1}{4^2} \right] = \frac{15}{16} \quad \dots(i)$$

For first line of Paschen series

$$\frac{1}{\lambda_2} = R \left[ \frac{1}{3^2} - \frac{1}{4^2} \right] = \frac{7}{144} R \quad \dots(ii)$$

From equation (i) and (ii), we get

$$\frac{\lambda_1}{\lambda_2} = \frac{7 \times 16}{144 \times 15} = \frac{7}{135}$$

3. (d): The resolving power of electron microscope is more than optical microscope. The de Broglie wavelength of the electrons ( $\approx 0.1 \text{ \AA}$ ) emitted from the electron gun is much less than the wavelength of visible light (380nm to 700 nm).

4. (b): Centripetal force,  $F = \frac{mv^2}{R}$  ... (i)

According to question,  $F = \frac{K}{R^3}$  ... (ii)

Where  $K$  is a proportional constant.

From equation (i) and (ii),

$$\Rightarrow \frac{mv^2}{R} = \frac{K}{R^3} \Rightarrow v = \sqrt{\frac{K}{m}} \cdot \frac{1}{R}$$

$$\therefore T = \frac{2\pi R}{v} = \frac{2\pi R}{\sqrt{\frac{K}{m}}} R \sqrt{m} = 2\pi \sqrt{\frac{m}{K}} R^2 \Rightarrow T \propto R^2$$

5. (b): Energy gap,  $E = 1.9 \text{ eV}$

$$\text{As } E = \frac{hc}{\lambda} \Rightarrow \lambda = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{1.9 \times 1.6 \times 10^{-19}}$$

$$\Rightarrow \lambda \approx 654 \text{ nm}$$


Wavelength of red colour range from 620 – 750 nm.

6. (d): As the two sheets are in series, so rate of flow of heat is same from both sheets.

$$\therefore H = \frac{\theta_1 - \theta}{R_1} = \frac{\theta - \theta_2}{R_2}$$

$$\Rightarrow \theta_1 R_2 - \theta R_2 = \theta R_1 - \theta_2 R_1$$

$$\Rightarrow \frac{\theta_1 R_2 + \theta_2 R_1}{R_1 + R_2} = \theta$$



7. (b): Bulk Modulus,  $K = \frac{-PV}{\Delta V}$


As mass is constant, so

$$\rho V = \text{constant} \Rightarrow V = \frac{1}{\rho}$$

$$\frac{\Delta V}{V} = -\frac{\Delta \rho}{\rho} \quad \text{put in (i)}$$

$$K = \frac{P\rho}{\Delta \rho} \Rightarrow \Delta \rho = \frac{P\rho}{K}$$

8. (d):  $a = \frac{\sqrt{3}}{2}L$ , Let us assume an elementary portion of length  $dx$  at distance  $x$  from  $O$ .



Electric field at point  $P$  due to this elementary portion is

$$dE = \frac{K\lambda d\theta}{r^2} = \frac{K\lambda dx}{r^2}$$

$$dE = \frac{K\lambda \cdot a \sec^2 \theta d\theta}{a^2 \sec^2 \theta}$$

$$dE_x = dE \cos \theta d\theta = \frac{K\lambda}{a} \cos \theta d\theta$$

$$dE_y = dE \sin \theta \quad (\text{They balance each other})$$

$$\therefore E = \int dE_x = \frac{K\lambda}{a} \int_{-\theta_1}^{\theta_2} \cos \theta d\theta = \frac{K\lambda}{a} [\sin \theta]_{-\theta_1}^{\theta_2}$$

$$\Rightarrow E = \frac{K\lambda}{a} [\sin \theta_2 + \sin \theta_1] = \frac{K\lambda}{a} \left[ \frac{L}{2r} + \frac{L}{2r} \right]$$

$$\Rightarrow E = \frac{K\lambda \cdot 2L}{a \cdot 2r} = \frac{K\lambda L}{a \cdot r} = \frac{KQ}{a \cdot \sqrt{\frac{L^2}{4} + \frac{3L^2}{4}}}$$

$$E = \frac{KQ \times 2}{\sqrt{3}L^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{2Q}{\sqrt{3}L^2} \Rightarrow E = \frac{Q}{2\sqrt{3}\pi\epsilon_0 L^2}$$

9. (c): According to Kepler's second law, areal velocity of planet is constant.

10. (\*): When connected in parallel,  $C_{eq} = C_1 + C_2$

When connected in series,  $C_{eq} = \frac{C_1 C_2}{C_1 + C_2}$

$$\text{Here, } C_1 + C_2 = \frac{15}{4} \frac{C_1 C_2}{C_1 + C_2}$$

$$\Rightarrow \frac{(C_1 + C_2)^2}{C_1 C_2} = \frac{15}{4} \Rightarrow \frac{C_1^2 + C_2^2 + 2C_1 C_2}{C_1 C_2} = \frac{15}{4}$$

$$\Rightarrow \frac{C_1}{C_2} + \frac{C_2}{C_1} = \frac{15}{4} - 2$$

$$\text{Let } \frac{C_2}{C_1} = x \Rightarrow \frac{1}{x} + x = \frac{7}{4}$$

$$\Rightarrow 4x^2 - 7x + 4 = 0$$

$$D = b^2 - 4ac = 49 - 64 < 0$$

No solution exist.

$$11. (d): W = \alpha^2 \beta e^{-\frac{\beta x^2}{KT}}$$

$$\text{As, } \frac{\beta x^2}{KT} = \text{dimensionless} \Rightarrow [\beta] = \frac{[KT]}{[x^2]} = \frac{ML^2 T^{-2}}{L^2}$$

Dimensions of  $(\alpha^2 \beta)$  = Dimension of work

$$\Rightarrow \alpha^2 \times [MT^{-2}] = [ML^2 T^{-2}]$$


$$\Rightarrow \alpha = [L]$$

12. (c): Gravitational field due to ring at a distance  $x$  is

$$E = G \cdot \frac{mx}{(R^2 + x^2)^{3/2}}$$

Force on sphere,

$$F = ME$$





$$F = \frac{MGmx}{(R^2 + x^2)^{3/2}}$$

$$\Rightarrow F = \frac{GMm \cdot \sqrt{8R}}{(R^2 + 8R^2)^{3/2}}$$

$$\Rightarrow F = \frac{GMm \cdot \sqrt{8R}}{9^{3/2} \cdot R^3} = \frac{GMm\sqrt{8}}{3^3 \cdot R^2}$$

$$\Rightarrow F = \frac{\sqrt{8}}{27} \cdot \frac{GMm}{R^2}$$

13. (b) : Here current,  $i = i_1 \sin \omega t + i_2 \cos \omega t$

$$\Rightarrow i = \sqrt{i_1^2 + i_2^2} \left[ \frac{i_2 \cos \omega t}{\sqrt{i_1^2 + i_2^2}} + \frac{i_1 \sin \omega t}{\sqrt{i_1^2 + i_2^2}} \right]$$

$$\Rightarrow i = \sqrt{i_1^2 + i_2^2} [\sin \alpha \cos \omega t + \cos \alpha \sin \omega t]$$

$$\Rightarrow i = \sqrt{i_1^2 + i_2^2} \sin(\omega t + \alpha) = i_0 \sin(\omega t + \alpha)$$

$$\Rightarrow i_{rms} = \frac{i_0}{\sqrt{2}} = \sqrt{\frac{i_1^2 + i_2^2}{2}}$$

14. (c) : Let the number of water drops =  $N$   
Volume remains constant

$$\therefore \frac{4}{3} \pi r^3 \cdot N = \frac{4}{3} \pi R^3 \Rightarrow R^3 = Nr^3$$

Energy released =  $T \times 4\pi r^2 N - T \times 4\pi R^2$

$$U = \left( T \times 4\pi r^2 \cdot \frac{R^3}{r^3} - T \cdot 4\pi R^2 \right) \frac{1}{J}$$

$$\therefore \frac{U}{\text{Volume}} = \frac{1}{J} \frac{T \times 4\pi R^2 \left[ \frac{1}{r} R - 1 \right]}{\frac{4}{3} \pi R^3}$$

$$\Rightarrow \frac{U}{\text{Volume}} = \frac{3T}{J} \left[ \frac{1}{r} - \frac{1}{R} \right]$$

15. (b) :  $g' = g \left( 1 - \frac{d}{R} \right) = \frac{R-d}{R}$

$$g' = -\frac{y}{R} g$$

$$\text{So, } g' = \frac{-g}{R} y = -\omega^2 y$$

$$\omega^2 = \frac{g}{R}$$

$$T = \frac{2\pi}{\omega} = \frac{2\pi\sqrt{R}}{\sqrt{g}}$$

$$T = 2\pi\sqrt{\frac{R}{g}}$$

16. (a) : When two springs are connected in series,


$$\text{equivalent spring constant, } K_S = \frac{K_1 K_2}{K_1 + K_2}$$

$$\Rightarrow K_S = \frac{K_1^2}{2K_1} = \frac{K_1}{2}$$

$\therefore$  Time period,

$$T = 2\pi\sqrt{\frac{m}{K_1}} \text{ and } T' = 2\pi\sqrt{\frac{m}{K_S}} = 2\pi\sqrt{\frac{2m}{K_1}}$$

$$\Rightarrow T' = \sqrt{2}T$$

17. (c) :  $m < M$   


(After collision)

Using conservation of linear momentum

$$Mu = MV + mv$$

...(i)

By, definition of coefficient of restitution

$$e = \frac{V - v}{0 - u} = 1 \Rightarrow V = v - u \text{ putting in equation (i)}$$

$$\Rightarrow Mu = M(v - u) + mv$$

$$v = \frac{2Mu}{M + m} \text{ if } M \gg m; v = 2u$$

18. (a) : Here,  $f = 40 \text{ cm}$ ,  $h = 100 \text{ cm}$ ,  $h' = 25 \text{ cm}$

As orientation is same, the image is virtual and smaller, mirror has to be convex and image formed on opposite side.

19. (b) : Given,  $d = 2 \text{ mm}$ ;  $D = 1 \text{ m}$ ;  $\lambda = 500 \text{ nm}$

$$\text{Fringe width, } \beta = \frac{D\lambda}{d} = \frac{1 \times 500 \times 10^{-9}}{2 \times 10^{-3}} = 2.5 \times 10^{-4} \text{ m}$$

$$\Rightarrow \beta = 0.25 \text{ mm}$$

20. (a) : It is a balanced Wheatstone bridge, therefore resistance between E and D can be removed.

So,

$$\frac{1}{R_{eq}} = \frac{1}{2R} + \frac{1}{2R} = \frac{1}{R}$$

$$\Rightarrow R_{AB} = R_{eq} = R$$

21. (300) :  $q = 20 \text{ C}$ ;  $V = 15 \text{ V}$

Work done =  $qV = 20 \times 15 = 300 \text{ J}$

22. (137) :  $P = 1000 \text{ W}$ ;  $r = 2 \text{ m}$ ;  $\eta = 1.25\%$

$$\text{Intensity, } I = \frac{1}{2} \epsilon_0 C E_0^2 \quad \dots(i)$$

$$I = \frac{P}{\text{area}} \times \eta \quad \dots(ii)$$

From equation (i) and (ii),

$$\frac{1000}{4\pi(2)^2} \times \frac{1.25}{100} = \frac{1}{2} \times 8.854 \times 10^{-12} \times 3 \times 10^8 E_0^2$$



$$E_0 = 13.7 \text{ V/m}$$

As question,  $x \times 10^{-1} \text{ V/m} = 13.7 \text{ V/m}$

$$x = 137$$

23. (1215) :

$$\mu = 0.135 \text{ g/cm} = \frac{0.135 \times 10^{-1}}{10^{-2}} \text{ kg/m} = 0.0135 \text{ kg/m}$$



$$y = -0.21 \sin(x + 30t) \quad \dots(i)$$

On comparing equation (i) with  $y = A \sin(kx + \omega t)$ , we get  
 $\omega = 30, k = 1$

$$v = \frac{\omega}{k} = 30$$

$$v = \sqrt{\frac{T}{\mu}} \Rightarrow 30 = \sqrt{\frac{T}{0.0135}}$$

$$T = 12.15 \text{ N}$$

As per question,  $T = x \times 10^{-2} \text{ N} = 12.15 \text{ N}$

$$x = 1215$$

**24. (20) :**  $D_2$  is in reverse bias, so no current flows through this branch.

The total resistance,

$$R = 50 + 130 + 120 = 300 \Omega$$

$$i = \frac{V}{R} = \frac{6}{300} = 0.02 \text{ A} = 20 \text{ mA}$$

**25. (33.3) :**  $A_{\max} = 16, A_{\min} = 8$

$$A_m = \frac{A_{\max} - A_{\min}}{2} = \frac{16 - 8}{2} = 4$$

$$A_c = \frac{A_{\max} + A_{\min}}{2} = \frac{16 + 8}{2} = 12$$

$$\mu = \frac{A_m}{A_c} = \frac{4}{12}$$

As per question,  $x \times 10^{-2} = \frac{4}{12}$

$$x = \frac{400}{12} = 33.3$$

**26. (492) :** When lift is at rest

$$N = mg$$

$$= 60 \times 10 = 600 \text{ N}$$

When lift moves with

downward acceleration

$$N' = m(g - a) = 60(10 - 18)$$

$$= 492 \text{ N}$$

**27. (500) :**  $a = 10 \text{ m/s}^2; m = 2 \text{ kg}$

$$v = u + at = 0 + 10 \times 10 = 100 \text{ m/s}$$

Also,  $\frac{1}{2}mv^2 = F \cdot S$

$$\frac{1}{2} \times 2 \times (100)^2 = 20 S$$

$$S = 500 \text{ m}$$



**28. (3.33) :** Coefficient of friction,  $\mu = \frac{1}{3\sqrt{3}}$

$$m = \sqrt{3} \text{ kg}$$

$$N = mg + F \sin 60^\circ \quad \dots(i)$$

$$\mu N = F \cos 60^\circ \text{ (for at rest)} \dots(ii)$$

From equation (i) and (ii), we get



$$\frac{1}{3\sqrt{3}} \left( mg + F \frac{\sqrt{3}}{2} \right) = \frac{F}{2}$$

$$\frac{1}{3\sqrt{3}} \left( \sqrt{3} \times 10 + F \frac{\sqrt{3}}{2} \right) = \frac{F}{2}$$

$$\frac{10}{3} + \frac{F}{6} = \frac{F}{2}$$

$$F = 10 \text{ N}$$

As per question,  $10 = 3x$

$$\Rightarrow x = 3.33$$

**29. (400) :** Quality factor,  $Q = \frac{\omega L}{R}$  ;

$$\frac{Q'}{Q} = \frac{L'R}{R'L} = \frac{2L}{L} \cdot \frac{R \times 2}{R} = 4$$

$$Q' = 4 \times 100 = 400$$

**30. (25.5) :**  $V_1 = 4.5 \text{ L}; V_2 = 5.5 \text{ L}$

$$n_1 = 3 \quad n_2 = 4$$

$$P_1 = 2 \text{ atm} \quad P_2 = 3 \text{ atm}$$

According to the conservation of energy

$$K_1 + K_2 = K$$

$$\frac{3}{2}n_1RT_1 + \frac{3}{2}n_2RT_2 = \frac{3}{2}(n_1 + n_2)RT$$

$$\therefore PV = nRT$$

$$\frac{3}{2}P_1V_1 + \frac{3}{2}P_2V_2 = \frac{3}{2}P(V_1 + V_2)$$

$$P = 2.55 \text{ atm}$$

As per question  $x \times 10^{-1} = 2.55$


$$x = 25.5$$

**31. (c) :** Angular nodes =  $l$

Radial nodes =  $n - l - 1$


Orbital	Angular nodes	Radial node
4f	$l = 3$	$4 - 3 - 1 = 0$
4d	$l = 2$	$4 - 2 - 1 = 1$
5d	$l = 2$	$5 - 2 - 1 = 2$
3p	$l = 1$	$3 - 1 - 1 = 1$

**32. (c) :** Neoprene is a polymer of chloroprene.




**33. (d) :**  $2\text{NH}_3 + \text{H}_2\text{O} + \text{CO}_2 \longrightarrow (\text{NH}_4)_2\text{CO}_3$

(A)



(B)



(C)


**34. (a) :** Due to inert pair effect,  $\text{Pb}^{4+}$  is strong oxidising agent. As given that compound A is used in lead storage batteries hence it is  $\text{PbO}_2$ .

35. (c) : As we move from left to right in a period, ionisation enthalpy increases. But half-filled and fully-filled orbitals have higher ionisation enthalpy than the next upcoming element.

Hence, the correct order of ionisation enthalpy is

	B	<	Be	<	O	<	N
E.C. :	$1s^2 2s^2 2p^1$		$1s^2 2s^2$		$1s^2 2s^2 2p^4$		$1s^2 2s^2 2p^3$
$\Delta_i H$ :	801		899		1314		1402

36. (c) :  $\text{CH}_3 - \text{CH} = \text{CH} - \text{Br} \xrightarrow[\text{-HBr}]{\text{NaNH}_2} \text{CH}_3 - \text{C} \equiv \text{CH}$




37. (c) : Kernite ( $\text{Na}_2\text{B}_4\text{O}_7 \cdot 4\text{H}_2\text{O}$ ) (A-ii)  
 Cassiterite ( $\text{SnO}_2$ ) (B-i)  
 Calamine ( $\text{ZnCO}_3$ ) (C-iv)  
 Cryolite ( $\text{Na}_3\text{AlF}_6$ ) (D-iii)

38. (d) : Vitamin K is a group of vitamins that the body needs for blood clotting, helping wounds to heal. On the other hand, vitamin E delays the blood clotting.

39. (a) : *o*-Nitrophenol has intramolecular hydrogen bonding thus, it is steam volatile. Molecular association is not possible in *o*-nitrophenol hence it has low melting point.

40. (a) :  $\text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 + 3\text{SO}_2 \longrightarrow$   
 (X)  $\text{K}_2\text{SO}_4 + \text{Cr}_2(\text{SO}_4)_3 + \text{H}_2\text{O}$   
 (Y) (Green coloured)

41. (a) : Ion-dipole interactions can also result in hydrogen bonding. Fluorine is most electronegative and forms symmetrical H-bonds.




42. (b) : Chloroform (b.p. 334 K) and aniline (b.p. 457 K) have a big difference in their boiling point. Hence, the mixture can be separated by simple distillation. Statement-II is also true.

43. (b) : Ozone in stratosphere helps us to protect from UV rays. But in troposphere it generates photochemical smog.

44. (d) : Viscosity of  $\text{D}_2\text{O}$  is greater than  $\text{H}_2\text{O}$ .



45. (d) : Pr, Nd, Tb and Dy exhibit +4 state, but only in oxides  $\text{MO}_2$ .  $\text{Yb}^{2+}$ , which has  $f^{14}$  configuration does not form oxide  $\text{MO}_2$ .

46. (c) : As B reacts with hydroxylamine, so it is a carbonyl compound, but as it does not give Tollen's test hence it is a ketone not an aldehyde. Thus, the reactions are summarised as



47. (a) : As given, the product formed with benzene sulphonyl chloride is insoluble in alkaline solution hence the amine is secondary amine. Amine is prepared by ammonolysis of ethyl chloride thus it should contain  $\text{CH}_3\text{CH}_2\text{NH}-$  group. So the amine could be  $\text{CH}_3\text{CH}_2\text{CH}_2\text{NHCH}_2\text{CH}_3$ .

48. (d) : Carius method is used in estimation of sulphur or halogen in organic compound.



51. (200) :  $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$

Given,  $\Delta S^\circ = 2T$

$\Delta G^\circ = \Delta H^\circ - T(2T)$

Minimum temperature can be calculated when  $\Delta G^\circ = 0$

$$2T^2 = \Delta H^\circ = 80000$$

$$T^2 = 40000 \text{ or } T = 200 \text{ K}$$

52. (18) :  $\text{SO}_2 + 2\text{NaOH} \longrightarrow \text{Na}_2\text{SO}_3 + \text{H}_2\text{O}$

$$\text{Number of moles of SO}_2 = \frac{0.224 \times 1}{0.0821 \times 298} = 0.0092 \text{ mol}$$

Number of moles of NaOH = 0.01 mol

NaOH is the limiting reagent

Hence, number of moles of  $\text{Na}_2\text{SO}_3 = 0.005 \text{ mol}$

$$\text{Mole fraction of Na}_2\text{SO}_3 = \frac{0.005}{2.005} = 0.0025$$

$$i = 3$$

$$\text{Lowering in vapour pressure} = P_{\text{H}_2\text{O}}^0 \times \frac{n_{\text{Na}_2\text{SO}_3}}{n_{\text{H}_2\text{O}}} \times i$$

$$= 24 \times \frac{0.005}{2.005} \times 3 = \frac{24 \times 5 \times 3}{2005} = 0.18$$


$$\text{Lowering in vapour pressure} = 18 \times 10^{-2} \text{ mm Hg}$$

$$\therefore x = 18$$

$$53. (50) : E_{af} = 30 \text{ kJ/mol}$$

$$\Delta E = -20 \text{ kJ/mol}$$


For exothermic reaction



$$\Delta E = E_{af} - E_{ab}$$

$$-20 = 30 - E_{ab}$$

$$\Rightarrow E_{ab} = 50 \text{ kJ mol}^{-1}$$



Initial	1	0	0
At eqm.	$1 - \alpha$	$\alpha$	$2\alpha$

$$\text{Moles at equilibrium} = 1 + 2\alpha$$

$$PV = nRT$$

$$1.9 \times 25 = n \times 0.082 \times 300$$

$$n = 1.93 \text{ or } \alpha = 0.465$$

$$K_p = \frac{P_A P_B^2}{P_{AB_2}} = \frac{\left(\frac{0.465}{1.93} \times 1.9\right) \left(\frac{0.93}{1.93} \times 1.9\right)^2}{\left(\frac{0.535}{1.93} \times 1.9\right)}$$

$$K_p = 0.727 \text{ or } 72.7 \times 10^{-2}$$

55. (8) : It has eight significant figures.


$$56. (1) : P(V_m - b) = RT$$

$$PV_m - Pb = RT$$

$$\frac{PV_m}{RT} = 1 + \frac{Pb}{RT} \Rightarrow Z = 1 + \frac{Pb}{RT} ; \left(\frac{dZ}{dP}\right)_T = 0 + \frac{b}{RT} \times 1$$

$$\text{Hence, } x = 1.$$

57. (25) : For the balanced equation,



We can say that 1 mole of  $\text{MnO}_4^-$  require 5 F electricity.


Hence, 5 moles of  $\text{MnO}_4^-$  will require 25 F of electricity.

$$58. (2) : \text{Moles of oxygen} = \frac{3.12}{32} = 0.0975$$


$$\text{Volume of oxygen} = \frac{nRT}{P} = \frac{0.0975 \times 0.0821 \times 300}{1} = 2.40 \text{ L}$$

1.2 g of Pt metal adsorbs = 2.40 L oxygen

$$1 \text{ g of Pt metal will adsorb} = \frac{2.4}{1.2} \text{ L oxygen} \\ = 2 \text{ L oxygen}$$



Oxidation state of Cr in  $\text{K}_2\text{CrO}_4$  or  $\text{CrO}_4^{2-}$  is 6.



No bridging CO group.

$$61. (b) : \text{Let } A = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$$

$$\therefore A^2 = \begin{bmatrix} a^2 + b^2 & b(a+c) \\ b(a+c) & b^2 + c^2 \end{bmatrix}$$

$$\text{Since, } a^2 + b^2 + b^2 + c^2 = 1$$

$$\Rightarrow a^2 + 2b^2 + c^2 = 1, \text{ where } a, b, c \in I$$

$$\therefore b = 0 \text{ and } a^2 + c^2 = 1$$

$$\text{At } a = 0, c = \pm 1$$

$$\text{At } c = 0, a = \pm 1$$

So, number of possible matrices is 4.

$$62. (c) : \text{Let } I = \int_{-\pi/2}^{\pi/2} \frac{\cos^2 x}{1+3^x} dx \quad \dots (i)$$

$$= \int_{-\pi/2}^{\pi/2} \frac{\cos^2(-x)}{1+3^{(-x)}} dx \quad \left[ \because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

$$\therefore I = \int_{-\pi/2}^{\pi/2} \frac{\cos^2 x}{1+3^{-x}} dx \quad \dots (ii)$$

Adding (i) and (ii), we get

$$2I = \int_{-\pi/2}^{\pi/2} \frac{\cos^2 x}{1+3^x} dx + \int_{-\pi/2}^{\pi/2} \frac{\cos^2 x}{1+3^{-x}} dx$$

$$= 2 \int_0^{\pi/2} \left( \frac{\cos^2 x}{1+3^x} + \frac{3^x \cdot \cos^2 x}{1+3^x} \right) dx$$

$$\Rightarrow I = \int_0^{\pi/2} \cos^2 x dx = \frac{1}{2} \int_0^{\pi/2} (1 + \cos 2x) dx$$

$$= \frac{1}{2} \left[ x + \frac{\sin 2x}{2} \right]_0^{\pi/2} = \frac{1}{2} \left[ \frac{\pi}{2} \right] = \frac{\pi}{4}$$

$$63. (a) : \text{We have, } \sum_{n=1}^{100} \int_{n-1}^n e^{x-[x]} dx$$

$$= \sum_{n=1}^{100} \int_{n-1}^n e^{\{x\}} dx \quad [\because x - [x] = \{x\}]$$

$$= \sum_{n=1}^{100} \int_0^1 e^x dx \quad [\because \text{period of } \{x\} \text{ is } 1]$$

$$= 100 \left[ e^x \right]_0^1 = 100(e-1)$$

64. (b) : We have,  $\lim_{h \rightarrow 0} 2 \left\{ \frac{\frac{\sqrt{3}}{2} \sin\left(\frac{\pi}{6} + h\right) - \frac{1}{2} \cos\left(\frac{\pi}{6} + h\right)}{\sqrt{3}h \left( \frac{\sqrt{3}}{2} \cosh - \frac{1}{2} \sinh \right)} \right\}$

$$= \lim_{h \rightarrow 0} 2 \left\{ \frac{\cos \frac{\pi}{6} \sin\left(\frac{\pi}{6} + h\right) - \sin \frac{\pi}{6} \cos\left(\frac{\pi}{6} + h\right)}{\sqrt{3}h \left( \cos \frac{\pi}{6} \cosh - \sin \frac{\pi}{6} \sinh \right)} \right\}$$

$$= \lim_{h \rightarrow 0} 2 \left\{ \frac{\sin\left(\frac{\pi}{6} + h - \frac{\pi}{6}\right)}{\sqrt{3}h \cos\left(\frac{\pi}{6} + h\right)} \right\}$$

$$= \lim_{h \rightarrow 0} \frac{2}{\sqrt{3}} \left( \frac{\sinh}{h} \right) \frac{1}{\cos\left(\frac{\pi}{6} + h\right)}$$

$$= \frac{2}{\sqrt{3} \cos \frac{\pi}{6}} = \frac{2}{\sqrt{3} \cdot \frac{\sqrt{3}}{2}} = \frac{4}{3}$$

65. (d) : Given,  $\frac{dx}{dt} = \lambda x$

(where  $\lambda$  is a constant of proportionality)

Now, when  $t = 0$ ,  $x = 1000$  and

when  $t = 2$ ,  $x = 1000 + 20\%$  of  $1000 = 1000 + 200 = 1200$

$$\Rightarrow \int_{1000}^{1200} \frac{dx}{x} = \lambda \int_0^2 dt \Rightarrow [\log_e x]_{1000}^{1200} = \lambda [t]_0^2$$

$$\Rightarrow \log_e \left( \frac{1200}{1000} \right) = 2\lambda \Rightarrow \lambda = \frac{1}{2} \log_e \left( \frac{6}{5} \right)$$

Also,  $\int_{1000}^{2000} \frac{dx}{x} = \frac{1}{2} \log_e \left( \frac{6}{5} \right) \int_0^k dt$

$$\Rightarrow \log_e \left( \frac{2000}{1000} \right) = \frac{1}{2} \log_e \left( \frac{6}{5} \right) \cdot \frac{k}{\log_e \left( \frac{6}{5} \right)}$$

$$\Rightarrow \log_e 2 = \frac{k}{2} \Rightarrow \frac{k}{\log_e 2} = 2 \Rightarrow \left( \frac{k}{\log_e 2} \right)^2 = 4$$

66. (b) : Let  $\frac{\sin^{-1} x}{a} = \frac{\cos^{-1} x}{b} = \frac{\tan^{-1} y}{c} = \lambda$  (say)

$$\Rightarrow \sin^{-1} x = a\lambda, \cos^{-1} x = b\lambda, \tan^{-1} y = c\lambda$$

$$\therefore (a+b)\lambda = \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

$$\Rightarrow \frac{\pi}{a+b} = 2\lambda$$

$$\text{Now, } \cos \left( \frac{\pi c}{a+b} \right) = \cos(2\lambda c) = \cos(2 \tan^{-1} y)$$

$$\text{Let } \tan^{-1} y = \theta \Rightarrow y = \tan \theta$$

$$\therefore \cos \left( \frac{\pi c}{a+b} \right) = \cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1 - y^2}{1 + y^2}$$

67. (c) : Let  $A(1, 5, 35)$ ,  $B(7, 5, 5)$ ,  $C(1, \lambda, 7)$  and  $D(2\lambda, 1, 2)$  be the given points.

For the points to be coplanar,

$$\begin{vmatrix} 6 & 0 & -30 \\ 0 & \lambda - 5 & -28 \\ 2\lambda - 1 & -4 & -33 \end{vmatrix} = 0$$

$$\Rightarrow 6[-33(\lambda - 5) - 112] - 30[-(\lambda - 5)(2\lambda - 1)] = 0$$

$$\Rightarrow -33\lambda + 53 + 5(2\lambda^2 - 11\lambda + 5) = 0$$

$$\Rightarrow 10\lambda^2 - 88\lambda + 78 = 0 \Rightarrow 5\lambda^2 - 44\lambda + 39 = 0$$

$$\therefore \text{Required sum} = 44/5$$

68. (b) : Let the coin be tossed  $n$  times.

$$\text{Also, } P(H) = P(T) = \frac{1}{2}$$

Since,  $P(7 \text{ heads}) = P(9 \text{ heads})$

$$\Rightarrow {}^nC_7 \left( \frac{1}{2} \right)^7 \left( \frac{1}{2} \right)^{n-7} = {}^nC_9 \left( \frac{1}{2} \right)^9 \left( \frac{1}{2} \right)^{n-9}$$

$$\Rightarrow {}^nC_7 = {}^nC_9 \Rightarrow n = 7 + 9 = 16$$

$$\therefore P(2 \text{ heads}) = {}^{16}C_2 \left( \frac{1}{2} \right)^2 \left( \frac{1}{2} \right)^{14} = \frac{16 \times 15}{2 \times 1} \cdot \frac{1}{2^{16}} = \frac{15}{2^{13}}$$

69. (b) : Since,  $\vec{a}$  and  $\vec{b}$  are perpendicular.

$$\therefore \vec{a} \cdot \vec{b} = 0 \quad \dots (i)$$

Now,  $\vec{a} \times (\vec{a} \times (\vec{a} \times (\vec{a} \times \vec{b})))$

$$= \vec{a} \times (\vec{a} \times ((\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}))$$

$$= \vec{a} \times (\vec{a} \times (-|\vec{a}|^2 \vec{b}))$$

$$= -|\vec{a}|^2 (\vec{a} \times (\vec{a} \times \vec{b})) = -|\vec{a}|^2 (-|\vec{a}|^2 \vec{b})$$

$$= |\vec{a}|^4 \vec{b}$$

[using (i)]

70. (d) : We have,  $|f(x) - f(y)| \leq |(x - y)^2|$

$$\Rightarrow \left| \frac{f(x) - f(y)}{x - y} \right| \leq |x - y|$$

$$\Rightarrow \lim_{h \rightarrow 0} \left| \frac{f(y+h) - f(y)}{h} \right| \leq 0$$

$$\Rightarrow |f'(y)| \leq 0 \Rightarrow f'(y) = 0$$

$$\Rightarrow f(y) = \text{constant}$$

$$\text{As } f(0) = 1, \therefore f(y) = 1 \Rightarrow f(x) = 1$$

$$\therefore f(x) > 0 \forall x \in R$$

71. (d) : Given,  $T_2 + T_6 = \frac{25}{2} \Rightarrow ar + ar^5 = \frac{25}{2} \quad \dots (i)$

$$\text{Also, } T_3 \cdot T_5 = 25$$



$$\Rightarrow ar^2 ar^4 = 25 \Rightarrow a^2 r^6 = 25 \Rightarrow ar^3 = 5$$

Dividing (i) by (ii), we get

$$\frac{r+r^5}{r^3} = \frac{5}{2} \Rightarrow 2+2r^4 = 5r^2$$

$$\Rightarrow 2r^4 - 5r^2 + 2 = 0 \Rightarrow r^2 = 2, \frac{1}{2}$$

$$\therefore r^2 = 2 \quad \left[ \text{Rejecting } r^2 = \frac{1}{2} \right]$$

$$\text{Now, } T_4 + T_6 + T_8 = ar^3 + ar^5 + ar^7 \\ = ar^3(1 + r^2 + r^4) = 5(1 + 2 + 4) = 35$$

$$72. \text{ (b) : Let } S = 1 + \frac{2}{3} + \frac{7}{3^2} + \frac{12}{3^3} + \dots \infty$$

$$\text{and } \frac{1}{3}S = \frac{1}{3} + \frac{2}{3^2} + \frac{7}{3^3} + \frac{12}{3^4} + \dots \infty$$

Subtracting (ii) from (i), we get

$$\frac{2}{3}S = 1 + \frac{1}{3} + \frac{5}{3^2} + \frac{5}{3^3} + \dots \infty$$

$$\Rightarrow \frac{2}{3}S = \frac{4}{3} + \left( \frac{5}{3^2} + \frac{5}{3^3} + \dots \infty \right)$$


$$\Rightarrow \frac{2}{3}S = \frac{4}{3} + \left( \frac{5}{3^2} \right) \left( \frac{1}{1 - \frac{1}{3}} \right) = \frac{4}{3} + \frac{5}{6} = \frac{13}{6}$$

$$\therefore S = \frac{13}{6} \times \frac{3}{2} = \frac{13}{4}$$

$$73. \text{ (c) : } AB = \sqrt{(2-1)^2 + (2-1)^2} = \sqrt{2} \text{ units,}$$

$$BC = \sqrt{(3-1)^2 + (0-1)^2} = \sqrt{5} \text{ units and}$$

$$CA = \sqrt{(3-2)^2 + (0-2)^2} = \sqrt{5} \text{ units}$$



As  $BC = AC$

$\therefore \triangle ABC$  is isosceles.

74. (c) : General term,

$$T_{r+1} = {}^{10}C_r (tx^{1/5})^{10-r} \left( \frac{(1-x)^{1/10}}{t} \right)^r$$

Since, it is independent of  $t$ .

$$\therefore 10 - r - r = 0 \Rightarrow r = 5$$

$$\therefore T_6 = {}^{10}C_5 x(1-x)^{1/2}$$

$$\text{Now, } \frac{d}{dx}(T_6) = {}^{10}C_5 \left[ (1-x)^{1/2} - \frac{x}{2\sqrt{1-x}} \right] = 0$$

$$\dots \text{ (ii) } \Rightarrow 2(1-x) - x = 0 \Rightarrow 2 - 3x = 0 \Rightarrow x = 2/3$$

$$\therefore \text{Maximum value of } T_6 = {}^{10}C_5 \left( \frac{2}{3} \right) \left( \frac{1}{3} \right)^{1/2} = \frac{2 \times 10!}{3\sqrt{3} (5!)^2}$$

75. (c) : We have,  $P_1 : x + 5y + 7z = 3$

$$P_2 : x - 3y - z = 5$$

$$P_3 : x + 5y + 7z = 5/2$$

So,  $P_1$  and  $P_3$  are parallel as direction ratios of normal are same.

76. (b) : Equivalence class of  $(1, -1)$  is a circle with centre  $(0, 0)$  and radius  $\sqrt{2}$ .

$$\text{i.e. } x^2 + y^2 = 2$$

$$\therefore \text{Required set is } S = \{(x, y) \mid x^2 + y^2 = 2\}$$

$\dots \text{ (i)}$

$\dots \text{ (ii)}$

$$77. \text{ (d) : We have, } y = \frac{1}{2}x^4 - 5x^3 + 18x^2 - 19x$$

$$\text{Slope} = \frac{dy}{dx} = 2x^3 - 15x^2 + 36x - 19$$

$$\Rightarrow \frac{d^2y}{dx^2} = 6x^2 - 30x + 36$$

$$\text{For the slope to be maximum or minimum, } \frac{d^2y}{dx^2} = 0$$

$$\Rightarrow 6(x^2 - 5x + 6) = 0 \Rightarrow x = 2, 3$$

$$\text{Now, } \frac{d^3y}{dx^3} = 6(2x - 5)$$

$$\text{At } x = 2, \frac{d^3y}{dx^3} = -6 < 0$$

$\therefore$  Slope is maximum at  $x = 2$ .

$$\text{At } x = 2, y = \frac{1}{2}(2)^4 - 5(2)^3 + 18(2)^2 - 19(2)$$

$$= 8 - 40 + 72 - 38 = 2$$

$\therefore$  Required point is  $(2, 2)$ .

78. (d) : Case-I : Digits can be 1, 1, 1, 1, 2, 3

$$\therefore \text{Required number of ways} = \frac{7!}{5!} = 7 \times 6 = 42$$

Case-II : Digits can be 1, 1, 1, 1, 2, 2, 2

$$\therefore \text{Required number of ways} = \frac{7!}{4!3!} = \frac{7 \times 6 \times 5}{3 \times 2} = 35$$

Hence, total number of ways =  $42 + 35 = 77$

$$79. \text{ (d) : Let } \Delta = \begin{vmatrix} (a+1)(a+2) & (a+2) & 1 \\ (a+2)(a+3) & (a+3) & 1 \\ (a+3)(a+4) & (a+4) & 1 \end{vmatrix}$$


Applying  $R_2 \rightarrow R_2 - R_1$  and  $R_3 \rightarrow R_3 - R_1$

$$\Delta = \begin{vmatrix} (a+1)(a+2) & (a+2) & 1 \\ 2a+4 & 1 & 0 \\ 4a+10 & 2 & 0 \end{vmatrix}$$

$$= 2(2a+4) - (4a+10) = 4a+8-4a-10 = -2$$

80. (c) : Given,  $OB = 13$  units and  $OA = 1$  unit

$\therefore AB = 12$  units



Let C be the centre of circle.

Thus,  $PC = \frac{13}{2} = 6.5$  units and  $AC = 5.5$  units

Now, in  $\triangle PAC$ ,  $PA^2 = PC^2 - CA^2$

$$\Rightarrow PA = \sqrt{(6.5)^2 - (5.5)^2} = \sqrt{42.25 - 30.25} = \sqrt{12} \text{ units}$$

$$\therefore PQ = 2\sqrt{12} \text{ units}$$

$$\begin{aligned} \therefore \text{Area of } \triangle PQB &= \frac{1}{2} \times AB \times PQ \\ &= \frac{1}{2} \times 12 \times 2\sqrt{12} = 24\sqrt{3} \text{ sq. units} \end{aligned}$$

$$81. (2) : \text{We have, } y^2 = a \left( x + \frac{\sqrt{a}}{2} \right) \quad \dots (i)$$

$$\Rightarrow 2yy_1 = a$$

From (i) and (ii), we get

$$y^2 = 2yy_1 \left( x + \frac{\sqrt{2yy_1}}{2} \right)$$

$$\Rightarrow y^2 = 2xyy_1 + \frac{(2yy_1)^{3/2}}{2} \Rightarrow y^2 - 2xyy_1 = 2^{1/2}(yy_1)^{3/2}$$

Squaring both sides, we get

$$(y^2 - 2xyy_1)^2 = 2y^3 y_1^3 \Rightarrow (y - 2xy_1)^2 = 2yy_1^3$$

$\therefore$  Order = 1 and degree = 3

Required difference =  $3 - 1 = 2$

82. (45) : We have,

$$30 \binom{30}{0} + 29 \binom{30}{1} + \dots + 2 \binom{30}{28} + 1 \binom{30}{29}$$

$$= 30 \binom{30}{30} + 29 \binom{30}{29} + \dots + 2 \binom{30}{2} + 1 \binom{30}{1}$$

$$= \sum_{r=1}^{30} r \cdot \binom{30}{r} = \sum_{r=1}^{30} r \cdot \frac{30}{r} \binom{29}{r-1}$$

$$= 30 \left[ \binom{29}{0} + \binom{29}{1} + \binom{29}{2} + \dots + \binom{29}{29} \right]$$

$$= 30 \cdot 2^{29} = 15 \times 2^{30} = n \cdot 2^m \text{ (given)}$$

$$\therefore n + m = 15 + 30 = 45$$

83. (1) : Given equation is

$$e^{\sin y} \cos y \frac{dy}{dx} + e^{\sin y} \cos x = \cos x \quad \dots (i)$$

$$\text{Put } e^{\sin y} = t \Rightarrow e^{\sin y} \cos y \frac{dy}{dx} = \frac{dt}{dx}$$

$\therefore$  (i) becomes,  $\frac{dt}{dx} + t \cos x = \cos x$ , which is a L.D.E.

$$\text{Here, I.F.} = e^{\int \cos x dx} = e^{\sin x}$$

$\therefore$  Solution is given by

$$t \cdot e^{\sin x} = \int \cos x \cdot e^{\sin x} dx \Rightarrow e^{\sin y} e^{\sin x} = e^{\sin x} + c$$

$$\text{At } x = 0, y = 0 \Rightarrow c = 0$$

$$\therefore e^{\sin y} = 1 \Rightarrow \sin y = 0 \Rightarrow y = 0, \forall x$$

$$\therefore 1 + y \left( \frac{\pi}{6} \right) + \frac{\sqrt{3}}{2} y \left( \frac{\pi}{3} \right) + \frac{1}{\sqrt{2}} y \left( \frac{\pi}{4} \right) = 1 + 0 + 0 + 0 = 1$$

$$84. (8) : \overline{AB} = \hat{i} + 5\hat{j} - 6\hat{k}$$

$$\overline{PQ} = (4 - \lambda)\hat{i} - 4\hat{j} + \hat{k}$$

Since,  $\overline{AB} \perp \overline{PQ}$ .

$$\therefore \overline{AB} \cdot \overline{PQ} = 0$$

$$(\hat{i} + 5\hat{j} - 6\hat{k}) \cdot ((4 - \lambda)\hat{i} - 4\hat{j} + \hat{k}) = 0$$

$$\Rightarrow 4 - \lambda - 20 - 6 = 0 \Rightarrow \lambda = -22$$

$$\begin{aligned} \therefore \left( \frac{\lambda}{11} \right)^2 - \frac{4\lambda}{11} - 4 &= \left( \frac{-22}{11} \right)^2 - 4 \left( \frac{-22}{11} \right) - 4 \\ &= 4 + 8 - 4 = 8 \end{aligned}$$

$$85. (1) : \text{We have, } \log_4 (x - 1) = \log_2 (x - 3)$$

$$\Rightarrow \frac{1}{2} \log_2 (x - 1) = \log_2 (x - 3)$$

$$\Rightarrow \log_2 (x - 1)^{1/2} = \log_2 (x - 3) \Rightarrow (x - 1)^{1/2} = x - 3$$

Squaring both sides, we get

$$(x - 1) = x^2 + 9 - 6x$$

$$\Rightarrow x^2 - 7x + 10 = 0 \Rightarrow x = 2, 5$$

$\therefore x = 5$  as  $x = 2$  does not satisfy the domain of given equation

$\therefore$  Number of solution is 1.

$$86. (11) : \text{We have, } 3\sin x + 4\cos x = k + 1$$

$$\Rightarrow (k + 1) \in \left[ -\sqrt{3^2 + 4^2}, \sqrt{3^2 + 4^2} \right]$$

$$\Rightarrow (k + 1) \in [-5, 5] \Rightarrow k \in [-6, 4]$$

$\therefore$  Number of integral values of  $k$  is 11.

$$87. (1) : \text{We have, } \sqrt{3} \cos^2 x - (\sqrt{3} - 1) \cos x - 1 = 0$$

$$\Rightarrow \sqrt{3} \cos^2 x - \sqrt{3} \cos x + \cos x - 1 = 0$$

$$\Rightarrow \sqrt{3} \cos x (\cos x - 1) + 1(\cos x - 1) = 0$$


$$\Rightarrow (\cos x - 1)(\sqrt{3} \cos x + 1) = 0$$

$$\Rightarrow \cos x = 1 \text{ or } \cos x = -\frac{1}{\sqrt{3}} \left( \text{not possible as } x \in \left[ 0, \frac{\pi}{2} \right] \right)$$

$$\therefore \cos x = 1 \Rightarrow x = 0$$

$\therefore$  Number of solutions is 1.

$$88. (3) : \text{We have, } x^3 - 2x^2 + 2x - 1 = 0 \quad \dots (i)$$



Since,  $x = 1$  satisfies the given equation.

$\therefore x - 1$  is a factor of (i)

$$\therefore (x^3 - 2x^2 + 2x - 1) = (x - 1)(x^2 - x + 1)$$

$$\therefore x = 1, \frac{1 \pm i\sqrt{3}}{2}$$

i.e.,  $x = 1, -\omega^2, -\omega$

$\therefore$  Sum of  $162^{\text{th}}$  power of roots

$$= (1)^{162} + (-\omega^2)^{162} + (-\omega)^{162}$$


$$= 1 + \omega^{324} + \omega^{162} = 1 + (\omega^3)^{108} + (\omega^3)^{54} = 1 + 1 + 1 = 3.$$

89. (4) : We have,  $y = ||x - 1| - 2|$

$$= \begin{cases} x - 3 & , \quad x \geq 3 \\ 3 - x & , \quad 1 \leq x \leq 3 \\ x + 1 & , \quad -1 \leq x \leq 1 \\ -1 - x & , \quad x \leq -1 \end{cases}$$

$$\therefore \text{ Required area } = \int_{-1}^1 (x + 1) dx + \int_1^3 (3 - x) dx$$

$$= \left[ \frac{x^2}{2} + x \right]_{-1}^1 + \left[ 3x - \frac{x^2}{2} \right]_1^3 = 0 + 2 + 6 - 4 = 4 \text{ sq. units}$$



90. (2) : Let  $I = \int_0^{\pi} |\sin 2x| dx$

$$= 2 \int_0^{\pi/2} \sin 2x dx$$

$$\left[ \because \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx \text{ if } f(2a - x) = f(x) \right]$$

$$= 2 \left[ \frac{-\cos 2x}{2} \right]_0^{\pi/2} = 2$$

