

A framework to drive a proof or refutation of the Collatz Conjecture

The problem

Given the function defined for the natural numbers:

$$f(n) = (3n+1)/2 \text{ if } n \text{ is odd } n/2 \text{ if } n \text{ is even (} \\ [/f(n) =]$$

And its kth application $f^k(n)$. By definition $f^0(n) = n$

Notice that for odd branches we are using the shortcut version provided that $3n+1$, being n odd is guaranteed to be even and the next step is always dividing by two.

Let's define that a number n is *reducible* if it exist a finite k so that $f^k(n)=1$

By brute force we know that all the first checked natural numbers are reducible. As for today this has been proved until 2^{68} (David Bařina <https://github.com/hellpig/collatz>).

The Collatz conjecture states that: all natural numbers are reducible.

Toolbox

Powers of three

We can relate a power of 3 with lesser powers of 3

Given that for any base b and natural power n :

$$b^n - 1 = (b-1) * \sum_{0 \leq i < n} (b^i) \\ b^n = 1 + (b-1) * \sum_{0 \leq i < n} (b^i)$$

For $b = 3$:

$$3^n = 1 + 2 * \sum_{0 \leq i < n} (3^i) \\ 3^n = 1 + 2 * (3^{n-1} + 3^{n-2} \dots + 3 + 1)$$

This is also equivalent to those formulas:

$$(3^n - 1)/2 = (3^{n-1} + 3^{n-2} \dots + 3 + 1) \\ (3^n - 1)/2 = \sum_{0 \leq i < n} (3^i)$$

$$\sum_{0 \leq k < n} (3^k) = (3^n - 1) / 2$$

$$(3^n + 1)/2 = 1 + (3^{n-1} + 3^{n-2} \dots + 3 + 1) \\ (3^n + 1)/2 = 1 + \sum_{0 \leq i < n} (3^i)$$

We can also relate powers of 3 in terms of powers of 2 by using the binomial theorem

```

3^n = (2 + 1)^n
3^n = sum[0<=i<=n] ( 2^i * n! / (n-i)! / i! )
3^n = sum[0<=i<=n] ( 2^i * bincoef(i,n) )

```

Boolean with integer arithmetics

Enable us to integrate boolean conditions into natural numbers expressions.
Useful to eliminate formula branching.

Let's define a boolean integer as $B \in [0,1]$

Being a,b,c... boolean integers, we can represent boolean operations with integer algebra like this:

- not: $\text{not } a = 1-a$
- and: $a \text{ and } b = (a*b)$
 - Properties:
 - * $a*1 = a$
 - * $a*0 = 0$
 - * $a*a = a$ — because both 1 and 0 multiplied by themselves return themselves
 - * $a(1-a) = a - aa = a - a = 0$
- or: $a \text{ or } b = (a + b - a*b)$
 - Properties:
 - * $a \text{ or } 1 = a + 1 - 1*a = 1$
 - * $a \text{ or } 0 = a + 0 - 0*a = a$
 - * $a \text{ or } a = a + a - a*a = a$
 - * $a \text{ or not } a = a + (1-a) - a * (1-a) = a + 1 - a - a + a*a = 1$

Other derived operators

- xor: $a \text{ xor } b = a(1-b) + b(1-a) = a-ab+b-ab = a+b-2ab = aa +bb -2ab = (a-b)^2$
 - $a \text{ xor } 0 = (a-0)^2 = a$
 - $a \text{ xor } 1 = (a-1)^2 = aa +1 - 2a = 1-a = \text{not } a$
 - $a \text{ xor } a = (a-a)^2 = 0$
 - $a \text{ xor not } a = (a - (1-a))^2 = (2a-1)^2 = 4a + 1 - 4*a = 1$
- eq: $a \text{ eq } b = ab + (1-a)(1-b) = 2ab -a -b +1 = 1 - (a-b)^2 = \text{not } (a \text{ xor } b)$

Be careful that $a+b-ab$ is an OR while $a+b-2ab$ is an XOR.

Those operations ensure a closure among booleans integers. Meaning that while the operands are 0 or 1, the result will be also 0 or 1.

So, how to use those boolean expressions inside an algebraic formula? We can make multiplication factors and addition terms optional.

Being x and y natural numbers and 'a' a boolean condition represented as integer,

- conditional addition of a term x: $a*x + y$
- conditional multiplication by a factor x: $x^a * y$

Oddity of algebraic expressions

Oddity is a function that returns a boolean integer, 1 if the number is odd. Also useful to eliminate branching.

Being a and b integer expressions:

```
odd(1) = 1
odd(2*a) = 0
odd(a+b) = odd(a) xor odd(b) = odd(a) + odd(b) - 2*odd(a)*odd(b) = (odd(a)-odd(b))^2
odd(a*b) = odd(a) and odd(b) = odd(a)*odd(b)
```

Pair terms can be ignored for oddity:

```
odd(2*a + b) = odd(2*a) xor odd(b) = 0 xor odd(b) = odd(b)
odd((1+2*a)**b) = 1
odd((2*a)**b) = (b!=0)    --- TODO: which opp gives this?
```

Unbranched formula

Unbranched formula for the generator function:

```
f(n) = 1/2 * (n*3^odd(n) + odd(n) )
```

Thus we can define the series recursively by:

```
f0(n) = n
fk+1(n) = 1/2 * ( fk(n) * 3^(odd(fk(n))) + odd(fk(n)) )
```

Another formulation is:

```
fk+1(n) = 1/2 * ( fk(n) * (1 + 2*odd(fk(n))) + odd(fk(n)) )
```

Which can be convenient to extract factors of two.

Lets define Ok as odd(fk(n)). Then we can express both formulations as

```
fk+1(n) = (3^Ok * fk(n) + Ok) / 2
```

Or also:

```
fk+1(n) = ((1+2*Ok) * fk(n) + Ok) / 2
```

For compactness, we will omit (n), so fk = fk(n)

```
fk+1 = (3^Ok * fk + Ok) / 2    # Ok exponential form
fk+1 = (fk + 2*Ok*fk + Ok) / 2 # Ok factor form
```

Additive/Subtractive view

```
fk+1 - fk =
= (fk + 2*Ok*fk + Ok) / 2 - fk    # Using Ok factor form
= (-fk + 2*Ok*fk + Ok) / 2        # fk inside 1/2
    Odd: (-fk + 2*fk + 1) / 2 = fk/2 + 1/2
```

Even: $-fk/2$

fk odd : $+fk/2 + 1/2$

fk even: $-fk/2$

Conclusión: Depending on the oddity of the previous result, we are adding or subtracting half of the sequence value, rounding up for odds.

Strategies

Hypothesis: Exists a first natural number A that it is not reducible.

Strategy 1: If exists a first non reducible natural A implies that any $n < A$ is reducible. Because A is not reducible, $f_k(A) \geq A$ for all k . Having $f_k(A) < A$ will contradict the hypothesis. If the hypothesis is true, it could lead to a search algorithm for A .

Suposition: The oddness of the $f_k(n)$ just depends on the lower $k+1$ bits of n .

Strategy 2: Being A a finite number, at a given k all remaining bits are zero. Because the oddity of $f_k(n)$ is inverted depending on the k th bit, maybe we could find a pattern for which appending 0 bits gets higher and higher or cyclic.

Strategy 3: Constructive. Requires demonstrating that k th bit of A controls oddity of $f_k(x)$. Instead of starting with every number and apply the function to reduce it. Start on 1 and invert f so that we can get to that number by the odd and even formula.

Solution structure

Theorem: All $f^k(n)$ can be expressed as $(an+b)/c$ being a and c extrictly positive integers, and b positive integer.

Proof:

$f^0(n) = n$; ($a_0=1$, $b_0=0$, $c_0=1$)

Given that $f^k(n)$ can be expressed as $(ak*n + bk)/ck$,
can $f^{k+1}(n)$ be expressed as $(a'*n+b')/c'$?

if $f^k(n)$ is odd: $f^{k+1}(n) = (3*a*n + 3*b + c)/2*c$
($a'=3*a$, $b'=3*b+c$, $c'=2*c$)

if $f^k(n)$ is even: $f^{k+1}(n) = (a*n + b) /2*c$
($a'=a$, $b'=b$, $c'=2*c$)

qvd

In single branch

$$f^{k+1}(n) = (a + 2^k a, b + 2^k b + 0_k, 2^k c)$$

$$\begin{aligned} a' &= (2^k + 1) * a \\ b' &= b + b * 2^k + c * 0_k \\ c' &= 2^k c \end{aligned}$$

$$\begin{aligned} a_{k+1} &= \prod_{i=0..k} 3^{0_i} = 3^{\sum_{i=0..k} 0_i} \\ b_{k+1} &= b_k * 3^{0_k} + c_k + 0_k = b_k * 3^{0_k} + 2^{k-1} + 0_k \\ c_{k+1} &= 2^k \end{aligned}$$

Empirical observations

Being n_k the k th bit of n binary base representation. And $N_k = N \gg k$, this is the integer division by 2^k .

All developed solutions have the form:

$$f_k(n) = 3^{B_k} N_k + C_k$$

Where B_k and C_k depend only on bits n_{k-1} to n_0 . $0 \leq B_k \leq k$ and $0 \leq C_k < 3^{B_k}$. Indeed $\max(C_k) = 3^{B_k-1}$ and happens when the B_k *higher processed* bits are 1 (Caution this has been observed just for the first 5 levels)

Lets try to demonstrate those observations.

All solutions as $f_k(n) = 3^{B_k} N_k + C_k$

Hypothesis: solutions can be represented as:

$$f_k(n) = N_k * 3^{B_k} + C_k$$

So that:

$$\begin{aligned} 0 &\leq B_k \leq k \\ 0 &\leq C_k < 3^{B_k} \end{aligned}$$

$$\begin{aligned} \text{for } k=0, \quad 0_0 &= n_0 \\ f_0(n) &= N_0; \quad \text{thus } B_0 = 1, \quad C_0 = 0 \end{aligned}$$

$$\begin{aligned} \text{for } k=1 \\ f_1(n) &= (3^{n_0} * N_0 + n_0) / 2 \\ f_1(n) &= (3^{n_0} * (2*N_1 + n_0) + n_0) / 2 && \text{--- Expand } N_0 = 2*N_1 + n_0 \\ f_1(n) &= (3^{n_0} * 2*N_1 + 3^{n_0} * n_0 + n_0) / 2 && \text{-- distribute } 3^{n_0} \\ f_1(n) &= (3^{n_0} * 2*N_1 + n_0 * 3^{n_0} + n_0) / 2 && \text{-- reorder factors} \\ f_1(n) &= (3^{n_0} * 2*N_1 + n_0 * 3^{n_0} + n_0) / 2 && \text{-- } 3^{n_0} = 1 + 2*n_0 \text{ for } n_0 \in (0,1) \\ f_1(n) &= (3^{n_0} * 2*N_1 + n_0 * (1 + 2*n_0) + n_0) / 2 && \text{-- } 3^{n_0} = 1 + 2*n_0 \text{ for } n_0 \in (0,1) \\ f_1(n) &= (3^{n_0} * 2*N_1 + 2*n_0 * 2*n_0) / 2 && \text{-- } 3^{n_0} = 1 + 2*n_0 \text{ for } n_0 \in (0,1) \\ f_1(n) &= (3^{n_0} * 2*N_1 + 4*n_0) / 2 && \text{-- } n_0^2 = n_0 \text{ for } n_0 \in (0,1) \\ f_1(n) &= 3^{n_0} * N_1 + 2*n_0 && \text{-- divide by 2} \end{aligned}$$

$$f_1(n) = N_1 + 2^{n_0}N_1 + 2^{n_0} \quad \text{---} \quad 3^{n_0} = 2^{n_0} + 1$$

$$O_1 = \text{odd}(f_1(n)) = \text{odd}(N_1 + 2^{n_0}N_1 + 2^{n_0}) = \text{odd}(N_1) = n_1$$

So for $k=1$, $B_1=n_0$ and $C_1 = 2^{n_0}$

$$0 \leq B_1 = n_0 \leq k = 1$$

$$0 \leq C_1 = 2^{n_0} < 3^{n_0} = 1+2^{n_0}$$

Now, suposing that:

$$f_k = N_k \cdot 3^{B_k} + C_k$$

$$0 \leq B_k \leq k$$

$$0 \leq C_k < 3^{B_k}$$

Let's demonstrate that:

$$f_{k+1}(n) = (N_{k+1}) \cdot 3^{B_{k+1}} + C_{k+1}$$

$$0 \leq B_k \leq B_{k+1} \leq k+1$$

$$0 \leq C_{k+1} < 3^{B_{k+1}}$$

$$O_k = \text{odd}(f_k)$$

$$= \text{odd}(N_k \cdot 3^{B_k} + C_k)$$

$$= \text{odd}(C_k) + \text{odd}(N_k \cdot 3^{B_k}) - 2 \cdot \text{odd}(C_k) \cdot \text{odd}(N_k \cdot 3^{B_k}) \quad \text{--- exclusive or}$$

$$= \text{odd}(C_k) + \text{odd}(N_k) - 2 \cdot \text{odd}(C_k) \cdot \text{odd}(N_k) \quad \text{---} \quad \text{odd}(N_k \cdot 3^{B_k}) = \text{odd}(N_k)$$

$$= \text{odd}(C_k) + n_k - 2 \cdot n_k \cdot \text{odd}(C_k) \quad \text{---} \quad \text{odd}(N_k) = \text{odd}(2 \cdot N_{k+1} + n_k) = n_k$$

$$f_{k+1} = (3^{O_k} \cdot f_k + O_k) / 2 \quad \text{--- Single branch formula}$$

$$f_{k+1} = (3^{O_k} \cdot (N_k \cdot 3^{B_k} + C_k) + O_k) / 2 \quad \text{---} \quad f_k = N_k \cdot 3^{B_k} + C_k$$

$$f_{k+1} = (3^{O_k} \cdot N_k \cdot 3^{B_k} + 3^{O_k} \cdot C_k + O_k) / 2 \quad \text{--- distribute}$$

$$f_{k+1} = (3^{(B_k+O_k)} \cdot N_k + C_k \cdot 3^{O_k} + O_k) / 2 \quad \text{--- adding exponents}$$

$$f_{k+1} = (3^{(B_k+O_k)} \cdot (2 \cdot N_{k+1} + n_k) + C_k \cdot 3^{O_k} + O_k) / 2 \quad \text{---} \quad N_k = 2 \cdot N_{k+1} + n_k$$

$$f_{k+1} = (3^{(B_k+O_k)} \cdot 2 \cdot N_{k+1} + 3^{(O_k+B_k)} \cdot n_k + C_k \cdot 3^{O_k} + O_k) / 2 \quad \text{--- distribute}$$

$$f_{k+1} = 3^{(B_k+O_k)} \cdot N_{k+1} + (3^{(O_k+B_k)} \cdot n_k + C_k \cdot 3^{O_k} + O_k) / 2 \quad \text{--- divide } N_{k+1} \text{ term}$$

$$B_{k+1} = B_k + O_k$$

$$B_{k+1} = B_k + n_k + \text{Odd}(C_k) - 2 \cdot n_k \cdot \text{Odd}(C_k)$$

$$2 \cdot C_{k+1} =$$

$$= 3^{(B_k+O_k)} \cdot n_k + C_k \cdot 3^{O_k} + O_k \quad \text{--- from } f_{k+1} \text{ expression}$$

$$= 3^{O_k} \cdot 3^{B_k} \cdot n_k + C_k \cdot 3^{O_k} + O_k \quad \text{--- split powers}$$

$$= (2 \cdot O_k + 1) \cdot 3^{B_k} \cdot n_k + C_k \cdot (2 \cdot O_k + 1) + O_k \quad \text{---} \quad 3^{O_k} = 2 \cdot O_k + 1$$

$$= 2 \cdot O_k \cdot 3^{B_k} \cdot n_k + 3^{B_k} \cdot n_k + C_k \cdot 2 \cdot O_k + C_k + O_k \quad \text{--- distribute}$$

$$= 2 \cdot (\text{odd}(C_k) + n_k - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot 3^{B_k} \cdot n_k + 3^{B_k} \cdot n_k + C_k \cdot 2 \cdot (\text{odd}(C_k) + n_k - 2 \cdot n_k \cdot \text{odd}(C_k)) + O_k$$

$$= \quad \text{--- just reorder}$$

$$+ 2 \cdot n_k \cdot 3^{B_k} \cdot (\text{odd}(C_k) + n_k - 2 \cdot n_k \cdot \text{odd}(C_k))$$

$$+ 3^{B_k} \cdot n_k$$

$$+ C_k \cdot 2 \cdot (\text{odd}(C_k) + n_k - 2 \cdot n_k \cdot \text{odd}(C_k))$$

$$+ C_k$$

$$\begin{aligned}
& + (\text{odd}(\text{Ck}) + \text{nk} - 2*\text{nk}*\text{odd}(\text{Ck})) \\
= & \quad \text{--- distribute} \\
& + 2*\text{nk}*3^{\text{Bk}}*(\text{odd}(\text{Ck})) \\
& + 2*\text{nk}*3^{\text{Bk}}*(\text{nk}) \\
& + 2*\text{nk}*3^{\text{Bk}}*(-2*\text{nk}*\text{odd}(\text{Ck})) \\
& + 3^{\text{Bk}}*\text{nk} \\
& + \text{Ck}^2*(\text{odd}(\text{Ck})) \\
& + \text{Ck}^2*(\text{nk}) \\
& + \text{Ck}^2*(-2*\text{nk}*\text{odd}(\text{Ck})) \\
& + \text{Ck} \\
& + \text{odd}(\text{Ck}) \\
& + \text{nk} \\
& - 2*\text{nk}*\text{odd}(\text{Ck}) \\
= & \quad \text{--- distribute} \\
& + 2*\text{nk}*3^{\text{Bk}}*\text{odd}(\text{Ck}) \\
& + 2*\text{nk}*3^{\text{Bk}}*\text{nk} \\
& - 4*\text{nk}*3^{\text{Bk}}*\text{nk}*\text{odd}(\text{Ck}) \\
& + 3^{\text{Bk}}*\text{nk} \\
& + 2*\text{Ck}*\text{odd}(\text{Ck}) \\
& + 2*\text{Ck}*\text{nk} \\
& - 4*\text{Ck}*\text{nk}*\text{odd}(\text{Ck}) \\
& + \text{Ck} \\
& + \text{odd}(\text{Ck}) \\
& + \text{nk} \\
& - 2*\text{nk}*\text{odd}(\text{Ck}) \\
= & \quad \text{--- nk*nk = nk} \\
& + 2*\text{nk}*3^{\text{Bk}}*\text{odd}(\text{Ck}) \\
& - 4*\text{nk}*3^{\text{Bk}}*\text{odd}(\text{Ck}) \\
& + 2*\text{nk}*3^{\text{Bk}} \\
& + 3^{\text{Bk}}*\text{nk} \\
& + 2*\text{Ck}*\text{odd}(\text{Ck}) \\
& + 2*\text{Ck}*\text{nk} \\
& - 4*\text{Ck}*\text{nk}*\text{odd}(\text{Ck}) \\
& + \text{Ck} \\
& + \text{odd}(\text{Ck}) \\
& + \text{nk} \\
& - 2*\text{nk}*\text{odd}(\text{Ck}) \\
= & \quad \text{--- grouping factors} \\
& - 2*\text{nk}*3^{\text{Bk}}*\text{odd}(\text{Ck}) \\
& + 3*\text{nk}*3^{\text{Bk}} \\
& + \text{nk} \\
& + 2*\text{Ck}*\text{nk} \\
& - 4*\text{Ck}*\text{nk}*\text{odd}(\text{Ck}) \\
& + 2*\text{Ck}*\text{odd}(\text{Ck}) \\
& + \text{Ck} \\
& + \text{odd}(\text{Ck})
\end{aligned}$$

$$\begin{aligned}
& - 2nk \cdot \text{odd}(Ck) \\
= & \quad \text{--- grouping factors} \\
& + nk \cdot 3^{Bk} \cdot (3 - 2 \cdot \text{odd}(Ck)) \\
& + nk \\
& + 2Ck \cdot (nk + \text{odd}(ck) - 2nk \cdot \text{odd}(Ck)) \\
& + Ck \\
& + \text{odd}(Ck) \\
& - 2nk \cdot \text{odd}(Ck)
\end{aligned}$$

In order to endup with a natural number 2^{Ck+1} should be even:

$$\begin{aligned}
& \text{odd}(2^{0k} \cdot 3^{Bk} \cdot nk + 3^{Bk} \cdot nk + Ck \cdot 2^{0k} + Ck + 0k) = \\
= & \text{odd}(3^{Bk} \cdot nk + Ck + 0k) \quad \text{--- Removed even terms} \\
= & \text{odd}(3^{Bk} \cdot nk + Ck + \text{odd}(Ck) + nk - 2nk \cdot \text{odd}(Ck)) \quad \text{--- } 0k = \text{odd}(Ck) + nk - 2nk \cdot \text{odd}(Ck) \\
= & \text{odd}(Ck + \text{odd}(Ck) - 2nk \cdot \text{odd}(Ck)) \quad \text{--- } \text{odd}(3^{Bk} \cdot nk + nk) = 0 \\
= & \text{odd}(Ck + \text{odd}(Ck)) \quad \text{--- pair term ignored} \\
= & \text{odd}(0) \quad \text{--- } \text{odd}(Ck + \text{odd}(Ck)) = 0 \\
= & 0 \quad (\text{qvd})
\end{aligned}$$

$$\begin{aligned}
Ck+1 = & (\text{--- grouping factors} + nk \cdot 3^{Bk} \cdot (3 - 2 \cdot \text{odd}(Ck)) + 2Ck \cdot (nk + \text{odd}(ck)) \\
& - 2nk \cdot \text{odd}(Ck) \\
& + Ck + nk + \text{odd}(Ck) - 2nk \cdot \text{odd}(Ck)) / 2
\end{aligned}$$

By cases nk , $\text{odd}(Ck)$.

$$\begin{aligned}
nk=0; \text{odd}(Ck)=0; Bk+1=Bk \cdot (1+2^{0k}) & = Bk \\
2^{Ck+1} & = \\
= & \\
& + nk \cdot 3^{Bk} \cdot (3 - 2 \cdot \text{odd}(Ck)) \\
& + 2Ck \cdot (nk + \text{odd}(ck) - 2nk \cdot \text{odd}(Ck)) \\
& + Ck \\
& + nk \\
& + \text{odd}(Ck) \\
& - 2nk \cdot \text{odd}(Ck) \\
= & Ck
\end{aligned}$$

$$Ck+1 = Ck/2 < Ck < 3^{Bk} = 3^{Bk+1}$$

$$\begin{aligned}
nk=0; \text{odd}(Ck)=1; Bk+1 & = Bk + 1 \\
2^{Ck+1} & = \\
= & \\
& + nk \cdot 3^{Bk} \cdot (3 - 2 \cdot \text{odd}(Ck)) \\
& + 2Ck \cdot (nk + \text{odd}(ck) - 2nk \cdot \text{odd}(Ck)) \\
& + Ck \\
& + nk \\
& + \text{odd}(Ck) \\
& - 2nk \cdot \text{odd}(Ck) \\
= & 3 \cdot Ck + 1
\end{aligned}$$

$$\begin{aligned}
2*C_k+1 &= 3*C_k + 1 <? 2*3*3^{B_k} \\
3*C_k &<? 2*3*3^{B_k} - 1 \\
C_k &<? 2*3^{B_k} - 1/3 \\
C_k &< 3^{B_k} <! 2*3^{B_k} - 1/3
\end{aligned}$$

$$\begin{aligned}
&nk=1; \text{ odd}(C_k)=0 \\
2*C_k+1 &= \\
&= \\
&\quad + nk*3^{B_k} * (3 - 2*\text{odd}(C_k)) \\
&\quad + 2*C_k*(nk + \text{odd}(C_k) - 2*nk*\text{odd}(C_k)) \\
&\quad + C_k \\
&\quad + nk \\
&\quad + \text{odd}(C_k) \\
&\quad - 2*nk*\text{odd}(C_k) \\
&= \\
&\quad + 3^{B_k} * 3 \\
&\quad + 3*C_k \\
&\quad + 1
\end{aligned}$$

$$\begin{aligned}
2*C_k+1 &= 3*3^{B_k} + 3*C_k + 1 <? 2*3*3^{B_k} \\
3*C_k + 1 &<? 3*3^{B_k} \\
C_k &<? 3^{B_k} - 1/3
\end{aligned}$$

$$\begin{aligned}
&nk=1; \text{ odd}(C_k)=1 \\
2*C_k+1 &= \\
&= \\
&\quad + nk*3^{B_k} * (3 - 2*\text{odd}(C_k)) \\
&\quad + 2*C_k*(nk + \text{odd}(C_k) - 2*nk*\text{odd}(C_k)) \\
&\quad + C_k \\
&\quad + nk \\
&\quad + \text{odd}(C_k) \\
&\quad - 2*nk*\text{odd}(C_k) \\
&= \\
&\quad + 3^{B_k} \\
&\quad + C_k \\
2*C_k+1 &= 3^{B_k} + C_k <? 2*3^{B_k} \\
3^{B_k} + C_k &<? 2*3^{B_k} \\
C_k &<! 3^{B_k}
\end{aligned}$$

Thus, it's demonstrated that for every k:

$$fk(n) = N_k*3^{B_k} + C_k$$

Where:

$$0 \leq B_k \leq k$$

$$0 \leq C_k < 3^{B_k}$$

Ck Oddity

From the previous demonstration we got an expression of what feeds Cks from nks

$$\begin{aligned}
 Ck+1 &= \\
 &= (\\
 &\quad + nk \cdot 3^{Bk} * (3 - 2 \cdot \text{odd}(Ck)) \\
 &\quad + 2 \cdot Ck * (nk + \text{odd}(ck) - 2 \cdot nk \cdot \text{odd}(Ck)) \\
 &\quad + Ck \\
 &\quad + nk \\
 &\quad + \text{odd}(Ck) \\
 &\quad - 2 \cdot nk \cdot \text{odd}(Ck) \\
 &\quad) / 2
 \end{aligned}$$

It would be nice to have a generalization of Ck oddity

$$\begin{aligned}
 \text{odd}(Ck+1) &= \\
 &\text{odd}(\\
 &\quad + nk \cdot 3^{Bk} * (3 - 2 \cdot \text{odd}(Ck)) \\
 &\quad + 2 \cdot Ck * (nk + \text{odd}(ck) - 2 \cdot nk \cdot \text{odd}(Ck)) \\
 &\quad + Ck \\
 &\quad + nk \\
 &\quad + \text{odd}(Ck) \\
 &\quad - 2 \cdot nk \cdot \text{odd}(Ck) \\
 &\quad) / 2)
 \end{aligned}$$

Again by cases:

$$\begin{aligned}
 \text{odd}(Ck) &= 0; \quad nk = 0 \\
 \text{odd}(Ck+1) &= \\
 &= \\
 &\quad \text{odd}(\\
 &\quad \quad + nk \cdot 3^{Bk} * (3 - 2 \cdot \text{odd}(Ck)) \\
 &\quad \quad + 2 \cdot Ck * (nk + \text{odd}(ck) - 2 \cdot nk \cdot \text{odd}(Ck)) \\
 &\quad \quad + Ck \\
 &\quad \quad + nk \\
 &\quad \quad + \text{odd}(Ck) \\
 &\quad \quad - 2 \cdot nk \cdot \text{odd}(Ck) \\
 &\quad) / 2) \\
 &= \quad \quad \quad \text{--- } nk = 0 \\
 &\quad \text{odd}(\\
 &\quad \quad + 2 \cdot Ck \cdot \text{odd}(ck) \\
 &\quad \quad + Ck \\
 &\quad \quad + \text{odd}(Ck) \\
 &\quad) / 2) \\
 &= \quad \quad \quad \text{--- } \text{odd}(Ck) = 0 \\
 &\quad \text{odd}(
 \end{aligned}$$

```

      + Ck
    ) /2)
=      --- simplify
      odd(Ck/2)

odd(Ck) = 0; nk = 1
odd(Ck+1) =
=
  odd((
    + nk*3^Bk * (3 - 2*odd(Ck))
    + 2*Ck*(nk + odd(ck) - 2*nk*odd(Ck))
    + Ck
    + nk
    + odd(Ck)
    - 2*nk*odd(Ck)
  ) /2)
=      --- odd(Ck) = 0
  odd((
    + nk*3^Bk * (3)
    + 2*Ck*(nk)
    + Ck
    + nk
  ) /2)
=      --- nk = 1
  odd((
    + 3^Bk * (3)
    + 2*Ck
    + Ck
    + 1
  ) /2)

=      --- even outside half
  odd(
    Ck + Ck/2 +
    (
      + 3^Bk * (3)
      + 1
    ) /2
  )

=      --- Ck being even does not affect overall oddity
  odd(
    Ck/2 +
    (
      + 3^Bk * (3)
      + 1
    )
  )

```

```

    ) /2
  )
=      --- factor of
  odd(
    Ck/2 +
    (
      + 3^(Bk + 1)
      + 1
    ) /2
  )

odd(Ck) = 1; nk = 0
odd(Ck+1) =
=
  odd((
    + nk*3^Bk * (3 - 2*odd(Ck))
    + 2*Ck*(nk + odd(ck) - 2*nk*odd(Ck))
    + Ck
    + nk
    + odd(Ck)
    - 2*nk*odd(Ck)
  ) /2)

odd(Ck) = 1; nk = 1
odd(Ck+1) =
=
  odd((
    + nk*3^Bk * (3 - 2*odd(Ck))
    + 2*Ck*(nk + odd(ck) - 2*nk*odd(Ck))
    + Ck
    + nk
    + odd(Ck)
    - 2*nk*odd(Ck)
  ) /2)

```

The last significative bit

Let's define the upper Nk and lower bits Lk so that:

```

N0 = 2**k * Nk + Lk
Lk = sum(ni * 2**i for i in range(k))
Nk = sum(ni+k * 2**i for i in range(n-k))

```

For any finite number there exists a position of the most significative bit k so that nk=1 and ni=0 for any i>k.

Also Nk=1, Ni = 0 for i>k.

$$fk(N0) = Nk * 3^{Bk} + Ck$$

$$fk(N0) = 3^{Bk} + Ck$$

$$fk(Lk) = Ck$$

Let's be $N0$ the first unreductible natural number. Because $Lk = N0 - 2^k < N0$, both Lk and Ck are reductible

$$fk+1 = (Nk+1)*3^{Bk} + Ck+1 = Ck+1$$

$$2*fk+1 = 2*Ck+1 =$$

$$\begin{aligned}
&= \\
&\quad + nk*3^{Bk} * (3 - 2*odd(Ck)) \\
&\quad + nk \\
&\quad + 2*Ck*(nk + odd(ck) - 2*nk*odd(Ck)) \\
&\quad + Ck \\
&\quad + odd(Ck) \\
&\quad - 2*nk*odd(Ck) \\
&= \quad \text{----- } nk = 1 \\
&\quad + 3^{Bk} * (3 - 2*odd(Ck)) \\
&\quad + 1 \\
&\quad + 2*Ck*(1 + odd(ck) - 2*odd(Ck)) \\
&\quad + Ck \\
&\quad + odd(Ck) \\
&\quad - 2*odd(Ck) \\
&= \quad \text{----- } nk = 1 \\
&\quad + 3^{Bk} * (3 - 2*odd(Ck)) \\
&\quad + 1 \\
&\quad + 2*Ck*(1 - odd(Ck)) \\
&\quad + Ck \\
&\quad - odd(Ck) \\
&= \quad \text{----- split terms} \\
&\quad + 3^{Bk} * (3 - 2*odd(Ck)) \\
&\quad + 1 \\
&\quad + 3*Ck \\
&\quad - 2*Ck*odd(Ck)) \\
&\quad - odd(Ck)
\end{aligned}$$

$$\text{For } odd(Ck) = 1$$

$$2*fk+1 = 3^{Bk} + Ck$$

$$fk+1 = (3^{Bk} + Ck)/2$$

$$fk+1 = \sum_{0 \leq i < Bk} (3^i) + (Ck + 1)/2$$

$$\text{For } odd(Ck) = 0$$

$$2*fk+1 = 3*3^{Bk} + 3*Ck + 1$$

$$fk+1 = (3*3^{Bk} + 3*Ck + 1)/2$$

Prunning outcomes

An outcome gets pruned whenever exists a k so that:

$$2^k * N_k + L_k > 3^{B_k} * N_k + C_k$$

$$N_k (2^k - 3^{B_k}) > C_k - L_k$$

Beyond the most significative bit:

$$C_k < L_k = N_0 \quad \text{## Nothing new apparently}$$

For the most significative bit

$$N_k=1$$

$$(2^k - 3^{B_k}) > (C_k - L_k)$$

For the less significative: $N_k > 1$

$$N_k > (C_k - L_k) / (2^k - 3^{B_k})$$

Because N_k Can be discarded whenever the right side is negative: $-L_k \leq C_k$ and $2^k \geq 3^{B_k} - L_k \geq C_k$ and $2^k \leq 3^{B_k}$

$$3^h / 2^n < 2^h$$

$$\ln(3^h / 2^n) < \ln(2^h)$$

$$\ln(3^h) - \ln(2^n) < \ln(2^h)$$

$$h \cdot \ln(3) - n \cdot \ln(2) < h \cdot \ln(2)$$

$$h \cdot \ln(3) < h \cdot \ln(2) - n \cdot \ln(2)$$

$$h \cdot \ln(3) < (h-n) \cdot \ln(2)$$

$$B = A \gg 1$$

$$\text{A Even, } A = 2 \cdot B \quad A:B0 \quad f^1(A) = B/2 = B$$

$$\text{A Odd, } A = 2B + 1 \quad A:B1 \quad f^1(A) = (3A+1)/2 = (6B+3+1)/2 = 3B + 2 \quad \text{So: } f^1(A) = 3 \cdot B + 2 \text{ same oddity as B}$$

$$C = B \gg 1$$

$$\text{B Even, } B = 2 \cdot C$$

$$A:C01$$

$$f^2(A) = f^1(3 \cdot B + 2) = (6 \cdot C + 2)/2 = 3 \cdot C + 1$$

$$\text{So: } f^2(A) = 3 \cdot C + 1 \quad (\text{opposite oddity than C})$$

$$D = C \gg 1$$

$$\text{C Even, } C = 2 \cdot D$$

$$A:D001$$

$$f^3(A) = f^1(3 \cdot C + 1) = f^1(6 \cdot D + 1) = (18 \cdot D + 4)/2 = 9 \cdot D + 2$$

$$\text{So: } f^3(A) = 9 \cdot D + 2 \quad (\text{same oddity})$$

C Odd, $C = 2*D + 1$
A:D101
 $f^3(A) = f^1(3*C + 1) = f^1(6*D + 4) = 3*D + 2$
So: $f^3(A) = 3*D + 2$ (same oddity)

B Odd, $B = 2*C + 1$
A:C11
 $f^2(A) = f^1(3*B + 2) = (9*B+6+1)/2 = (9*(2*C + 1)+6+1)/2 = 9*C + 8$
So: $f^2(A) = 9*C + 8$ (same oddity as C)

$D = C \gg 1$

C Even, $C = 2*D$
A:D011
 $f^3(A) = F^1(9*C + 8) = F^1(18*D+8) = (18D+8)/2 = 9*D + 4$
 $f^3(A) = 9*D + 4$ (same oddity as C)

C Odd, $C = 2*D + 1$
A:D111
 $f^3(A) = F^1(9*C + 8) = F^1(18*D+17) = (3*18D+3*17 +1)/2 = 27*D + 26$
 $f^3(A) = 27*D + 26$ (same oddity as C)

- Let be a_k the kth bit of the binary representation of A.
- Let be $r_k = A \gg (k)$ (the integer division of A by the n power of two)

If r_0 is even, a_0 is 0, then $f^1(r_0) = r_1 = A/2 < A$, thus imposible. Thus r_0 is odd. $A = 2 r_1 + 1$, $a_0=1$ A odd means that bit 0 is 1. And then $f^1(A) = 3A+1/2 = (6B+3+1)/2 = 3B + 2$

$a_1=0$ B even? If B was even, then $f^2(A)=(3A+1)/4 \geq A$ (for $A>1$) and A would be reducible. So B is odd, lets $B=2C+1$; $A = 2(2C+1)+1 = 4C+3$ B odd means that bit 1 is 1. $f^2(A) = 3((3A+1)/2)+1 = (9A+3)/2+1 = (9A+5)/2 = 18C + 16$ so even $f^3(A) = (9A+5)/4 = 9C+8$

$aX+b$

a even, b even -> even a even, b odd -> odd a odd, b even -> whatever X is a odd, b odd -> whatever X is not

Visualization: We will get a binary search tree, where each levels decide a bit of the number. We can prune the tree every time the function evaluates less than A. We evolve an expression for the $f^n(A)$ in terms of A to get this comparison. We evolve an equivalent expresion for $f^n(A)$ in terms of the undecided bits. If A is a finite number, beyond a given n, the undecided bits are always zero. Which pattern makes that even with zeros as undecided bits it keeps always $\geq A$?

Note: If we can demonstrate that in order not getting under A, we have to add bits for ever then we are demonstrating that the number is infinite.

k even: $f^3(n_0) = (3n_0+1)/4 < n_0$

k odd: $k=2k'+1$ $f^3(n_0) = 3((3n_0+1)/2)+1 = (9n_0+3)/2 + 1 = (9n_0+5)/2$
 $f^3(n_0) = 3(3k+2)+1 = 9k + 7 = 18k'$

Because n_0 is odd, n_0 reduces to $3n_0+1 = 6k+3+1 = 6k + 4$ (even) Being even reduces to $3k+2$

Constructive aproach

$f_{\text{odd}}^{-1} = (2n-1)/3$ $f_{\text{even}}^{-1} = 2n$

```

1
  1/3 X
  2 10
    1 loop
    4 100
      7/3 X
      8 1000
        5 101
          3 11
            5/3 X
            6 110
              11/3 X
              12 1100 ...
                10 1010
                  19/3 X
                  20 10100
                    13 1101 ...
                    40 101000 ...
                      16 10000
                        31/3 X
                        32 100000
                          21 10101 ...
                          41/3 X
                          42 101010
                            85/3 C
                            84 1010100
                              64 1000000 ...

```

Limiting factor

N_0 will be reductible if for any k

$$N_0 > 3^{BkNk + Ck Nk^2} k + \sum (n_i \cdot 2^i) (0 \leq i < k) > 3^{Bk} Nk + Ck$$

Expansion depending on the least significative bits

```

0 N1 + 0
  00 N2 + 0
    000 N3 + 0
      0000 N4 + 0
        00000 N5 + 0
          000000 N6 + 0
            100000 3*N6 + 2
              10000 3*N5 + 2
                010000 3*N6 + 1
                  110000 9*N6 + 8
                    1000 3*N4 + 2
                      01000 3*N5 + 1
                        001000 9*N6 + 2
                          101000 3*N6 + 2
                            11000 9*N5 + 8
                              011000 9*N6 + 4
                                111000 27*N6 + 26
                                  100 3*N3 + 2
                                    0100 3*N4 + 1
                                      00100 9*N5 + 2
                                        000100 9*N6 + 1
                                          100100 27*N6 + 17
                                            10100 3*N5 + 2
                                              010100 3*N6 + 1
                                                110100 9*N6 + 8
                                                  1100 9*N4 + 8
                                                    01100 9*N5 + 4
                                                      001100 9*N6 + 2
                                                        101100 27*N6 + 20
                                                          11100 27*N5 + 26
                                                            011100 27*N6 + 13
                                                              111100 81*N6 + 80
                                                                10 3*N2 + 2
                                                                  010 3*N3 + 1
                                                                    0010 9*N4 + 2
                                                                      00010 9*N5 + 1
                                                                          000010 27*N6 + 2
                                                                              100010 9*N6 + 5
                                                                                  10010 27*N5 + 17
                                                                                      010010 81*N6 + 26
                                                                                          110010 27*N6 + 22
                                                                                            1010 3*N4 + 2
                                                                                              01010 3*N5 + 1
                                                                                                  001010 9*N6 + 2

```

```

101010 3*N6 + 2
11010 9*N5 + 8
011010 9*N6 + 4
111010 27*N6 + 26
110 9*N3 + 8
0110 9*N4 + 4
00110 9*N5 + 2
000110 9*N6 + 1
100110 27*N6 + 17
10110 27*N5 + 20
010110 27*N6 + 10
110110 81*N6 + 71
1110 27*N4 + 26
01110 27*N5 + 13
001110 81*N6 + 20
101110 27*N6 + 20
11110 81*N5 + 80
011110 81*N6 + 40
111110 243*N6 + 242
1 3*N1 + 2
01 3*N2 + 1
001 9*N3 + 2
0001 9*N4 + 1
00001 27*N5 + 2
000001 27*N6 + 1
100001 81*N6 + 44
10001 9*N5 + 5
010001 27*N6 + 8
110001 9*N6 + 7
1001 27*N4 + 17
01001 81*N5 + 26
001001 81*N6 + 13
101001 243*N6 + 161
11001 27*N5 + 22
011001 27*N6 + 11
111001 81*N6 + 74
101 3*N3 + 2
0101 3*N4 + 1
00101 9*N5 + 2
000101 9*N6 + 1
100101 27*N6 + 17
10101 3*N5 + 2
010101 3*N6 + 1
110101 9*N6 + 8
1101 9*N4 + 8
01101 9*N5 + 4

```

```

001101 9*N6 + 2
101101 27*N6 + 20
11101 27*N5 + 26
011101 27*N6 + 13
111101 81*N6 + 80
11 9*N2 + 8
011 9*N3 + 4
0011 9*N4 + 2
00011 9*N5 + 1
000011 27*N6 + 2
100011 9*N6 + 5
10011 27*N5 + 17
010011 81*N6 + 26
110011 27*N6 + 22
1011 27*N4 + 20
01011 27*N5 + 10
001011 27*N6 + 5
101011 81*N6 + 56
11011 81*N5 + 71
011011 243*N6 + 107
111011 81*N6 + 76
111 27*N3 + 26
0111 27*N4 + 13
00111 81*N5 + 20
000111 81*N6 + 10
100111 243*N6 + 152
10111 27*N5 + 20
010111 27*N6 + 10
110111 81*N6 + 71
1111 81*N4 + 80
01111 81*N5 + 40
001111 81*N6 + 20
101111 243*N6 + 182
11111 243*N5 + 242
011111 243*N6 + 121
111111 729*N6 + 728

```

Demonstrated facts

$f_k = N_k \cdot 3^{B_k} + C_k$

where

$N_k = N_0 \gg k$

$0 \leq C_k < 3^{B_k}$

$0 \leq B_k \leq k$

$B_k = \sum_{0 \leq i < k} (\text{odd}(f_i))$

C_k also depends only on the k th lower bits of N_0 ($N_0 - 2^k * N_k$)
Numbers sharing lower k bits share also the oddity of the k th first terms

$$B_{k+1} = B_k + \text{odd}(n_k + C_k)$$

$$C_{k+1} = \left(\begin{aligned} &+ n_k 3^{B_k} * (3 - 2 \cdot \text{odd}(C_k)) \\ &+ C_k * (1 + 2 \cdot \text{odd}(n_k + C_k)) \\ &+ \text{odd}(n_k + C_k) \end{aligned} \right) / 2$$

$$\begin{aligned} 2 \cdot C_{k+1} &= (2 \cdot 0_k + 1) \cdot 3^{B_k} \cdot n_k + C_k \cdot (2 \cdot 0_k + 1) + 0_k && \text{--- } 3^{0_k} = 2 \cdot 0_k + 1 \\ &= 2 \cdot 0_k \cdot 3^{B_k} \cdot n_k + 3^{B_k} \cdot n_k + C_k \cdot 2 \cdot 0_k + C_k + 0_k && \text{--- distribute} \\ C_{k+1} &= 0_k \cdot (n_k \cdot 3^{B_k} + C_k) + (n_k \cdot 3^{B_k} + C_k + 0_k) / 2 && \text{--- divide by 2} \\ C_{k+1} &= 0_k \cdot (n_k \cdot 3^{B_k} + C_k) + (n_k \cdot 3^{B_k} + C_k + 0_k) / 2 && \text{--- } 0_k = n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k) \\ C_{k+1} &= (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot (n_k \cdot 3^{B_k} + C_k) + (n_k \cdot 3^{B_k} + C_k + n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \\ C_{k+1} &= && \text{--- split in lines} \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot (n_k \cdot 3^{B_k} + C_k) \\ &+ (n_k \cdot 3^{B_k} + C_k + n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \\ C_{k+1} &= && \text{--- distribute first term} \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot n_k \cdot 3^{B_k} \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot C_k \\ &+ (n_k \cdot 3^{B_k} + C_k + n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \\ C_{k+1} &= && \text{--- } n_k \cdot n_k = n_k \\ &+ (1 + \text{odd}(C_k) - 2 \cdot \text{odd}(C_k)) \cdot n_k \cdot 3^{B_k} \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot C_k \\ &+ (n_k \cdot 3^{B_k} + C_k + n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \\ C_{k+1} &= && \text{--- } n_k \cdot n_k = n_k \\ &+ (1 - \text{odd}(C_k)) \cdot n_k \cdot 3^{B_k} \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot C_k \\ &+ (n_k \cdot 3^{B_k} + C_k + n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \\ C_{k+1} &= && \text{--- group } 3^{B_k} + 1 \\ &+ (1 - \text{odd}(C_k)) \cdot n_k \cdot 3^{B_k} \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot C_k \\ &+ (n_k \cdot (3^{B_k} + 1) + \text{odd}(C_k) + C_k - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \\ C_{k+1} &= && \text{--- split } (3^{B_k} + 1) / 2 \text{ term} \\ &+ (1 - \text{odd}(C_k)) \cdot n_k \cdot 3^{B_k} \\ &+ n_k \cdot (3^{B_k} + 1) / 2 \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot C_k \\ &+ (\text{odd}(C_k) + C_k - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \\ C_{k+1} &= && \text{--- } (3^{B_k} + 1) / 2 = 1 + \sum_{0 \leq i < k} (3^{B_i}) \\ &+ (1 - \text{odd}(C_k)) \cdot n_k \cdot 3^{B_k} \\ &+ n_k \cdot \sum_{0 \leq i < k} (3^{B_i}) \\ &+ (n_k + \text{odd}(C_k) - 2 \cdot n_k \cdot \text{odd}(C_k)) \cdot C_k \\ &+ (\text{odd}(C_k) + C_k - 2 \cdot n_k \cdot \text{odd}(C_k)) / 2 \end{aligned}$$

```

Ck+1 =
+ (1 - odd(Ck))*nk*3^Bk
+ nk*sum([0<=i<k](3^Bi)
+ (nk + odd(Ck) -2*nk*odd(Ck))*Ck
- nk*odd(Ck)
+ (odd(Ck) + Ck)/2
Ck+1 =
+ nk*sum([0<=i<k+1-odd(Ck)](3^Bi)
+ (nk + odd(Ck) -2*nk*odd(Ck))*Ck
+ (odd(Ck) + Ck)/2
- nk*odd(Ck)
Ck+1 =
+ (nk*3^(k+1-odd(Ck)) -nk)/2
+ (nk + odd(Ck) -2*nk*odd(Ck))*Ck
+ (odd(Ck) + Ck)/2
- nk*odd(Ck)

```

--- extract pair from division
--- extract pair from division
--- $\text{sum}(3^k)[0 \leq k < n] = (3^n - 1) / 2$