

Bayesian Age-Period-Cohort Prediction

Volker Schmid


2026-08-05

Prediction

Because period and cohort effects are modelled as random walks, they can be extrapolated as a continuation of these trends, which lets `predict_apc()` forecast cases for years beyond the observed data. For a first introduction to `bamp()`, see *vignette("bamp", package="bamp")*; for other model specifications, see *vignette("modeling", package="bamp")*.

We use the bundled data example, which covers ten years:

```
data(apc)
plot(cases[,1],type="l",ylim=range(cases), ylab="cases", xlab="year", main="cases per age group")
for (i in 2:8)lines(cases[,i], col=i)
```



To see how well the forecast holds up against real data, we fit the model on only the first nine years and predict the tenth, so it can be compared to the cases actually observed that year.


```
model0 <- bamp(cases[-10,], population[-10,], age="rw1", period="rw1", cohort="rw1",
              periods_per_agegroup = 5)
```

`predict_apc()` extends the fitted model by the given number of periods. With `update = TRUE` the forecast is merged back into `model0` (as `model0$predicted`) instead of being returned as a separate object:

```
model0 <- predict_apc(object=model0, periods=1, population=population, update = TRUE)
```


The forecast for year 10 (dashed lines: 90% credible interval; solid line: median) lines up well with the true total cases (points), which were held out of the fit:

```
ts.plot(t(model0$predicted$cases_period), lty=c(2,1,2))
points(apply(cases,1,sum), pch=19)
```



The same credible intervals extend the period and cohort effects themselves into year 10:

```
ts.plot(t(model0$predicted$period), lty=c(2,1,2))
```



```
ts.plot(t(model0$predicted$cohort), lty=c(2,1,2))
```

