Companion software for "Volker Ziemann, *Physics and Finance, Springer*, 2021" (https://link.springer.com/book/10.1007/978-3-030-63643-2)

Delta hedging (Section 5.4)

Volker Ziemann, 211115, CC-BY-SA-4.0

With this example we provide the opportunity to explore delta hedging and, in particular, recreate the plots shown in Figure 5.3 in the book. The simulation is based on following the value of a call option over 252 trading days as the stock value evolves with annual return ρ and volatility σ . Note that we only take σ into account when calculating the value of the option, but not when evolving the stock S in order to make the development more transparent.

The four sliders can be used to adjust the parameters of the simulation. In particular the return ρ , volatility σ , the risk-free rate r_f , and the strike price K.

In the next section we define the duration of the simulation T0 and the number of trading days N, as well as the initial stock price S0. then we allocate a number of arrays to store the stock price at each trading day, the value of the hedge, the amount of money to borrow from the bank, and the value of the money, originally received when selling the call option. We also prepare an array KK to display the constant value of the agreed-upon strike price

At the start of the year, we calculate the value of the call option c0, the fraction of stocks Δ = delta needed for the hedge, which we borrow from the bank.

Now we iterate through the 252 trading days and at the end of every daym we calculate the remaining time dt until maturity, the interest we have to pay for the borrowed money, the value of the money we received when selling the option, as well as the new stock value (with random variation turned off!).

```
S(i)=S(i-1)*exp(rho/N);%*(1+sig*randn/sqrt(N)); % stocks growth rate
```


Given the updated stock value S(i), we calculate the new hedging fraction delta, calculate the new value of the hedge, and figure out how much money we have to borrow, which is the difference in the values of the hedge and the inetest we need to pay for the borrowed money.

```
[c,delta]=black_scholes_call(S(i),K,dt,rf,sig); % calculate delta
hedge(i)=delta*S(i); % optimal hedge
borrow(i)=borrow(i-1)+hedge(i)-hedge(i-1)+interest; % borrow to adjust hedge
end
```

Once we reach maturity after $\mathbb N$ trading days, we work out how much the hedging actually costs. If the option is *in the money*, our costs is the boorowed money, which includes the handed-over stock, to the holder of the option, minus the agreed-upon strike price K that I receive from the option holder. If the option is out of the money, the hedge at the end is zero, but I had incurred interest payments along the way. Note that the cost calculated here is has to be compared to the value of the money received for the option c0 that has increased in value at the risk-free rate.

Finally we plot the stock value S as a solid black line, the strike price K as a dotted black line, the hedge in red and the borrowed money as a blue dashed line.

```
nn=1:N;
plot(nn,S,'k-',nn,KK,'k:',nn,borrow,'b--',nn,hedge,'r-.','LineWidth',2);
xlim([0,255]);
xlabel('Trading days')
ylabel('S,K,\Delta S,Borrowed')
legend('Stock S','Strike K','Borrowed','Hedge \Delta S')
```



The values set by the sliders at the start lead to the upper plot in Figure 5.3, while in the lower plot κ was increased to 1.2, such that the hedge can be reduced to zero as it cecomes more and more obvious at we do not have to hand over a stock at maturity.

Appendix

The function black_scholes_call() receives the initial stock value S0, the strike price K, the time until maturity dt, the risk-free rate rf, and the volatility sig as input and returns the value of a call option c, its delta and the two other Greeks gamma and theta, which are discussed in Section 6.1.

```
function [c,delta,gamma,theta]=black_scholes_call(S0,K,dt,rf,sig)
N=@(z)0.5*erfc(-z/sqrt(2));
d1=(log(S0./K)+(rf+0.5.*sig.^2).*dt)./(sig.*sqrt(dt));
d2=(log(S0./K)+(rf-0.5.*sig.^2).*dt)./(sig.*sqrt(dt));
c=S0.*N(d1)-K.*exp(-rf.*dt).*N(d2);
delta=N(d1);
gamma=exp(-0.5*d1.^2)./(S0.*sqrt(2*pi*sig.^2.*dt));
theta=-0.5*(sig.^2).*(S0.^2).*gamma-rf.*K.*exp(-rf.*dt).*N(d2);
end
```