

# ECON 899A - Problem Set 7

Alex von Hafften\*

10/27/2021

1. Derive the following asymptotic moments associated with  $m_3(x)$ : mean, variance, first order autocorrelation. Furthermore, compute  $\nabla_b g(b_0)$ . Which moments are informative for estimating  $b$ ?

With  $|\rho_0| < 1$ , the stochastic process  $\{x_t\}$  is stationary, so  $E[x_t] = E[x_{t-1}]$  and  $Var[x_t] = Var[x_{t-1}]$ :<sup>1</sup>

$$E[x_t] = E[\rho_0 x_{t-1} + \varepsilon_t] = \rho_0 E[x_{t-1}] = \rho_0 E[x_t]$$

$$\implies E[x_t] = 0$$

$$Var[x_t] = Var[\rho_0 x_{t-1} + \varepsilon_t] = \rho_0^2 Var[x_{t-1}] + \sigma_0^2 = \rho_0^2 Var[x_t] + \sigma_0^2$$

$$\implies Var[x_t] = \frac{\sigma_0^2}{1 - \rho_0^2}$$

$$Cov[x_t, x_{t-1}] = Cov[\rho_0 x_{t-1} + \varepsilon_t, x_{t-1}] = \rho_0 Cov[x_{t-1}, x_{t-1}] + Cov[\varepsilon_t, x_{t-1}] = \rho_0 Var[x_t] = \frac{\sigma_0^2 \rho_0}{1 - \rho_0^2}$$

---

\*This problem set is for ECON 899A Computational Economics taught by Dean Corbae with assistance from Philip Coyle at UW-Madison. I worked on this problem set with a study group of Michael Nattinger, Sarah Bass, and Xinxin Hu.

<sup>1</sup>Alternatively, we can write  $x_t$  in terms of  $x_0$ ,  $\{\varepsilon_i\}_{i=1}^t$ , and  $\rho_0$ :

$$\begin{aligned} x_t &= \rho_0 x_{t-1} + \varepsilon_t \\ &= \rho_0(\rho_0 x_{t-2} + \varepsilon_{t-1}) + \varepsilon_t \\ &= \rho_0^2 x_{t-2} + \varepsilon_t + \rho_0 \varepsilon_{t-1} \\ &= \rho_0^2(\rho_0 x_{t-3} + \varepsilon_{t-2}) + \varepsilon_t + \rho_0 \varepsilon_{t-1} \\ &= \rho_0^3 x_{t-3} + \varepsilon_t + \rho_0 \varepsilon_{t-1} + \rho_0^2 \varepsilon_{t-2} \\ &= \rho_0^t x_0 + \sum_{i=1}^t \rho_0^{t-i} \varepsilon_i \\ &= \sum_{i=1}^t \rho_0^{t-i} \varepsilon_i \\ E[x_t] &= E\left[\sum_{i=1}^t \rho_0^{t-i} \varepsilon_i\right] = \sum_{i=1}^t \rho_0^{t-i} E[\varepsilon_i] = 0 \end{aligned}$$

Thus, the asymptotic moments associated with  $m_3(x)$  are:

$$\mu(x) = E[m_3(x)] = \begin{bmatrix} 0 \\ \frac{\sigma_0^2}{1-\rho_0^2} \\ \frac{\sigma_0^2 \rho_0}{1-\rho_0^2} \end{bmatrix}$$

To calculate the Jacobian, we compute the derivative of the moment conditions with respect to each parameter:


$$\begin{aligned} \frac{\partial}{\partial \rho} \left( \frac{\sigma^2}{1-\rho^2} \right) &= \frac{\sigma^2(-2\rho) - (1-\rho^2)(0)}{(1-\rho^2)^2} \\ &= \frac{-2\rho\sigma^2}{(1-\rho^2)^2} \\ \frac{\partial}{\partial \sigma} \left( \frac{\sigma^2}{1-\rho^2} \right) &= \frac{2\sigma}{1-\rho^2} \\ \frac{\partial}{\partial \rho} \left( \frac{\sigma^2 \rho}{1-\rho^2} \right) &= \frac{\sigma^2 \rho(-2\rho) - (1-\rho^2)\sigma^2}{(1-\rho^2)^2} \\ &= \frac{-\sigma^2(1+\rho^2)}{(1-\rho^2)^2} \\ \frac{\partial}{\partial \sigma} \left( \frac{\sigma^2 \rho}{1-\rho^2} \right) &= \frac{2\sigma \rho}{1-\rho^2} \end{aligned}$$

Each cell of the Jacobian is the negative of the derivative of the moment condition:

$$\Rightarrow \nabla_b g(b_0) = \begin{bmatrix} 0 & 0 \\ \frac{2\rho_0\sigma_0^2}{(1-\rho_0^2)^2} & \frac{-2\sigma_0}{1-\rho_0^2} \\ \frac{\sigma_0^2(1+\rho_0^2)}{(1-\rho_0^2)^2} & \frac{-2\sigma_0\rho_0}{1-\rho_0^2} \end{bmatrix}$$

Variance and first order autocorrelation are informative for estimating  $b$ .

2. Simulate a series of “true” data of length  $T = 200$  using (1). We will use this to compute  $M_T(x)$ .




$$M_T(x) = \begin{bmatrix} -0.0764 \\ 1.3953 \\ 0.6916 \end{bmatrix}$$

3. Set  $H = 10$  and simulate  $H$  vectors of length  $T = 200$  random variables  $e_t$  from  $N(0, 1)$ . We will use this to compute  $M_{TH}(y(b))$ . Store these vectors. You will use the same vector of random variables throughout the entire exercise. Since this exercise requires you to estimate  $\sigma^2$ , you want to change the variance of  $e_t$  during the estimation. You can simply use  $\sigma e_t$  when the variance is  $\sigma^2$ .
4. We will start by estimating the  $\ell = 2$  vector  $b$  for the just identified case where  $m_2$  uses mean and variance. Given what you found in part (1), do you think there will be a problem? In general we would not know whether this case would be a problem, so hopefully the standard error of the estimate of  $b$  as well as the  $J$  test will tell us something.

From part (1), I found that in expectation the mean is zero, so the problem is that the mean is not informative for estimating  $b$ .

- (a) Set  $W = I$  and graph in three dimensions, the objective function (3) over  $\rho \in [0.35, 0.65]$  and  $\sigma \in [0.8, 1.2]$ . Obtain an estimate of  $b$  by using  $W = I$  in (4) using `fminsearch`. Report  $\hat{b}_{TH}^1$ .



$$\hat{b}_{TH}^1 = \begin{bmatrix} 0.5320 \\ 1.0115 \end{bmatrix}$$

- (b) Set  $i(T) = 4$ . Obtain an estimate of  $W^*$ . Using  $\hat{W}_{TH}^* = \hat{S}_{TH}^{-1}$  in (4), obtain an estimate of  $\hat{b}_{TH}^2$ . Report  $\hat{b}_{TH}^2$ .

...

- (c) To obtain standard errors, compute numerically  $\nabla_b g_T(\hat{b}_{TH}^2)$  defined in (6). Report the values of  $\nabla_b g_T(\hat{b}_{TH}^2)$ . Next, obtain the  $\ell \times \ell$  variance-covariance matrix of  $\hat{b}_{TH}^2$  as in (7). Finally, what are the standard errors defined in (8)? How can we use the information on  $\nabla_b g_T(\hat{b}_{TH}^2)$  to think about local identification?

...

- (d) Since we are in the just identified case, the  $J$  test should be zero (on a computer this may be not be exact). However, given the identification issues in this particular case where we use mean and variance, the  $J$  test may not be zero. Compute the value of the  $J$  test:

$$T \frac{H}{1+H} \times J_{TH}(\hat{b}_{TH}^2) \rightarrow \chi^2$$

noting that in this just identified case  $n - \ell = 0$  degrees of freedom recognizing that there really is not distribution.

...

5. Next we estimating the  $\ell = 2$  vector  $b$  for the just identified case where  $m_2$  uses the variance and autocorrelation. Given what you found in part (1), do you now think there will be a problem? If not, hopefully the standard error of the estimate of  $b$  as well as the  $J$  test will tell us something. For this case, perform steps (a)-(d) above.

(a)

...

(b)

...

(c)

...

(d)

...

6. Next, we will consider the overidentified case where  $m_3$  uses the mean, variance and autocorrelation. For this case, perform steps (a)-(d) above.

(a)

...

(b)

...

(c)

...

(d)

...

- (e) Bootstrap the the finite sample distribution of the estimators by repeatedly drawing  $\varepsilon_t$  and  $e_t^h$  from  $N(0, 1)$  for  $t = 1, \dots, T$  and  $h = 1, \dots, H$ . Compute  $(\hat{b}_{TH}^1, \hat{b}_{TH}^2)$ .

...