

ECON 709B - Problem Set 1

Alex von Hafften*

11/11/2020

1. 2.1 - 2.2¹

2.1 Find $E[E[E[Y|X_1, X_2, X_3]|X_1, X_2]|X_1]$.

Using the law of iterated expectations,

$$E[E[E[Y|X_1, X_2, X_3]|X_1, X_2]|X_1] = E[E[Y|X_1, X_2]|X_1] = E[Y|X_1]$$

2.2 If $E[Y|X] = a + bX$, find $E[YX]$ as a function of moments of X .

Using the law of iterated expectations,

$$E[YX] = E[E[YX|X]] = E[XE[Y|X]] = E[X(a + bX)] = E[aX + bX^2] = aE[X] + bE[X^2]$$

2. 2.3 Prove conclusion (4) of Theorem 2.4.

If $E|Y| < \infty$ then for any function $h(x)$ such that $E|h(X)e| < \infty$ then $E[h(X)e] = 0$.

Proof: Using the law of iterated expectations, Theorem 2.3, and conclusion (1) (i.e., $E[e|X] = 0$),

$$E[h(X)e] = E[E[h(X)e|X]] = E[h(X)E[e|X]] = E[h(X)(0)] = E[0] = 0$$

□

*I worked on this problem set with a study group of Michael Nattinger, Andrew Smith, and Ryan Mather. I also discussed problems with Emily Case, Sarah Bass, and Danny Edgel.

¹These problems come from *Econometrics* by Bruce Hansen, revised on October 23, 2020.

3. 2.4 Suppose that the random variables Y and X only take the values 0 and 1, and have the following joint probability distribution

	$X = 0$	$X = 1$
$Y = 0$.1	.2
$Y = 1$.4	.3

Find $E[Y|X]$, $E[Y^2|X]$ and $\text{var}[Y|X]$ for $X = 0$, $X = 1$.

$$\begin{aligned}
 E[Y|X = 0] &= (1)P[Y = 1|X = 0] + (0)P[Y = 0|X = 0] = (1)(.4)/(.5) = .8 \\
 E[Y|X = 1] &= (1)P[Y = 1|X = 1] + (0)P[Y = 0|X = 1] = (1)(.3)/(.5) = .6 \\
 E[Y^2|X = 0] &= (1)^2P[Y = 1|X = 0] + (0)^2P[Y = 0|X = 0] = (1)^2(.4)/(.5) = .8 \\
 E[Y^2|X = 1] &= (1)^2P[Y = 1|X = 1] + (0)^2P[Y = 0|X = 1] = (1)^2(.3)/(.5) = .6 \\
 \text{var}[Y|X = 0] &= E[Y^2|X = 0] - (E[Y|X = 0])^2 = (.8) - (.8)^2 = 0.16 \\
 \text{var}[Y|X = 1] &= E[Y^2|X = 1] - (E[Y|X = 1])^2 = (.6) - (.6)^2 = 0.24
 \end{aligned}$$

4. 2.5 (c) Show that $\sigma^2(X)$ is the best predictor of e^2 given X . Show that $\sigma^2(X)$ minimizes the mean-squared error and is thus the best predictor.

.....

5. 2.8 Suppose that Y is discrete-valued, taking values only on the non-negative integers, and the conditional distribution of Y given $X = x$ is Poisson: $P[Y = j|X = x] = \frac{\exp(-x'\beta)(x'\beta)^j}{j!}$, $j = 0, 1, 2, \dots$. Compute $E[Y|X]$ and $\text{var}[Y|X]$. Does this justify a linear regression model of the form $Y = X'\beta + e$?²

Using the hint, we know that $E[Y|X] = x'\beta$ and $\text{var}[Y|X] = x'\beta$.

Yes, this justifies a linear regression model because $E[e|X] = E[Y - X'\beta|X] = E[Y|X] - E[X'\beta|X] = x'\beta - x'\beta = 0$.

6. 2.10 - 2.14 Explain your answers.

2.10 If $Y = X\beta + e$, $X \in \mathbb{R}$, and $E[e|X] = 0$, then $E[X^2e] = 0$.

True, based on the law of iterated expectation:

$$E[X^2e] = E[E[X^2e|X]] = E[X^2E[e|X]] = E[X^2(0)] = E[0] = 0$$

2.11 If $Y = X\beta + e$, $X \in \mathbb{R}$, and $E[Xe] = 0$, then $E[X^2e] = 0$.

False, for a counter example, assume $X \sim N(0, 1)$ and e is a degenerate random variable equal to 1. Notice that $E[Xe] = E[X] = 0$ and $E[X^2e] = E[X^2] = 1$.

2.12 If $Y = X'\beta + e$, and $E[e|X] = 0$, then e is independent of X .

False, for a counter example, assume $X = (X_1, \dots, X_k)'$ and $U = (U_1, \dots, U_k)'$ where X_1, \dots, X_k and U_1, \dots, U_k are independently distributed standard normal. Let $e = X'U$. Thus, conditional on X , e is distributed $N(0, X'X)$, so $E[e|X] = 0$, but X and e are not independent.

2.13 If $Y = X'\beta + e$, and $E[Xe] = 0$, then $E[e|X] = 0$.

False, for a counter example, assume $X = (X_1, \dots, X_k)'$ where $X_1, \dots, X_k \sim N(0, 1)$ and e is a degenerate random variable equal to 1. Notice that $E[Xe] = E[X] = 0$ and $E[e|X] = E[e] = 1$.

²Hint: $P[Y = j] = \frac{\exp(-\lambda)(\lambda)^j}{j!}$, then $E[Y] = \lambda$ and $\text{var}[Y] = \lambda$.

2.14 If $Y = X'\beta + e$, and $E[e|X] = 0$, and $E[e^2|X] = \sigma^2$, then e is independent of X .
False, ...

7. 2.16 Let X and Y have the joint density $f(x, y) = \frac{3}{2}(x^2 + y^2)$ on $0 \leq x \leq 1, 0 \leq y \leq 1$. Compute the coefficients of the best linear predictor $Y = \alpha + \beta X + e$. Compute the conditional expectation $m(x) = E[Y|X = x]$. Are the best linear predictor and conditional expectation different?

Best Linear Predictor (BLP):

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy = \int_0^1 \frac{3}{2}(x^2 + y^2) dy = \frac{3}{2} \left[x^2 y + \frac{y^3}{3} \right]_{y=0}^1 = \frac{3}{2}x^2 + \frac{1}{2}$$

$$E[X] = \int_0^1 x \left(\frac{3}{2}x^2 + \frac{1}{2} \right) dx = \left[\frac{3}{8}x^4 + \frac{1}{4}x^2 \right]_0^1 = \frac{5}{8}$$

$$E[X^2] = \int_0^1 x^2 \left(\frac{3}{2}x^2 + \frac{1}{2} \right) dx = \left[\frac{3}{10}x^5 + \frac{1}{6}x^3 \right]_0^1 = \frac{7}{15}$$

Since the joint distribution is symmetric, $E[Y] = \frac{5}{8}$ and $E[Y^2] = \frac{7}{15}$.

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy \frac{3}{2}(x^2 + y^2) dx dy \\ &= \frac{3}{2} \int_0^1 \int_0^1 x^3 y + xy^3 dx dy \\ &= \frac{3}{2} \int_0^1 \left[\frac{x^4 y}{4} + \frac{x^2 y^3}{2} \right]_{x=0}^1 dy \\ &= \frac{3}{2} \int_0^1 \frac{y}{4} + \frac{y^3}{2} dy \\ &= \frac{3}{2} \left[\frac{y^2}{8} + \frac{y^4}{8} \right]_0^1 \\ &= \frac{3}{8} \end{aligned}$$

$$\begin{aligned} S(\alpha, \beta) &= E[(Y - \alpha - \beta X)^2] \\ &= E[Y^2 - \alpha Y - \beta XY - \alpha Y + \alpha^2 + \alpha\beta X - \beta XY + \alpha\beta X + \beta^2 X^2] \\ &= E[Y^2] - \alpha E[Y] - \beta E[XY] - \alpha E[Y] + \alpha^2 + \alpha\beta E[X] - \beta E[XY] + \alpha\beta E[X] + \beta^2 E[X^2] \\ &= \frac{7}{15} - \frac{5}{8}\alpha - \frac{3}{8}\beta - \frac{5}{8}\alpha + \alpha^2 + \frac{5}{8}\alpha\beta - \frac{3}{8}\beta + \frac{5}{8}\alpha\beta + \frac{7}{15}\beta^2 \\ &= \frac{7}{15} - \frac{5}{4}\alpha - \frac{3}{4}\beta + \alpha^2 + \frac{5}{4}\alpha\beta + \frac{7}{15}\beta^2 \end{aligned}$$

FOC $[\alpha]$:

$$0 = -\frac{5}{4} + 2\alpha + \frac{5}{4}\beta \implies \beta = 1 - \frac{8}{5}\alpha$$

FOC $[\beta]$:


$$0 = -\frac{3}{4} + \frac{5}{4}\alpha + \frac{14}{15}\beta \implies 45 = 75\alpha + 56\beta$$

$$45 = 75\alpha + 56\left(1 - \frac{8}{5}\alpha\right) \implies \alpha = \frac{55}{73} \implies \beta = \frac{-15}{73}$$

Conditional Expectation Function (CEF):

$$f_{Y|X}(y|x) = \frac{f(x, y)}{f_X(x)} = \frac{\frac{3}{2}(x^2 + y^2)}{\frac{3}{2}x^2 + \frac{1}{2}} = \frac{3x^2 + 3y^2}{3x^2 + 1}$$

$$m(x) = E[Y|X = x] = \int_{-\infty}^{\infty} y f_{Y|X}(y|x) dy = \int_0^1 \frac{3x^2 y + 3y^3}{3x^2 + 1} dy = \left[\frac{\frac{3}{2}x^2 y^2 + \frac{3}{4}y^4}{3x^2 + 1} \right]_{y=0}^1 = \frac{6x^2 + 3}{12x^2 + 4}$$



8. 4.1 - 4.6

4.1 For some integer k , set $\mu_k = E[Y^k]$.

(a) Construct an estimator $\hat{\mu}_k$ for μ_k .

$$\hat{\mu}_k = \frac{1}{n} \sum_{i=0}^n Y_i^k$$

(b) Show that $\hat{\mu}_k$ is unbiased for μ_k .

$$E[\hat{\mu}_k] = E\left[\frac{1}{n} \sum_{i=0}^n Y_i^k\right] = \frac{1}{n} \sum_{i=0}^n E[Y_i^k] = \frac{1}{n} \sum_{i=0}^n \mu_k = \mu_k$$

(c) Calculate the variance of $\hat{\mu}_k$, say $\text{var}[\hat{\mu}_k]$. What assumption is needed for $\text{var}[\hat{\mu}_k]$ to be finite?

We need to assume that $\mu_{2k} < \infty$ for $\text{var}[\hat{\mu}_k]$ to be finite:

$$\text{var}[\hat{\mu}_k] = \text{var}\left[\frac{1}{n} \sum_{i=0}^n Y_i^k\right] = \frac{1}{n^2} \sum_{i=0}^n \text{var}[Y_i^k] = \frac{1}{n^2} \sum_{i=0}^n (E[Y_i^{2k}] - E[Y_i^k]^2) = \frac{1}{n^2} \sum_{i=0}^n (\mu_{2k} - \mu_k^2) = \frac{\mu_{2k} - \mu_k^2}{n}$$

(d) Propose an estimator of $\text{var}[\hat{\mu}_k]$.

$$\frac{\hat{\mu}_{2k} - \hat{\mu}_k^2}{n} = \frac{\sum_{i=0}^n Y_i^{2k} - (\sum_{i=0}^n Y_i^k)^2}{n}$$

4.2 Calculate $E[(\bar{Y} - \mu)^3]$, the skewness of \bar{y} . Under what conditions is it zero?

...

4.3 Explain the difference between \bar{Y} and μ . Explain the difference between $n^{-1} \sum_{i=1}^n X_i X_i'$ and $E X_i X_i'$.

...

4.4 True or False. If $Y_i = X_i \beta + e_i$, $X_i \in \mathbb{R}$, $E[x_i | X_i] = 0$, and \hat{e}_i is the OLS residual from the regression of Y_i on X_i , then $\sum_{i=1}^n X_i^2 \hat{e}_i = 0$.

...

4.5 Prove (4.15) and (4.16).

$$(4.15) \quad E[\hat{\beta} | X] = \beta$$

...

$$(4.16) \quad \text{var}[\hat{\beta} | X] = (X'X)^{-1}(X'\Omega X)(X'X)^{-1}$$

...

4.6 Prove Theorem 4.5.

Theorem 4.5 Generalized Gauss-Markov

In the linear regression model (Assumption 4.2) and $\Omega > 0$, if $\tilde{\beta}$ is a linear unbiased estimator of β then $\text{var}[\tilde{\beta} | X] \geq (X'\Omega^{-1}X)^{-1}$

...