

ECON 703 - PS 3

Alex von Hafften*

8/28/2020

(1) Let (X, d) be a nonempty complete metric space. Suppose an operator $T : X \rightarrow X$ satisfies $d(T(x), T(y)) < d(x, y)$ for all $x \neq y, x, y \in X$. Prove or disprove that T has a fixed point. Compare with the Contraction Mapping Theorem.

I prove that T has a fixed point.

Proof: Define $\beta_{xy} := \frac{d(T(x), T(y))}{d(x, y)}$ and $\beta := \max \beta_{xy}$ for all $x \neq y, x, y \in X$. Notice that $d(T(x), T(y)) < d(x, y) \implies \beta_{xy} < 1$. Thus, $d(T(x), T(y)) = \beta_{xy}d(x, y) \leq \beta d(x, y)$. By the convergence mapping theorem, T has a fixed point. \square

(2) Does there exist a countable set, which is compact?

Yes.

Proof: By definition, \mathbb{N} is countable. \mathbb{N} is a finite open cover for \mathbb{N} in metric space (\mathbb{N}, d_E) because \mathbb{N} is open in (\mathbb{N}, d_E) and $\mathbb{N} \subset \mathbb{N}$. Thus, \mathbb{N} is compact in (\mathbb{N}, d_E) . \square

(3) Prove that the function $f(x) = \cos^2(x)e^{5-x-x^2}$ has a maximum on \mathbb{R} .

Proof: Using the extreme value theorem, I show that f attains a maximum on \mathbb{R} by (i) finding $C \subset \mathbb{R}$ where $\exists x \in C$ such that $f(x) > 1$ and $f(y) < 1 \forall y \in C^c$, (ii) showing C is compact, and (iii) f is continuous on \mathbb{R} . Define $g(x) = \cos^2(x)$, $h_1(x) = e^x$, $h_2(x) = 5 - x - x^2$, so that $f(x) = g(x)h_1(h_2(x))$.

For (i), define $C = [(-\sqrt{21} - 1)/2, (\sqrt{21} - 1)/2]$. For $0 \in C$, $f(0) = \cos^2(0)e^{5-0-0^2} = e^5 > 1$. Since $0 \leq \cos^2(x) \leq 1$, $f(x) \leq h_1(h_2(x))$ for all $x \in \mathbb{R}$. Furthermore, if $h_2(x) < 0 \implies h_1(h_2(x)) < 1$. Because $h_2(0) = 5 - 0 - 0^2 = 5$ and the quadratic roots of h_2 are $x = (\sqrt{21} - 1)/2$ and $x = (-\sqrt{21} - 1)/2$, $f(x) \leq h_1(h_2(x)) < 1$ for all $x \in C^c$. Thus, the maximum of $f(x)$ must occur on C .

For (ii), as a closed interval of \mathbb{R} , C is compact by the Heine-Borel theorem.

For (iii), since g , h_1 , and h_2 are continuous, f is continuous.

Thus, by the extreme value theorem, f attains a maximum on C and therefore on \mathbb{R} . \square

*I worked on this problem set with a study group of Michael Nattinger, Andrew Smith, Tyler Welch, and Ryan Mather. I also discussed problems with Emily Case, Sarah Bass, and Danny Edgel.

(4) Suppose you have two maps of Wisconsin: one large and one small. You put the large one on top of the small one, so that the small one is completely covered by the large one. Prove that it is possible to pierce the stack of those two maps in a way that the needle will go through exactly the same (geographical) points on both maps.

Proof: Draw Cartesian planes on both maps such that the Wisconsin Capital Building in Madison is at the origin. For the larger map, scale the x coordinate such that moving one unit right corresponds to the point on the map that is one inch east of the Capital Building on the map and scale the y coordinate such that moving one unit up corresponds to the point on the map that is one inch north of the Capital Building on the map. Scale the Cartesian plane on the smaller map similarly.

Define β as the ratio of the miles between geographical points per inch on the larger map to the miles between geographical points per inch on the smaller map. Notice that $\beta < 1$. Define $A \subset \mathbb{R} \times \mathbb{R}$ as the closed set of points on the larger map that are on or within Wisconsin's borders. Notice that the metric space (A, d_E) is complete because A is closed. Define operator $T : A \rightarrow \mathbb{R} \times \mathbb{R}$ such that, for $x \in A$, (x_1, x_2) on the larger map and $(T(x_1), T(x_2))$ on the smaller map represent the same geographical point. Notice that, for all $x, y \in A, x \neq y$, $d_E(x, y) = \beta d_E(T(x), T(y))$. Thus, T is a contraction with modulus $\beta < 1$. Thus, T has a fixed point where $x^* = T(x^*)$, where you can pierce the stack of maps at the same geographical point. \square