

ECON 710B - Problem Set 7

Alex von Hafften*

3/23/2021

13.1

Take the model:

$$\begin{aligned}Y &= X'\beta + e \\E[Xe] &= 0 \\e^2 &= Z'\gamma + \eta \\E[Z\eta] &= 0\end{aligned}$$

Find the method of moments estimators $(\hat{\beta}, \hat{\gamma})$ for (β, γ) .

The moment conditions are:

$$\begin{aligned}&\begin{pmatrix} E[Xe] \\ E[Z\eta] \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ \Rightarrow &\begin{pmatrix} E[X(Y - X'\beta)] \\ E[Z((Y - X'\beta)^2 - Z'\gamma)] \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ \Rightarrow &\begin{pmatrix} E[g_1(\beta, \gamma)] \\ E[g_2(\beta, \gamma)] \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ &\text{where } g_1(\beta, \gamma) = XY - XX'\beta, \\ &\quad g_2(\beta, \gamma) = Z(Y - X'\beta)^2 - ZZ'\gamma\end{aligned}$$

Replacing with the sample moment:

$$\begin{aligned}\frac{1}{n} \sum_{i=1}^n (X_i Y_i - X_i X_i' \hat{\beta}) &= 0 \Rightarrow \hat{\beta} = \left(\frac{1}{n} \sum_{i=1}^n X_i X_i' \right)^{-1} \left(\frac{1}{n} \sum_{i=1}^n X_i Y_i \right) \\ \frac{1}{n} \sum_{i=1}^n (Z_i (Y_i - X_i' \hat{\beta})^2 - Z_i Z_i' \hat{\gamma}) &= 0 \Rightarrow \hat{\gamma} = \left(\frac{1}{n} \sum_{i=1}^n Z_i Z_i' \right)^{-1} \left(\frac{1}{n} \sum_{i=1}^n Z_i (Y_i - X_i' \hat{\beta})^2 \right)\end{aligned}$$

*I worked on this problem set with a study group of Michael Nattinger, Andrew Smith, and Ryan Mather. I also discussed problems with Sarah Bass, Emily Case, Danny Edgel, and Katherine Kwok.

13.2

Take the model $Y = X'\beta + e$ with $E[e|Z] = 0$. Let β_{gmm} be the GMM estimator using the weight matrix $W_n = (Z'Z)^{-1}$. Under the assumption $E[e^2|Z] = \sigma^2$ show that

$$\sqrt{n}(\hat{\beta}_{gmm} - \beta) \rightarrow_d N(0, \sigma^2(Q'M^{-1}Q)^{-1})$$

where $Q = E[ZX']$ and $M = E[ZZ']$.

We can rewrite $\hat{\beta}_{gmm}$ as:

$$\begin{aligned}\hat{\beta}_{gmm} &= (X'ZW_nZ'X)^{-1}(X'ZW_nZ'Y) \\ &= (X'Z(nW_n)Z'X)^{-1}(X'Z(nW_n)Z'Y) \\ &= (X'ZV_nZ'X)^{-1}(X'ZV_nZ'Y)\end{aligned}$$

where $V_n = (n^{-1}Z'Z)^{-1}$. Notice that

$$n^{-1}Z'Z \rightarrow_p E[Z'Z]$$

by law of large numbers, so by CMT:

$$V_n = (n^{-1}Z'Z)^{-1} \rightarrow_p E[Z'Z]^{-1} \equiv W$$

Notice that $M = W^{-1}$. If $E[e^2|Z] = \sigma^2$, then

$$\Omega = E[ZZ'e^2] = E[ZZ'E[e^2|Z]] = \sigma^2 E[ZZ'] = \sigma^2 M = \sigma^2 W^{-1}$$

By Theorem 13.3, we know that $\sqrt{n}(\hat{\beta}_{gmm} - \beta) \rightarrow_d N(0, V_\beta)$ where

$$\begin{aligned}V_\beta &= (Q'WQ)^{-1}(Q'W\Omega WQ)(Q'WQ)^{-1} \\ &= (Q'WQ)^{-1}(Q'W\sigma^2 W^{-1}WQ)(Q'WQ)^{-1} \\ &= \sigma^2(Q'WQ)^{-1}(Q'WQ)(Q'WQ)^{-1} \\ &= \sigma^2(Q'WQ)^{-1} \\ &= \sigma^2(Q'M^{-1}Q)^{-1}\end{aligned}$$

13.3

Take the model $Y = X'\beta + e$ with $E[Ze] = 0$. Let $\tilde{e} = Y - X'\hat{\beta}$ where $\hat{\beta}$ is consistent for β (e.g. a GMM estimator with some weight matrix). An estimator of the optimal GMM weight matrix is

$$\hat{W} = \left(\frac{1}{n} \sum_{i=1}^n Z_i Z_i' \tilde{e}_i^2 \right)^{-1}$$

Show that $\hat{W} \rightarrow_p \Omega^{-1}$ where $\Omega = E[ZZ'e^2]$.

$$\frac{1}{n} \sum_{i=1}^n \tilde{e}_i^2 = \frac{1}{n} \sum_{i=1}^n (Y_i - X_i' \hat{\beta})^2 = \frac{1}{n} \sum_{i=1}^n Y_i^2 - 2 \frac{1}{n} \sum_{i=1}^n Y_i X_i' \hat{\beta} + \frac{1}{n} \sum_{i=1}^n X_i' X_i \hat{\beta} \hat{\beta}' \rightarrow_p E[Y^2] - 2E[YX'\beta] + E[X'X\beta'\beta] = E[(Y - X'\beta)^2]$$

By the continuous mapping theorem and the weak law of large numbers.

...

13.4

In the linear model estimated by GMM with general weight matrix W the asymptotic variance of $\hat{\beta}_{gmm}$ is

$$V = (Q'WQ)^{-1}Q'W\Omega WQ(Q'WQ)^{-1}$$

(a) Let V_0 be this matrix when $W = \Omega^{-1}$. Show that $V_0 = (Q'\Omega^{-1}Q)^{-1}$.

$$\begin{aligned} V_0 &= (Q'\Omega^{-1}Q)^{-1}Q'\Omega^{-1}\Omega\Omega^{-1}Q(Q'\Omega^{-1}Q)^{-1} \\ &= (Q'\Omega^{-1}Q)^{-1}Q'\Omega^{-1}Q(Q'\Omega^{-1}Q)^{-1} \\ &= (Q'\Omega^{-1}Q)^{-1} \end{aligned}$$

(b) We want to show that for any W , $V - V_0$ is positive semi-definite (for then V_0 is the smaller possible covariance matrix and $W = \Omega^{-1}$ is the efficient weight matrix). To do this start by finding matrices A and B such that $V = A'\Omega A$ and $V_0 = B'\Omega B$.

$$\begin{aligned} V &= (Q'WQ)^{-1}Q'W\Omega WQ(Q'WQ)^{-1} \\ &= A'\Omega A \\ A &:= WQ(Q'WQ)^{-1} \\ A' &= (WQ(Q'WQ)^{-1})' \\ &= ((Q'WQ)')^{-1}Q'W' \\ &= (Q'WQ)^{-1}Q'W \end{aligned}$$

Since W is symmetric $\implies Q'WQ$ is symmetric.

$$\begin{aligned} V_0 &= (Q'\Omega^{-1}Q)^{-1}Q'\Omega^{-1}\Omega\Omega^{-1}Q(Q'\Omega^{-1}Q)^{-1} \\ &= B'\Omega B \\ B &:= \Omega^{-1}Q(Q'\Omega^{-1}Q)^{-1} \\ B' &= (\Omega^{-1}Q(Q'\Omega^{-1}Q)^{-1})' \\ &= (Q'\Omega^{-1}Q)^{-1}Q'\Omega^{-1} \end{aligned}$$

13.11

As a continuation of Exercise 12.7 derive the efficient GMM estimator using the instrument $Z = (XX2)'$. Does this differ from 2SLS and/or OLS?

13.13

- (a) (See appendix A10 pg 972; Theorem A.4 (4): By the spectral decomposition, $A = H\Lambda H'$ where $H'H = I_k$ and Λ is diagonal with non-negative diagonal elements. All diagonal elements of Λ are strictly positive if (and only if) $A > 0$.)

13.18

13.19

13.28

17.15

In this exercise you will replicate and extend the empirical work reported in Arellano and Bond (1991) and Blundell and Bond (1998). Arellano-Bond gathered a dataset of 1031 observations from an unbalanced panel of 140 U.K. companies for 1976-1984 and is in the datafile **AB1991** on the textbook webpage. The variables we will be using are log employment (N), log real wages (W), and log capital (K). See the description file for definitions.

- (a) Estimate the panel AR(1) $K_{it} = \alpha K_{it-1} + u_i + v_t + \varepsilon_{it}$ using Arellano-Bondone-step GMM with clustered standard errors. Note that the model includes year fixed effects.

...

- (b) Re-estimate using Blundell-Bondone-step GMM with clustered standard errors.

...

- (c) Explain the difference in the estimates. ...