

ECON 709B - Problem Set 1

Alex von Hafften*

11/11/2020

1. 2.1 - 2.2¹

2.1 Find $E[E[E[Y|X_1, X_2, X_3]|X_1, X_2]|X_1]$.

By the law of iterated expectations,

$$E[E[E[Y|X_1, X_2, X_3]|X_1, X_2]|X_1] = E[E[Y|X_1, X_2]|X_1] = E[Y|X_1]$$

2.2 If $E[Y|X] = a + bX$, find $E[XY]$ as a function of moments of X .

By the conditioning theorem,

$$E[XY] = E[E[XY|X]] = E[XE[Y|X]] = E[X(a + bX)] = E[aX + bX^2] = aE[X] + bE[X^2]$$

2. 2.3 Prove conclusion (4) of Theorem 2.4.

If $E|Y| < \infty$ then for any function $h(x)$ such that $E|h(X)e| < \infty$ then $E[h(X)e] = 0$.

Proof: Let h be a function such that $E|h(X)e| < \infty$. By the conditioning theorem and conclusion (1) of Theorem 2.4 (i.e., $E[e|X] = 0$),

$$E[h(X)e] = E[E[h(X)e|X]] = E[h(X)E[e|X]] = E[h(X)(0)] = E[0] = 0$$

□

*I worked on this problem set with a study group of Michael Nattinger, Andrew Smith, and Ryan Mather. I also discussed problems with Emily Case, Sarah Bass, and Danny Edgel.

¹These problems come from *Econometrics* by Bruce Hansen, revised on October 23, 2020.

3. 2.4 Suppose that the random variables Y and X only take the values 0 and 1, and have the following joint probability distribution

	$X = 0$	$X = 1$
$Y = 0$.1	.2
$Y = 1$.4	.3

Find $E[Y|X]$, $E[Y^2|X]$ and $\text{var}[Y|X]$ for $X = 0$, $X = 1$.

$$\begin{aligned}
E[Y|X = 0] &= (1)P[Y = 1|X = 0] + (0)P[Y = 0|X = 0] = (1)(.4)/(.5) = .8 \\
E[Y|X = 1] &= (1)P[Y = 1|X = 1] + (0)P[Y = 0|X = 1] = (1)(.3)/(.5) = .6 \\
E[Y^2|X = 0] &= (1)^2P[Y = 1|X = 0] + (0)^2P[Y = 0|X = 0] = (1)^2(.4)/(.5) = .8 \\
E[Y^2|X = 1] &= (1)^2P[Y = 1|X = 1] + (0)^2P[Y = 0|X = 1] = (1)^2(.3)/(.5) = .6 \\
\text{var}[Y|X = 0] &= E[Y^2|X = 0] - (E[Y|X = 0])^2 = (.8) - (.8)^2 = 0.16 \\
\text{var}[Y|X = 1] &= E[Y^2|X = 1] - (E[Y|X = 1])^2 = (.6) - (.6)^2 = 0.24
\end{aligned}$$

4. 2.5 (c) Show that $\sigma^2(X)$ is the best predictor of e^2 given X . Show that $\sigma^2(X)$ minimizes the mean-squared error and is thus the best predictor.

For $S(X)$ some predictor of e^2 given X :

$$\begin{aligned}
E[(e^2 - S(X))^2] &= E[(e^2 - \sigma^2(X) + \sigma^2(X) - S(X))^2] \\
&= E[(e^2 - \sigma^2(X))^2] + 2E[(e^2 - \sigma^2(X))(\sigma^2(X) - S(X))] + E[(\sigma^2(X) - S(X))^2]
\end{aligned}$$

The middle term is zero:

$$\begin{aligned}
E[(e^2 - \sigma^2(X))(\sigma^2(X) - S(X))] &= E[E[(e^2 - \sigma^2(X))(\sigma^2(X) - S(X))|X]] \\
&= E[E[(e^2 - \sigma^2(X))|X](\sigma^2(X) - S(X))] \\
&= E[(E[e^2|X] - E[\sigma^2(X)|X])(\sigma^2(X) - S(X))] \\
&= E[(\sigma^2(X) - \sigma^2(X))(\sigma^2(X) - S(X))] \\
&= 0
\end{aligned}$$

Thus,

$$E[(e^2 - S(X))^2] = E[(e^2 - \sigma^2(X))^2] + E[(\sigma^2(X) - S(X))^2]$$

The choice of S does not change the first term and the second term is minimized when $S(X) = \sigma^2(X)$. Thus, $\sigma^2(X)$ is the best predictor of e^2 .

5. 2.8 Suppose that Y is discrete-valued, taking values only on the non-negative integers, and the conditional distribution of Y given $X = x$ is Poisson: $P[Y = j|X = x] = \frac{\exp(-x'\beta)(x'\beta)^j}{j!}, j = 0, 1, 2, \dots$. Compute $E[Y|X]$ and $\text{var}[Y|X]$. Does this justify a linear regression model of the form $Y = X'\beta + e$?²

Using the hint, we know that $E[Y|X] = x'\beta$ and $\text{var}[Y|X] = x'\beta$.

Yes, this justifies a linear regression model because $E[e|X] = E[Y - X'\beta|X] = E[Y|X] - E[X'\beta|X] = x'\beta - x'\beta = 0$.

6. 2.10 - 2.14 Explain your answers.

2.10 If $Y = X\beta + e$, $X \in \mathbb{R}$, and $E[e|X] = 0$, then $E[X^2e] = 0$.

True, by the conditioning theorem:

$$E[X^2e] = E[E[X^2e|X]] = E[X^2E[e|X]] = E[X^2(0)] = E[0] = 0$$

2.11 If $Y = X\beta + e$, $X \in \mathbb{R}$, and $E[Xe] = 0$, then $E[X^2e] = 0$.

False, for a counter example, assume $X \sim N(0, 1)$ and e is a degenerate random variable equal to 1. Notice that $E[Xe] = E[X] = 0$ and $E[X^2e] = E[X^2] = 1$.

2.12 If $Y = X'\beta + e$, and $E[e|X] = 0$, then e is independent of X .

False, for a counter example, assume $X = (X_1, \dots, X_k)'$ and $U = (U_1, \dots, U_k)'$ where X_1, \dots, X_k and U_1, \dots, U_k are independently distributed standard normal. Let $e = X'U$. Thus, conditional on X , e is distributed $N(0, X'X)$, so $E[e|X] = 0$, but X and e are not independent.

2.13 If $Y = X'\beta + e$, and $E[Xe] = 0$, then $E[e|X] = 0$.

False, for a counter example, assume $X = (X_1, \dots, X_k)'$ where $X_1, \dots, X_k \sim N(0, 1)$ and e is a degenerate random variable equal to 1. Notice that $E[Xe] = E[X] = 0$ and $E[e|X] = E[e] = 1$.

2.14 If $Y = X'\beta + e$, and $E[e|X] = 0$, and $E[e^2|X] = \sigma^2$, then e is independent of X .

False, for a counter example, assume $X = (X_1, \dots, X_k)'$ where $X_i \sim N(0, 1)$ and $Z = (Z_1, \dots, Z_k)$ where $Z_i \sim N(1, \sigma^2/x_i^2)$. Thus, $E[Z_i|X_i] = 1$ and $\text{var}[Z_i|X_i] = \sigma^2/x_i^2$. Define $Y := X'Z$. Notice that $E[Y|X] = E[X'Z|X] = X'E[Z|X] = \sum_{i=1}^n X_i$, so $\beta = (1, \dots, 1)'$. Define $e := Y - E[Y|X] = Y - \sum_{i=1}^n X_i = X'(Z - 1)$. Notice that $E[e|X] = E[Y - E[Y|X]|X] = 0$ and $E[e^2|X] = E[X'(Z - 1)(Z - 1)'X|X] = X'E[(Z - 1)(Z - 1)'|X]X = X'E[(Z - 1)(Z - 1)'|X]X = X'\text{var}[Z|X]X = X'(X'\sigma^2X)^{-1}X = \sigma^2$. However, X and e are not independent.

²Hint: $P[Y = j] = \frac{\exp(-\lambda)(\lambda)^j}{j!}$, then $E[Y] = \lambda$ and $\text{var}[Y] = \lambda$.

7. 2.16 Let X and Y have the joint density $f(x, y) = \frac{3}{2}(x^2 + y^2)$ on $0 \leq x \leq 1, 0 \leq y \leq 1$. Compute the coefficients of the best linear predictor $Y = \alpha + \beta X + e$. Compute the conditional expectation $m(x) = E[Y|X = x]$. Are the best linear predictor and conditional expectation different?

Best Linear Predictor (BLP):

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy = \int_0^1 \frac{3}{2}(x^2 + y^2) dy = \frac{3}{2} \left[x^2 y + \frac{y^3}{3} \right]_{y=0}^1 = \frac{3}{2}x^2 + \frac{1}{2}$$

$$E[X] = \int_0^1 x \left(\frac{3}{2}x^2 + \frac{1}{2} \right) dx = \left[\frac{3}{8}x^4 + \frac{1}{4}x^2 \right]_0^1 = \frac{5}{8}$$

$$E[X^2] = \int_0^1 x^2 \left(\frac{3}{2}x^2 + \frac{1}{2} \right) dx = \left[\frac{3}{10}x^5 + \frac{1}{6}x^3 \right]_0^1 = \frac{7}{15}$$

Since the joint distribution is symmetric, $E[Y] = \frac{5}{8}$ and $E[Y^2] = \frac{7}{15}$.

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy \frac{3}{2}(x^2 + y^2) dx dy \\ &= \frac{3}{2} \int_0^1 \int_0^1 x^3 y + xy^3 dx dy \\ &= \frac{3}{2} \int_0^1 \left[\frac{x^4 y}{4} + \frac{x^2 y^3}{2} \right]_{x=0}^1 dy \\ &= \frac{3}{2} \int_0^1 \frac{y}{4} + \frac{y^3}{2} dy \\ &= \frac{3}{2} \left[\frac{y^2}{8} + \frac{y^4}{8} \right]_0^1 \\ &= \frac{3}{8} \end{aligned}$$

$$\begin{aligned} S(\alpha, \beta) &= E[(Y - \alpha - \beta X)^2] \\ &= E[Y^2 - \alpha Y - \beta XY - \alpha Y + \alpha^2 + \alpha\beta X - \beta XY + \alpha\beta X + \beta^2 X^2] \\ &= E[Y^2] - \alpha E[Y] - \beta E[XY] - \alpha E[Y] + \alpha^2 + \alpha\beta E[X] - \beta E[XY] + \alpha\beta E[X] + \beta^2 E[X^2] \\ &= \frac{7}{15} - \frac{5}{8}\alpha - \frac{3}{8}\beta - \frac{5}{8}\alpha + \alpha^2 + \frac{5}{8}\alpha\beta - \frac{3}{8}\beta + \frac{5}{8}\alpha\beta + \frac{7}{15}\beta^2 \\ &= \frac{7}{15} - \frac{5}{4}\alpha - \frac{3}{4}\beta + \alpha^2 + \frac{5}{4}\alpha\beta + \frac{7}{15}\beta^2 \end{aligned}$$

FOC $[\alpha]$:

$$0 = -\frac{5}{4} + 2\alpha + \frac{5}{4}\beta \implies \beta = 1 - \frac{8}{5}\alpha$$

FOC $[\beta]$:


$$0 = -\frac{3}{4} + \frac{5}{4}\alpha + \frac{14}{15}\beta \implies 45 = 75\alpha + 56\beta$$

$$45 = 75\alpha + 56\left(1 - \frac{8}{5}\alpha\right) \implies \alpha = \frac{55}{73} \implies \beta = \frac{-15}{73}$$

Conditional Expectation Function (CEF):

$$f_{Y|X}(y|x) = \frac{f(x, y)}{f_X(x)} = \frac{\frac{3}{2}(x^2 + y^2)}{\frac{3}{2}x^2 + \frac{1}{2}} = \frac{3x^2 + 3y^2}{3x^2 + 1}$$

$$m(x) = E[Y|X = x] = \int_{-\infty}^{\infty} y f_{Y|X}(y|x) dy = \int_0^1 \frac{3x^2 y + 3y^3}{3x^2 + 1} dy = \left[\frac{\frac{3}{2}x^2 y^2 + \frac{3}{4}y^4}{3x^2 + 1} \right]_{y=0}^1 = \frac{6x^2 + 3}{12x^2 + 4}$$



8. 4.1 - 4.6

4.1 For some integer k , set $\mu_k = E[Y^k]$.

(a) Construct an estimator $\hat{\mu}_k$ for μ_k .

$$\hat{\mu}_k := \frac{1}{n} \sum_{i=1}^n Y_i^k$$

(b) Show that $\hat{\mu}_k$ is unbiased for μ_k .

$$E[\hat{\mu}_k] = E\left[\frac{1}{n} \sum_{i=1}^n Y_i^k\right] = \frac{1}{n} \sum_{i=1}^n E[Y_i^k] = \frac{1}{n} \sum_{i=1}^n \mu_k = \mu_k$$

(c) Calculate the variance of $\hat{\mu}_k$, say $\text{var}[\hat{\mu}_k]$. What assumption is needed for $\text{var}[\hat{\mu}_k]$ to be finite?

We need to assume that $|\mu_{2k}| < \infty$ for $\text{var}[\hat{\mu}_k]$ to be finite:

$$\text{var}[\hat{\mu}_k] = \text{var}\left[\frac{1}{n} \sum_{i=1}^n Y_i^k\right] = \frac{1}{n^2} \sum_{i=1}^n \text{var}[Y_i^k] = \frac{1}{n^2} \sum_{i=1}^n (E[Y_i^{2k}] - E[Y_i^k]^2) = \frac{1}{n^2} \sum_{i=1}^n (\mu_{2k} - \mu_k^2) = \frac{\mu_{2k} - \mu_k^2}{n}$$

(d) Propose an estimator of $\text{var}[\hat{\mu}_k]$.

$$\frac{\hat{\mu}_{2k} - \hat{\mu}_k^2}{n} = \frac{n^{-1} \sum_{i=1}^n Y_i^{2k} - (n^{-1} \sum_{i=1}^n Y_i^k)^2}{n}$$

4.2 Calculate $E[(\bar{Y} - \mu)^3]$, the skewness of \bar{y} . Under what conditions is it zero?

$$\begin{aligned}
E[(\bar{Y} - \mu)^3] &= E\left[\left(\left(\frac{1}{n} \sum_{i=1}^n Y_i\right) - \mu\right)^3\right] \\
&= E\left[\left(\frac{1}{n} \left(\sum_{i=1}^n Y_i - n\mu\right)\right)^3\right] \\
&= E\left[\left(\frac{1}{n} \left(\sum_{i=1}^n (Y_i - \mu)\right)\right)^3\right] \\
&= \frac{1}{n^3} E\left[\left(\sum_{i=1}^n (Y_i - \mu)\right)^3\right] \\
&= \frac{1}{n^3} E\left[\sum_{i=1}^n (Y_i - \mu)^3 + 3 \sum_{i=1}^n \sum_{j=1; j \neq i}^n (Y_i - \mu)^2 (Y_j - \mu) + \sum_{i=1}^n \sum_{j=1; j \neq i}^n \sum_{k=1; k \neq i; k \neq j}^n (Y_i - \mu)(Y_j - \mu)(Y_k - \mu)\right] \\
&= \frac{1}{n^3} \left(\sum_{i=1}^n E[(Y_i - \mu)^3] + 3 \sum_{i=1}^n \sum_{j=1; j \neq i}^n E[(Y_i - \mu)^2] E[Y_j - \mu] \right. \\
&\quad \left. + \sum_{i=1}^n \sum_{j=1; j \neq i}^n \sum_{k=1; k \neq i; k \neq j}^n E[Y_i - \mu] E[Y_j - \mu] E[Y_k - \mu] \right) \\
&= \frac{1}{n^3} \left(\sum_{i=1}^n E[(Y_i - \mu)^3] \right) \\
&= \frac{1}{n^3} \left(n E[(Y_i - \mu)^3] \right) \\
&= \frac{E[(Y_i - \mu)^3]}{n^2}
\end{aligned}$$

The skewness of \bar{y} is zero if the skewness of Y_i is zero ($E[(Y_i - \mu)^3] = 0$). The skewness of \bar{y} approaches zero as n gets large.

4.3 Explain the difference between \bar{Y} and μ . Explain the difference between $n^{-1} \sum_{i=1}^n X_i X_i'$ and $E[X_i X_i']$.

The difference between \bar{Y} and μ is \bar{Y} is a statement about a sample and μ is a statement about a population. Namely, \bar{Y} is the sample mean and μ is the population mean. Similarly, $n^{-1} \sum_{i=1}^n X_i X_i'$ is the sample variance and $E[X_i X_i']$ is the population variance.

4.4 True or False. If $Y_i = X_i\beta + e_i$, $X_i \in \mathbb{R}$, $E[e_i|X_i] = 0$, and \hat{e}_i is the OLS residual from the regression of Y_i on X_i , then $\sum_{i=1}^n X_i^2 \hat{e}_i = 0$.

False. Counter example with simulated data:

```
beta <- 2
x <- runif(n = 100)
e <- rnorm(n=100)

y <- x * beta + e

beta_hat <- as.numeric((y %*% x)/(x %*% x))
print(beta_hat)

## [1] 2.118199
e_hat <- y - x * beta_hat
print(sum(x^2 %*% e_hat))

## [1] -0.7489966
```

4.5 Prove (4.15) and (4.16).

$$(4.15) \ E[\hat{\beta}|X] = \beta$$

Since $E[Y|X] = E[X\beta + e|X] = E[X\beta|X] + E[e|X] = X\beta$,

$$E[\hat{\beta}|X] = E[(X'X)^{-1}X'Y|X] = (X'X)^{-1}X'E[Y|X] = (X'X)^{-1}X'X\beta = \beta$$

$$(4.16) \ \text{var}[\hat{\beta}|X] = (X'X)^{-1}(X'\Omega X)(X'X)^{-1}$$

Since $\text{var}[Y|X] = \text{var}[X\beta + e|X] = \text{var}[e|X] = \Omega$,

$$\begin{aligned} \text{var}[\hat{\beta}|X] &= E[(\hat{\beta} - \beta)(\hat{\beta} - \beta)'|X] \\ &= E[((X'X)^{-1}X'e)((X'X)^{-1}X'e)'|X] \\ &= E[(X'X)^{-1}(X'ee'X)(X'X)^{-1}|X] \\ &= (X'X)^{-1}(X'E[ee'|X]X)(X'X)^{-1} \\ &= (X'X)^{-1}(X'\text{var}[e|X]X)(X'X)^{-1} \\ &= (X'X)^{-1}(X'\Omega X)(X'X)^{-1} \end{aligned}$$

4.6 Prove Generalized Gauss-Markov Theorem (Theorem 4.5): In the linear regression model (Assumption 4.2) and $\Omega > 0$, if $\tilde{\beta}$ is a linear unbiased estimator of β then $\text{var}[\tilde{\beta}|X] \geq (X'\Omega^{-1}X)^{-1}$

Let A be any $n \times k$ function of X of such that $A'X = I_k$. The estimator $\tilde{\beta} = A'Y$ is unbiased: $E[\tilde{\beta}|X] = E[A'Y|X] = E[A'(X\beta + e)|X] = E[A'X\beta|X] + E[A'e|X] = \beta$. The variance of $\tilde{\beta}$ is $\text{var}[\tilde{\beta}|X] = A'\Omega A$