

Two questions

1. Correlation between ITW and topology of the eye?
2. Can we group BT samples based on PH that correspond with size?
 - Persistent homology is a good choice, measures shape not size.
 - H_1 encodes loops in volume, H_2 encodes voids in volume.

- Two groups, one consisting of *Bombus terrestris* and one of others

- Two groups, one consisting of *Bombus terrestris* and one of others
- We compute *persistent homology* for the cornea of each sample (less noisy)

- Two groups, one consisting of *Bombus terrestris* and one of others
- We compute *persistent homology* for the cornea of each sample (less noisy)
- We create a distance matrix that tells us how close each sample is to the other (topologically, ITW)

- Two groups, one consisting of *Bombus terrestris* and one of others
- We compute *persistent homology* for the cornea of each sample (less noisy)
- We create a distance matrix that tells us how close each sample is to the other (topologically, ITW)
- Mantel test each group separately with PH matrix and ITW matrix


- Two groups, one consisting of *Bombus terrestris* and one of others
- We compute *persistent homology* for the cornea of each sample (less noisy)
- We create a distance matrix that tells us how close each sample is to the other (topologically, ITW)
- Mantel test each group separately with PH matrix and ITW matrix
- Cluster *Bombus terrestris* based on best metric from Mantel test

Mantel test results

Group	Metric	$H_1\rho$	H_1 p-value	$H_2\rho$	H_2 p-value
<i>BT</i>	Bottleneck	-0.069	0.84	0.86	0.0083
<i>BT</i>	Wasserstein	0.59	0.053	0.49	0.080
<i>Others</i>	Bottleneck	0.22	0.030	0.33	0.0073
<i>Others</i>	Wasserstein	0.23	0.023	0.26	0.013

H_2 with bottleneck distance shows significant results for *Bombus terrestris* and the other group of insects.

Clustering on H_2 (bottleneck)



- Supports the hypothesis that ITW and topology of eye are related
- Other dimensions/metrics were not as correlated, but do we want strong correlation?