

Chapter 4

Functions

Discrete Structures for Computing on September 14, 2017

Nguyen An Khuong, Tran Tuan Anh, Le Hong Trang
Faculty of Computer Science and Engineering
University of Technology - VNUHCM
nakhuong@hcmut.edu.vn



Contents

① Functions

② One-to-one and Onto Functions

③ Sequences and Summation

④ Recursion

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto Functions

Sequences and Summation

Recursion

Course outcomes

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Course learning outcomes

L.O.1	Understanding of logic and discrete structures
	L.O.1.1 – Describe definition of propositional and predicate logic L.O.1.2 – Define basic discrete structures: set, mapping, graphs
L.O.2	Represent and model practical problems with discrete structures
	L.O.2.1 – Logically describe some problems arising in Computing
	L.O.2.2 – Use proving methods: direct, contrapositive, induction L.O.2.3 – Explain problem modeling using discrete structures
L.O.3	Understanding of basic probability and random variables
	L.O.3.1 – Define basic probability theory L.O.3.2 – Explain discrete random variables
L.O.4	Compute quantities of discrete structures and probabilities
	L.O.4.1 – Operate (compute/ optimize) on discrete structures L.O.4.2 – Compute probabilities of various events, conditional ones, Bayes theorem

Introduction

- Each student is assigned a grade from set $\{0, 0.1, 0.2, 0.3, \dots, 9.9, 10.0\}$ at the end of semester

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Introduction

- Each student is assigned a grade from set $\{0, 0.1, 0.2, 0.3, \dots, 9.9, 10.0\}$ at the end of semester
- Function is extremely important in mathematics and computer science



Introduction

- Each student is assigned a grade from set $\{0, 0.1, 0.2, 0.3, \dots, 9.9, 10.0\}$ at the end of semester
- Function is extremely important in mathematics and computer science
 - linear, polynomial, exponential, logarithmic,...

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Introduction

- Each student is assigned a grade from set $\{0, 0.1, 0.2, 0.3, \dots, 9.9, 10.0\}$ at the end of semester
- Function is extremely important in mathematics and computer science
 - linear, polynomial, exponential, logarithmic,...
- Don't worry! For discrete mathematics, we need to understand functions at a basic set theoretic level

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$



Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f



Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f
- B : **codomain** (*miền giá trị*) of f



Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f
- B : **codomain** (*miền giá trị*) of f
- For each $a \in A$, if $f(a) = b$



Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f
- B : **codomain** (*miền giá trị*) of f
- For each $a \in A$, if $f(a) = b$
 - b is an **image** (*ảnh*) of a



Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f
- B : **codomain** (*miền giá trị*) of f
- For each $a \in A$, if $f(a) = b$
 - b is an **image** (*ảnh*) of a
 - a is **pre-image** (*ngược ảnh*) of $f(a)$



Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f
- B : **codomain** (*miền giá trị*) of f
- For each $a \in A$, if $f(a) = b$
 - b is an **image** (*ảnh*) of a
 - a is **pre-image** (*ngược ảnh*) of $f(a)$
- **Range** of f is the set of all images of elements of A



Function

Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .

- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f
- B : **codomain** (*miền giá trị*) of f
- For each $a \in A$, if $f(a) = b$
 - b is an **image** (*ảnh*) of a
 - a is **pre-image** (*ngược ảnh*) of $f(a)$
- **Range** of f is the set of all images of elements of A
- f **maps** (*ánh xạ*) A to B




Function


Definition

Let A and B be nonempty sets. A **function** f from A to B is an assignment of **exactly one** element of B to each element of A .


- $f : A \rightarrow B$
- A : **domain** (*miền xác định*) of f
- B : **codomain** (*miền giá trị*) of f
- For each $a \in A$, if $f(a) = b$
 - b is an **image** (*ảnh*) of a
 - a is **pre-image** (*ngược ảnh*) of $f(a)$
- **Range** of f is the set of all images of elements of A
- f **maps** (*ánh xạ*) A to B



Example




Example



Example:



Example




Example:

- y is an image of d



Example



Example:

- y is an image of d
- c is a pre-image of z



Example

Example

What are domain, codomain, and range of the function that assigns grades to students includes: student A: 5, B: 3.5, C: 9, D: 5.2, E: 4.9?

Example

Let $f : \mathbb{Z} \rightarrow \mathbb{Z}$ assign the the square of an integer to this integer. What is $f(x)$? Domain, codomain, range of f ?



Example

Example

What are domain, codomain, and range of the function that assigns grades to students includes: student A: 5, B: 3.5, C: 9, D: 5.2, E: 4.9?

Example

Let $f : \mathbb{Z} \rightarrow \mathbb{Z}$ assign the the square of an integer to this integer. What is $f(x)$? Domain, codomain, range of f ?

- $f(x) = x^2$
- **Domain**: set of all integers
- **Codomain**: Set of all integers
- **Range of f** : $\{0, 1, 4, 9, \dots\}$



Example

Example

What are domain, codomain, and range of the function that assigns grades to students includes: student A: 5, B: 3.5, C: 9, D: 5.2, E: 4.9?

Example

Let $f : \mathbb{Z} \rightarrow \mathbb{Z}$ assign the the square of an integer to this integer. What is $f(x)$? Domain, codomain, range of f ?

- $f(x) = x^2$
- **Domain**: set of all integers
- **Codomain**: Set of all integers
- **Range of f** : $\{0, 1, 4, 9, \dots\}$



Add and multiply real-valued functions

Definition

Let f_1 and f_2 be functions from A to \mathbb{R} . Then $f_1 + f_2$ and $f_1 f_2$ are also functions from A to \mathbb{R} defined by

$$(f_1 + f_2)(x) = f_1(x) + f_2(x)$$

$$(f_1 f_2)(x) = f_1(x) f_2(x)$$



Add and multiply real-valued functions



Definition

Let f_1 and f_2 be functions from A to \mathbb{R} . Then $f_1 + f_2$ and $f_1 f_2$ are also functions from A to \mathbb{R} defined by

$$(f_1 + f_2)(x) = f_1(x) + f_2(x)$$

$$(f_1 f_2)(x) = f_1(x) f_2(x)$$

Example

Let $f_1(x) = x^2$ and $f_2(x) = x - x^2$. What are the functions $f_1 + f_2$ and $f_1 f_2$?

$$(f_1 + f_2)(x) = f_1(x) + f_2(x) = x^2 + x - x^2 = x$$

$$(f_1 f_2)(x) = f_1(x) f_2(x) = x^2(x - x^2) = x^3 - x^4$$

Image of a subset

Definition

Let $f : A \rightarrow B$ and $S \subseteq A$. The **image** of S :

$$f(S) = \{f(s) \mid s \in S\}$$




Image of a subset

Definition

Let $f : A \rightarrow B$ and $S \subseteq A$. The **image** of S :

$$f(S) = \{f(s) \mid s \in S\}$$





Image of a subset

Definition

Let $f : A \rightarrow B$ and $S \subseteq A$. The **image** of S :

$$f(S) = \{f(s) \mid s \in S\}$$



$$f(\{a, b, c, d\}) = \{x, y, z\}$$



Definition

A function f is **one-to-one** or **injective** (*đơn ánh*) if and only if

$$\forall a \forall b (f(a) = f(b) \rightarrow a = b)$$




One-to-one



Definition

A function f is **one-to-one** or **injective** (*đơn ánh*) if and only if

$$\forall a \forall b (f(a) = f(b) \rightarrow a = b)$$




- Is $f : \mathbb{Z} \rightarrow \mathbb{Z}, f(x) = x + 1$ one-to-one?
- Is $f : \mathbb{Z} \rightarrow \mathbb{Z}, f(x) = x^2$ one-to-one?



Definition

$f : A \rightarrow B$ is **onto** or **surjective** (*toàn ánh*) if and only if

$$\forall b \in B, \exists a \in A : f(a) = b$$




- Is $f : \mathbb{Z} \rightarrow \mathbb{Z}, f(x) = x + 1$ onto?
- Is $f : \mathbb{Z} \rightarrow \mathbb{Z}, f(x) = x^2$ onto?

One-to-one and onto (bijection)

Definition

$f : A \rightarrow B$ is **bijective** (one-to-one correspondence) (*song ánh*) if and only if f is **injective** and **surjective**



- Let f be the function from $\{a, bc, d\}$ to $\{1, 2, 3, 4\}$ with $f(a) = 4$, $f(b) = 2$, $f(c) = 1$, $f(d) = 3$. Is f a bijection?



Example

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents


Functions

One-to-one and Onto
Functions



Sequences and
Summation

Recursion




Example







Example



Example



Example




Inverse function (Hàm ngược)

Definition

Let $f : A \rightarrow B$ be a **bijection** then the **inverse of f** is the function $f^{-1} : B \rightarrow A$ defined by

$$\text{if } f(a) = b \text{ then } f^{-1}(b) = a$$

A one-to-one correspondence is call **invertible** (*khả nghịch*) because we can define the inverse of this function.



Example

Example

$A = \{a, b, c\}$ and $B = \{1, 2, 3\}$ with

$$f(a) = 2 \quad f(b) = 3 \quad f(c) = 1$$

f is invertible and its inverse is

$$f^{-1}(1) = c \quad f^{-1}(2) = a \quad f^{-1}(3) = b$$



Example

Example

$A = \{a, b, c\}$ and $B = \{1, 2, 3\}$ with

$$f(a) = 2 \quad f(b) = 3 \quad f(c) = 1$$

f is invertible and its inverse is

$$f^{-1}(1) = c \quad f^{-1}(2) = a \quad f^{-1}(3) = b$$

Example

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ with $f(x) = x^2$. If f invertible?



Example

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x + 1$$



Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x + 1$$

$$f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$$

$$f^{-1}(x) = \frac{x - 1}{2}$$



Function Composition



Definition

Given a pair of functions $g : A \rightarrow B$ and $f : B \rightarrow C$. Then the **composition** (*hợp thành*) of f and g , denoted $f \circ g$ is defined by

$$f \circ g : A \rightarrow C$$

$$f \circ g(a) = f(g(a))$$

Example

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents


Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Example



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents



Functions

One-to-one and Onto
Functions




Sequences and
Summation

Recursion

Example



Example



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Graphs of Functions

Example


The graph of $f(x) = x^2$ from \mathbb{Z} to \mathbb{Z} .



Graphs of Functions

Example


The graph of $f(x) = x^2$ from \mathbb{Z} to \mathbb{Z} .



Graphs of Functions

Example

The graph of $f(x) = x^2$ from \mathbb{Z} to \mathbb{Z} .



Definition

Let f be a function from the set A to the set B . The **graph** of the function f is the set of ordered pairs $\{(a, b) \mid a \in A \text{ and } f(a) = b\}$.



Important Functions

Definition

Floor function (*hàm sàn*) of x ($\lfloor x \rfloor$): the largest integer $\leq x$
 $\lfloor \frac{1}{2} \rfloor = 0, \lfloor 3.1 \rfloor = 3, \lfloor 7 \rfloor = 7$



Important Functions

Definition

Floor function (*hàm sàn*) of x ($\lfloor x \rfloor$): the largest integer $\leq x$

$$\lfloor \frac{1}{2} \rfloor = 0, \lfloor 3.1 \rfloor = 3, \lfloor 7 \rfloor = 7$$

Ceiling function (*hàm trần*) of x ($\lceil x \rceil$): the smallest integer $\geq x$

$$\lceil \frac{1}{2} \rceil = 1, \lceil 3.1 \rceil = 4, \lceil 7 \rceil = 7$$



Important Functions

Definition

Floor function (*hàm sàn*) of x ($\lfloor x \rfloor$): the largest integer $\leq x$

$$\lfloor \frac{1}{2} \rfloor = 0, \lfloor 3.1 \rfloor = 3, \lfloor 7 \rfloor = 7$$

Ceiling function (*hàm trần*) of x ($\lceil x \rceil$): the smallest integer $\geq x$

$$\lceil \frac{1}{2} \rceil = 1, \lceil 3.1 \rceil = 4, \lceil 7 \rceil = 7$$

Bảng: Properties (n is an integer, x is a real number)

$$(1a) \quad \lfloor x \rfloor = n \text{ iff } n \leq x < n + 1$$

$$(1b) \quad \lceil x \rceil = n \text{ iff } n - 1 < x \leq n$$

$$(1c) \quad \lfloor x \rfloor = n \text{ iff } x - 1 < n \leq x$$

$$(1d) \quad \lceil x \rceil = n \text{ iff } x \leq n < x + 1$$



Important Functions

Definition

Floor function (*hàm sàn*) of x ($\lfloor x \rfloor$): the largest integer $\leq x$

$$\lfloor \frac{1}{2} \rfloor = 0, \lfloor 3.1 \rfloor = 3, \lfloor 7 \rfloor = 7$$

Ceiling function (*hàm trần*) of x ($\lceil x \rceil$): the smallest integer $\geq x$

$$\lceil \frac{1}{2} \rceil = 1, \lceil 3.1 \rceil = 4, \lceil 7 \rceil = 7$$

Bảng: Properties (n is an integer, x is a real number)

$$(1a) \quad \lfloor x \rfloor = n \text{ iff } n \leq x < n + 1$$

$$(1b) \quad \lceil x \rceil = n \text{ iff } n - 1 < x \leq n$$

$$(1c) \quad \lfloor x \rfloor = n \text{ iff } x - 1 < n \leq x$$

$$(1d) \quad \lceil x \rceil = n \text{ iff } x \leq n < x + 1$$

$$(2) \quad x - 1 < \lfloor x \rfloor \leq \lceil x \rceil < x + 1$$



Important Functions

Definition

Floor function (*hàm sàn*) of x ($\lfloor x \rfloor$): the largest integer $\leq x$

$$\lfloor \frac{1}{2} \rfloor = 0, \lfloor 3.1 \rfloor = 3, \lfloor 7 \rfloor = 7$$

Ceiling function (*hàm trần*) of x ($\lceil x \rceil$): the smallest integer $\geq x$

$$\lceil \frac{1}{2} \rceil = 1, \lceil 3.1 \rceil = 4, \lceil 7 \rceil = 7$$

Bảng: Properties (n is an integer, x is a real number)

$$(1a) \quad \lfloor x \rfloor = n \text{ iff } n \leq x < n + 1$$

$$(1b) \quad \lceil x \rceil = n \text{ iff } n - 1 < x \leq n$$

$$(1c) \quad \lfloor x \rfloor = n \text{ iff } x - 1 < n \leq x$$

$$(1d) \quad \lceil x \rceil = n \text{ iff } x \leq n < x + 1$$

$$(2) \quad x - 1 < \lfloor x \rfloor \leq \lceil x \rceil < x + 1$$

$$(3a) \quad \lfloor -x \rfloor = -\lceil x \rceil$$

$$(3b) \quad \lceil -x \rceil = -\lfloor x \rfloor$$



Important Functions

Definition

Floor function (*hàm sàn*) of x ($\lfloor x \rfloor$): the largest integer $\leq x$

$$\lfloor \frac{1}{2} \rfloor = 0, \lfloor 3.1 \rfloor = 3, \lfloor 7 \rfloor = 7$$

Ceiling function (*hàm trần*) of x ($\lceil x \rceil$): the smallest integer $\geq x$

$$\lceil \frac{1}{2} \rceil = 1, \lceil 3.1 \rceil = 4, \lceil 7 \rceil = 7$$

Bảng: Properties (n is an integer, x is a real number)

$$(1a) \quad \lfloor x \rfloor = n \text{ iff } n \leq x < n + 1$$

$$(1b) \quad \lceil x \rceil = n \text{ iff } n - 1 < x \leq n$$

$$(1c) \quad \lfloor x \rfloor = n \text{ iff } x - 1 < n \leq x$$

$$(1d) \quad \lceil x \rceil = n \text{ iff } x \leq n < x + 1$$

$$(2) \quad x - 1 < \lfloor x \rfloor \leq \lceil x \rceil < x + 1$$

$$(3a) \quad \lfloor -x \rfloor = -\lceil x \rceil$$

$$(3b) \quad \lceil -x \rceil = -\lfloor x \rfloor$$

$$(4a) \quad \lfloor x + n \rfloor = \lfloor x \rfloor + n$$

$$(4b) \quad \lceil x + n \rceil = \lceil x \rceil + n$$



Sequences

What are the rule of these sequences (*dãy*)?

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Sequences

What are the rule of these sequences (*dãy*)?

Example

1, 3, 5, 7, 9, ...

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Sequences

What are the rule of these sequences (*dãy*)?

Example

$$1, 3, 5, 7, 9, \dots \quad a_n = 2n - 1$$



Sequences

What are the rule of these sequences (*dãy*)?

Example

$$1, 3, 5, 7, 9, \dots \quad a_n = 2n - 1$$

Example

$$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$$



Sequences

What are the rule of these sequences (*dãy*)?

Example

$$1, 3, 5, 7, 9, \dots \quad a_n = 2n - 1$$

Example

$$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots \quad a_n = \frac{1}{2^{n-1}}$$



Sequences

What are the rule of these sequences (dãy)?

Example

$1, 3, 5, 7, 9, \dots$ $a_n = 2n - 1$
Arithmetic sequence (cấp số cộng)

Example

$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$ $a_n = \frac{1}{2^{n-1}}$



Sequences

What are the rule of these sequences (dãy)?

Example

$1, 3, 5, 7, 9, \dots$ $a_n = 2n - 1$
Arithmetic sequence (cấp số cộng)

Example

$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$ $a_n = \frac{1}{2^{n-1}}$
Geometric sequence (cấp số nhân)



Sequences

What are the rule of these sequences (dãy)?

Example

1, 3, 5, 7, 9, ... $a_n = 2n - 1$
Arithmetic sequence (cấp số cộng)

Example

$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$ $a_n = \frac{1}{2^{n-1}}$
Geometric sequence (cấp số nhân)

Example

$\{a_n\}$ 5, 11, 17, 23, 29, 35, 41, 47, ...



Sequences

What are the rule of these sequences (dãy)?

Example

$1, 3, 5, 7, 9, \dots$ $a_n = 2n - 1$
Arithmetic sequence (cấp số cộng)

Example

$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$ $a_n = \frac{1}{2^{n-1}}$
Geometric sequence (cấp số nhân)

Example

$\{a_n\}$ $5, 11, 17, 23, 29, 35, 41, 47, \dots$ $a_n = 6n - 1$



Sequences

What are the rule of these sequences (dãy)?

Example

1, 3, 5, 7, 9, ... $a_n = 2n - 1$
Arithmetic sequence (cấp số cộng)

Example

$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$ $a_n = \frac{1}{2^{n-1}}$
Geometric sequence (cấp số nhân)

Example

$\{a_n\}$ 5, 11, 17, 23, 29, 35, 41, 47, ... $a_n = 6n - 1$
 $\{b_n\}$ 1, 7, 25, 79, 241, 727, 2185, ...



Sequences

What are the rule of these sequences (dãy)?

Example

1, 3, 5, 7, 9, ... $a_n = 2n - 1$
Arithmetic sequence (cấp số cộng)

Example

$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$ $a_n = \frac{1}{2^{n-1}}$
Geometric sequence (cấp số nhân)

Example

$\{a_n\}$	5, 11, 17, 23, 29, 35, 41, 47, ...	$a_n = 6n - 1$
$\{b_n\}$	1, 7, 25, 79, 241, 727, 2185, ...	$b_n = 3^n - 2$



Recurrence Relations

Example

$$\{a_n\} \quad 5, 11, 17, 23, 29, 35, 41, 47, \dots$$

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Recurrence Relations

Example

$\{a_n\}$ 5, 11, 17, 23, 29, 35, 41, 47, ...

$a_n = a_{n-1} + 6$ for $n = 2, 3, 4, \dots$ and $a_1 = 5$



Recurrence Relations

Example

$\{a_n\}$ 5, 11, 17, 23, 29, 35, 41, 47, ...

$a_n = a_{n-1} + 6$ for $n = 2, 3, 4, \dots$ and $a_1 = 5$

Recurrence relations: công thức truy hồi



Recurrence Relations

Example

$\{a_n\}$ 5, 11, 17, 23, 29, 35, 41, 47, ...

$a_n = a_{n-1} + 6$ for $n = 2, 3, 4, \dots$ and $a_1 = 5$

Recurrence relations: công thức truy hồi

Definition (Fibonacci Sequence)

Initial condition: $f_0 = 0$ and $f_1 = 1$

$f_n = f_{n-1} + f_{n-2}$ for $n = 2, 3, 4, \dots$



Recurrence Relations

Example

$\{a_n\}$ 5, 11, 17, 23, 29, 35, 41, 47, ...

$a_n = a_{n-1} + 6$ for $n = 2, 3, 4, \dots$ and $a_1 = 5$

Recurrence relations: công thức truy hồi

Definition (Fibonacci Sequence)

Initial condition: $f_0 = 0$ and $f_1 = 1$

$f_n = f_{n-1} + f_{n-2}$ for $n = 2, 3, 4, \dots$

Example

Find the Fibonacci numbers f_2, f_3, f_4, f_5 and f_6



Recurrence Relations

Example

$\{a_n\}$ 5, 11, 17, 23, 29, 35, 41, 47, ...

$a_n = a_{n-1} + 6$ for $n = 2, 3, 4, \dots$ and $a_1 = 5$

Recurrence relations: công thức truy hồi

Definition (Fibonacci Sequence)

Initial condition: $f_0 = 0$ and $f_1 = 1$

$f_n = f_{n-1} + f_{n-2}$ for $n = 2, 3, 4, \dots$

Example

Find the Fibonacci numbers f_2, f_3, f_4, f_5 and f_6

$$f_2 = f_1 + f_0 = 1 + 0 = 1$$

$$f_3 = f_2 + f_1 = 1 + 1 = 2$$

$$f_4 = f_3 + f_2 = 2 + 1 = 3$$

$$f_5 = f_4 + f_3 = 3 + 2 = 5$$

$$f_6 = f_5 + f_4 = 5 + 3 = 8$$



Exercise (1)

Initial deposit: \$10,000

Interest: 11%/year, **compounded** annually (*lãi suất kép*)

After 30 years, how much do you have in your account?



Exercise (1)

Initial deposit: \$10,000

Interest: 11%/year, **compounded** annually (*lãi suất kép*)

After 30 years, how much do you have in your account?

Solution:

Let P_n be the amount in the account after n years. The sequence $\{P_n\}$ satisfies the recurrence relation

$$P_n = P_{n-1} + 0.11P_{n-1} = (1.11)P_{n-1}.$$

The initial condition is $P_0 = 10,000$



Exercise (1)

Initial deposit: \$10,000

Interest: 11%/year, **compounded** annually (*lãi suất kép*)

After 30 years, how much do you have in your account?

Solution:

Let P_n be the amount in the account after n years. The sequence $\{P_n\}$ satisfies the recurrence relation

$$P_n = P_{n-1} + 0.11P_{n-1} = (1.11)P_{n-1}.$$

The initial condition is $P_0 = 10,000$

Step 1. Solve the recurrence relation (iteration technique)

$$P_1 = (1.11)P_0$$

$$P_2 = (1.11)P_1 = (1.11)^2 P_0$$

$$P_3 = (1.11)P_2 = (1.11)^3 P_0$$

$$\vdots$$

$$P_n = (1.11)P_{n-1} = (1.11)^n P_0.$$



Exercise (1)

Initial deposit: \$10,000

Interest: 11%/year, **compounded** annually (*lãi suất kép*)

After 30 years, how much do you have in your account?

Solution:

Let P_n be the amount in the account after n years. The sequence $\{P_n\}$ satisfies the recurrence relation

$$P_n = P_{n-1} + 0.11P_{n-1} = (1.11)P_{n-1}.$$

The initial condition is $P_0 = 10,000$

Step 1. Solve the recurrence relation (iteration technique)

$$P_1 = (1.11)P_0$$

$$P_2 = (1.11)P_1 = (1.11)^2 P_0$$

$$P_3 = (1.11)P_2 = (1.11)^3 P_0$$

$$\vdots$$

$$P_n = (1.11)P_{n-1} = (1.11)^n P_0.$$

Step 2. Calculate

$$P_{30} = (1.11)^{30} 10,000 = \$228,922.97.$$



Exercise (2)

What is the 2012th number in the sequence $\{x_n\}$: 1, 2, 2, 3, 3, 3, 4, 4, 4, 4, 5, 5, 5, 5, 5, 6,...





Exercise (2)

What is the **2012th number** in the sequence $\{x_n\}$: 1, 2, 2, 3, 3, 3, 4, 4, 4, 4, 5, 5, 5, 5, 5, 6,...

Solution:

In this sequence, integer 1 appears once, the integer 2 appears twice, the integer 3 appears three times, and so on. Therefore integer n appears n times in the sequence.

We can prove that (**try it!**)

$$\sum_{i=1}^n i = 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

and can easily calculate that

$$\sum_{i=1}^{62} i = 1953$$

so the next 63 numbers (until 2016) is 63.

Therefore, 2012th number in the sequence is 63.

Theorem

If a and r are real numbers and $r \neq 0$, then

$$\sum_{j=0}^n ar^j = \begin{cases} \frac{ar^{n+1}-a}{r-1} & \text{if } r \neq 1 \\ (n+1)a & \text{if } r = 1. \end{cases}$$





Theorem

If a and r are real numbers and $r \neq 0$, then

$$\sum_{j=0}^n ar^j = \begin{cases} \frac{ar^{n+1}-a}{r-1} & \text{if } r \neq 1 \\ (n+1)a & \text{if } r = 1. \end{cases}$$

Chứng minh.

Let $S_n = \sum_{j=0}^n ar^j$.

$$\begin{aligned} rS_n &= r \sum_{j=0}^n ar^j \\ &= \sum_{j=0}^n ar^{j+1} \\ &= \sum_{k=1}^{n+1} ar^k \\ &= \left(\sum_{k=0}^n ar^k \right) + (ar^{n+1} - a) \\ &= S_n + (ar^{n+1} - a) \end{aligned}$$

Solving for S_n shows that if $r \neq 1$, then $S_n = \frac{ar^{n+1}-a}{r-1}$

If $r = 1$, then $S_n = \sum_{j=0}^n a = (n+1)a$



Recursion

Definition (Recurrence Relation)

An equation that **recursively defines** a sequence.



Recursion

Definition (Recurrence Relation)

An equation that **recursively defines** a sequence.

Definition (Recursion (đệ quy))

The act of defining an object (usually a function) in terms of that object itself.




Recursion

Definition (Recurrence Relation)

An equation that **recursively defines** a sequence.

Definition (Recursion (đệ quy))

The act of defining an object (usually a function) in terms of that object itself.



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Recursive Algorithms

Definition

An algorithm is called **recursive** if it solves a problem by reducing it to an instance of the same problem with smaller input.



Recursive Algorithms

Definition

An algorithm is called **recursive** if it solves a problem by reducing it to an instance of the same problem with smaller input.

Example

Give a recursive algorithm for computing $n!$, where n is a nonnegative integer.



Recursive Algorithms

Definition

An algorithm is called **recursive** if it solves a problem by reducing it to an instance of the same problem with smaller input.

Example

Give a recursive algorithm for computing $n!$, where n is a nonnegative integer.

Solution. We base on the recursive definition of $n!$:
 $n! = n \cdot (n - 1)!$ and $0! = 1$.



Recursive Algorithms



Definition

An algorithm is called **recursive** if it solves a problem by reducing it to an instance of the same problem with smaller input.

Example

Give a recursive algorithm for computing $n!$, where n is a nonnegative integer.

Solution. We base on the recursive definition of $n!$:
 $n! = n \cdot (n - 1)!$ and $0! = 1$.

```
procedure factorial( $n$ : nonnegative integer)
if  $n = 0$  then return 1
else return  $n \cdot \text{factorial}(n - 1)$ 
{output is  $n!$ }
```

Algorithms for Fibonacci Numbers

Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Algorithms for Fibonacci Numbers

Recursive Algorithm

```
procedure fibonacci( $n$ : nonnegative integer)
if  $n = 0$  then return 0
else if  $n = 1$  then return 1
else return fibonacci( $n-1$ ) + fibonacci( $n-2$ )
{output is fibonacci( $n$ )}
```



Algorithms for Fibonacci Numbers

Recursive Algorithm

```
procedure fibonacci( $n$ : nonnegative integer)
if  $n = 0$  then return 0
else if  $n = 1$  then return 1
else return fibonacci( $n-1$ ) + fibonacci( $n-2$ )
{output is fibonacci( $n$ )}
```

Iterative Algorithm

```
procedure iterative fibonacci( $n$ : nonnegative integer)
if  $n = 0$  then return 0
else
     $x := 0$ 
     $y := 1$ 
    for  $i := 1$  to  $n - 1$ 
         $z := x + y$ 
         $x := y$ 
         $y := z$ 
    return  $y$ 
{output is the  $n$ th Fibonacci number}
```



Tower of Hanoi

There is a tower in Hanoi that has three pegs mounted on a board together with 64 gold disks of different sizes.

Initially, these disks are placed on the first peg in order of size, with the largest on the borrom.



Tower of Hanoi

There is a tower in Hanoi that has three pegs mounted on a board together with 64 gold disks of different sizes.

Initially, these disks are placed on the first peg in order of size, with the largest on the bottom.

The rules:

- ① Move one at a time from one peg to another
- ② A disk is never placed on top of a smaller disk



Tower of Hanoi

There is a tower in Hanoi that has three pegs mounted on a board together with 64 gold disks of different sizes.

Initially, these disks are placed on the first peg in order of size, with the largest on the bottom.

The rules:

- ① Move one at a time from one peg to another
- ② A disk is never placed on top of a smaller disk

Goals: all the disks on the third peg in order of size.



Tower of Hanoi

There is a tower in Hanoi that has three pegs mounted on a board together with 64 gold disks of different sizes.

Initially, these disks are placed on the first peg in order of size, with the largest on the bottom.

The rules:


- ① Move one at a time from one peg to another
- ② A disk is never placed on top of a smaller disk

Goals: all the disks on the third peg in order of size.

The myth says that **the world will end** when they finish the puzzle.



Tower of Hanoi – 64 Discs



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Tower of Hanoi – 1 Disc



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto Functions

Sequences and Summation

Recursion


Tower of Hanoi – 1 Disc



Moved disc from peg 1 to peg 3.



Tower of Hanoi – 1 Disc



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Tower of Hanoi – 2 Discs



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents


Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion


Tower of Hanoi – 2 Discs



Moved disc from peg 1 to peg 2.



Tower of Hanoi – 2 Discs



Moved disc from peg 1 to peg 3.




Tower of Hanoi – 2 Discs



Moved disc from peg 2 to peg 3.



Tower of Hanoi – 2 Discs



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Tower of Hanoi – 3 Discs



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents


Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion


Tower of Hanoi – 3 Discs



Moved disc from peg 1 to peg 3.




Tower of Hanoi – 3 Discs



Moved disc from peg 1 to peg 2.




Tower of Hanoi – 3 Discs



Moved disc from peg 3 to peg 2.




Tower of Hanoi – 3 Discs



Moved disc from peg 1 to peg 3.



Tower of Hanoi – 3 Discs



Moved disc from peg 2 to peg 1.



Tower of Hanoi – 3 Discs



Moved disc from peg 2 to peg 3.




Tower of Hanoi – 3 Discs



Moved disc from peg 1 to peg 3.



Tower of Hanoi – 3 Discs



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents


Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Tower of Hanoi – 4 Discs



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents


Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion


Tower of Hanoi – 4 Discs



Moved disc from peg 1 to peg 2.



Tower of Hanoi – 4 Discs



Moved disc from peg 1 to peg 3.




Tower of Hanoi – 4 Discs



Moved disc from peg 2 to peg 3.




Tower of Hanoi – 4 Discs



Moved disc from peg 1 to peg 2.




Tower of Hanoi – 4 Discs



Moved disc from peg 3 to peg 1.




Tower of Hanoi – 4 Discs



Moved disc from peg 3 to peg 2.




Tower of Hanoi – 4 Discs



Moved disc from peg 1 to peg 2.




Tower of Hanoi – 4 Discs



Moved disc from peg 1 to peg 3.




Tower of Hanoi – 4 Discs



Moved disc from peg 2 to peg 3.




Tower of Hanoi – 4 Discs



Moved disc from peg 2 to peg 1.




Tower of Hanoi – 4 Discs



Moved disc from peg 3 to peg 1.




Tower of Hanoi – 4 Discs



Moved disc from peg 2 to peg 3.




Tower of Hanoi – 4 Discs



Moved disc from peg 1 to peg 2.



Tower of Hanoi – 4 Discs



Moved disc from peg 1 to peg 3.




Tower of Hanoi – 4 Discs



Moved disc from peg 2 to peg 3.



Tower of Hanoi – 4 Discs



Functions

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Functions

One-to-one and Onto
Functions

Sequences and
Summation

Recursion

Tower of Hanoi

Algorithm

```
procedure hanoi( $n$ , A, B, C)
if  $n = 1$  then move the disk from A to C
else
    call hanoi( $n - 1$ , A, C, B)
    move disk  $n$  from A to C
    call hanoi( $n - 1$ , B, A, C)
```

Recurrence Relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2H(n-1) + 1 & \text{if } n > 1. \end{cases}$$

Recurrence Solving

$$H(n) = 2^n - 1$$

If one move takes 1 second, for $n = 64$

$$\begin{aligned} 2^{64} - 1 &\approx 2 \times 10^{19} \text{ sec} \\ &\approx 500 \text{ billion years!} \end{aligned}$$

