

Chapter 5

Relations

Discrete Structures for Computing on September 20, 2017

Nguyen An Khuong, Tran Tuan Anh, Le Hong Trang
Faculty of Computer Science and Engineering
University of Technology - VNUHCM
nakhuong@hcmut.edu.vn



Contents

① Properties of Relations

② Combining Relations

③ Representing Relations

④ Closures of Relations

⑤ Types of Relations

Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Course outcomes



Contents

Properties of Relations

Combining Relations

Representing Relations


Closures of Relations

Types of Relations

Course learning outcomes

L.O.1	Understanding of logic and discrete structures
	L.O.1.1 – Describe definition of propositional and predicate logic L.O.1.2 – Define basic discrete structures: set, mapping, graphs
L.O.2	Represent and model practical problems with discrete structures
	L.O.2.1 – Logically describe some problems arising in Computing
	L.O.2.2 – Use proving methods: direct, contrapositive, induction
	L.O.2.3 – Explain problem modeling using discrete structures
L.O.3	Understanding of basic probability and random variables
	L.O.3.1 – Define basic probability theory
	L.O.3.2 – Explain discrete random variables
L.O.4	Compute quantities of discrete structures and probabilities
	L.O.4.1 – Operate (compute/ optimize) on discrete structures
	L.O.4.2 – Compute probabilities of various events, conditional ones, Bayes theorem

Introduction



Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents


Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations



Function?





Definition

Let A and B be sets. A **binary relation** (*quan hệ hai ngôi*) from a set A to a set B is a set

$$R \subseteq A \times B$$

Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations



Definition

Let A and B be sets. A **binary relation** (*quan hệ hai ngôi*) from a set A to a set B is a set

$$R \subseteq A \times B$$

- Notations:

Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations



Definition

Let A and B be sets. A **binary relation** (*quan hệ hai ngôi*) from a set A to a set B is a set

$$R \subseteq A \times B$$

- Notations:

$$(a, b) \in R \longleftrightarrow aRb$$

Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations



Definition

Let A and B be sets. A **binary relation** (*quan hệ hai ngôi*) from a set A to a set B is a set

$$R \subseteq A \times B$$

- Notations:

$$(a, b) \in R \longleftrightarrow aRb$$

- **n-ary relations?**

Contents

Properties of Relations

Combining Relations

Representing Relations


Closures of Relations

Types of Relations

Example

Example


Let $A = \{a, b, c\}$ be the set of students, $B = \{l, c, s, d\}$ be the set of the available optional courses. We can have relation R that consists of pairs (a, b) , where a is a student enrolled in course b .



Example

Example

Let $A = \{a, b, c\}$ be the set of students, $B = \{l, c, s, d\}$ be the set of the available optional courses. We can have relation R that consists of pairs (a, b) , where a is a student enrolled in course b .




$$R = \{(a, l), (a, s), (a, g), (b, c), (b, s), (b, g), (c, l), (c, g)\}$$



Example

Example

Let $A = \{a, b, c\}$ be the set of students, $B = \{l, c, s, d\}$ be the set of the available optional courses. We can have relation R that consists of pairs (a, b) , where a is a student enrolled in course b .



$$R = \{(a, l), (a, s), (a, g), (b, c), (b, s), (b, g), (c, l), (c, g)\}$$

R	l	c	s	g
a	x		x	x
b		x	x	x
c	x			x





- Is a function a relation?

Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Functions as Relations

- Is a function a relation?
- **Yes!**



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Functions as Relations

- Is a function a relation?
- **Yes!**
- $f : A \rightarrow B$



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

- Is a function a relation?

- **Yes!**

- $f : A \rightarrow B$

$$R = \{(a, b) \mid b = f(a)\}$$

Functions as Relations

- Is a relation a function?



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Functions as Relations

- Is a relation a function?
- **No**



Contents

Properties of Relations

Combining Relations


Representing Relations

Closures of Relations

Types of Relations

Functions as Relations

- Is a relation a function?
- **No**



Contents

Properties of Relations

Combining Relations


Representing Relations

Closures of Relations

Types of Relations

Functions as Relations

- Is a relation a function?
- **No**



- Relations are a **generalization** of functions



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Relations on a Set

Definition

A **relation on the set A** is a relation from A to A .



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Relations on a Set

Definition

A **relation on the set A** is a relation from A to A .

Example

Let A be the set $\{1, 2, 3, 4\}$. Which ordered pairs are in the relation $R = \{(a, b) \mid a \text{ divides } b\}$ (*a là ước số của b*)?



Relations on a Set

Definition

A **relation on the set A** is a relation from A to A .

Example

Let A be the set $\{1, 2, 3, 4\}$. Which ordered pairs are in the relation $R = \{(a, b) \mid a \text{ divides } b\}$ (a là ước số của b)?

Solution:

$$R = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}$$

R	1	2	3	4
1	x	x	x	x
2		x		x
3			x	
4				x



Properties of Relations

Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Properties of Relations



Reflexive (phản xạ)	$xRx, \forall x \in A$
-------------------------------	------------------------

Properties of Relations



Reflexive (phản xạ)	$xRx, \forall x \in A$
Symmetric (đối xứng)	$xRy \rightarrow yRx, \forall x, y \in A$

Properties of Relations



Reflexive (phản xạ)	$xRx, \forall x \in A$
Symmetric (đối xứng)	$xRy \rightarrow yRx, \forall x, y \in A$
Antisymmetric (phản đối xứng)	$(xRy \wedge yRx) \rightarrow x = y, \forall x, y \in A$

Properties of Relations



Reflexive (phản xạ)	$xRx, \forall x \in A$
Symmetric (đối xứng)	$xRy \rightarrow yRx, \forall x, y \in A$
Antisymmetric (phản đối xứng)	$(xRy \wedge yRx) \rightarrow x = y, \forall x, y \in A$
Transitive (bắc cầu)	$(xRy \wedge yRz) \rightarrow xRz, \forall x, y, z \in A$

Example

Example

Consider the following relations on $\{1, 2, 3, 4\}$:

$$R_1 = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 4), (4, 1), (4, 4)\},$$

$$R_2 = \{(1, 1), (1, 2), (2, 1)\},$$

$$R_3 = \{(1, 1), (1, 2), (1, 4), (2, 1), (2, 2), (3, 3), (4, 1), (4, 4)\},$$

$$R_4 = \{(2, 1), (3, 1), (3, 2), (4, 1), (4, 2), (4, 3)\},$$

$$R_5 = \{(3, 4)\}$$



Example

Example

Consider the following relations on $\{1, 2, 3, 4\}$:

$$R_1 = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 4), (4, 1), (4, 4)\},$$

$$R_2 = \{(1, 1), (1, 2), (2, 1)\},$$

$$R_3 = \{(1, 1), (1, 2), (1, 4), (2, 1), (2, 2), (3, 3), (4, 1), (4, 4)\},$$

$$R_4 = \{(2, 1), (3, 1), (3, 2), (4, 1), (4, 2), (4, 3)\},$$

$$R_5 = \{(3, 4)\}$$

Solution:

- Reflexive: R_3



Example

Example

Consider the following relations on $\{1, 2, 3, 4\}$:

$$R_1 = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 4), (4, 1), (4, 4)\},$$

$$R_2 = \{(1, 1), (1, 2), (2, 1)\},$$

$$R_3 = \{(1, 1), (1, 2), (1, 4), (2, 1), (2, 2), (3, 3), (4, 1), (4, 4)\},$$

$$R_4 = \{(2, 1), (3, 1), (3, 2), (4, 1), (4, 2), (4, 3)\},$$

$$R_5 = \{(3, 4)\}$$

Solution:

- Reflexive: R_3
- Symmetric: R_2, R_3



Example

Example

Consider the following relations on $\{1, 2, 3, 4\}$:

$$R_1 = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 4), (4, 1), (4, 4)\},$$

$$R_2 = \{(1, 1), (1, 2), (2, 1)\},$$

$$R_3 = \{(1, 1), (1, 2), (1, 4), (2, 1), (2, 2), (3, 3), (4, 1), (4, 4)\},$$

$$R_4 = \{(2, 1), (3, 1), (3, 2), (4, 1), (4, 2), (4, 3)\},$$

$$R_5 = \{(3, 4)\}$$

Solution:

- Reflexive: R_3
- Symmetric: R_2, R_3
- Antisymmetric: R_4, R_5



Example

Example

Consider the following relations on $\{1, 2, 3, 4\}$:

$$R_1 = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 4), (4, 1), (4, 4)\},$$

$$R_2 = \{(1, 1), (1, 2), (2, 1)\},$$

$$R_3 = \{(1, 1), (1, 2), (1, 4), (2, 1), (2, 2), (3, 3), (4, 1), (4, 4)\},$$

$$R_4 = \{(2, 1), (3, 1), (3, 2), (4, 1), (4, 2), (4, 3)\},$$

$$R_5 = \{(3, 4)\}$$

Solution:

- Reflexive: R_3
- Symmetric: R_2, R_3
- Antisymmetric: R_4, R_5
- Transitive: R_4, R_5



Example

Example

What is the properties of the **divides** (ước số) relation on the set of positive integers?



Example

Example

What is the properties of the **divides** (ước số) relation on the set of positive integers?

Solution:

- $\forall a \in \mathbb{Z}^+, a \mid a$: **reflexive**
- $1 \mid 2$, but $2 \nmid 1$: **not symmetric**
- $\exists a, b \in \mathbb{Z}^+, (a \mid b) \wedge (b \mid a) \rightarrow a = b$: **antisymmetric**
- $a \mid b \Rightarrow \exists k \in \mathbb{Z}^+, b = ak; b \mid c \Rightarrow \exists l \in \mathbb{Z}^+, c = bl$. Hence, $c = a(kl) \Rightarrow a \mid c$: **transitive**



Example



Example

What are the properties of these relations on the set of integers:

$$R_1 = \{(a, b) \mid a \leq b\}$$

$$R_2 = \{(a, b) \mid a > b\}$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\}$$

Combining Relations

Because relations from A to B are **subsets** of $A \times B$, two relations from A to B can be combined in any way two sets can be combined.



Combining Relations

Because relations from A to B are **subsets** of $A \times B$, two relations from A to B can be combined in any way two sets can be combined.

Example

Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 3, 4\}$. List the combinations of relations $R_1 = \{(1, 1), (2, 2), (3, 3)\}$ and $R_2 = \{(1, 1), (1, 2), (1, 3), (1, 4)\}$.



Combining Relations

Because relations from A to B are **subsets** of $A \times B$, two relations from A to B can be combined in any way two sets can be combined.

Example

Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 3, 4\}$. List the combinations of relations $R_1 = \{(1, 1), (2, 2), (3, 3)\}$ and $R_2 = \{(1, 1), (1, 2), (1, 3), (1, 4)\}$.

Solution: $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$ and $R_2 - R_1$.



Combining Relations

Because relations from A to B are **subsets** of $A \times B$, two relations from A to B can be combined in any way two sets can be combined.

Example

Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 3, 4\}$. List the combinations of relations $R_1 = \{(1, 1), (2, 2), (3, 3)\}$ and $R_2 = \{(1, 1), (1, 2), (1, 3), (1, 4)\}$.

Solution: $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$ and $R_2 - R_1$.

Example

Let A and B be the set of all students and the set of all courses at school, respectively. Suppose $R_1 = \{(a, b) \mid a \text{ has taken the course } b\}$ and $R_2 = \{(a, b) \mid a \text{ requires course } b \text{ to graduate}\}$. What are the relations $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 \oplus R_2$, $R_1 - R_2$, $R_2 - R_1$?



Composition of Relations

Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Composition of Relations



Definition

Let R be **relations** from A to B and S be from B to C . Then the **composite** (*hợp thành*) of S and R is

$$S \circ R = \{(a, c) \in A \times C \mid \exists b \in B (aRb \wedge bSc)\}$$

Composition of Relations



Definition

Let R be **relations** from A to B and S be from B to C . Then the **composite** (*hợp thành*) of S and R is

$$S \circ R = \{(a, c) \in A \times C \mid \exists b \in B (aRb \wedge bSc)\}$$

Example

$$R = \{(0, 0), (0, 3), (1, 2), (0, 1)\}$$

$$S = \{(0, 0), (1, 0), (2, 1), (3, 1)\}$$

Composition of Relations



Definition

Let R be **relations** from A to B and S be from B to C . Then the **composite** (*hợp thành*) of S and R is

$$S \circ R = \{(a, c) \in A \times C \mid \exists b \in B (aRb \wedge bSc)\}$$

Example

$$R = \{(0, 0), (0, 3), (1, 2), (0, 1)\}$$

$$S = \{(0, 0), (1, 0), (2, 1), (3, 1)\}$$

$$S \circ R =$$

Composition of Relations



Definition

Let R be **relations** from A to B and S be from B to C . Then the **composite** (*hợp thành*) of S and R is

$$S \circ R = \{(a, c) \in A \times C \mid \exists b \in B (aRb \wedge bSc)\}$$

Example

$$R = \{(0, 0), (0, 3), (1, 2), (0, 1)\}$$

$$S = \{(0, 0), (1, 0), (2, 1), (3, 1)\}$$

$$S \circ R = \{(0, 0), (0, 1), (1, 1)\}$$

Power of Relations

Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations



Definition

Let R be a relation on the set A . The **powers** (*lũy thừa*) $R^n, n = 1, 2, 3, \dots$ are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R.$$



Definition

Let R be a relation on the set A . The **powers** (*lũy thừa*) $R^n, n = 1, 2, 3, \dots$ are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R.$$

Example

Let $R = \{(1, 1), (2, 1), (3, 2), (4, 3)\}$. Find the powers $R^n, n = 2, 3, 4, \dots$



Definition

Let R be a relation on the set A . The **powers** (*lũy thừa*) $R^n, n = 1, 2, 3, \dots$ are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R.$$

Example

Let $R = \{(1, 1), (2, 1), (3, 2), (4, 3)\}$. Find the powers $R^n, n = 2, 3, 4, \dots$

Solution:

$$R^2 = \{(1, 1), (2, 1), (3, 1), (4, 2)\}$$



Definition

Let R be a relation on the set A . The **powers** (*lũy thừa*) $R^n, n = 1, 2, 3, \dots$ are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R.$$

Example

Let $R = \{(1, 1), (2, 1), (3, 2), (4, 3)\}$. Find the powers $R^n, n = 2, 3, 4, \dots$

Solution:

$$R^2 = \{(1, 1), (2, 1), (3, 1), (4, 2)\}$$

$$R^3 = \{(1, 1), (2, 1), (3, 1), (4, 1)\}$$



Definition

Let R be a relation on the set A . The **powers** (*lũy thừa*) $R^n, n = 1, 2, 3, \dots$ are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R.$$

Example

Let $R = \{(1, 1), (2, 1), (3, 2), (4, 3)\}$. Find the powers $R^n, n = 2, 3, 4, \dots$

Solution:

$$R^2 = \{(1, 1), (2, 1), (3, 1), (4, 2)\}$$

$$R^3 = \{(1, 1), (2, 1), (3, 1), (4, 1)\}$$

$$R^4 = \{(1, 1), (2, 1), (3, 1), (4, 1)\}$$



Definition

Let R be a relation on the set A . The **powers** (*lũy thừa*) $R^n, n = 1, 2, 3, \dots$ are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R.$$

Example

Let $R = \{(1, 1), (2, 1), (3, 2), (4, 3)\}$. Find the powers $R^n, n = 2, 3, 4, \dots$

Solution:

$$R^2 = \{(1, 1), (2, 1), (3, 1), (4, 2)\}$$

$$R^3 = \{(1, 1), (2, 1), (3, 1), (4, 1)\}$$

$$R^4 = \{(1, 1), (2, 1), (3, 1), (4, 1)\}$$

...

Representing Relations Using Matrices

Definition

Suppose R is a relation from $A = \{a_1, a_2, \dots, a_m\}$ to $B = \{b_1, b_2, \dots, b_n\}$, R can be represented by the **matrix** $\mathbf{M}_R = [m_{ij}]$, where

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$



Representing Relations Using Matrices

Definition

Suppose R is a relation from $A = \{a_1, a_2, \dots, a_m\}$ to $B = \{b_1, b_2, \dots, b_n\}$, R can be represented by the **matrix** $\mathbf{M}_R = [m_{ij}]$, where

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$

Example

R is relation from $A = \{1, 2, 3\}$ to $B = \{1, 2\}$. Let $R = \{(2, 1), (3, 1), (3, 2)\}$, the matrix for R is

$$\mathbf{M}_R = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 1 & 1 \end{bmatrix}$$



Representing Relations Using Matrices

Definition

Suppose R is a relation from $A = \{a_1, a_2, \dots, a_m\}$ to $B = \{b_1, b_2, \dots, b_n\}$, R can be represented by the **matrix** $\mathbf{M}_R = [m_{ij}]$, where

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$

Example

R is relation from $A = \{1, 2, 3\}$ to $B = \{1, 2\}$. Let $R = \{(2, 1), (3, 1), (3, 2)\}$, the matrix for R is

$$\mathbf{M}_R = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 1 & 1 \end{bmatrix}$$

Determine whether the relation has certain properties (reflexive, symmetric, antisymmetric,...)



Representing Relations Using Digraphs



Definition

Suppose R is a relation in $A = \{a_1, a_2, \dots, a_m\}$, R can be represented by the **digraph** (đồ thị có hướng) $G = (V, E)$, where

$$V = A$$
$$(a_i, a_j) \in E \text{ if } (a_i, a_j) \in R$$

Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Representing Relations Using Digraphs



Definition


Suppose R is a relation in $A = \{a_1, a_2, \dots, a_m\}$, R can be represented by the **digraph** (đồ thị có hướng) $G = (V, E)$, where

$$V = A$$
$$(a_i, a_j) \in E \text{ if } (a_i, a_j) \in R$$

Example

Given a relation on $A = \{1, 2, 3, 4\}$,
 $R = \{(1, 1), (1, 3), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (4, 1)\}$
Draw corresponding digraph.

Resulting digraph





Definition

The **closure** (*bao đóng*) of relation R with respect to **property** P is the relation S that

- i. **contains** R
- ii. **has** property P
- iii. is **contained in any** relation satisfying (i) and (ii).



Definition

The **closure** (*bao đóng*) of relation R with respect to **property** P is the relation S that

- i. **contains** R
- ii. **has** property P
- iii. is **contained in any** relation satisfying (i) and (ii).

S is the “smallest” relation satisfying (i) & (ii)

Reflexive Closure

Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Reflexive Closure



Example

Let $R = \{(a, b), (a, c), (b, d), (d, c)\}$

The **reflexive closure** of R

$\{(a, b), (a, c), (b, d), (d, c), (a, a), (b, b), (c, c), (d, d)\}$

Reflexive Closure



Example

Let $R = \{(a, b), (a, c), (b, d), (d, c)\}$

The **reflexive closure** of R

$\{(a, b), (a, c), (b, d), (d, c), (a, a), (b, b), (c, c), (d, d)\}$


$$R \cup \Delta$$

where

$$\Delta = \{(a, a) \mid a \in A\}$$

diagonal relation (*quan hệ đường chéo*).

Reflexive Closure



Symmetric Closure

Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Symmetric Closure

Example

Let $R = \{(a, b), (a, c), (b, d), (c, a), (d, e)\}$

The **symmetric closure** of R

$\{(a, b), (a, c), (b, d), (c, a), (d, e), (b, a), (d, b), (e, d)\}$



Symmetric Closure



Example

Let $R = \{(a, b), (a, c), (b, d), (c, a), (d, e)\}$

The **symmetric closure** of R

$\{(a, b), (a, c), (b, d), (c, a), (d, e), (b, a), (d, b), (e, d)\}$


$$R \cup R^{-1}$$

where

$$R^{-1} = \{(b, a) \mid (a, b) \in R\}$$

inverse relation (*quan hệ ngược*).

Symmetric Closure



Transitive Closure

Relations

Nguyen An Khuong,
Tran Tuan Anh, Le
Hong Trang



Contents

Properties of Relations

Combining Relations

Representing Relations

Closures of Relations

Types of Relations

Transitive Closure



Example

Let $R = \{(a, b), (a, c), (b, d), (d, e)\}$

The **transitive closure** of R

$\{(a, b), (a, c), (b, d), (d, e), (a, d), (b, e), (a, e)\}$

Transitive Closure



Example


Let $R = \{(a, b), (a, c), (b, d), (d, e)\}$

The **transitive closure** of R

$\{(a, b), (a, c), (b, d), (d, e), (a, d), (b, e), (a, e)\}$

$$\bigcup_{n=1}^{\infty} R^n$$

Transitive Closure



Equivalence Relations

Definition

A relation on a set A is called an **equivalence relation** (*quan hệ tương đương*) if it is **reflexive**, **symmetric** and **transitive**.



Equivalence Relations

Definition

A relation on a set A is called an **equivalence relation** (*quan hệ tương đương*) if it is **reflexive**, **symmetric** and **transitive**.

Example (1)

The relation $R = \{(a, b) | a \text{ and } b \text{ are in the same provinces}\}$ is an equivalence relation. a is **equivalent** to b and vice versa, denoted $a \sim b$.



Equivalence Relations

Definition

A relation on a set A is called an **equivalence relation** (*quan hệ tương đương*) if it is **reflexive**, **symmetric** and **transitive**.

Example (1)

The relation $R = \{(a, b) | a \text{ and } b \text{ are in the same provinces}\}$ is an equivalence relation. a is **equivalent** to b and vice versa, denoted $a \sim b$.

Example (2)

$$R = \{(a, b) \mid a = b \vee a = -b\}$$

R is an equivalence relation.



Equivalence Relations

Definition

A relation on a set A is called an **equivalence relation** (*quan hệ tương đương*) if it is **reflexive**, **symmetric** and **transitive**.

Example (1)

The relation $R = \{(a, b) | a \text{ and } b \text{ are in the same provinces}\}$ is an equivalence relation. a is **equivalent** to b and vice versa, denoted $a \sim b$.

Example (2)

$$R = \{(a, b) \mid a = b \vee a = -b\}$$

R is an equivalence relation.

Example (3)

$$R = \{(x, y) \mid |x - y| < 1\}$$

Is R an equivalence relation?



Example

Example (Congruence Modulo m - Đồng dư modulo m)

Let m be a positive integer with $m > 1$. Show that the relation

$$R = \{(a, b) \mid a \equiv b \pmod{m}\}$$

is an equivalence relation on the set of integers.





Definition

Let R be an **equivalence relation** on the set A . The set of all elements that are related to an element a of A is called the **equivalence class** (*lớp tương đương*) of a , denoted by

$$[a]_R = \{s \mid (a, s) \in R\}$$

Equivalence Classes



Definition

Let R be an **equivalence relation** on the set A . The set of all elements that are related to an element a of A is called the **equivalence class** (*lớp tương đương*) of a , denoted by

$$[a]_R = \{s \mid (a, s) \in R\}$$

Example

The equivalence class of “Thủ Đức” for the equivalence relation “in the same provinces” is { “Thủ Đức”, “Gò Vấp”, “Bình Thạnh”, “Quận 10”, ... }

Example

Example

What are the equivalence classes of 0, 1, 2, 3 for congruence modulo 4?



Example

Example

What are the equivalence classes of 0, 1, 2, 3 for congruence modulo 4?

Solution:

$$[0]_4 = \{\dots, -8, -4, 0, 4, 8, \dots\}$$

$$[1]_4 = \{\dots, -7, -3, 1, 5, 9, \dots\}$$

$$[2]_4 = \{\dots, -6, -2, 2, 6, 10, \dots\}$$

$$[3]_4 = \{\dots, -5, -1, 3, 7, 11, \dots\}$$



Equivalence Relations and Partitions



Theorem

Let R be an equivalence relation on a set A . These statements for elements a and b of A are equivalent:

- i aRb
- ii $[a] = [b]$
- iii $[a] \cap [b] \neq \emptyset$

Example 1

Example

Suppose that $S = \{1, 2, 3, 4, 5, 6\}$. The collection of sets $A_1 = \{1, 2, 3\}$, $A_2 = \{4, 5\}$, and $A_3 = \{6\}$ forms a partition of S , because these sets are disjoint and their union is S



Example 1

Example

Suppose that $S = \{1, 2, 3, 4, 5, 6\}$. The collection of sets $A_1 = \{1, 2, 3\}$, $A_2 = \{4, 5\}$, and $A_3 = \{6\}$ forms a partition of S , because these sets are disjoint and their union is S .

The equivalence classes of an equivalence relation R on a set S form a **partition** of S .



Example 1

Example


Suppose that $S = \{1, 2, 3, 4, 5, 6\}$. The collection of sets $A_1 = \{1, 2, 3\}$, $A_2 = \{4, 5\}$, and $A_3 = \{6\}$ forms a partition of S , because these sets are disjoint and their union is S .

The equivalence classes of an equivalence relation R on a set S form a **partition** of S .

Every partition of a set can be used to form an **equivalence relation**.



Example 2




Example

Divides set of all cities and towns in Vietnam into set of 64 provinces. We know that:

- there are no provinces with no cities or towns
- no city is in more than one province
- every city is accounted for



Example 2



Example

Divides set of all cities and towns in Vietnam into set of 64 provinces. We know that:


- there are no provinces with no cities or towns
- no city is in more than one province
- every city is accounted for

Definition

A **partition** of a Vietnam is a collection of non-overlapping non-empty subsets of Vietnam (provinces) that, together, make up all of Vietnam.



Relation in a Partition




- We divided based on relation

$$R = \{(a, b) | a \text{ and } b \text{ are in the same provinces}\}$$



Relation in a Partition




- We divided based on relation

$$R = \{(a, b) | a \text{ and } b \text{ are in the same provinces}\}$$

- “Thủ Đức” is related (equivalent) to “Gò Vấp”



Relation in a Partition



- We divided based on relation

$$R = \{(a, b) | a \text{ and } b \text{ are in the same provinces}\}$$

- “Thủ Đức” is related (equivalent) to “Gò Vấp”
- “Đà Lạt” is **not** related (not equivalent) to “Long Xuyên”



Partial Order Relations

- Order words such that x comes before y in the dictionary



Partial Order Relations

- Order words such that x comes before y in the dictionary
- Schedule projects such that x must be completed before y



Partial Order Relations

- Order words such that x comes before y in the dictionary
- Schedule projects such that x must be completed before y
- Order set of integers, where $x < y$



Partial Order Relations

- Order words such that x comes before y in the dictionary
- Schedule projects such that x must be completed before y
- Order set of integers, where $x < y$



Partial Order Relations

- Order words such that x comes before y in the dictionary
- Schedule projects such that x must be completed before y
- Order set of integers, where $x < y$

Definition

A relation R on a set S is called a **partial ordering** (có thứ tự bộ phận) if it is **reflexive**, **antisymmetric** and **transitive**. A set S together with a partial ordering R is called a partially ordered set, or **poset** (tập có thứ tự bộ phận), and is denoted by (S, R) or (S, \preceq) .



Partial Order Relations

- Order words such that x comes before y in the dictionary
- Schedule projects such that x must be completed before y
- Order set of integers, where $x < y$

Definition

A relation R on a set S is called a **partial ordering** (có thứ tự bộ phận) if it is **reflexive**, **antisymmetric** and **transitive**. A set S together with a partial ordering R is called a partially ordered set, or **poset** (tập có thứ tự bộ phận), and is denoted by (S, R) or (S, \preceq) .

Example

- (\mathbb{Z}, \geq) is a poset



Partial Order Relations

- Order words such that x comes before y in the dictionary
- Schedule projects such that x must be completed before y
- Order set of integers, where $x < y$

Definition


A relation R on a set S is called a **partial ordering** (có thứ tự bộ phận) if it is **reflexive**, **antisymmetric** and **transitive**. A set S together with a partial ordering R is called a partially ordered set, or **poset** (tập có thứ tự bộ phận), and is denoted by (S, R) or (S, \preceq) .

Example

- (\mathbb{Z}, \geq) is a poset
- Let S a set, $(P(S), \subseteq)$ is a poset



Example



Totally Order Relations

Example

In the poset $(\mathbb{Z}^+, |)$, 3 and 9 are **comparable** (so sánh được), because $3 \mid 9$, but 5 and 7 are not, because $5 \nmid 7$ and $7 \nmid 5$.



Totally Order Relations

Example

In the poset $(\mathbb{Z}^+, |)$, 3 and 9 are **comparable** (so sánh được), because $3 \mid 9$, but 5 and 7 are not, because $5 \nmid 7$ and $7 \nmid 5$.

→ That's why we call it **partially** ordering.



Totally Order Relations

Example

In the poset $(\mathbb{Z}^+, |)$, 3 and 9 are **comparable** (so sánh được), because $3 \mid 9$, but 5 and 7 are not, because $5 \nmid 7$ and $7 \nmid 5$.

→ That's why we call it **partially** ordering.

Definition

If (S, \preceq) is a poset and every two elements of S are comparable, S is called a **totally ordered** (có thứ tự toàn phần). A totally ordered set is also called a **chain** (dây xích).



Totally Order Relations



Example

In the poset $(\mathbb{Z}^+, |)$, 3 and 9 are **comparable** (so sánh được), because $3 \mid 9$, but 5 and 7 are not, because $5 \nmid 7$ and $7 \nmid 5$.

→ That's why we call it **partially** ordering.

Definition

If (S, \preccurlyeq) is a poset and every two elements of S are comparable, S is called a **totally ordered** (có thứ tự toàn phần). A totally ordered set is also called a **chain** (dây xích).

Example

The poset (\mathbb{Z}, \leq) is totally ordered.



Definition

- a is **maximal** (*cực đại*) in the poset (S, \preceq) if there is no $b \in S$ such that $a \prec b$.



Definition

- a is **maximal** (*cực đại*) in the poset (S, \preceq) if there is no $b \in S$ such that $a \prec b$.
- a is **minimal** (*cực tiểu*) in the poset (S, \preceq) if there is no $b \in S$ such that $b \prec a$.



Definition

- a is **maximal** (*cực đại*) in the poset (S, \preceq) if there is no $b \in S$ such that $a \prec b$.
- a is **minimal** (*cực tiểu*) in the poset (S, \preceq) if there is no $b \in S$ such that $b \prec a$.

Maximal & Minimal Elements



Definition

- a is **maximal** (*cực đại*) in the poset (S, \preceq) if there is no $b \in S$ such that $a \prec b$.
- a is **minimal** (*cực tiểu*) in the poset (S, \preceq) if there is no $b \in S$ such that $b \prec a$.

Example

Which elements of the poset $(\{2, 4, 5, 10, 12, 20, 25\}, |)$ are minimal and maximal?

Maximal & Minimal Elements




Definition

- a is **maximal** (*cực đại*) in the poset (S, \preceq) if there is no $b \in S$ such that $a \prec b$.
- a is **minimal** (*cực tiểu*) in the poset (S, \preceq) if there is no $b \in S$ such that $b \prec a$.

Example

Which elements of the poset $(\{2, 4, 5, 10, 12, 20, 25\}, |)$ are minimal and maximal?



Greatest Element & Least Element

Definition

- a is the **greatest element** (*lớn nhất*) of the poset (S, \preceq) if $b \preceq a$ for all $b \in S$.





Definition

- a is the **greatest element** (*lớn nhất*) of the poset (S, \preceq) if $b \preceq a$ for all $b \in S$.
- a is the **least element** (*nhỏ nhất*) of the poset (S, \preceq) if $a \preceq b$ for all $b \in S$.



Definition

- a is the **greatest element** (*lớn nhất*) of the poset (S, \preceq) if $b \preceq a$ for all $b \in S$.
- a is the **least element** (*nhỏ nhất*) of the poset (S, \preceq) if $a \preceq b$ for all $b \in S$.

Greatest Element & Least Element



Definition

- a is the **greatest element** (*lớn nhất*) of the poset (S, \preceq) if $b \preceq a$ for all $b \in S$.
- a is the **least element** (*nhỏ nhất*) of the poset (S, \preceq) if $a \preceq b$ for all $b \in S$.

The greatest and least element are **unique** if it exists.

Greatest Element & Least Element



Definition

- a is the **greatest element** (*lớn nhất*) of the poset (S, \preceq) if $b \preceq a$ for all $b \in S$.
- a is the **least element** (*nhỏ nhất*) of the poset (S, \preceq) if $a \preceq b$ for all $b \in S$.

The greatest and least element are **unique** if it exists.

Example

Let S be a set. In the poset $(P(S), \subseteq)$, the least element is \emptyset and the greatest element is S .

Upper Bound & Lower Bound

Definition

Let $A \subseteq (S, \preccurlyeq)$.

- If u is an element of S such that $a \preccurlyeq u$ for all elements $a \in A$, then u is called an **upper bound** (*cận trên*) of A .



Upper Bound & Lower Bound

Definition

Let $A \subseteq (S, \preccurlyeq)$.

- If u is an element of S such that $a \preccurlyeq u$ for all elements $a \in A$, then u is called an **upper bound** (*cận trên*) of A .
- If l is an element of S such that $l \preccurlyeq a$ for all elements $a \in A$, then l is called a **lower bound** (*cận dưới*) of A .



Upper Bound & Lower Bound

Definition

Let $A \subseteq (S, \preccurlyeq)$.

- If u is an element of S such that $a \preccurlyeq u$ for all elements $a \in A$, then u is called an **upper bound** (*cận trên*) of A .
- If l is an element of S such that $l \preccurlyeq a$ for all elements $a \in A$, then l is called a **lower bound** (*cận dưới*) of A .




Upper Bound & Lower Bound

Definition

Let $A \subseteq (S, \preccurlyeq)$.

- If u is an element of S such that $a \preccurlyeq u$ for all elements $a \in A$, then u is called an **upper bound** (*cận trên*) of A .
- If l is an element of S such that $l \preccurlyeq a$ for all elements $a \in A$, then l is called a **lower bound** (*cận dưới*) of A .




Upper Bound & Lower Bound

Definition

Let $A \subseteq (S, \preccurlyeq)$.

- If u is an element of S such that $a \preccurlyeq u$ for all elements $a \in A$, then u is called an **upper bound** (*cận trên*) of A .
- If l is an element of S such that $l \preccurlyeq a$ for all elements $a \in A$, then l is called a **lower bound** (*cận dưới*) of A .



Example

- Subset A does **not** have upper bound and lower bound.




Upper Bound & Lower Bound

Definition

Let $A \subseteq (S, \preccurlyeq)$.

- If u is an element of S such that $a \preccurlyeq u$ for all elements $a \in A$, then u is called an **upper bound** (*cận trên*) of A .
- If l is an element of S such that $l \preccurlyeq a$ for all elements $a \in A$, then l is called a **lower bound** (*cận dưới*) of A .



Example

- Subset A does **not** have upper bound and lower bound.
- The upper bound of B are 20, 40 and the lower bound is 2.

