Nucleated Self-Excitation: Conformational Propagation and Self-Sustained Excitation in Proteins, and a Universal Mechanism

Abstract

In protein science, nucleation-dependent polymerization and prion-type conformational propagation are two established phenomena: the former characterized by lag phases and seed-dose effects, the latter by conformational replication without nucleic acid templates. Yet the mechanism they share—the question of *what makes a conformational excitation self-sustaining*—lacks an operational criterion: "nucleation" in the protein literature is a phenomenon word (it names the existence of a lag phase), not a mechanism word. This paper derives a complete reasoning chain from the atomic level: conformational states as multiple local minima on the energy landscape → the **duality of the excited state** (a misfolded molecule is simultaneously an attenuator and an exciter) → competition of two rates (excitation transfer β against spontaneous decay γ) → crossing one (β>γ gives self-sustained propagation, a gain chain) → population coherence → macroscopic nucleation (a shape transition from soluble fluctuation to aggregate or phase-separated condensate). On this chain we give operational definitions and a three-layer criterion: mechanism (rate ratio crossing one), morphology (residence-time ratio crossing threshold), and maintenance (dual survival conditions—structural residence and replication commitment). Quantitative evidence includes the mechanistic reinterpretation of lag phase and seeding, ice as a physical fuse (four experimental strands), a two-stage carbon–protein fuse decomposition, and a contrast with autocatalytic-set theory (topological connectivity versus dynamical rate-ratio). The generalization section shows that the same criterion holds across chemical replication, ecological colonization, and epidemic spread ($R_0=\beta/\gamma$ being the established epidemiological form), with three published operational model cards as domain instances. Physical wave phenomena are outside the scope of this paper and will be treated separately.

Keywords: nucleated self-excitation; template nucleation; conformational propagation; rate ratio; amyloid aggregation; prion; self-sustainment

---

1 Introduction

1.1 Two established phenomena

In the study of protein misfolding and aggregation, nucleation-dependent polymerization is textbook-established: polymerization kinetics display a pronounced lag phase; addition of preformed fibril seeds shortens or abolishes the lag; seeds are conformation-specific—seeds of one conformation preferentially accelerate aggregation into that same conformation. Prion research supplies the stronger fact: conformational information can replicate without nucleic acids—a misfolded conformational state induces the native conformer to convert to the same misfolded state through physical contact, with complete experimental systems in mammals (transmissible spongiform encephalopathies) and fungi (yeast prion inheritance).

1.2 The gap

The shared mechanistic core of these phenomena can be posed as one question: **what makes a conformational excitation self-sustaining?** The existing vocabulary stops at two levels: the phenomenon level ("nucleation" names the lag phase—it answers *what happens*) and the phenomenological level (kinetic fits of rate constants—they answer *how fast*). What is missing is an operational criterion at the mechanism level: a physical condition for self-sustainment, a moment-criterion for ignition, and a discriminant for survival after ignition.

1.3 Route of this paper

We derive the reasoning chain from the atomic level (§2), give operational definitions and the three-layer criterion on the chain (§3), test against protein-domain quantitative evidence and fuse experiments (§4), generalize to three domains and state the universal mechanism (§5), and give testable readouts (§6). Physical wave phenomena are not discussed.

---

2 The reasoning chain from the atomic level

Each step below is established physical chemistry or a direct consequence thereof. **Nucleated self-excitation is not induced from biological phenomena; it grows out of the atomic level.**

Step 1 (conformational states). A protein possesses multiple local minima on its conformational energy landscape; each minimum corresponds to a (meta)stable conformational state, and inter-state conversion requires crossing an activation barrier. This is the standard picture of energy-landscape theory.

Step 2 (the duality of the excited state). Consider a molecule in a high-energy conformational state (e.g., a β-rich misfolded state). It is simultaneously two things:

- an **attenuator**: the high-energy state tends to return to lower energy, and cellular degradation channels (proteasome, autophagy) further remove it—spontaneous departure rate γ;
- an **exciter**: it can, through physical contact (a templating interface), induce a soluble molecule to convert into the same high-energy conformational state—excitation-transfer rate β.

The excited state is at once a loss source and a gain source. This is the atomic definition of *self-excitation*: the excitation of the system is carried by the already-excited.

Step 3 (competition of two rates). Define the rate ratio $R = \beta/\gamma$: the expected number of new molecules excited per excited molecule during its lifetime, against its expected dissipation.

Step 4 (crossing one). For $R<1$ the excitation chain decays (each generation shrinks; fluctuation). $R=1$ is critical. **For $R>1$ the excitation propagates self-sustainingly**—each excited molecule on average excites more than one successor;

the gain chain expands exponentially. The lag phase is the accumulation period before R crosses one; the seed-dose effect is the forward shift of the crossing time by the number of injected excitations.

Step 5 (population coherence → macroscopic nucleation). Molecular-scale excitation-transfer events accumulate into macroscopic ordered structure (fibrils, phase-separated condensates): the system converts from molecular fluctuation (short residence) to macroscopic structure (long residence)—**the nucleation divide**. We note the boundary honestly: R>1 is a *necessary* condition for this conversion, not a sufficient one. Spatial coherence into a supracritical nucleus additionally requires the classical-nucleation conditions (surface-energy barrier, critical nucleus size, competition among nuclei—Ostwald regimes); the rate-ratio criterion governs whether excitation is self-sustaining, not whether a given fluctuation clears the barrier. The full condition is layered: R>1 (this paper) plus barrier crossing (classical nucleation theory).

Step 6 (the cross-domain bridge). This structure—medium units excited, exciting neighbors, gain against decay, collective mode established—does not depend on the specifics of protein chemistry. It is generalized in §5 to ecology, epidemiology and phase separation; the corresponding structure for physical waves is treated in a companion paper.

---

3 Definitions

The definitions are the formal closures of the chain steps. **All four are given here in operational form for the first time (candidate-level definitions).**

Definition 1 (template excitation) (Step 2, exciter face). The mechanism by which conformational state X, through physical contact, induces a soluble molecule to convert into the same state X. Operational criteria: (i) seed conformation-specificity; (ii) measurable dose–lag relation.

Definition 2 (rate ratio) (Step 3). $R = \beta/\gamma$, excitation-transfer rate over spontaneous-decay rate—the expected number of successors per excited molecule. The expression is isomorphic to the basic reproduction number $R_0=\beta/\gamma$ of epidemiology (§5: the established epidemiological form of the same criterion). *Operationalization*: β = the bimolecular templating rate constant of the conversion interface times the available soluble-substrate concentration (s^{-1}); γ = the effective clearance rate, the sum of spontaneous conformational relaxation and degradation flux (s^{-1}); both are s^{-1} and their ratio is dimensionless. *Constancy window*: R is treated as constant over the substrate-unexhausted regime; substrate depletion (β falling as the soluble pool drains) and secondary nucleation (effective β growing with aggregate surface) are explicit boundary conditions outside the constant-R window—the branching-process prediction of §4.1 applies within the window.

Definition 3 (nucleated self-excitation) (Step 4). Template excitation propagating self-sustainingly at $R \geq 1$. **Crossing one = ignition (fuse event: any mechanism that carries R past one is a fuse).**

Definition 4 (the nucleation divide; three-layer criterion) (Step 5). The shape transition from soluble fluctuation to aggregate or phase-separated condensate:

- **Main criterion (morphological layer)**: the residence-time ratio τ/T crosses a threshold—soluble fluctuation $\tau \ll T$ (molecular excited states reside for less than the system response time), aggregate/condensate $\tau \gg T$ (macroscopic structure resides long);
- **Mechanism note (occurrence layer)**: the transition is driven by the rate ratio crossing one (fuse event);
- **Maintenance note (downstream layer)**: survival after the divide requires the **dual survival conditions**—structural residence and replication commitment—jointly. Four quadrants: both = nucleus survives; structure without commitment = silent nucleus (dormant seed); commitment without structure = reconstruction product; neither = dispersal.

The three layers compose one spectrum: R<1 (decay region, fluctuation) → crossing one (ignition) → R>1 with both conditions (survival region, macroscopic structure). The mechanism gives the cause, the morphology the reading (morphology is the integral display of mechanism), maintenance the persistence condition.

---

4 Quantitative evidence in the protein domain

4.1 Lag phase and seed dose: mechanistic reinterpretation

The two standard observables of nucleation-dependent polymerization acquire mechanism readings: the lag phase is the accumulation period while the population R rises to one (excitation chains decay each generation; the excitation number accumulates by fluctuation); the seed-dose effect is the direct forward shift of the crossing time by injected excitations. **Consistency check**: any intervention that changes β or γ (temperature, conformational stabilizers, chaperones, degradation flux, concentration) should move the lag phase in the direction dictated by R=β/γ. We flag honestly that this direction is shared by all nucleation-elongation models and is a consistency check, not a discriminating prediction.

Framework-specific quantitative prediction: under the simple branching-process approximation, the post-crossing excitation number grows as $N(t) \approx S_0\, e^{(\beta-\gamma)\,t}$ (S_0 = initial seed number). The detection time—the lag phase—therefore obeys $t_{\text{lag}} = \dfrac{1}{\beta-\gamma}\ln(N_{\text{det}}/S_0) = \dfrac{1}{\gamma\,(R-1)}\ln(N_{\text{det}}/S_0)$. The framework thus predicts: (i) the lag phase is *linear in* $\ln(S_0)$ with slope $1/(\beta-\gamma)$; (ii) interventions that change γ at fixed R move the slope by 1/γ; (iii) the slope·intercept product yields β and γ separately. This log-linearity with a mechanistically interpretable slope is not a generic consequence of nucleation-elongation phenomenology (which fits lag vs. S_0 with free exponents) and constitutes the framework's discriminating quantitative face. (a framework-level prediction, to be tested against data)

4.1b First-round data test of the quantitative prediction

The log-linearity prediction of §4.1 has been tested against published seed-titration data (digitized from original figures; sources and extraction notes in the accompanying test report). Four quantitative series across four systems:

Series	System	n	R^2 (t vs ln S_0)	R^2 (t vs S_0)	ΔAIC	β−γ (h^{-1})
D1	prion RT-QuIC (Wilham 2010)	7	0.934	0.365	−15.9	1.52
D2	IAPP, 30% DOPS (Elenbaas 2022)	4	0.998	0.800	−18.3	3.45
D3	IAPP, 70% DOPS (Elenbaas 2022)	3	0.964	—	+1.6	10.90
D4	tau K18 (Carlsson 2026)	5	0.962	0.612	−11.6	0.37

Readings: (i) the log-linear form holds across all four systems with no counterexample (3/4 series ΔAIC≤−10 strongly favoring log over raw; D3's narrow 15× dynamic range is structurally undiscriminating, not a counterexample); (ii) slope inversion gives β−γ = 0.2–10.9 h^{-1}, positive throughout as the crossing-one requirement demands; (iii) **independent cross-validation**: from Elenbaas 2022's own globally fitted rate constants (secondary-nucleation theory, $\lambda=\sqrt{(2k_+k_2m_0^2)}$) one obtains $\lambda_{theory}=2.80$ h^{-1} against our slope inversion $\lambda_{fit}=3.45$ h^{-1}—ratio 1.23 (second condition 1.74, same order and correct direction: anionic lipid ↑ → λ ↑). Honest limits: β and γ enter only as their difference (the framework's in-principle separation via slope×intercept awaits controlled single-parameter interventions); digitization precision bounds R^2 from above; open first-hand Aβ42 and yeast-prion titration series were not obtained. A fifth system (α-synuclein, Vaneyck 2023) supports log-linearity qualitatively per the authors' own statement.

4.2 Ice as a physical fuse: four experimental strands

The operational definition of a fuse (any mechanism that carries R past one) finds experimental support in the ice–water two-phase system (Trinks et al. 2005; Kanavarioti et al. 2001; Attwater et al. 2010): growing ice crystals exclude solutes, forming concentrated liquid films at the ice–water interface—reactant concentration raises β directly, carrying dilute systems (R<1) across one. Four strands of readings: RNA replication yields at low temperature (−18°C) reaching orders of magnitude above dilute controls; replication enhancement for poly(A) templates of the 400-nt class; protection against hydrolysis; compartmentalization (multiple ice crystals as independent reactors). **Ice is a fuse, not a reactant: it changes R, not the chemistry.**

4.3 The carbon–protein two-stage decomposition

The fuse concept unfolds in the origin-of-life context: amino-acid polymerization (expansion of sequence space—laying the fuel) is the first half of the fuse; the first self-catalytic replicator reaching R≥1 (e.g., autocatalytic RNA) is the second half. **The minimal definition of life under this framework is the first molecular replication cycle to cross one.**

4.4 Contrast with autocatalytic-set theory

Kauffman's autocatalytic-set theory gives a connectivity transition of the polymer-catalysis graph (catalytic closure appearing at a diversity threshold)—a topological formulation. The present mechanism gives the rate ratio crossing one—a dynamical formulation. The two are isomorphism candidates (both depict the emergence of collective self-sustainment) but differ in criterion: connectivity is a graph-theoretic (static) condition; the rate ratio is a rate (dynamic) condition, continuously tunable by temperature, concentration, intervention. **Operational distinction (narrowed)**: RAF theory itself distinguishes catalytic closure from kinetic feasibility (closure is an existence condition; realized growth requires sufficient catalytic levels—the Hordijk–Kauffman–Steel results on required levels of catalysis). We therefore do not claim the existence of connected-but-not-sustaining systems as a discriminating prediction. The narrowed contrast is: the rate ratio R is *continuously tunable* (temperature, concentration, intervention) and localizes an *ignition moment* (crossing one), whereas the graph-theoretic condition is binary and not parametrized by rates—an operability difference, not an existence difference. Artificial cross-catalytic pair experiments (fuse lit by the experimenters; self-sustained exponential amplification with hour-scale doubling) constitute the mid-station of the four quadrants—rate-ratio crossing attained, spontaneity not (the fuse was lit by the experimenters), providing a positive instance for the R criterion.

---

5 Generalization: three domains and universality

5.1 Domain invariance of the criterion

The form $R=\beta/\gamma$ is verbatim invariant across four domains: chemical replication (polymerization/decomposition), ecological colonization (birth/death— invasion biology), epidemic spread (transmission/removal—**$R_0=\beta/\gamma$, the established epidemiological name for this same mathematical quantity**), and information adoption (forwarding/forgetting). This is not analogy; it is one quantity instantiated across domains.

5.2 Three operational model cards (published; cross-referenced)

Three domain instances of this framework have been published separately as operational model cards (same delivery format, different domains): template nucleation (conformational-propagation domain), colonization nucleation (ecological domain), epidemic nucleation (spread domain). This paper supplies their unified theoretical layer; the cards supply domain-level operationalization with externally preregistered refutation clauses.

5.3 Phase separation and information nucleation (abridged)

The nucleation kinetics of biomolecular condensates (liquid–liquid phase separation) fall under the same framework (concentration crossing threshold; condensate residence). Information-domain nucleation (technology adoption, rumor spread, concept

formation) is supported by a source-checked alignment study (archived with the authors).

5.4 Statement of the universal mechanism

Nucleated self-excitation is a cross-domain isomorphic mechanism; the three-layer criterion is its unified test. The core is the duality of Step 2 (excited state = attenuator × exciter): wherever a medium admits this duality, the full structure of nucleated self-excitation follows.

---

6 Testability

Four channels of testable readouts:

1. **Rate-ratio micro-perturbation**: interventions on β/γ move lag phase and nucleation rate in the R-predicted direction—the qualitative behavioral criterion (§4.1 prediction);
2. **Connected-but-not-sustaining discriminating experiment**: constructing catalytic closure with R<1 (§4.4 operational distinction)—the fork between this framework and the connectivity criterion;
3. **Four-quadrant maintenance instances**: silent nuclei (dormant seeds—structure without commitment) and reconstruction products (commitment without structure)—seed storage/reactivation experimental faces;
4. **A silicon-domain instance in service (labeled as such)**: in a parameterized memory system, entry residence time extends with resonant access (an engineering display of structural residence × replication commitment, the half-life extension mechanism)—a non-biological instance note.

---

7 Scope note

The applicability of this framework to physical wave phenomena (electromagnetic and acoustic propagation) will be treated in a companion paper.

Appendices

Appendix A (formalization, selected): Lean assets of the nucleation line—no-presupposition items and dual-survival-condition formalizations. **Appendix B (observational origin)**: the chain originates in the following observation propositions (verbatim-anchored, archived with the authors): the universality proposition of self-excitation (from interstice layer to medium layer); “any mechanism that carries the replication-rate/decomposition-rate ratio past one is a fuse”; the dual survival conditions of nuclei. **Appendix C (glossary)**: nucleated self-excitation / template excitation / rate-ratio crossing one / fuse / dual survival conditions / silent nucleus / residence-time ratio, etc.

References

1. Harper, J. D. & Lansbury, P. T., Jr. (1997). Models of amyloid seeding in Alzheimer's disease and scrapie: mechanisms of amyloid fibril growth. *Annu. Rev. Biochem.* 66, 385–407. doi:10.1146/annurev.biochem.66.1.385
2. Prusiner, S. B. (1982). Novel proteinaceous infectious particles cause scrapie. *Science* 216(4542), 136–144. doi:10.1126/science.6801762
3. Wickner, R. B. (1994). [URE3] as an altered URE2 protein: evidence for a prion analog in *S. cerevisiae*. *Science* 264(5158), 566–569. doi:10.1126/science.7909170
4. Kauffman, S. A. (1986). Autocatalytic sets of proteins. *J. Theor. Biol.* 119(1), 1–24. doi:10.1016/S0022-5193(86)80047-9
5. Kauffman, S. A. (1993). *The Origins of Order: Self-Organization and Selection in Evolution*. Oxford University Press. doi:10.1093/oso/9780195079517.001.0001
6. Lincoln, T. A. & Joyce, G. F. (2009). Self-sustained replication of an RNA enzyme. *Science* 323(5918), 1229–1232. doi:10.1126/science.1167856
7. Trinks, H., Schröder, W. & Biebricher, C. K. (2005). Ice and the origin of life. *Orig. Life Evol. Biosph.* 35(5), 429–445. doi:10.1007/s11084-005-5009-1
8. Kanavarioti, A., Monnard, P.-A. & Deamer, D. W. (2001). Eutectic phases in ice facilitate nonenzymatic nucleic acid synthesis. *Astrobiology* 1(3), 271–281. doi:10.1089/15311070152757465
9. Attwater, J. et al. (2010). Ice as a protocellular medium for RNA replication. *Nat. Commun.* 1, 76. doi:10.1038/ncomms1076
10. Anderson, R. M. & May, R. M. (1991). *Infectious Diseases of Humans: Dynamics and Control*. Oxford University Press. doi:10.1093/oso/9780198545996.001.0001
11. Three operational model cards of this group: NMC lineage, doi:10.5281/zenodo.23613761; NMC operational model card, doi:10.5281/zenodo.23610599.
12. Wilham, J. M. et al. (2010). Rapid end-point quantitation of prion seeding activity with sensitivity comparable to bioassays. *PLoS Pathog.* 6(12), e1001217. doi:10.1371/journal.ppat.1001217
13. Elenbaas, B. O. et al. (2022). Membrane composition modulates IAPP amyloidogenesis. *Biochemistry* 61, 1229–1240. doi:10.1021/acs.biochem.2c00184
14. Carlsson, P. et al. (2026). Seed-dose titration in tau K18 aggregation. *Communications Chemistry*. doi:10.1038/s42004-026-02108-1
15. Vaneyck, T. A. et al. (2023). α-Synuclein seed titration. *J. Phys. Chem. B*. doi:10.1021/acs.jpcb.2c08326

EN v0.2 release-cleaned draft from CN v0.2 (verification round passed: 15 direct DOI checks, 4 errata corrected). ~7,000 words.