The Enantiomeric Blind Spot of Organic Geochemistry: Mirror-Image Biomarkers in Petroleum Are Structurally Invisible to the Standard Workflow

---

Abstract

Steranes and hopanes are among the most widely used biomarkers in petroleum geology and organic geochemistry. The standard analytical workflow resolves their stereochemistry only at the level of **diastereomers** — for example, the 20S/(20S+20R) ratio of C_{29} ααα-steranes as a maturity parameter. Compounds that are **enantiomers** of the standard biomarkers — mirror-image molecules with all chiral centers inverted — are structurally invisible to this workflow: on non-chiral chromatographic phases they co-elute with their conventional counterparts and produce indistinguishable mass spectra. The incompatibility is structurally general, and the recent literature confirms it in practice: reviewing enantiomeric analysis of hydrocarbon biomarkers, Leseigneur et al. (2022) note that cyclodextrin-based chiral phases exist, yet even the pristane/phytane enantiomer pair "have been resolved only by achiral columns so far in petroleum geochemistry," and list enantioselective chromatography of isoprenoid alkanes for the search of molecular biosignatures on Mars as a prospect. Sterane-scale biomarkers in geological samples have not been reached at all. Enantioselective methods are mature and remain actively developed — cyclodextrin-based chiral stationary phases have been in analytical use since the early 1990s (Menges & Armstrong, 1991; Keim et al., 1991) and are the subject of current methodological reviews (Betzenbichler et al., 2022) — but have never been applied to sterane-scale biomarkers in real geological samples; Leseigneur et al. (2022) list exactly this application as a future prospect. The result is an undiscussed methodological blind spot: the mirror-image biomarker hypothesis (Davies et al., 2009), which proposes mirror steranes as the most credible geological signature of an alternative biochemistry, has been neither tested nor falsified in the seventeen years since its publication. A documented survey of its citation trail (185 citing works; Appendix A) finds no analytical study among them — the silence is consistent with the output of an instrument that cannot see the signal. We (i) analyze the structural cause of the blind spot as a dimension analysis of the chromatography–mass spectrometry coupling; (ii) reconstruct the mismatch history of the hypothesis and the tool; and (iii) specify a minimal test — chiral GC-MS re-analysis of archived samples — requiring zero new field sampling (though non-trivial method development), with pre-registered criteria distinguishing the conventional biological baseline from a mirror-biomarker signal.

Keywords: biomarkers; steranes; enantiomers; chiral chromatography; shadow biosphere; methodological blind spot

---

1. Introduction

Biomarker stereochemistry occupies a curious position in organic geochemistry. On the one hand, stereochemistry is the discipline's most celebrated information channel: the isomerization of steranes and hopanes under thermal stress underwrites the maturity parameters that remain standard tools of petroleum systems analysis more than forty years after their introduction (Seifert & Moldowan, 1978; Peters, Walters & Moldowan, 2005). On the other hand, the stereochemical information extracted is selective by construction. Geological steranes carry multiple stereocenters, and the standard workflow resolves only one class of stereoisomeric relationship: **diastereomerism** — differences in configuration at some, but not all, chiral centers.

The complementary relationship, **enantiomerism** — complete mirror inversion at all chiral centers — is not a marginal case. Terrestrial biology builds its sterane skeletons with a single handedness, so any biology built on the opposite handedness would deposit biomarkers that are enantiomers of ours: molecules identical in every bulk property, identical in mass spectra, differing only in the sense of chirality. Davies et al. (2009) argued that such mirror biomarkers are a particularly credible geological signature of a shadow biosphere — an alternative biochemistry coexisting with, or predating, our own.

The question this paper poses is not whether mirror life exists. It is a question about a tool. Does the standard biomarker workflow — gas chromatography coupled to mass spectrometry (GC-MS) of the m/z 217 (steranes) and m/z 191 (hopanes) ions on non-chiral columns — have the resolving power to detect an enantiomeric biomarker signal if it were present? We show the answer is no, that this limitation is acknowledged in the recent literature but has never been treated as a problem, and that a minimal remedy has existed, unpublished in application, for over thirty years.

2. The blind spot: a dimension analysis of the standard workflow

2.1 What the workflow resolves

The standard biomarker analysis is a coupled instrument: a chromatographic dimension and a mass-spectrometric dimension. The chromatographic dimension (a non-chiral stationary phase, standard since the foundational work of the late 1970s) separates compounds by boiling point and molecular shape — properties shared by a molecule and its mirror image. The mass-spectrometric dimension fragments ions by mass-to-charge — again identical for enantiomers, which differ only in the sign of chiral configuration. The identified "stereochemical" information of the standard workflow therefore comes entirely from the diastereomeric component of the configuration: epimerization at C-14, C-17 and C-20 of steranes shifts retention times slightly (diastereomers differ in shape), and the ratios of the resulting peaks serve as maturity indices.

2.2 What it cannot resolve

An enantiomer of a given sterane differs from its conventional counterpart at **every** stereocenter simultaneously. Pairwise, the two molecules are diastereomer-free with

respect to each other: they are mirror images. Their retention times on a non-chiral phase are identical, and their 70 eV electron-impact mass spectra are identical. They **co-elute into a single peak** whose measured area is the sum of both enantiomers. No post-processing of the standard chromatogram can recover the split — the information was never encoded in the measurement. Formally: the measurement operator (retention dimension × m/z dimension) is invariant under exchange of the two enantiomers, so no function of its output can distinguish them; enantiomeric composition is recoverable only by coupling in a chiral interaction, under which the symmetry is broken.

This is a structural point, not an instrumental one: the blind spot is a property of the dimensionality of the coupled measurement. A detection system resolves a property only if that property maps to a distinguishable state of some detector dimension. Chirality maps to a distinguishable state only under chiral interaction (a chiral stationary phase, a chiral derivatizing agent, circularly polarized light). The standard workflow contains no chiral interaction at any stage. Nor is the lost information recoverable indirectly: the diastereomeric ratios the workflow does measure are functions of the conventional (non-mirror) epimerization system, and no combination of them constrains the abundance of a co-eluting mirror component — the mirror contribution enters only as an additive, unattributed term in the peak areas.

2.3 The limitation is acknowledged — but not treated as a gap

Leseigneur et al. (2022), reviewing enantiomeric hydrocarbon analysis, state in plain terms that cyclodextrin-based chiral phases exist, yet even the pristane/phytane pair "have been resolved only by achiral columns so far in petroleum geochemistry," and flag enantioselective applications (isoprenoid alkanes; molecular biosignatures on Mars) as prospects. The extension to sterane-scale biomarkers is not their claim but the structural argument of §§2.1–2.2: chirality maps to no detector dimension of the standard workflow, whatever the compound class — the argument that blinds the workflow to pristane/phytane enantiomers blinds it to enantiomeric steranes identically. What increases at sterane scale is the difficulty of the *remedy* — more stereocenters to resolve and elution temperatures further from standard chiral-phase operating windows (§4) — a fact about the cure, not about the disease. To our knowledge, no published work has treated this as a gap with consequences — that is, asked what signals the field is structurally unable to see as a result. The present paper is an attempt to draw that consequence.

3. A hypothesis that no tool has ever addressed

3.1 The mirror biomarker proposal

Davies et al. (2009), in a programmatic paper on signatures of a shadow biosphere, identified mirror-image steranes — the enantiomeric counterparts of conventional fossil biomarkers — as a particularly credible geological signature of alternative biochemistry: the molecules would be structurally familiar (hence analyzable by the same extraction and identification protocols), yet diagnostic of a different handedness of biochemistry.

3.2 Seventeen years without a test

The proposal is now seventeen years old. A documented survey of its citation trail (OpenAlex, 185 citing works, retrieved 2026-11-02; protocol and classification in Appendix A) finds that subsequent references occur in reviews, programmatic discussions, risk analyses and philosophy of science — and not a single analytical study: no citing work reports, or proposes to report, an enantioselective measurement of geological biomarkers. The 2024 technical assessment of risks from mirror-life research (Adamala et al., 2024, 38 co-signatories, *Science*) discusses synthesis and containment of mirror organisms at length; the geological detection question — whether a mirror biosphere has already left a record — is not engaged. The asymmetry is striking: the hypothesis concerns the *past* (a signal that would already be sitting in archived cores and oil fractions), while the mirror-life discussion concerns the *future* (organisms that do not yet exist). The past-facing version is testable with existing instrument concepts and existing samples. It has not been tested.

We emphasize what this silence does and does not show. An analysis that cannot distinguish enantiomers cannot report their absence either — so the seventeen-year silence is *consistent with* the expected output of a blind instrument. It is not, by itself, proof that instrumental blindness is the operative cause: a seventeen-year gap can also follow from a low perceived prior, funding priorities, or the programmatic character of the original proposal, which specified no detection method. What the survey does establish is narrower and sufficient for our purpose: the testable question has never been tested, and no result — positive or negative — exists anywhere in the trail. The blind-spot analysis of §2 explains how that is possible; the test of §5 is what would end the condition.

3.3 The mismatch as a category

The mirror-biomarker case instantiates a general category of scientific gap: **a signal proposed in one community's conceptual vocabulary that falls outside the resolving dimensions of the tools that community standardly deploys**. No individual decision anywhere in the history is an error — the non-chiral workflow was built to answer diastereomer-scale questions (maturity, correlation, source), and answers them superbly. The gap arises at the community level: when a new question (mirror biochemistry) is posed in terms of a property (enantiomeric handedness) that the deployed toolset does not resolve, the question remains open by default while appearing, from inside the workflow, to be closed.

4. The method exists; the application is missing

Enantioselective gas chromatography is mature and remains an active field: cyclodextrin-based chiral stationary phases — permethylated and functionalized cyclodextrins — have been in routine use for enantiomer separation since the late 1980s, with gas-chromatographic implementations already in place by 1991 (Keim, Köhnes & Meltzow, 1991; Menges & Armstrong, 1991 in HPLC mode), and the phase family is still the subject of current methodological development and review (Betzenbichler et al., 2022). Enantioselective analysis is in fact routine in *adjacent* fields — the environmental literature routinely resolves chiral pollutants

and traces their enantioselective degradation (e.g., brominated flame retardants; Badea et al., 2016) — which makes the absence specific: it is not enantioselective measurement that is missing from geoscience, but its application to geological biomarkers. What has never been demonstrated is precisely that extension — to **sterane-scale biomarkers in real geological samples** (crude oil fractions, rock extracts, archived core material). Leseigneur et al. (2022) flag the adjacent prospect (enantioselective analysis of isoprenoid alkanes for Mars biosignature search); the sterane extension goes beyond what any published work has reported.

The gap between standards and samples is real and should not be understated. Steranes are large, low-volatility, high-temperature-eluting molecules; cyclodextrin phases historically tolerate narrower temperature windows than the non-chiral polysiloxanes used in biomarker work, and the high-temperature stability of chiral phases at biomarker elution temperatures is a genuine methodological risk. This is, however, a risk profile of exactly the kind the field routinely retires for questions it cares about — and the first step is modest: **standards-level validation of enantiomer separation for C_{27}–C_{29} steranes**, which has, remarkably, not been published even at the standard level for the full sterane suite. (Menges & Armstrong, 1991 resolved related chiral hydrocarbons; a dedicated sterane-suite demonstration remains to be reported.)

5. The minimal test

The structural point of §2 implies that the mirror-biomarker question can be closed with **zero new field sampling**: the relevant extracts and oil fractions already exist in departmental and survey archives, in the very vials that produced the conventional biomarker datasets. Zero new sampling is not zero new work — the standards validation of (i), chiral-phase high-temperature stability, and confirmation of archive access are real method development; the claim is only that no new geological material is required. A minimal test program has four components.

(i) Standards validation. Demonstrate baseline separation (or quantify co-elution) of enantiomeric C_{27}–C_{29} sterane standards on current-generation cyclodextrin or replacement chiral phases at biomarker elution temperatures. This step calibrates what "resolution" means in practice, fixes the detection limit at the mirror retention position, and is a precondition for interpreting any sample result.

(ii) Sample re-analysis: a two-signal readout. Re-run archived biomarker fractions on the validated chiral phase. The readout must keep two signals separate, because they answer different questions:

- *Signal A — the enantiomeric composition of the conventional sterane itself.* The conventional biological configuration is expected to dominate at low maturity (biology's handedness), drifting toward the racemic equilibrium under thermal maturation. This signal is partly known physics, and serves as the in-run baseline and maturity cross-check.
- *Signal B — the mirror component.* The mirror-image sterane is a distinct compound on a chiral phase; its abundance relative to Signal A is the shadow-biosphere quantity. The discovery case is an excess at the mirror position above

the maturation-predicted baseline of Signal A — a peak that the non-chiral trace cannot, even in principle, have shown.

Three outcomes, read on both signals jointly: (a) mirror component below the validated detection limit at all sterane windows → the quantitative upper bound of (iv); (b) Signal A racemic and no distinct mirror peak → maturation equilibration or abiotic origin (the two are not separable by this measurement alone, and the result must be reported with that ambiguity explicit); (c) reproducible excess at the mirror position → the discovery case, which proceeds directly to the protocol of (iii).

(iii) Pre-registered criteria and contamination control. Because a positive result would be extraordinary, the decision criteria should be fixed before sample analysis: retention-index matching on the chiral phase, co-injection with standards where available, and an explicit contamination protocol. We note the field's recent experience with an extraordinary life-claim that dissolved under contamination skepticism (the GFAJ-1 episode): an enantiomeric excess claim will face the same structural skepticism, and the analysis design should anticipate it — contamination by conventional (non-mirror) biomarkers cannot mimic a mirror signal, which is an epistemic advantage of this particular test: the contamination direction is one-sided.

(iv) The value of a negative. Most probably, the test returns an upper bound: mirror steranes constitute less than D% of the total (conventional + mirror) peak area at a given sterane elution window, where D is the detection-limit fraction fixed by the validated chiral-phase signal-to-noise at the mirror retention position (standards validation, (i)). Even this is a new scientific quantity — the first published constraint on mirror biomarker abundance in the geological record — and it converts a seventeen-year undiscussed gap into a bounded question. In a field where the alternative-biochemistry hypothesis currently has no instrumental constraint at all, a well-designed upper bound is progress.

A side observation sharpens the asymmetry: the Mars Organic Molecule Analyzer (MOMA) instrument on the ExoMars rover carries a chiral GC column precisely because enantiomeric excesses are regarded as a strong biosignature on Mars. The Earth case differs in an ironic direction: the samples are here, the instrument concepts are here, and the biomarker community's standard workflow — for the structural reasons of §2 — has simply never pointed a chiral phase at its own archives.

6. Discussion: blind spots of this general form

The enantiomeric blind spot is one instance of a general pattern worth naming: **the resolving dimensions of a mature workflow are set by the questions that built it**. Signals defined along dimensions that the founding questions did not include are not rejected by the community — they are rendered invisible at the instrument level, which is stronger and quieter: no data point ever contradicts the status quo, because no data point could. The standard remedy is cheap by comparison to the usual costs of science: an audit of which dimensions the current instruments do not encode, cross-referenced against signals proposed in adjacent literatures (here: astrobiology's shadow-biosphere program) that live exactly on those dimensions.

The mirror-biomarker case has an unusual feature that makes it a clean test bed for

the general pattern: the proposed signal is *one exact mirror step away* from the field's most-measured molecules. Nothing about the extraction, handling, or identification changes; a single column swap in a single re-analysis converts an invisible dimension into a measured one. Few structural blind spots in science come with so small a turning radius. We suggest the turn is worth making.

Data availability

The citation-trail survey of Appendix A was retrieved from OpenAlex on 2026-11-02 (work identifier W2088537647; 185 citing works); the classification protocol is stated in Appendix A and the raw list is available from the corresponding author.

Appendix A: Citation-trail survey of Davies et al. (2009)

Source: OpenAlex, works citing "Signatures of a Shadow Biosphere" (Davies et al., 2009; doi:10.1089/ast.2008.0251), retrieved 2026-11-02. **Total**: 185 citing works. **Protocol**: (1) full listing retrieved via the OpenAlex API (filter: cites this work); (2) classification by work type and keyword scan of titles and venues for analytical-methods terms (chromatograph*, enantioselect*, enantiom*, chiral, sterane, hopane, GC-MS, mass spectrometry, in combination with petroleum/oil/crude/geochem*) and earth-science terms (geochem*, petroleum, crude oil, sediment, core, bitumen, kerogen).

Results. By type: articles 105, book chapters 54, reviews 7, preprints 7, paratexts 5, other 3, books 2, conference papers 2. By classification: the keyword scan returns three candidate hits, none an analytical geochemistry application: a theoretical origin-of-life study of chiral polymerization (2012); a SPIE conference abstract on alternate-chirality utilization in microbial communities (2010); and the present authors' own preprint of this paper (Zenodo, 2026). The single earth-science-venue hit is a methodological-caution paper on ancient DNA in marine sediments (2019), not an enantioselective biomarker measurement. All 29 citing works from 2022 onward are reviews, programmatic papers, mission-concept studies or philosophy-of-science discussions. **Conclusion**: within the retrievable citation trail, no work reports or proposes an enantioselective analytical measurement of geological biomarkers; the negative existence claim of §3.2 is documented to the limit of this protocol (which cannot exclude unindexed or industrial-internal work — a limit we state explicitly).

References

- Adamala, K. P., et al. (2024). Confronting risks of mirror life. *Science* 386(6728), 1351–1353. doi:10.1126/science.ads9158 (38 co-signatories)
- Badea, S.-L., Niculescu, V.-C., Ionete, R.-E. & Eljarrat, E. (2016). Advances in enantioselective analysis of chiral brominated flame retardants. *Science of the Total Environment* 566–567, 1120–1130. doi:10.1016/j.scitotenv.2016.05.148
- Betzenbichler, G., Huber, L., Kräh, S., Morkos, M.-L. K., Siegle, A. F. & Trapp, O. (2022). Chiral stationary phases and applications in gas chromatography. *Chirality* 34(5), 732–759. doi:10.1002/chir.23427

- Davies, P. C. W., Benner, S. A., Cleland, C. E., Lineweaver, C. H., McKay, C. P. & Wolfe-Simon, F. (2009). Signatures of a shadow biosphere. *Astrobiology* 9(2), 241–249. doi:10.1089/ast.2008.0251
- Keim, W., Köhnes, A. & Meltzow, W. (1991). Enantiomer separation by gas chromatography on cyclodextrin chiral stationary phases. *Journal of High Resolution Chromatography* 14, 507–529. doi:10.1002/jhrc.1240140803
- Leseigneur, G., Filippi, J.-J., Baldovini, N. & Meierhenrich, U. J. (2022). Absolute configuration of aliphatic hydrocarbon enantiomers identified by gas chromatography: theorized application for isoprenoid alkanes and the search of molecular biosignatures on Mars. *Symmetry* 14(2), 326. doi:10.3390/sym14020326
- Mackenzie, A., Brassell, S., Eglinton, G. & Maxwell, J. (1982). Chemical fossils: the geological fate of steroids. *Science* 217, 491–504.
- Menges, R. A. & Armstrong, D. W. (1991). Chiral separations using native and functionalized cyclodextrin-bonded stationary phases in high-pressure liquid chromatography. In *Chiral Separations* (ACS Symposium Series 471), 67–100. doi:10.1021/bk-1991-0471.ch004
- Peters, K. E., Walters, C. C. & Moldowan, J. M. (2005). *The Biomarker Guide*, 2nd ed. Cambridge University Press.
- Seifert, W. K. & Moldowan, J. M. (1978). Applications of steranes, terpanes and monoaromatics to the maturation, migration and source of crude oils. *Geochimica et Cosmochimica Acta* 42, 77–95.