

Real Analysis Exercise Sheet

1. Let X be an infinite set. For $p, q \in X$, define

$$d(p, q) = \begin{cases} 1 & \text{if } p \neq q, \\ 0 & \text{if } p = q. \end{cases}$$

- (a) Prove that this is a metric.
 - (b) Which subsets of the resulting metric space are open?
 - (c) Which are closed?
 - (d) Which are compact? (Definition. A metric space (X, d) is called compact if every sequence in X has a convergent subsequence whose limit lies in X ; a subset $Y \subset X$ is called compact if the subspace $(Y, d|_{Y \times Y})$ is compact).
 - (e) Prove that a sequence $\{x_n\} \subset X$ converges to $x \in X$ in the discrete metric if and only if there exists $N \in \mathbb{N}$ such that $x_n = x$ for all $n \geq N$.
2. Let (X, d) be a metric space. Show that $d'(x, y) = \sqrt{d(x, y)}$ is also a metric on X , and that the open sets for d' are the same as the open sets for d .
3. Consider \mathbb{R} with the standard metric. Let $E \subset \mathbb{R}$ be a subset which has no limit points. Show that E is at most countable.
4. Let (X, d) be a compact metric space, and $f : X \rightarrow X$ a map such that $d(f(x), f(y)) < d(x, y)$ for all $x \neq y$. Prove that there exists a point x such that $f(x) = x$. Hint: how small can $d(x, f(x))$ get? Comment: this is an example of a fixed point theorem, very popular in various sciences for showing the existence of equilibria and such.
5. Let (X, d) be a metric space.

- (a) Prove the following:

- (i) For all $x, y, z \in X$

$$|d(y, z) - d(z, x)| \leq d(x, y).$$

- (ii) For any $n \in \mathbb{N}$ and any $x_1, x_2, \dots, x_n \in X$

$$d(x_1, x_n) \leq d(x_1, x_2) + d(x_2, x_3) + \dots + d(x_{n-1}, x_n).$$

- (b) For $x \in \mathbb{R}$ and $y \in \mathbb{R}$ define

- (i) $d_1(x, y) = (x - y)^2$
- (ii) $d_2(x, y) = \sqrt{|x - y|}$
- (iii) $d_3(x, y) = |x^2 - y^2|$
- (iv) $d_4(x, y) = |x - 2y|$

Determine, for each of these, whether it is a metric or not.

6. (a) In a metric space (X, d) , prove that if a sequence (x_n) converges, then its limit is unique. That is, show that if $x_n \rightarrow x$ and $x_n \rightarrow y$, then $x = y$.
- (b) Prove that a subset E of a metric space (X, d) is closed if and only if

$$E = \overline{E},$$

where \overline{E} denotes the closure of E .

7. Let \mathbb{R}^2 be equipped with the Taxicab metric/Manhattan metric defined by

$$d((x_1, y_1), (x_2, y_2)) = |x_1 - x_2| + |y_1 - y_2|.$$

- (a) Prove that d is a metric.
- (b) Draw the open balls $B_1((0, 0))$, $B_2((0, 0))$, and $B_{\frac{1}{2}}((0, 0))$ in \mathbb{R}^2 .

- (c) On the same axes, sketch the corresponding Euclidean balls (under the usual metric) of the same radii.
- (d) What geometric shape do the taxicab balls form?
- (e) Are these taxicab balls open sets in the usual metric on Euclidean space? Prove that these two metrics are equivalent.
- (f) Let

$$A = \{(x, y) \in \mathbb{R}^2 : d((x, y), (1, 1)) \geq 2\}.$$

Sketch the set A. Is it closed in the taxicab metric?

- 8. (a) Let us consider the set $X = \mathbb{R} \setminus \{\sqrt{2}\}$ equipped with the usual metric $d(x, y) = |x - y|$. Is (X, d) a complete metric space? Justify your answer with appropriate reasoning.
- (b) Let \mathbb{R} be the set of real numbers with the usual metric $d(x, y) = |x - y|$.
 - (i) Show that the sequence $x_n = \frac{1}{n}$ is a Cauchy sequence in \mathbb{R} .
 - (ii) Show that the sequence $x_n = n$ is not a Cauchy sequence in \mathbb{R} .
- 9. Let $C[0, 1]$ be the set of all real-valued continuous functions on $[0, 1]$. Define:

$$d_1(f, g) = \int_0^1 |f(x) - g(x)| dx.$$

Define a sequence $\{f_n\} \subset C[0, 1]$ by:

$$f_n(x) = \begin{cases} 0, & 0 \leq x \leq \frac{1}{2} - \frac{1}{n} \\ n(x - (\frac{1}{2} - \frac{1}{n})), & \frac{1}{2} - \frac{1}{n} < x < \frac{1}{2} \\ 1, & \frac{1}{2} \leq x \leq 1 \end{cases}$$

- (a) Draw the graph of each function f_n and verify that $f_n \in C[0, 1]$ by observing that each f_n is piecewise linear and continuous.
- (b) Prove that f_n pointwise converge to f where

$$f(x) = \begin{cases} 0, & 0 \leq x < \frac{1}{2}, \\ 1, & \frac{1}{2} \leq x \leq 1. \end{cases}$$
- (c) Verify that $\{f_n\}$ is a Cauchy sequence in $(C[0, 1], d_1)$.
- (d) Is $(C[0, 1], d_1)$ a complete metric space?
- 10. Let l^2 be the set of all sequences $x = \{x_n\}$ over the real numbers \mathbb{R} such that $\sum_{n \geq 1} x_n^2$ converges. Prove that $d(x, y) = \sqrt{\sum_{n \geq 1} (x_n - y_n)^2}$ gives a metric on l^2 .
- 11. Let (X, d) be a metric space and A be a non-empty subset of X. For $x \in X$, we define

$$d(x, A) = \inf\{d(x, a) : a \in A\}.$$

- (a) Show that $d(x, A) = 0$ if and only if x lies in the closure of A.
- (b) Show that if A is compact, then $d(x, A) = d(x, a)$ for some $a \in A$.
- (c) The ϵ -neighborhood of A is defined to be

$$N(A; \epsilon) = \{x \in X : d(x, A) < \epsilon\}.$$

Prove that $N(A; \epsilon)$ is the union of the open balls $B_\epsilon(a)$ for $a \in A$.

- (d) If A is compact and U is an open set containing A, prove that there exists $\epsilon > 0$ such that

$$A \subset N(A; \epsilon) \subset U.$$

Is this fact still true if A is just a closed set?

- 12. Let (X, d) be a metric space. Then verify the following properties:

- (a) The union of any collection $\{G_\alpha\}$ of open sets is open; that is, $\bigcup_\alpha G_\alpha$ is open.
 - (b) The intersection of any collection $\{F_\alpha\}$ of closed sets is closed; that is, $\bigcap_\alpha F_\alpha$ is closed.
 - (c) The intersection of any finite collection G_1, \dots, G_n of open sets is open; that is, $\bigcap_{i=1}^n G_i$ is open.
 - (d) The union of any finite collection F_1, \dots, F_n of closed sets is closed; that is, $\bigcup_{i=1}^n F_i$ is closed.
13. In vector spaces, metrics are usually defined in terms of norms which measure the length of a vector. If V is a vector space defined over \mathbb{R} , then a norm is a function from vectors to real numbers, denoted by $\|\cdot\|$, satisfying:
- (i) $\|x\| \geq 0$ and $\|x\| = 0 \iff x = 0$,
 - (ii) For any $\lambda \in \mathbb{R}$, $\|\lambda x\| = |\lambda| \|x\|$,
 - (iii) $\|x + y\| \leq \|x\| + \|y\|$.

Prove that every norm defines a metric.

14. Let M be a metric space with metric d . Show that d_1 defined by

$$d_1(x, y) = \frac{d(x, y)}{1 + d(x, y)}$$

is also a metric on M . Observe that M itself is bounded in this metric.

15. Let A and B be two subsets of a metric space M . Recall that A° , the interior of A , is the set of interior points of A . Prove the following:

$$\text{a) } A^\circ \cup B^\circ \subseteq (A \cup B)^\circ, \quad \text{b) } A^\circ \cap B^\circ = (A \cap B)^\circ$$

Give an example of two subsets A and B of the real line such that $A^\circ \cup B^\circ \neq (A \cup B)^\circ$.

16. Let A be a subset of a metric space M . Recall that \bar{A} , the closure of A , is the union of A and its limit points. Recall that a point belongs to the boundary of A , ∂A , if every open ball centered at the point contains points of A and points of A^c , the complement of A . Prove that:

- (a) $\partial A = \bar{A} \cap \bar{A}^c$
- (b) $p \in \partial A \iff p$ is in \bar{A} , but not in A° (symbolically, $\partial A = \bar{A} \setminus A^\circ$)
- (c) ∂A is a closed set
- (d) A is closed $\iff \partial A \subseteq A$

17. Show that, in \mathbb{R}^n with the usual (Euclidean) metric, the closure of the open ball $B_R(p)$, $R > 0$, is the closed ball

$$\{q \in \mathbb{R}^n : d(p, q) \leq R\}.$$

Give an example of a metric space for which the corresponding statement is false.

18. Prove directly from the definition that the set $K \subseteq \mathbb{R}$ given by

$$K = \{0, 1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots\}$$

is compact.

19. Let K be a compact subset of a metric space M , and let $\{\mathcal{U}_\alpha\}_{\alpha \in I}$ be an open cover of K . Show that there is a positive real number δ with the property that for every $x \in K$ there is some $\alpha \in I$ with

$$B_\delta(x) \subseteq \mathcal{U}_\alpha$$

20. Let M be a non-empty set, and let d be a real-valued function of ordered pairs of elements of M satisfying

- (a) $d(x, y) = 0 \iff x = y$
- (b) $d(x, y) \leq d(x, z) + d(y, z)$.

Show that d is a metric on M .

21. Determine the boundaries of the following sets, $A \subseteq X$:

(i) $A = \mathbb{Q}, X = \mathbb{R}$

(ii) $A = \mathbb{R} \setminus \mathbb{Q}, X = \mathbb{R}$

(iii) $A = (\mathbb{Q} \times \mathbb{Q}) \cap B_R(0), X = \mathbb{R}^2$

22. Describe the interior of the Cantor set.

23. Let M be a metric space with metric d , and let d_1 be the metric defined above (in problem 2). Show that the two metric spaces $(M, d), (M, d_1)$ have the same open sets.