

# Bounding the Value of Extended Nonlocal Games

Theory Seminar

Vincent Russo

University of Waterloo

September 15, 2016

UNIVERSITY OF  
WATERLOO

IQC Institute for  
Quantum  
Computing

# Outline

Nonlocal games

Upper bounding nonlocal games


Extended nonlocal games

Upper bounding extended nonlocal games

# Nonlocal games

## Nonlocal games

A *nonlocal game* is a cooperative game played between *Alice* and *Bob* against a *referee*.



1. Question and answer sets:  $(\Sigma_A, \Sigma_B)$  and  $(\Gamma_A, \Gamma_B)$ ,
2. Distributions on question pairs:  $\pi : \Sigma_A \times \Sigma_B \rightarrow [0, 1]$ ,
3. A predicate  $V : \Gamma_A \times \Gamma_B \times \Sigma_A \times \Sigma_B \rightarrow \{0, 1\}$ , where

$$V(a, b|x, y) = \begin{cases} 1 & \text{if Alice and Bob win,} \\ 0 & \text{if Alice and Bob lose.} \end{cases}$$

# Strategies for nonlocal games

Alice and Bob could use different types of *strategies*:

- ▶ *Classical strategies*: Alice and Bob answer deterministically, determined by functions of  $f : \Sigma_A \rightarrow \Gamma_A$  and  $g : \Sigma_B \rightarrow \Gamma_B$ .
- ▶ *Quantum strategies*: Alice and Bob share a joint quantum system  $\rho \in \mathcal{D}(\mathcal{A} \otimes \mathcal{B})$  and allow their answers to be outcomes of measurements on this shared system.
- ▶ *Commuting measurement strategies*: Alice and Bob share a quantum system over a single Hilbert space  $\rho \in \mathcal{D}(\mathcal{H})$  and allow their answers to be outcomes of measurements on this system.

# Winning probabilities for Alice and Bob

For any type of strategy, the output probability distribution produced by Alice and Bob may be described by a *correlation operator*:

$$C \in L(\mathbb{R}^{\Sigma_A \times \Gamma_A}, \mathbb{R}^{\Sigma_B \times \Gamma_B}),$$

where  $C((x, a), (y, b))$  is the probability that Alice and Bob output  $(a, b)$  given questions  $(x, y)$ .

# Winning probabilities for Alice and Bob

For any type of strategy, the output probability distribution produced by Alice and Bob may be described by a *correlation operator*:

$$C \in L(\mathbb{R}^{\Sigma_A \times \Gamma_A}, \mathbb{R}^{\Sigma_B \times \Gamma_B}),$$

where  $C((x, a), (y, b))$  is the probability that Alice and Bob output  $(a, b)$  given questions  $(x, y)$ .

For a nonlocal game, the probability that Alice and Bob win for a given choice of  $x, y, a, b$  is

$$\sum_{x,y} \pi(x, y) \sum_{a,b} V(a, b|x, y) C((x, a), (y, b)).$$

# Values for nonlocal games

The *value* of a nonlocal game is the maximal winning probability for the players to win over all strategies of a specified type.

For a nonlocal game,  $G$ , we denote the classical and quantum values as

- ▶ Classical value:  $\omega(G)$ ,
- ▶ Commuting measurement value:  $\omega_c(G)$ ,
- ▶ Quantum value:  $\omega^*(G)$ .

The values obey the following relationship for any nonlocal game:

$$\omega(G) \leq \omega^*(G) \leq \omega_c(G).$$



# Quantum strategies for nonlocal games

A *quantum strategy* consists of finite-dimensional complex Euclidean spaces  $\mathcal{A}$  and  $\mathcal{B}$  as well as the following:

- ▶ Shared state:  $\rho \in \mathcal{A} \otimes \mathcal{B}$ .
- ▶ Measurements  $\{A_a^x\} \subset \text{Pos}(\mathcal{A})$  and  $\{B_b^y\} \subset \text{Pos}(\mathcal{B})$ .

# Quantum strategies for nonlocal games

A *quantum strategy* consists of finite-dimensional complex Euclidean spaces  $\mathcal{A}$  and  $\mathcal{B}$  as well as the following:

- ▶ Shared state:  $\rho \in \mathcal{A} \otimes \mathcal{B}$ .
- ▶ Measurements  $\{A_a^x\} \subset \text{Pos}(\mathcal{A})$  and  $\{B_b^y\} \subset \text{Pos}(\mathcal{B})$ .

Winning probability for a quantum strategy is given by:

$$\sum_{x,y} \pi(x,y) \sum_{a,b} V(a,b|x,y) \langle A_a^x \otimes B_b^y, \rho \rangle.$$

The *quantum value*, denoted as  $\omega^*(G)$ , is the supremum of the winning probability over all quantum strategies.

# Commuting measurement strategies for nonlocal games

A *commuting measurement strategy* consists of a single (possibly infinite-dimensional) complex Euclidean space  $\mathcal{H}$ , as well as the following:

- ▶ Shared state:  $\rho \in \mathcal{H}$ .
- ▶ Measurements  $\{A_a^x\} \subset \text{Pos}(\mathcal{H})$  and  $\{B_b^y\} \subset \text{Pos}(\mathcal{H})$ ,

such that  $[A_a^x, B_b^y] = 0$  for all  $x, y, a, b$ .

Winning probability for a commuting measurement strategy is given by:


$$\sum_{x,y} \pi(x,y) \sum_{a,b} V(a,b|x,y) \langle A_a^x B_b^y, \rho \rangle.$$

The *commuting measurement value*, denoted as  $\omega_c(G)$ , is the supremum of the winning probability over all commuting measurement strategies.

# Calculating the quantum value is hard

Want: Method to calculate the quantum value of a nonlocal game.

This figure shows representations of the space of correlations for fixed and finite number of possible questions and answers.




Unfortunately, the set  $\mathcal{Q}$  is a non-polyhedral set with an infinite number of extreme points.

## Upper bounding nonlocal games

## Upper bounds for nonlocal games

- ▶ The NPA hierarchy<sup>¶</sup> is a method of placing *upper bounds* on the *quantum value* of nonlocal games.
- ▶ Hierarchy of semidefinite programs is *guaranteed* to converge to the commuting measurement value for some finite level,  $\ell$  of the hierarchy.
- ▶ The commuting measurement value is an upper bound on the quantum value,  $\omega^*(G) \leq \omega_c(G)$ , for all nonlocal games,  $G$ .



---

<sup>¶</sup>[Navascués, Pironio, and Acín, (2008)]

# NPA theorem (Main idea)

- ▶ Finding a quantum state and measurements for a quantum strategy is a computationally difficult task.

# NPA theorem (Main idea)

- ▶ Finding a quantum state and measurements for a quantum strategy is a computationally difficult task.
- ▶ Instead then, let's think about a set of *weaker* conditions that correspond to a commuting measurement strategy.



# NPA theorem (Main idea)

- ▶ Finding a quantum state and measurements for a quantum strategy is a computationally difficult task.
- ▶ Instead then, let's think about a set of *weaker* conditions that correspond to a commuting measurement strategy.
- ▶ In the NPA hierarchy, each condition amounts to verifying the existence of a positive semidefinite matrix with structure that depends on algebraic properties satisfied by a commuting measurement strategy.

# NPA theorem (Main idea)

- ▶ Finding a quantum state and measurements for a quantum strategy is a computationally difficult task.
- ▶ Instead then, let's think about a set of *weaker* conditions that correspond to a commuting measurement strategy.
- ▶ In the NPA hierarchy, each condition amounts to verifying the existence of a positive semidefinite matrix with structure that depends on algebraic properties satisfied by a commuting measurement strategy.
- ▶ If any of these conditions are violated, we may conclude that there does not exist an adequate state and sets of measurements.

# NPA theorem (Main idea)

NPA states that there exists some matrix  $C^{(\ell)}$  that allows  $\omega_c(G)$  to be calculated by maximizing

$$\sum_{x,y,a,b} \pi(x,y) V(a,b|x,y) C^{(\ell)}((x,a),(y,b))$$

for some finite level,  $\ell$ , such that  $C^{(\ell)}$  satisfies certain linear constraints satisfied by a commuting measurement strategy.

- ▶ These linear constraints can be checked via a semidefinite program!
- ▶ We refer to  $C^{(\ell)}$  as an  $\ell$ -th order pseudo commuting measurement assemblage.
- ▶ The next few slides will describe how  $C^{(\ell)}$  is defined.

## Example

Consider a nonlocal game where  $|\Sigma_A| = |\Sigma_B| = |\Gamma_A| = |\Gamma_B| = 2$ .  
Let's compute  $C^{(1)}$ :

$$C^{(1)} = \left( \begin{array}{c|cccc|cccc} & \mathbb{1} & A_0^0 & \cdots & A_1^1 & B_0^0 & \cdots & B_1^1 \\ \hline \mathbb{1} & & & & & & & \\ A_0^0 & & & & & & & \\ \vdots & & & & & & & \\ A_1^1 & & & & & & & \\ \hline B_0^0 & & & & & & & \\ \vdots & & & & & & & \\ B_1^1 & & & & & & & \end{array} \right)$$

- ▶ Fill in matrix with products of row and column to generate entries.
- ▶ For each entry,  $Z$ , computed in this way, the entry refers to an inner product between  $Z$  and the shared state  $\rho$ , i.e.  $\langle Z, \rho \rangle$ .

## Example

Consider a nonlocal game where  $|\Sigma_A| = |\Sigma_B| = |\Gamma_A| = |\Gamma_B| = 2$ .  
Let's compute  $C^{(1)}$ :

$$C^{(1)} = \left( \begin{array}{c|cccc|cccc} & \mathbb{1} & A_0^0 & \cdots & A_1^1 & B_0^0 & \cdots & B_1^1 \\ \hline \mathbb{1} & \mathbb{1} & A_0^0 & \cdots & A_1^1 & B_0^0 & \cdots & B_1^1 \\ A_0^0 & A_0^0 & A_0^0 & \cdots & A_0^0 A_1^1 & A_0^0 B_0^0 & \cdots & A_0^0 B_1^1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ A_1^1 & A_1^1 & A_1^1 A_0^0 & \cdots & A_1^1 & A_1^0 B_0^0 & \cdots & A_1^1 B_1^1 \\ \hline B_0^0 & B_0^0 & B_0^0 A_0^0 & \cdots & B_0^0 A_1^1 & B_0^0 & \cdots & B_0^0 B_1^1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ B_1^1 & B_1^1 & B_1^1 A_0^0 & \cdots & B_1^1 A_1^1 & B_1^1 B_0^0 & \cdots & B_1^1 \end{array} \right)$$

# Strings

In order to index into  $C^{(\ell)}((x, a), (y, b))$ , we will consider strings.

Define alphabets

$$\Sigma_A = \Sigma_A \times \Gamma_A, \quad \Sigma_B = \Sigma_B \times \Gamma_B, \quad \Sigma^* = \{\epsilon\} \cup \Sigma_A \cup \Sigma_B.$$

For example, we can refer to operators (or products of operators) as tuples of concatenated strings. For example:

$$A_a^x \rightarrow (x, a), \quad \text{and} \\ A_{a_1}^{x_1} \cdots A_{a_k}^{x_k} \rightarrow (x_1, a_1) \cdots (x_k, a_k).$$

Similarly for Bob.

# Equivalence relations for strings

The measurements in a commuting measurement strategy are *projective* and they *commute*. This property can be conveyed in terms of a string relation:

For all strings  $s, t \in \Sigma^*$ ,

1. Projective:  $s\sigma t \sim s\sigma\sigma t$  for all  $\sigma \in \Sigma$
2. Commute:  $s\sigma\tau t \sim s\tau\sigma t$  for all  $\sigma \in \Sigma_A$  and  $\tau \in \Sigma_B$ .

# Admissible functions

The function

$$\phi : \Sigma^* \rightarrow \mathbb{C}$$

is *admissible* iff it satisfies the following conditions:

1. Measurements sum to identity:

$$\sum_a \phi(s(x, a)t) = \sum_b \phi(s(y, b)t) = \phi(st),$$

for all  $x, y \in X \times Y$ .

2. Something:

$$\phi(s(x, a)(x, a')t) = \phi(s(y, b)(y, b')t) = 0$$

3. For all  $s, t \in \Sigma^*$  where  $s \sim t$

$$\phi(s) = \phi(t).$$



## $\ell$ -th order admissible matrices

We call the matrix  $C^{(\ell)}$  an  $\ell$ -th order admissible matrix if

1. There exists an admissible function

$$\phi : \Sigma^{\leq 2\ell} \rightarrow \mathbb{C},$$

such that

$$C^{(\ell)}(s, t) = \phi(s^R t) \quad \forall s, t \in \Sigma^{\leq \ell},$$

2. Normalization:  $C^{(\ell)}(\epsilon, \epsilon) = 1$ ,
3.  $C^{(\ell)}$  is positive semidefinite.

## $\ell$ -th order pseudo commuting measurement assemblages

Define an  $\ell$ -th order pseudo commuting measurement assemblage

$$K : A \times B \times X \times Y \rightarrow \mathcal{L}(\mathbb{C}),$$


for which there exists an  $\ell$ -th order admissible matrix  $C^{(\ell)}$  such that

$$K(a, b|x, y) = C^{(\ell)}((x, a), (y, b)) \quad \forall x, y, a, b.$$

## Extended nonlocal games

## Extended nonlocal games

An *extended nonlocal game* is a nonlocal game where the *referee* also holds a quantum system that he measures provided by Alice and Bob.



1. Question and answer sets  $(\Sigma_A, \Sigma_B)$  and  $(\Gamma_A, \Gamma_B)$ .
2. Distribution on question pairs:  $\pi : \Sigma_A \times \Sigma_B \rightarrow [0, 1]$ .
3. A measurement operator  $V : \Gamma_A \times \Gamma_B \times \Sigma_A \times \Sigma_B \rightarrow \text{Pos}(\mathcal{R})$ .

# Extended nonlocal games: Winning and losing probabilities

At the end of the protocol, the referee has:

1. The state at the end of the protocol:

$$\rho_{a,b}^{x,y} \in \mathcal{D}(\mathcal{R}).$$

2. A measurement the referee makes on its part of the state  $\rho$ :

$$V(a, b|x, y) \in \text{Pos}(\mathcal{R}).$$

The respective winning and losing probabilities are given by

$$\left\langle V(a, b|x, y), \rho_{a,b}^{x,y} \right\rangle \quad \text{and} \quad \left\langle \mathbb{1} - V(a, b|x, y), \rho_{a,b}^{x,y} \right\rangle.$$

# Standard quantum strategies

A *standard quantum strategy* consists of finite-dimensional complex Euclidean spaces  $\mathcal{R}$ ,  $\mathcal{A}$ , and  $\mathcal{B}$  as well as the following:

- ▶ Shared state:  $\rho \in \mathcal{R} \otimes \mathcal{A} \otimes \mathcal{B}$ .
- ▶ Measurements:  $\{A_a^x\} \subset \text{Pos}(\mathcal{A})$ ,  $\{B_b^y\} \subset \text{Pos}(\mathcal{B})$ .

# Standard quantum strategies

A *standard quantum strategy* consists of finite-dimensional complex Euclidean spaces  $\mathcal{R}$ ,  $\mathcal{A}$ , and  $\mathcal{B}$  as well as the following:

- ▶ Shared state:  $\rho \in \mathcal{R} \otimes \mathcal{A} \otimes \mathcal{B}$ .
- ▶ Measurements:  $\{A_a^x\} \subset \text{Pos}(\mathcal{A})$ ,  $\{B_b^y\} \subset \text{Pos}(\mathcal{B})$ .

Winning probability for a standard quantum strategy is given by:

$$\sum_{x,y} \pi(x,y) \sum_{a,b} \left\langle V(a,b|x,y) \otimes A_a^x \otimes B_b^y, \rho \right\rangle$$

The *standard quantum value*, denoted as  $\omega^*(G)$ , is the supremum of the winning probability over all standard quantum strategies.


## Upper bounding extended nonlocal games



# Upper bounds for extended nonlocal games

*Extended NPA hierarchy:*

- ▶ Uses the same idea as the NPA hierarchy. (For  $\dim(\mathcal{R}) = 1$ , the NPA hierarchy is a special case.)
- ▶ Enables one to compute *upper bounds* on the *standard quantum value* for *extended nonlocal games*.
- ▶ Same idea as before, only now we need to take into account the actions of the referee.



# Commuting measurement strategies (for ENLG)

A *commuting measurement strategy* consists of a finite-dimensional complex Euclidean space  $\mathcal{H}$  as well as the following:

- ▶ Shared state:  $\rho \in \mathcal{R} \otimes \mathcal{H}$ .
- ▶ Measurements:  $\{A_a^x\} \subset \text{Pos}(\mathcal{H})$ ,  $\{B_b^y\} \subset \text{Pos}(\mathcal{H})$ ,  
where  $[A_a^x, B_b^y] = 0$  for all  $x, y, a, b$ .

## Commuting measurement strategies (for ENLG)

A *commuting measurement strategy* consists of a finite-dimensional complex Euclidean space  $\mathcal{H}$  as well as the following:

- ▶ Shared state:  $\rho \in \mathcal{R} \otimes \mathcal{H}$ .
- ▶ Measurements:  $\{A_a^x\} \subset \text{Pos}(\mathcal{H})$ ,  $\{B_b^y\} \subset \text{Pos}(\mathcal{H})$ ,  
where  $[A_a^x, B_b^y] = 0$  for all  $x, y, a, b$ .

The expected *pay-off* for a commuting measurement strategy is given by:

$$\sum_{(x,y) \in \Sigma_A \times \Sigma_B} \pi(x,y) \sum_{(a,b) \in \Gamma_A \times \Gamma_B} \left\langle V(a,b|x,y) \otimes A_a^x B_b^y, \rho \right\rangle$$

The *commuting measurement value*, denoted as  $\omega_c(G)$ , is the supremum of the pay-off over all commuting measurement strategies.

## Extended NPA theorem

There exists some matrix  $M^{(\ell)}$  that allows  $\omega_c(G)$  (where  $G$  is an ENLG) to be calculated by maximizing

$$\sum_{x,y,a,b} \pi(x,y) \left\langle V(a,b|x,y), M^{(\ell)}((x,a),(y,b)) \right\rangle$$

## Extended NPA hierarchy

Same idea, but now we're taking into account the referee, and therefore have a larger matrix.

For each  $\ell$ , now consider block matrices

$$M^{(\ell)} = \begin{pmatrix} M_{1,1}^{(\ell)} & \cdots & M_{1,m}^{(\ell)} \\ \vdots & \ddots & \vdots \\ M_{m,1}^{(\ell)} & \cdots & M_{m,m}^{(\ell)} \end{pmatrix}$$

where each block takes the form  $M_{i,j}^{(\ell)} : \Sigma^{\leq \ell} \times \Sigma^{\leq \ell} \rightarrow \mathbb{C}$ .

- ▶ Each submatrix has similar properties to what we saw for the NPA hierarchy.
- ▶ The overall matrix also has some structure, which is unique to this case.

Supplementary material:  
Extended nonlocal games

# Winning probability for standard quantum strategies

The winning probability is given by the following equation:

$$\sum_{x,y} \pi(x,y) \sum_{a,b} \frac{\langle V(a,b|x,y), \text{Tr}_{\mathcal{A} \otimes \mathcal{B}} (\mathbb{1}_{\mathcal{R}} \otimes A_a^x \otimes B_b^y) \rho \rangle}{\text{Tr}(\mathbb{1}_{\mathcal{R}} \otimes A_a^x \otimes B_b^y) \rho} \text{Tr}(\mathbb{1}_{\mathcal{R}} \otimes A_a^x \otimes B_b^y) \rho$$

The probabilities cancel giving

$$\sum_{x,y} \pi(x,y) \sum_{a,b} \text{Tr}(V(a,b|x,y) \text{Tr}_{\mathcal{A} \otimes \mathcal{B}} (\mathbb{1}_{\mathcal{R}} \otimes A_a^x \otimes B_b^y) \rho)$$

The trace operator slips past the  $\mathbb{1}_{\mathcal{R}}$  giving

$$\sum_{x,y} \pi(x,y) \sum_{a,b} \text{Tr}(V(a,b|x,y) \otimes (A_a^x \otimes B_b^y) \rho)$$

Writing the trace in terms of the inner product, we have that

$$\sum_{x,y} \pi(x,y) \sum_{a,b} \left\langle V(a,b|x,y) \otimes A_a^x \otimes B_b^y, \rho \right\rangle.$$

Supplementary material:  
Lower bounds for extended nonlocal games



## Lower bounds for extended nonlocal games

*Key idea:* Fixing measurements on one system yields the optimal measurements of the other system via an SDP¶

---

¶[Liang and Doherty (2007)]

## Lower bounds for extended nonlocal games

*Key idea:* Fixing measurements on one system yields the optimal measurements of the other system via an SDP¶

Iterative “see-saw” algorithm between two SDPs:

- ▶ SDP-1: Fix Bob’s measurements. Optimize over Alice’s measurements.
- ▶ SDP-2: Fix Alice’s measurements (from SDP-1). Optimize over Bob’s measurements.
- ▶ Repeat.

Not guaranteed to give optimal value, as the algorithm can get stuck in a local minimum.

---

¶[Liang and Doherty (2007)]

## Lower bounds for extended nonlocal games

Define  $\{\rho_a^x : x \in \Sigma_A, a \in \Gamma_A\} \subset \text{Pos}(\mathcal{R} \otimes \mathcal{B})$  as the residual states acting on the referee and Bob's systems and let

$$f = \sum_{(x,y) \in \Sigma_A \times \Sigma_B} \pi(x,y) \sum_{(a,b) \in \Gamma_A \times \Gamma_B} \left\langle V(a,b|x,y) \otimes B_b^y, \rho_a^x \right\rangle$$

Lower bound (SDP-1)

max:  $f$

s.t.:  $\sum_{a \in \Gamma_A} \rho_a^x = \tau,$   
 $\tau \in \text{D}(\mathcal{R} \otimes \mathcal{B}).$

Lower bound (SDP-2)

max:  $f$

s.t.:  $\sum_{b \in \Gamma_B} B_b^y = \mathbb{1}_B,$   
 $B_b^y \in \text{Pos}(\mathcal{B}).$

- Iterate between SDP-1 and SDP-2 until desired numerical precision is reached.