

1

Solve the system of linear congruences:

$$x \equiv 4 \pmod{11}$$

$$x \equiv 7 \pmod{12}$$

$$x \equiv -3 \pmod{13}$$

$N = 11 \cdot 12 \cdot 13 = 1716$. So, we must calculate m_n of the following

$$\frac{N}{11} = 156m_1 \equiv 2m_1 \equiv 1 \pmod{11}$$

$$\frac{N}{12} = 143m_2 \equiv 11m_2 \equiv 1 \pmod{12}$$

$$\frac{N}{13} = 132m_3 \equiv 2m_3 \equiv 1 \pmod{13}$$

q	r	m_1	y	q	r	m_2	y	q	r	m_3	y
	11	0	1		12	0	1		13	0	1
5	2	1	0	1	11	1	0	6	2	1	0
1	1	-5	1	1	1	-1	1	1	1	-6	1

Thus, we have $m_1 = 6$, $m_2 = 11$, and $m_3 = 7$. Now we can calculate the solution:

$$x \equiv 4 \cdot 156 \cdot 6 + 7 \cdot 143 \cdot 11 + (-3) \cdot 132 \cdot 7 \pmod{1716}$$

$$\equiv 3744 + 11011 - 2772 \pmod{1716}$$

$$\equiv 11983 \pmod{1716}$$

$$\equiv 1687 \pmod{1716}$$

2

Solve the system of linear congruences:

$$2x \equiv 1 \pmod{3}$$

$$3x \equiv 2 \pmod{5}$$

$$4x \equiv 3 \pmod{7}$$

$$5x \equiv 4 \pmod{11}$$

$N = 3 \cdot 5 \cdot 7 \cdot 11 = 1155$. Now, let's find the inverses of the given congruences.

q	r	m_1	y	q	r	m_2	y	q	r	m_3	y	q	r	m_4	y
	3	0	1		5	0	1		7	0	1		11	0	1
1	2	1	0	1	3	1	0	1	4	1	0	2	5	1	0
1	1	-1	1	2	1	-1	1	3	1	-1	1	1	1	-2	1

Thus, we found 2, 4, 6, and 9 respectively. Let's plug these in

$$x \equiv 1 \cdot 2 \equiv 2 \pmod{3}$$

$$x \equiv 2 \cdot 4 \equiv 3 \pmod{5}$$

$$x \equiv 3 \cdot 6 \equiv 4 \pmod{7}$$

$$x \equiv 4 \cdot 9 \equiv 36 \equiv 3 \pmod{11}$$

Now, we must solve for m_n in

$$\frac{N}{3} = 385m_1 \equiv m_1 \equiv 1 \pmod{3}$$

$$\frac{N}{5} = 231m_2 \equiv m_2 \equiv 1 \pmod{5}$$

$$\frac{N}{7} = 165m_3 \equiv 4m_3 \equiv 1 \pmod{7}$$

$$\frac{N}{11} = 105m_4 \equiv 6m_4 \equiv 1 \pmod{11}$$

q	r	m_1	y	q	r	m_2	y	q	r	m_3	y	q	r	m_4	y
	3	0	1		5	0	1		7	0	1		11	0	1
1	2	1	0	1	3	1	0	1	4	1	0	2	5	1	0
1	1	-1	1	1	2	-1	1	1	3	-1	1	1	1	-2	1
		1	-1			1	-1	1	1	2	-1			2	-2

Thus, we have $m_1 = 1$, $m_2 = 1$, $m_3 = 2$, and $m_4 = 2$. Now we can calculate the solution:

$$\begin{aligned}
 x &\equiv 2 \cdot 385 \cdot 1 + 3 \cdot 231 \cdot 1 + 4 \cdot 165 \cdot 2 + 3 \cdot 105 \cdot 2 \pmod{1155} \\
 &\equiv 770 + 693 + 1320 + 630 \pmod{1155} \\
 &\equiv 3413 \pmod{1155} \\
 &\equiv 839 \pmod{1155}
 \end{aligned}$$

3

Solve the system of linear congruences:

$$x \equiv 2 \pmod{2}$$

$$x \equiv 4 \pmod{6}$$

$$x \equiv 2 \pmod{14}$$

$$x \equiv 10 \pmod{15}$$

We see that for any pair of congruences in the system, the GCD of the moduli divides the difference of the residues. Thus, a solution exists. Let's start from the first congruence working

our way down.

$$x \equiv 2 \pmod{2} \quad (1)$$

$$x = 2 + 2k_1 \text{ for } k_1 \in \mathbb{Z}$$

$$2 + 2k_1 \equiv 4 \pmod{6} \quad (2)$$

$$2k_1 \equiv 2 \pmod{6}$$

$$k_1 \equiv 1 \pmod{3}$$

$$k_1 = 1 + 3k_2 \text{ for } k_2 \in \mathbb{Z}$$

$$x = 2 + 2(1 + 3k_2)$$

$$x = 4 + 6k_2$$

$$4 + 6k_2 \equiv 2 \pmod{14} \quad (3)$$

$$6k_2 \equiv -2 \pmod{14}$$

$$3k_2 \equiv 6 \pmod{7}$$

$$3k_2 = 6 + 7k_3 \text{ for } k_3 \in \mathbb{Z}$$

$$x = 4 + 2(6 + 7k_3)$$

$$x = 16 + 14k_3$$

$$16 + 14k_3 \equiv 10 \pmod{15} \quad (4)$$

$$14k_3 \equiv -6 \pmod{15}$$

$$14k_3 \equiv 9 \pmod{15}$$

$$-k_3 \equiv 9 \pmod{15}$$

$$k_3 \equiv 6 \pmod{15}$$

$$k_3 = 6 + 15k_4 \text{ for } k_4 \in \mathbb{Z}$$

$$x = 16 + 14(6 + 15k_4)$$

$$x = 100 + 210k_4$$

$$x \equiv 100 \pmod{210} \quad (5)$$

4

Prove that there are arbitrarily long strings of consecutive integers in which every one of the numbers is divisible by a square (other than 1!).

First, let $P = \{p_1, p_2, \dots, p_n\}$ be the set of all prime numbers where all p_i are unique. We have previously proven that there are infinite prime numbers, so $|P| = \infty$. Now, we want to find a string of n consecutive integers starting with x . Let's say that x is divisible by p_1^2 , $x+1$ is divisible by p_2^2 , \dots , and $x+n-1$ is divisible by p_n^2 . Now, let's rewrite in terms of congruences.

$$x \equiv 0 \pmod{p_1^2}$$

$$x+1 \equiv 0 \pmod{p_2^2}$$

$$\vdots$$

$$x+n-1 \equiv 0 \pmod{p_n^2}$$

Rearranging gives us:

$$x \equiv 0 \pmod{p_1^2}$$

$$x \equiv -1 \pmod{p_2^2}$$

$$\vdots$$

$$x \equiv -n+1 \pmod{p_n^2}$$

Since p_1, p_2, \dots, p_n are all unique primes, any pair of congruences has coprime moduli. Thus, we can use the Chinese Remainder Theorem. It tells us that there exists a solution to this system

of congruences. Since we can choose n to be arbitrarily large, we can construct arbitrarily long sequences of consecutive integers, each divisible by a square.

5

Find the least positive residue of $3^{984} \pmod{31}$.

Luckily, 31 is prime, so we can simply use Fermat's Little Theorem. From it, we know $3^{31-1} = 3^{30} \equiv 1 \pmod{31}$.

$$\begin{aligned} 3^{984} &\equiv 3^{30 \cdot 32 + 24} \pmod{31} \\ &\equiv (3^{30})^{32} \cdot 3^{24} \pmod{31} \\ &\equiv 1^{32} \cdot 3^{24} \pmod{31} \\ &\equiv 3^{24} \pmod{31} \end{aligned}$$

Now, let's compute $3^{24} \pmod{31}$ with successive squares.

$$\begin{aligned} 3^2 &\equiv 9 \pmod{31} \\ 3^4 &\equiv 81 \equiv 19 \pmod{31} \\ 3^8 &\equiv 361 \equiv 20 \pmod{31} \\ 3^{16} &\equiv 400 \equiv 28 \pmod{31} \end{aligned}$$

$$\begin{aligned} 3^{24} &\equiv 3^{16+8} \pmod{31} \\ &\equiv 3^{16} \cdot 3^8 \pmod{31} \\ &\equiv 28 \cdot 20 \pmod{31} \\ &\equiv 560 \pmod{31} \\ &\equiv 2 \pmod{31} \end{aligned}$$

Thus, the least possible residue of $3^{984} \pmod{31}$ is 2.

6

Find the least positive residue of $3^{984} \pmod{360}$.

$360 = 2^3 \cdot 3^2 \cdot 5$. We can then split this into congruences of 8, 9, and 5 and combine with the Chinese Remainder Theorem. Let's calculate the congruences.

$$\begin{aligned} 3^{984} &= 9^{492} \\ &\equiv 1^{492} \pmod{8} \\ &\equiv 1 \pmod{8} \end{aligned}$$

$$\begin{aligned} 3^{984} &= 9^{492} \\ &\equiv 0^{492} \pmod{9} \\ &\equiv 0 \pmod{9} \end{aligned}$$

By Fermat's Little Theorem, we can say $3^{5-1} = 3^4 \equiv 1 \pmod{5}$.

$$\begin{aligned} 3^{984} &= 3^{4 \cdot 246} \\ &= (3^4)^{246} \\ &\equiv 1^{246} \pmod{5} \\ &\equiv 1 \pmod{5} \end{aligned}$$

Thus, we have 3 congruences with coprime moduli.

$$x \equiv 1 \pmod{8}$$

$$x \equiv 0 \pmod{9}$$

$$x \equiv 1 \pmod{5}$$

Now, we can solve this using the Chinese remainder Theorem. $N = 8 \cdot 9 \cdot 5 = 360$. Then, we need to find all inverses m_n .

$$\frac{N}{8} = 45m_1 \equiv 5m_1 \equiv 1 \pmod{8}$$

$$\frac{N}{9} = 40m_2 \equiv 4m_2 \equiv 1 \pmod{9}$$

$$\frac{N}{5} = 72m_3 \equiv 2m_3 \equiv 1 \pmod{5}$$

q	r	m_1	y	q	r	m_2	y	q	r	m_3	y
	8	0	1		9	0	1		5	0	1
5	5	1	0	4	4	1	0	2	2	1	0
1	3	-1	1	2	1	-2	1	2	1	-2	1
1	2	2	-1								
1	1	-3	2								

Thus, we have $m_1 = 5$, $m_2 = 7$, and $m_3 = 3$. Now we can calculate the solution.

$$x \equiv 1 \cdot 45 \cdot 5 + 0 \cdot 40 \cdot 7 + 1 \cdot 72 \cdot 3 \pmod{360}$$

$$\equiv 225 + 0 + 216 \pmod{360}$$

$$\equiv 441 \pmod{360}$$

$$\equiv 81 \pmod{360}$$

Thus, the least possible residue of $3^{984} \pmod{360}$ is 81.