

λ -calculi and Church-Rosser property

Vincent Rahli

supervisors: Professor Fairouz Kamareddine and Doctor J. B. Wells

ULTRA group, MACS, Heriot Watt University

June 12, 2008

A bit of History

formalisation of Mathematics (and functions)

- ▶ First formalisation of the concept of function by Frege (1879, premises of the formalisation of Mathematics)
- ▶ Discovery of some **paradoxes** in Mathematics (around 1900).
- ▶ Functions can be applied to any function: reflexiveness.
- ▶ Formalization of the concept of **type** by Russel to restrict application of functions (1908).
- ▶ Formalisation of Mathematics: design of logical systems

A bit of History

Improvement of the formalisation of the concept of function

- ▶ Design of the **λ -calculus** by Church as part of a formal system for logic and functions (1932).
- ▶ The full system was inconsistent:
 - ▶ Church uses the type free λ -calculus to investigate functions (successful model for computation);
 - ▶ Church adds simple types ($\text{int}, \text{int} \rightarrow \text{int}$) to λ -calculus in a system with logical axioms to deal with logic and function.
- ▶ Functions are studied as **computational rules** rather than as sets of pairs.

The λ -Calculus

“The traditional theory of λ -calculus relies on β -reduction, that is the capture by a function of its argument followed by the process of substituting this argument to the places where it is used.”
(Hugo Herbelin)

The λ -Calculus

the terms

Let Var be a countably infinite set of variables and $x, y, z, f \in \text{Var}$.

$$M, N \in \Lambda ::= x \mid (\lambda x.M) \mid (MN)$$

The λ -Calculus

the terms

Let Var be a countably infinite set of variables and $x, y, z, f \in \text{Var}$.

$$M, N \in \Lambda ::= x \mid (\lambda x.M) \mid (MN)$$

Natural Numbers: Church Numerals

$$\underline{0} := \lambda f x. x$$

$$\underline{1} := \lambda f x. f x$$

$$\underline{2} := \lambda f x. f(f x)$$

$$\underline{3} := \lambda f x. f(f(f x))$$

$\underline{0} := \lambda f x. x$ means $\underline{0}$ is a notation for $\lambda f x. x$

The λ -Calculus

the computational process

the β -reduction is the compatible closure[†] of the following rule:

$$(\lambda x.M_1)M_2 \rightarrow_{\beta} M_1[x := M_2]$$

[†]if $M_1 \rightarrow_{\beta} M_2$ then $\lambda x.M_1 \rightarrow_{\beta} \lambda x.M_2$ and $M_1 N \rightarrow_{\beta} M_2 N$ and $N M_1 \rightarrow_{\beta} N M_2$

The λ -Calculus

the computational process

the β -reduction is the compatible closure[†] of the following rule:

$$(\lambda x.M_1)M_2 \rightarrow_{\beta} M_1[x := M_2]$$

Natural Numbers: Church Numerals

$$2 + 1 = 3 \quad \text{or} \quad 2 + 1 \rightarrow 3$$

$$\text{Succ} := \lambda y f x. f(y f x)$$

$$\text{Plus} := \lambda y. y \text{ Succ}$$

$$\text{Plus } \underline{2} \ \underline{1} \rightarrow_{\beta}^* \underline{3}$$

[†]if $M_1 \rightarrow_{\beta} M_2$ then $\lambda x.M_1 \rightarrow_{\beta} \lambda x.M_2$ and $M_1 N \rightarrow_{\beta} M_2 N$ and $N M_1 \rightarrow_{\beta} N M_2$

The λ -Calculus

interlude

$\text{Succ} := \lambda y f x. f(y f x)$ (term in the λ -calculus)

and not

$\{\langle 0, 1 \rangle, \langle 1, 2 \rangle, \langle 2, 3 \rangle, \dots\}$ (set of pairs)

The λ -Calculus

the theory λ

The theory λ : the set of terms Λ and the equivalence relation $=_\beta$

The equivalence relation:

$$M =_\beta M'$$

is defined as the symmetric, transitive and reflexive closure of:

$$M =_\beta M' \quad \text{if} \quad M \rightarrow_\beta M'$$


Is the theory λ consistent?

(λ is consistent iff there exists M, N such that $M =_\beta N$ is not provable in λ)

The Church-Rosser Property

in arithmetic

Intuition:



The Church-Rosser property is also called the confluence property.

The Church-Rosser Theorem

in the λ -calculus

Plus $\underline{2} \underline{1} \rightarrow_{\beta}^* \underline{3}$

Is that always the case?

Does $\text{Plus } \underline{2} \underline{1} \rightarrow_{\beta}^* \underline{n}$ for $n \in \mathbb{N} \setminus \{3\}$?

Theorem

The λ -calculus satisfies the Church-Rosser property

First, proved by Church and Rosser in 1936 for a restricted version of the λ -calculus [CR36].

Since then, proved by many people (the most common one by Tait and Martin-Löf using the parallel reductions method).

The Church-Rosser Theorem

consistency of the λ -calculus

Assume

for all $M_1, N_2 \in \Lambda$, $M_1 =_{\beta} M_2$

By the Church-Rosser theorem

there exists M_3 such that $M_1 \rightarrow_{\beta}^* M_3$ and $M_2 \rightarrow_{\beta}^* M_3$

But, for example:

$\underline{0} \not\rightarrow_{\beta} \underline{1}$ and $\underline{1} \not\rightarrow_{\beta} \underline{0}$

So


$\underline{0} \neq_{\beta} \underline{1}$

(recall: $\underline{0} := \lambda fx.x$ and $\underline{1} := \lambda fx.fx$)

The Church-Rosser Theorem

how do we prove that?

- ▶ Embedding into a simpler calculus (**parallel reduction**):
 - ▶ By modifying the reduction relation.
 - ▶ By modifying the set of terms
(for example: based on a subset of the set of terms of the λ -calculus).



Extensionality

the theory $\lambda + \text{ext}$

The extensional theory $\lambda + \text{ext}$ is defined as follows:

The theory λ is extended with the following rule:

$$M =_{\beta\eta} N \text{ if } Mx =_{\beta\eta} Nx$$

Is the theory $\lambda + \text{ext}$ consistent?

Extensionality

η -reduction

The η -reduction is the compatible closure of the rule:

$$\lambda x.Mx \rightarrow_{\eta} M, \quad \text{if } x \notin \text{fv}(M)$$

(example: $x \in \text{fv}(\lambda y.xy)$ and $y, z \notin \text{fv}(\lambda y.xy)$)

Intuition for this rule:

$$(\lambda x.Mx)N \rightarrow_{\beta} MN \text{ and } (\lambda x.Mx)N \rightarrow_{\eta} MN, \quad \text{if } x \notin \text{fv}(M)$$

The theory $\lambda + \mathbf{ext}$ is equal to λ extended with the axiom scheme:

$$\lambda x.Mx =_{\beta\eta} M, \quad \text{if } x \notin \text{fv}(M)$$

Theorem

The λ -calculus satisfies the Church-Rosser property w.r.t. the $\beta\eta$ -reduction.

Proved by:

- ▶ Hindley in 1974 [Hin74]
- ▶ Klop in 1980 [Klo80]
- ▶ Kamareddine and Rahli in 2008 [KR08]

Extensionality

what is extensionality useful for? - a completeness result

We say that M has a normal form if there exists N such that $M \rightarrow_{\beta\eta}^* N$ and for all N' , $N \not\rightarrow_{\beta\eta} N'$.

Theorem [Bar84]

Suppose M, N have a normal form. Then either $M =_{\beta\eta} N$ is provable in $\lambda + \mathbf{ext}$ or $\lambda + \mathbf{ext} + M =_{\beta\eta} N$ is inconsistent.

The theory $\lambda + \mathbf{ext} + M =_{\beta\eta} N$ is the theory $\lambda + \mathbf{ext}$ augmented with the axiom $M =_{\beta\eta} N$.

Extensionality

how do we prove the Church-Rosser theorem w.r.t. the $\beta\eta$ -reduction?

For example:


- ▶ Let $c, d \in \text{Var}$. Let $\bar{x} \in \text{Var}_{cd} = \text{Var} \setminus \{c, d\}$.
Let the parametrised set of terms $\Lambda_{cd}^{\beta\eta}$ be defined as follows:

$$\bar{M} \in \Lambda_{cd}^{\beta\eta} ::= \bar{x} \mid d(c\bar{M}) \mid d(\lambda\bar{x}.\bar{M}) \mid (\lambda\bar{x}.\bar{M}_1)\bar{M}_2 \mid c\bar{M}_1\bar{M}_2$$

- ▶ Why this calculus? To impose a control on the reductions.
- ▶ This calculus satisfies the Church-Rosser property w.r.t. the $\beta\eta$ -reduction.
(By **reducibility**: method based on a semantic argument.)
- ▶ With an embedding of the λ -calculus in this calculus we prove that the λ -calculus satisfies the Church-Rosser property w.r.t. the $\beta\eta$ -reduction.
(Using the **method of parallel reduction**.)


Extensionality

one of our proof scheme for both β - and $\beta\eta$ -reduction




Extensionality

one of our proof scheme for both β - and $\beta\eta$ -reduction



Extensionality

one of our proof scheme for both β - and $\beta\eta$ -reduction




Extensionality

one of our proof scheme for both β - and $\beta\eta$ -reduction

Confluence of developments:

confluence of the simple calculus (reducibility)




Extensionality

one of our proof scheme for both β - and $\beta\eta$ -reduction

Confluence of developments:


confluence of the simple calculus (reducibility)



Another proof of Church-Rosser [GK01]


scheme of the Ghilezan and Kunčák's proof

Our proof is a simplification and extension of the Ghilezan and Kunčák's proof of the Church-Rosser property w.r.t. the β -reduction:



A general proof scheme

Both of the previous proof methods follow this proof scheme:



Remark

For now, this proof scheme is suitable for the Church-Rosser property only. Earlier attempts to find a common proof scheme for different properties of the lambda-calculus failed.

What else? What next?

Some of the other projects in which the ULTRA group is involved

- ▶ **Semantics** [KNRW08b, KNRW08a]: Soundness and completeness of semantics based on realisability, w.r.t. some type systems (intersection type systems with expansion variables). **Project**: find a complete semantics for an intersection type system with expansion (the whole mechanism).
- ▶ **Reducibility** [KRW08, KR08]: Study and simplification of proofs of properties of the lambda-calculus. **Project**: find a general proof scheme suitable for different properties of the lambda-calculus (such as Church-Rosser/normalisation/standardisation).
- ▶ **Type Error Slicing** [HW04]: The aim is to accurately identify the location of a type error of a piece of code (for a SML-based programming language), by providing a minimal and necessary set of points in the piece of code (a slice). **Project**: Extension to a bigger language.



Henk P. Barendregt.

The Lambda Calculus: Its Syntax and Semantics.

North-Holland, revised edition, 1984.



Alonzo Church and John B. Rosser.

Some properties of conversion.

Transactions of the American Mathematical Society, 39(3):472–482, 1936.



Silvia Ghilezan and Viktor Kunčák.

Confluence of untyped lambda calculus via simple types.

Lecture Notes in Computer Science, 2202:38–??, 2001.



Roger Hindley.

An abstract church-rosser theorem. II. applications.

J. Symbolic Logic, 39:1–21, 1974.



Christian Haack and J. B. Wells.

Type error slicing in implicitly typed higher-order languages.

Sci. Comput. Program., 50(1-3):189–224, 2004.



Jan W. Klop.

Combinatory Reductions Systems.

PhD thesis, Mathematisch Centrum, Amsterdam, 1980.



Fairouz Kamareddine, Karim Nour, Vincent Rahli, and Joe B. Wells.

A complete realisability semantics for intersection types and arbitrary expansion variables.

Accepted at ICTAC 2008, 2008.



Fairouz Kamareddine, Karim Nour, Vincent Rahli, and Joe B. Wells.

Realisability semantics for intersection type systems and expansion variables.

Presented to ITRS '08, 4th Workshop on Intersection Types and Related Systems, Turin, Italy, 25 March 2008, 2008.



Fairouz Kamareddine and Vincent Rahli.

Simplified reducibility proofs of church-rosser for beta- and beta eta-reduction.

2008.



Fairouz Kamareddine, Vincent Rahli, and J. B. Wells.

Reducibility proofs in the λ -calculus.

Presented to ITRS '08, 4th Workshop on Intersection Types and Related Systems, Turin, Italy, 25 March 2008, 2008.