Genetic Contributing Factors for Depression

0:00 / 4:44



Genetic Contributing Factors for Depression

Epidemiologist Myrna Weissman and her colleagues examined the risk of developing depression in the children and the grandchildren of depressed parents (Weissman et al., 2005). Even those depressive disorders are known to run in families, children of depressed parents (like Rowena Wells) do develop depression. Weissman and her colleagues found that 40% of children with a depressed parent (like Rowena Wells) had developed depression by adulthood. Parental factors were also the risk for the depressive illness of the grandchildren. Children with a depressed parent and depressed grandparent had the highest risk for developing depression themselves (Goodman et al., 2011; Weissman et al., 2016).

A heritability rate of 37-50% for depression indicates that environmental factors also play a role in the development of depression (Sullivan et al., 2000). Environmental factors can be considered as anything that has a significant influence on both family, such as family, household, and neighborhood variables have relatively little efficacy on depression. Mother, it is unique environmental influences (environmental influences or one two hundred the effect) that are not shared by family members (Plomin & Daniels, 2011). Environmental factors can be considered as anything that influences or observation include extreme childhood experiences and victimization, such as the bullying (Kwong et al., 2019).


Epigenetics

There's excitement that gene expression is impacted by the environment. **Epigenetics** is the field of study that investigates the ways in which the environment controls the activations or deactivation of various parts of our genome. Environmental influences, such as stress, can influence the expression of genes, and no reverse: the level of gene expression can influence the likelihood that a person will develop depression (Nestler et al., 2016). In depression, methylation is the key. Additionally, it can environmental factor that has been found to activate parts of the genome related to depression and suicide ideation (Cecil Jones et al., 2016; Fetterman et al., 2018).

The mechanism of epigenetics can be thought of as three basic steps (Park et al., 2005; Dulcis et al., 2013):

1. First hormonal stressors make persistent changes to an individual's genome.
2. Genomic changes promote activation or deactivation of genes on the person's DNA, which promotes the expression or suppression of an important protein.
3. The protein influences mood and behavior outcomes of the brain.

The Mechanism of Epigenetics



Research on identical twins has demonstrated that epigenetics plays a role in who manifests disorders such as depression and who does not. Identical twins have identical genetic material (Bruder et al., 2008; Kato, 2016; Labrie et al., 2018). Epigenetics influences are expressed in identical twins as an effect only one-twentieth of the time.

Epigenetics has been used (Tonoyan et al., 2019) to explain how stressful experiences can be passed from one generation to the next. Epigenetic changes have been found in survivors and developed posttraumatic stress disorder–related epigenetic markers or genes that could even suggested links between parental trauma and offspring PTSD (Yehuda & Lehrner, 2018). Epigenetic changes have been found in controlled research in this area, evidence supporting the trans-generational transmission of trauma through epigenetics is inclusive is still not definitive (Yehuda & Lehrner, 2018) and more research is needed.

Mark as completed

Like Dislike Report an issue