← Back

Module Quiz: Depressive and Bipolar Disorders

Graded Assignment • 16 min

Due Sep 12, 11:59 PM IST

Your grade: 87.50%

Your latest: **87.50%** • Your highest: **87.50%** • To pass you need at least 80%. We keep your highest score.

Next item →

Instructions

1. **Which of these is a symptom of dysthymic disorder but not depressive disorder?** 1 / 1 point

○ Twice as common in women than men.
○ Difficulty in making decisions and poor concentration.
● Symptoms present most of the day, more days than not for at least two years with no break in symptoms lasting more than 2 months.
○ Difficulty sleeping or oversleeping.

✓ Nice work

Of the options, dysthymic disorder is characterized by symptoms that are persistent and lasting for at least two years. Difficulty sleeping or oversleeping, difficulty in making decisions, having poor concentration, and prevalence rates twice as common in women than men are all common symptoms to both dysthymic disorder and the spectrum of depressive disorders.

2. **When asked to identify and choose items, objects, and activities, individuals with depression** 1 / 1 point

○ can identify what they like but will make their choices at random.
○ cannot identify what they like and will not make any choice.
● can identify what they like but choose not to engage with them.
○ cannot identify what they like and will select them at random.

✓ Nice work

Anhedonia is a key symptom of depression, specifically, it is the inability to experience pleasure. Individuals with depression have difficulty motivating themselves to engage in activities that were once rewarding because they no longer find pleasure in them.

3. **Which of these symptoms are common to ordinary sadness?** 1 / 1 point

○ Maladaptive cognition
○ Anhedonia
● Negative emotional state
○ Psychological changes

✓ Nice work

Ordinary sadness is marked by a negative emotional state or sadness. The symptom of sadness is also common in all depressive disorders. However, depressive disorders have symptoms that go beyond sadness including maladaptive cognitions, physiological changes, and anhedonia. Importantly, ordinary sadness tends to be temporary. Depressive disorders are often long term and impact daily life.

4. **Which of these is an example of physiological changes common to depressive disorders?** 1 / 1 point

● Sleep disturbances
○ Emotional numbing
○ Thoughts of self-harm
○ Mania

✓ Nice work

Sleep disturbances, fatigue, and lethargy are physiological changs common to depressive disorders. Thoughts of self-harm and emotional numbing are examples of maladaptive cognition. Mania is not a symptom of depressive disorders.

5. **Bipolar disorders and depressive disorders are both [BLANK] disorders. The primary difference is that bipolar disorders include symptoms of [BLANK] mood states.** 1 / 1 point

○ personality; normal
○ phobic; normal
● mood; elevated
○ anxiety; elevated

✓ Nice work

Bipolar disorders differ from depressive disorders in that people with one of the bipolar disorders cycle between elevated and depressive mood states.

6. **While experiencing a manic episode Marion sends her partner a message. Her partner fails to respond in a timely manner because their phone was turned off. Marion becomes increasingly agitated, and starts to think that her partner is cheating on her. In this situation, Marion is exhibiting** 0 / 1 point

● paranoia.
○ hallucination.
○ delusion.
○ deprivation.

⊗ Try again

7. **Which disorder is difficult to diagnose because elevated and depressed moods fail to meet criteria for hypomania, mania, or a depressive episode?** 1 / 1 point

○ Dysthymic disorder
○ Bipolar I disorder
● Cyclothymic disorder
○ Bipolar II disorder

✓ Nice work

Cyclothymic disorder is difficult to diagnose because its symptoms are less severe and often fail to meet the criteria for mania, hypomania, or a depressive episode.

8. **A diagnosis of cyclothymic disorder requires patients to show symptoms over [BLANK], whereas a diagnosis of bipolar I or II disorders requires the presence of symptoms over [BLANK].** 1 / 1 point

○ days, years
○ years, months
● years, days
○ weeks, months

✓ Nice work

To receive a diagnosis of cyclothymic disorder, the symptoms must have persisted for at least two years. For bipolar disorders, patients show symptoms of one manic or mixed episode lasting at least one week.

9. **A diagnosis for bipolar II disorder requires at least [BLANK] depressive episode/s, and one [BLANK] episode.** 1 / 1 point

○ one, manic
○ two, hypomanic
○ two, manic
● one, hypomanic

✓ Nice work

A diagnosis for bipolar II disorder requires at least one depressive episode and one hypomanic episode, but without a history of manic or mixed episodes.

10. **Bipolar I disorder is defined by the experience of at least [BLANK] manic or mixed episode/s lasting at least [BLANK] week. Some episodes may be hypomanic or depressive.** 1 / 1 point

○ one, two
● one, one
○ two, one
○ two, two

✓ Nice work

Bipolar I disorder is defined by the experience of at least one manic or mixed episode lasting at least one week. Some episodes may be hypomanic or depressive. The typical course of the disorder is characterized by recurrent depressive and manic or mixed episodes. Hypomanic episodes are not necessary for a diagnosis of bipolar I disorder. Approximately three-quarters of patients with bipolar I disorder experience psychotic symptoms at some point in their lives.

11. **What percentage of the global adult population is estimated to experience a depressive disorder?** 1 / 1 point

○ 7%
○ 3%
○ 10%
● 5%

✓ Nice work

It is estimated that depressive disorders account for about 3.8% of the global population and 5% of the adult population.

12. **The global prevalence of depression is greater for adult men than adult women.** 1 / 1 point

● False
○ True

✓ Nice work

Of the adults, prevalence is estimated to be about 4% among adult men and 6% among adult women.

13. **Global prevalence rates for bipolar II disorder are estimated to be between** 0 / 1 point

○ 4–6%
● 0.5–2%
○ less than 0.5%
○ 2–4%

⊗ Try again

14. **Of those experiencing bipolar I disorder, depression is often the predominant episode experienced by men and mania is the predominant episode experienced by women.** 1 / 1 point

● False
○ True

✓ Nice work

Depression is often the predominant episode experienced by women whereas the more prominent episode is mania for men. In addition, substance abuse tends to be a common comorbid disorder for men but eating disorders tend to be more common for women.

15. **Bipolar patients with mixed features who develop the disorder earlier in life are less likely to be diagnosed with bipolar I disorder and have lower number of suicide attempts.** 1 / 1 point

○ True
● False

✓ Nice work

Research has discovered that bipolar patients with mixed features develop the disorder earlier in life, are more likely to be diagnosed with bipolar I disorder, and have a higher number of suicide attempts than bipolar patients without mixed symptoms.

16. **The global lifetime prevalence rates for cyclothymic disorder are reported to be** 1 / 1 point

○ 2–3%
○ 3–4%
● 1% or less
○ 1–2%

✓ Nice work

Lifetime prevalence for cyclothymic disorder is estimated to be 0.4%–1% globally and occurs equally often in men and women. However, because of a lack of effective diagnosis, prevalence rates may actually be higher.

^ **See feedback**

After you submit this assignment click "See feedback" to see how you did, along with helpful information that will reinforce the module concepts.