← Back **Module Quiz: Suicide**
Graded Assignment • 15 min

Due Sep 18, 11:59 PM IST

Your grade: 93.33%

Your latest: **93.33%** • Your highest: **93.33%** • To pass you need at least 80%. We keep your highest score.

Next item →

Instructions

1. **Nonsuicidal self injury (NSSI) behaviors are** 1 / 1 point

○ always a precursor to a suicide attempt.
○ considered very rare.
● a risk factor for suicidal behaviors.
○ not related to suicidal behaviors.

✓ Nice work

While the relationship between nonsuicidal self-injury and suicide attempts is somewhat complicated, it is clear that nonsuicidal self-injury is a risk factor for suicidal behaviors. It remains unclear whether nonsuicidal self-injury increases the likelihood of suicide, being suicidal increases the likelihood of nonsuicidal self-injury, or some third factor, such as a lack of emotion regulation ability or another underlying psychological disorder such as depression causes both nonsuicidal self-injury or suicidal behaviors.

2. **Based on global reports, which country has the highest overall suicide rate (per 100,000 individuals) in 2019?** 0 / 1 point

○ Brazil
○ Indonesia
○ The United States
● China

⊗ Try again

3. **Which of the following is true concerning gender and suicide?** 1 / 1 point

○ Women complete suicide at a higher rate than men.
● Women attempt suicide at a higher rate than men.
○ There is no difference in the rate of suicide between women and men.
○ Women typically use more lethal or violent means of suicide.

✓ Nice work

Rates of suicide attempts and suicide completion are heavily influenced by gender. Globally, women attempt suicide more often than men, yet suicide completion rates are over two times higher for men than women. Contributing to this discrepancy, men are more likely to employ more deadly or violent means of suicide such as firearms, whereas women are more likely to employ less deadly means like cutting and drug overdose.

4. **Teenagers are the age group at the greatest risk of dying by suicide.** 1 / 1 point

○ True
● False

✓ Nice work

Teenagers are frequently viewed as the age group at the greatest risk of dying by suicide, however, teenagers are among the least likely age groups to die by suicide. Suicide rates typically increase with age.

5. **In the United States, American Indian or Alaska Native individuals have the highest suicide rate.** 1 / 1 point

○ False
● True

✓ Nice work

In the United States, the American Indian or Alaska Native population experienced a much higher suicide rate compared to other groups in 2021, at 28.1 per 100,000 individuals, following by White individuals with a rate of 17.4.

6. **Among LGBT adolescents, which group is at the highest risk of suicide?** 1 / 1 point

○ Bisexual adolescents
○ Lesbian adolescents
● Transgender adolescents
○ Gay adolescent

✓ Nice work

A meta-analysis that included studies from 10 countries concluded that transgender adolescents had the highest risk out of the LGBT youth included in the sample, with rates that were 5.87 times higher than the average for all adolescents.

7. **Why are techniques, such as MRI, DTI, fMRI, PET, and SPECT, used to diagnose suicidality in patients?** 1 / 1 point

○ Because these CBT techniques are effective in treating patients expressing suicidal intentions.
○ Because there may be an epigenetic effect on the expression of a person's genes.
○ Because neurotransmitter imbalance is linked to suicidal behavior.
● Because some brain regions are noted to change in patients with suicidal behavior.

✓ Nice work

Because of strong evidence linking suicidal behavior with changes to brain regions, neuroimaging techniques, such as magnetic resonance imaging (MRI), diffusion tensor imaging (DTI), functional MRI (fMRI), positron emission tomography (PET), and single photon emission computed tomography (SPECT), are being explored as techniques for identifying patients at risk of suicide.

8. **Psychological disorders, such as depressive disorders, bipolar disorders, schizophrenia, and borderline personality disorder, account for about what percentage of people who die by suicide?** 1 / 1 point

○ 40%
● 90%
○ 60%
○ 75%

✓ Nice work

Depressive disorders, bipolar disorders, schizophrenia, borderline personality disorder, and other psychological disorders account for an estimated 90% of people who die by suicide with depression as the symptom most closely associated with suicide ideation, attempts, and completions.

9. **Which of these is considered a neurochemical stressor?** 1 / 1 point

○ Marital stress
● Serotonin imbalance
○ Natural disaster
○ Career upheaval

✓ Nice work

An imbalance in the neurotransmitter serotonin is a psychological stressor linked to depression and suicide.

10. **Which of these factors is the most common predictor of completed adolescent suicide?** 1 / 1 point

○ Adverse childhood experiences
○ Loss of a parent
● Parent-child discord
○ Parent divorce

✓ Nice work

The high incidence of suicidal behavior in families is not due entirely to biological factors. The quality of family relationships has been shown to predict adolescent suicidal behavior. Parent-child discord is the most common predictor of completed adolescent suicide.

11. **Which of these is a culturally determined risk factor for suicide?** 1 / 1 point

○ Climate change
● Marital status in men
○ Sexual abuse
○ Alcohol and opiate use

✓ Nice work

Marital status, whether a person is married, partnered, single, or divorced, has been investigated as a risk factor for suicide, but there are cultural and geographical differences that affect whether marital status is a protective factor against suicide or a risk factor for suicide.

12. **What is the name of the phenomenon by which the occurrence of one suicide contributes to increased suicide completions or attempts by another person?** 1 / 1 point

○ Mass cluster
○ Clustering
● Contagion
○ Point cluster

✓ Nice work

Suicide contagion is the phenomenon by which the occurrence of one suicide contributes to increased suicide completions or attempts by another person. The Centers for Disease Control in the United States suggested that suicide contagion primarily affects at-risk individuals who identify with the person who died by suicide.

13. **Suicidal prevention interventions have advocated limiting access to all but which of these?** 1 / 1 point

○ High places for jumping
○ Lethal medications
● Awareness programs
○ Firearms

✓ Nice work

Researchers have concluded that suicide can be prevented by limiting access to lethal means of dying by suicide, especially medications, access to high places for jumping, and firearms. Educating clinicians to recognize and begin to treat individuals with suicidal symptoms and promoting suicide awareness programs have also been acknowledged as important preventative strategies.

14. **Suicide prevention involves removing motivation and means, which involves all but which of these goals?** 1 / 1 point

○ Work with those exposed to a completed suicide to prevent suicide contagion
○ Prevent as many first attempts at suicide as possible
● Education and awareness programs for doctors
○ Treat patients who have attempted suicide to prevent a second attempt

✓ Nice work

For someone to die by suicide, they must have the motivation to end their life and the ability to end their life. Education and awareness programs for doctors are important for screening individuals at a high risk of suicide, but it does not remove motivation or means. Preventing first attempters, treating those who have attempted, and working with those exposed to the completed or attempted suicide are suggested strategies for removing both motive and means.

15. **Which of these practices are encouraged during the postvention process following a suicide within a school or community setting?** 1 / 1 point

○ Share suicide helpful resources through social media
○ Confirming the facts from a knowledgeable authority
○ Conduct small-group educational and psychological support sessions
● All of these are important in the postvention process.

✓ Nice work

All of these are important in the postvention process. It's important to take immediate action, confirm the facts, communicate with the support community, identify at-risk individuals through monitoring of social media and small-group support sessions, make support resources available to everyone and everywhere, and encourage responsible reporting.

^ **See feedback**

After you submit this assignment click "See feedback" to see how you did, along with helpful information that will reinforce the module concepts.