

CE-25

EE23BTECH11063 - Vemula Siddhartha

Question:

The following function is defined over the interval $[-L, L]$:

$$f(x) = px^4 + qx^5$$

It is expressed as a Fourier series,

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left\{ a_n \sin\left(\frac{\pi nx}{L}\right) + b_n \cos\left(\frac{\pi nx}{L}\right) \right\}$$

which options amongst the following are true?

- (a) $a_n, n = 1, 2, \dots, \infty$ depend on p
- (b) $a_n, n = 1, 2, \dots, \infty$ depend on q
- (c) $b_n, n = 1, 2, \dots, \infty$ depend on p
- (d) $b_n, n = 1, 2, \dots, \infty$ depend on q

Solution:

The complex exponential Fourier Series of $f(x)$ is,

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{\pi nx}{L}} \quad (1)$$

$$c_n = \frac{1}{2L} \int_{-L}^L f(x) e^{-j\frac{\pi nx}{L}} dx \quad (2)$$

$$= \frac{1}{2L} \int_{-L}^L (px^4 + qx^5) e^{-j\frac{\pi nx}{L}} dx \quad (3)$$

$$= \frac{pL^4}{2} (e^{j\pi n} - e^{-j\pi n}) \left(\frac{1}{j\pi n} + \frac{12}{(j\pi n)^3} + \frac{24}{(j\pi n)^5} \right) \\ - \frac{pL^4}{2} (e^{j\pi n} + e^{-j\pi n}) \left(\frac{4}{(j\pi n)^2} + \frac{24}{(j\pi n)^4} \right) \\ - \frac{qL^5}{2} (e^{j\pi n} + e^{-j\pi n}) \left(\frac{1}{j\pi n} + \frac{20}{(j\pi n)^3} + \frac{120}{(j\pi n)^5} \right) \\ + \frac{qL^5}{2} (e^{j\pi n} - e^{-j\pi n}) \left(\frac{5}{(j\pi n)^2} + \frac{60}{(j\pi n)^4} + \frac{120}{(j\pi n)^6} \right) \quad (4)$$

$$c_n = 0 - (pL^4)(-1)^n \left(-\frac{4}{(\pi n)^2} + \frac{24}{(\pi n)^4} \right) \\ - (qL^5)(-1)^n \left(-\frac{j}{\pi n} + \frac{20j}{(\pi n)^3} - \frac{120j}{(\pi n)^5} \right) + 0 \quad (5)$$

We know,

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left\{ a_n \sin\left(\frac{\pi nx}{L}\right) + b_n \cos\left(\frac{\pi nx}{L}\right) \right\} \quad (6)$$

Finding the Fourier Coefficient a_0 ,

$$a_0 = c_0 \quad (7)$$

$$\Rightarrow a_0 = \frac{1}{2L} \int_{-L}^L f(x) dx \quad (8)$$

$$= \frac{1}{2L} \int_{-L}^L (px^4 + qx^5) dx \quad (9)$$

$$= \frac{pL^4}{5} \quad (10)$$

Finding the Fourier Coefficients a_n ,

$$a_n = \frac{c_{-n} - c_n}{j} \quad (11)$$

$$\Rightarrow a_n = (-2qL^5)(-1)^n \left(\frac{1}{\pi n} - \frac{2}{(\pi n)^3} + \frac{120}{(\pi n)^5} \right) \quad (12)$$

Finding the Fourier Coefficients b_n ,

$$b_n = c_n + c_{-n} \quad (13)$$

$$\Rightarrow b_n = (2pL^4)(-1)^n \left(\frac{4}{(\pi n)^2} - \frac{24}{(\pi n)^4} \right) \quad (14)$$

Hence, options (b) and (c) are correct.




Fig. 4. Fourier Series Approximation of $f(x)$ for $p = 5, q = 2, L = 0.08$