1

10.5.3.9

EE23BTECH11063 - Vemula Siddhartha

Question:

If the sum of first 7 terms of an AP is 49 and that of 17 terms is 289, find the sum of first n terms.

Solution:

The sum of first r terms of an Arithmetic Progression (AP) x(r), whose first term is a and common difference is d is:

$$x(r) = \frac{r}{2} (2a + (r - 1) d)$$
 (1)

Let the given AP have first term a and common difference d.

Given, the sum of first 7 terms of the AP is 49.

$$x(7) = 49$$

$$49 = \frac{7}{2}(2a + (7 - 1)d)$$

$$49 = \frac{7}{2}(2a + 6d)$$

$$a + 3d = 7$$
(2)

Also given, the sum of first 17 terms of the AP is 289.

$$x(17) = 289$$

$$289 = \frac{17}{2} (2a + (17 - 1) d)$$

$$289 = \frac{17}{2} (2a + 16d)$$

$$a + 8d = 17$$
(3)

Subtracting equation 2 from equation 3 we get:

$$5d = 10$$

$$d = 2 \tag{4}$$

Substituting the value of d in equation 1 we get:

$$a + 6 = 7$$

$$a = 1 \tag{5}$$

The sum of first n terms of the AP is:

$$x(n) = \frac{n}{2} (2a + (n-1) d)$$

Substituting the values of a and d:

$$x(n) = \frac{n}{2} (2(1) + (n-1)(2))$$

$$x(n) = n (1 + n - 1)$$

$$x(n) = n^{2}$$
(6)

Applying z-transform:

$$X(z) = \sum_{n=1}^{\infty} (n^2) z^{-n}$$

$$X(z) = (1^2) z^{-1} + (2^2) z^{-2} + (3^2) z^{-3} + \dots$$

$$X(z) = z^{-1} + 4z^{-2} + 9z^{-3} + 16z^{-4} + \dots$$
 (7)

Multiplying the equation 7 with z^{-1} :

$$z^{-1}X(z) = z^{-2} + 4z^{-3} + 9z^{-4} + 16z^{-5} + \dots$$
 (8)

Subtracting equation 8 from equation 7:

$$X(z)(1-z^{-1}) = z^{-1} + 3z^{-2} + 5z^{-3} + 7z^{-4} + \dots$$
 (9)

Multiplying the equation 9 with z^{-1} :

$$X(z)(z^{-1} - z^{-2}) = z^{-2} + 3z^{-3} + 5z^{-4} + 7z^{-5} + \dots$$
(10)

Subtracting equation 10 from equation 9:

$$X(z)(1 - 2z^{-1} + z^{-2}) = z^{-1} + 2(z^{-2} + z^{-3} + z^{-4} + \dots)$$

$$X(z)(1 - z^{-1})^{2} = z^{-1} + 2\frac{(z^{-2})}{(1 - z^{-1})}$$

$$X(z) = \frac{z^{-1}(1 - z^{-1}) + 2z^{-2}}{(1 - z^{-1})^{3}}$$

$$X(z) = \frac{z^{-1}(1 + z^{-1})}{(1 - z^{-1})^{3}}$$
(11)