

TEXAS TECH
UNIVERSITY®

Cypress Ck nr Westfield

Spatial Hydrology

Submitted By:
Vahid Salahi

Submitted To:
Dr. Uddameri

February 10, 2017

Contents

Section 1: Watershed Delineation	4
Section 2: Soil Hydrological Group	11
Section 3: Land Use Land Cover	14
Section 4: Curve Number Calculation	18

List of Figures

1	Digital Elevation Model Clipped to the corresponding 8 digit HUC	5
2	Flow direction map based on DEM	6
3	Flow accumulation map	6
4	Flow accumulation map (close view of the station point)	7
5	Watershed associated with pour point	8
6	Digital Elevation Map with 30m resolution	9
7	Watershed (30m resolution)	10
8	SHG map of watershed using SSURGO	11
9	SHG map of watershed using STATSGO2	12
10	LULC map for the year 2006	14
11	LULC map for the year 2011	15
12	Landuse changes from 2006 to 2011	16
13	Reclassified LULC of 2011	19
14	Reclassified SHG using SSURGO delineation	20
15	Curve Number Map	23

List of Tables

1	Comparison of Watershed Areas	10
2	Soil Hydrological Groups of Watershed	13
3	Change in Land Use Land Cover from 2006 to 2011	17
4	Reclassification of LULC data	18
5	Curve Number	21

Section 1

Delineate the watershed for the USGS gaging station corresponding to USGS 08069000 Cypress Ck nr Westfield, TX.

1. Show the flow direction, flow accumulation, and the final watershed maps (not screenshots). At a minimum you map must contain a legend, a north arrow and a scale bar.

The Digital Elevation Model has been used to derive flow direction map. The following prepossessing steps have been done to obtain the proper DEM to start the analysis:

- Two DEM subcategories with 10 m resolution (1/3 arc) downloaded from <https://nationalmap.gov/> which cover the entire 8 digit HUC corresponding to station point.
- "UTM Zone 14 N" was used as projection system to re-project both DEM subcategories. For this study, all the data and maps was projected using "UTM Zone 14N".
- "*Mosaic to New Raster*" tool has been utilized to combine two DEMs to get one raster file.
- 8 digit HUC with number of 12040102 which is containing the station point, was used as "*Clip*" feature to constrain the DEM raster data based on.
- "*Fill*" tool was used to remove small imperfections.

The final map of digital elevation model is depicted as Figure 1.




Figure 1: Digital Elevation Model Clipped to the corresponding 8 digit HUC

Based on the corrected DEM, the flow direction and flow accumulation maps are generated, shown as Figure 2 and 3 respectively. The flow accumulation map looks very uniform with low accumulation for entire raster data because the number of cells with low accumulation is much higher than the number of cells with high accumulation. By zooming in, the streams of high accumulation cells can be seen. Figure 4 shows a closer view of the accumulation map corresponding to the vicinity area of the station point.




Figure 2: Flow direction map based on DEM




Figure 3: Flow accumulation map

Figure 4 shows that the station point is not located on any cell with high accumulation (stream). The distance of 25 m was measured from the station point to the near stream and used as the tolerance in "Snap Pour Point" tool. The location of Pour Point on the stream is shown on Figure 4 in blue color.




Figure 4: Flow accumulation map (close view of the station point)

"Watershed" tool has been used to delineate the watershed associated with the snap pour point. The output watershed is shown in Figure 5.




Figure 5: Watershed associated with pour point

2. Calculate the contributing drainage area. Does it match reasonably with that indicated by the USGS on their website.

The area of delineated watershed shown in Figure 5 is calculated as 268 mi^2 (for obtaining the area, watershed was converted from raster to polygon using "*Raster to Polygon*" tool and then the area of polygon has been calculated). USGS has indicated an area of 285 mi^2 for the corresponding watershed. Difference in area can be mainly due to pour point estimation. Here a 25 m distance has been chosen as the tolerance but watershed is significantly sensitive to the pour point and different results might be obtained by changing the pour point. There are some other parameters that can affect the watershed

delineation process such as the resolution of DEM and the projection used for analysis.

In the next part, a sensitivity analysis has been considered on the DEM resolution. A courser resolution of 30 m (1 arc) has been chosen for the same analysis and the results have been compared. Figure 6 shows the DEM prepared for analysis in 30 m resolution.




Figure 6: Digital Elevation Map with 30m resolution

The new delineated watershed based on the new DEM (Figure 6) has been shown in Figure 7. The difference between two delineated watershed cannot be distinguished visually and they are similar to each other. The corresponding area for the new watershed calculated to be 267.5 mi^2 , indicating almost no

change in area (half square mile difference). Decreasing resolution from 10 m to 30 m did not affect the watershed delineation result significantly.

The comparison of the delineated watersheds is summarized in Table 1. For the rest of analysis, the delineated watershed from 10 m resolution DEM has been taken to consideration.




Figure 7: Watershed (30m resolution)

Table 1: Comparison of Watershed Areas

Description	Area mi^2	% Difference to USGS Watershed
Watershed from USGS Website	285	
Watershed using DEM of 1/3 arc resolution	268	5.96 %
Watershed using DEM of 1 arc resolution	267.5	6.14 %

Section 2

Download or extract the following data pertinent to the watershed – Soil Hydrologic Group (SHG) using SSURGO and Soil Hydrologic Group (SHG) corresponding to STATSGO2

1. Develop SHG maps corresponding to these two datasets.

The SHG map for both cases of SSURGO and STATSGO2 are shown in Figure 8 and Figure 9 respectively. For SSURGO case, the SHG for two counties of Harris and Waller have been *clipped* to the watershed and then *merged* to create the complete coverage of SHG on watershed. To obtain SHG using STATSGO2, the data of Texas state has been clipped to the watershed.




Figure 8: SHG map of watershed using SSURGO




Figure 9: SHG map of watershed using STATSGO2

2. Discuss the difference between the two delineations.

As it can be seen from Figure 8 and 9, these two delineations are different. The SHG map developed from STATGO2 has lost many details of SHG data and only have three class of SHG. But, the SHG map developed by SSURGO has more details of SHG distribution on the watershed. The other difference that can be observed is the lack of consistency between to delineations. For instance, the southern west part of the watershed has different SHG of C and D in these two delineation. Also, we can see significant areas of watershed

in west are grouped as B in SSURGO, however, in STATGO2 delineation, there is very small area of group B which is located in the northern east. This failure in correspondence is another difference in these two methods of delineation. The SHG map from SSURGO delineation has been chosen for the rest of analysis. Table 2 has summarized the percentage area of each SHG corresponding to Both SSURGO and STATGO2. For SSURGO, B/D and C/D have been considered as B and C respectively. This reclassification has been taken with the assumption of pre-development analysis.

Table 2: Soil Hydrological Groups of Watershed

Soil Hydrological Group	% Area (SSURGO)	% Area (STATGO2)
A	1.3	0
B	17	0.1
C	65.1	81.3
D	16.6	18.7

Section 3

Download the Land Use Land Cover (LULC) maps for the watershed for the years 2006 and 2011.

1. Map the LULC for these two years.

The LULC map for entire US was obtained from USDA web soil survey. Figure 10 and 11 represent the LULC maps clipped to the delineated watershed for year 2006 and 2011 respectively. All the maps are projected to UTM zone 14N as well.




Figure 10: LULC map for the year 2006




Figure 11: LULC map for the year 2011

2. Make a separate map that shows the areas that have undergone land use changes within that 5 year period.

Figure 12 depicts all the places that LULC record has been changed from 2006 to 2011. This map has been obtained by comparison of LULC data corresponding two years of 2006 and 2011. "Raster Calculator" was used to obtain the places that LULC has been changed.




Figure 12: Landuse changes from 2006 to 2011

3. How has the land use changed in the watershed within the 5 year period? Which LULC has the most dominant change?

The total area of the watershed that has undergone land use changes is 12.89 mi^2 . This value was obtained by converting the raster data shown in Figure 12 to polygon and then calculating the area. Table 3 shows the change of LULC from year 2006 to 2011 for all individual LULC classes.

The class code presented in Table 3 are encoded according to the USGS land

Table 3: Change in Land Use Land Cover from 2006 to 2011

Class Code and Name	Area in 2006 (%)	Area in 2011 (%)	Change (%)
11 (Open Water)	0.51	0.72	0.21
21 (Developed, Open Space)	11.12	10.74	-0.38
22 (Developed, Low Intensity)	9.99	10.73	0.74
23 (Developed, Medium Intensity)	9.23	10.78	1.55
24 (Developed High Intensity)	1.65	2.26	0.61
31 (Barren Land)	0.92	0.89	-0.03
41 (Deciduous Forest)	2.07	1.84	-0.23
42 (Evergreen Forest)	3.39	2.8	-0.59
43 (Mixed Forest)	0.6	0.53	-0.07
52 (Shrub/Scrub)	1.96	2	0.03
71 (Grassland/Herbaceous)	1.8	1.91	0.11
81 (Pasture/Hay)	37.9	36.44	-1.46
82 (Cultivated Crops)	12.46	12.22	-0.24
90 (Woody Wetlands)	5.07	4.83	-0.24
95 (Emergent Herbaceous Wetlands)	1.32	1.32	0

cover institute (LCI). Based on this table, the highest change of LULC in the watershed is in Residential (Medium Intensity). 1.55 % of watershed area has been increased for LULC class of 23. Also, land use corresponding to class 81 which is indicating Pasture/Hay land use has been decreased by 1.46 %. These changes imply lower capacity of infiltration and higher capacity of runoff for the watershed.

Section 4

Use the 2011 LULC and SSURGO SHG datasets to develop a spatial distribution of the Curve Number within the watershed.

1. Show a map of the spatial distribution of the Curve Number within the watershed.

The 2011 LULC includes 15 classes of different land uses. Table 4 shows the reclassification of LULC in four major classes. This reclassification has been done to make the task easier.

Table 4: Reclassification of LULC data

Original LULC Classification		Reclassification	
Class Number	Description	New Class Num	Description
11	Open Water	1	Water
90	Woody Wetlands		
95	Emergent Herbaceous Wetlands		
21	Developed, Open Space	2	Residential
22	Developed, Low Intensity		
23	Developed, Medium Intensity		
24	Developed High Intensity		
41	Deciduous Forest	3	Forest
42	Evergreen Forest		
43	Mixed Forest		
31	Barren Land	4	Agriculture
52	Shrub/Scrub		
71	Grassland/Herbaceous		
81	Pasture/Hay		
82	Cultivated Crops		

Figure 13 shows the spatial distribution of reclassified LULC on the watershed. The reclassification has been done using "Reclassify" toll in ArcGIS and 2011 LULC rater data was used as input.




Figure 13: Reclassified LULC of 2011

Based on Figure 8, there are some regions that SHG are assigned as B/D or C/D. Assuming pre-development evaluation of the watershed, SHG of B and C was substituted for groups B/D and C/D respectively. The spatial distribution of the new SHG dataset is shown in Figure 14.




Figure 14: Reclassified SHG using SSURGO delineation

After preparation of LULC and SHG data, a fishnet of $50 \times 50\text{ m}$ cell size was created covering the watershed. *"Spatial Join"* tool was used to spatially join LULC and SHG data to each cell of fishnet. The LULC and SHG are going to be used in conjunction with curve numbers, to assign a curve number to each cell of fishnet. Table 5 summarize the curve numbers associated with pairs of LULCs and SHGs. Classification of Table 4 and 5 were done by using datasets presented in our text book and also a report with topic of *"Creating SCS Curve Number Grid using HEC-GeoHMS"* which was prepared by Venkatesh Merwade from Purdue University. This report is available online (<https://web.ics.purdue.edu/~vmerwade/education/cngrid.pdf>).

Table 5: Curve Number

LULC	Soil Hydrological Group			
	A	B	C	D
1 (Water)	100	100	100	100
2 (Residential)	77	81	86	91
3 (Forest)	30	58	71	78
4 (Agriculture)	67	77	83	87

The following code has been written in R to assign the curve numbers to the fishnet cells based of Table 5.

```

1 # Set work directory:
2 setwd("C:\\\\Users\\\\vahid\\\\Google Drive\\\\GraduateStudies\\\\Semester
3       6\\\\SurfaceWaterHydrology\\\\Module2\\\\Datasets\\\\Ass2\\\\Fishnet")
4
5 # fishnet attribute table including SHG and LULC attributes is read
6 # as "table"
7 table <- read.csv("Table.csv")
8
9 # Curve number table (Table 5 in report)
10 cn <- read.csv("CNtable.csv")
11
12
13 table$HydrolGrp <- as.character(table$HydrolGrp)
14 #missing values for Hydrological group in fishnet set as "E"
15 v <- replace(table$HydrolGrp, table$HydrolGrp==" ", "E")
16 #create new table by using modified missing values of SHG
17 table$HydrolGrp <- v
18
19
20 # creating a list named as "encode" to encode SHG and LULC as
21 # numbers
22 encode<-list()
23 encode[[ "A" ]]<-2
24 encode[[ "B" ]]<-3
25 encode[[ "C" ]]<-4
26 encode[[ "D" ]]<-5
27 encode[[ "E" ]]<-6
28
29 # "0" is LULC value for cells with missing values
30 encode[[ "0" ]]<-5
31 encode[[ "1" ]]<-1

```

```

32 encode[[ "2" ]]<-2
33 encode[[ "3" ]]<-3
34 encode[[ "4" ]]<-4
35
36
37 #modifying CN table by adding values of "0" curve number for cels
  with SHG of "E" or LULC of "0"
38 E<-rep(0,4)
39 cn<-cbind(cn,E)
40
41 u<-c(5,rep(0,5))
42 cn<-rbind(cn,u)
43
44
45 # the final CN results will be save in a vector named as "CN"
46 CN<-vector()
47 # c1 and c2 are vectors of SHG and LULC respectively. They are
  coerced as character to encode them to numbers in the next line
48 c1<-as.character(table$HydrolGrp)
49 c2<-as.character(table$gridcode)
50
51 # for loop to check SHG and LULC for each cell and assign a curve
  number based on 'cn' table
52 for(i in 1:nrow(table)){
53   temp1<-c1[i]
54   temp2<-c2[i]
55   a <- cn[encode[[temp2]], encode[[temp1]]]
56   CN[i] <- a
57 }
58
59 # add the new attribute of Curve Number (CN) to the origin table
60 table<-cbind(table,CN)

```

The processed fishnet dataset was taken back to ArcGIS including the new attribute of Curve Number (CN). Figure 15 shows the spatial distribution of curve number on the watershed.




Figure 15: Curve Number Map

2. Calculate the area-weighted curve number for your watershed. (Show your calculations/code).

The area-weighted curve number can be obtained using Equation 1:

$$\overline{CN} = \frac{\sum_{i=1}^n CN_i \times A_i}{\sum_{i=1}^n A_i} \quad (1)$$

Since we used a fishnet to obtain the CN distribution map, the area associated with each cell is same for all and equal to cell size. So the CN calculation can

be simplified as follows:

$$\overline{CN} = \frac{A_{cell} \times \sum_{i=1}^n CN_i}{A_{cell} \times n} = \frac{\sum_{i=1}^n CN_i}{n}$$

where n is the total number of cells in fishnet (279673 active cells for CN). Calling a query on curve number attribute in fishnet data results in CN summation equals to 23073022. Then we have:

$$\overline{CN} = \frac{\sum_{i=1}^n CN_i}{n} = \frac{23073022}{279673} \approx 82$$

It should be considered that average curve number of 82 has been obtained with assumptions made regards to reclassification of LULD and SHG data.