

BROWN

Meteorological and Behavioral Correlates of COVID-19 Transmissibility Across the United States



Venkatsai Bellala¹, Annabel Coyle², Linsey Marr³

¹Biomedical Engineering, Brown University

²Biological Sciences, Virginia Tech

³Civil and Environmental Engineering, Virginia Tech

Background

Respiratory viruses (i.e. SARS-CoV-2) primarily transmit through the air.

Instantaneous reproduction number (R_t): Expected infections by one individual at a given time and a measure of transmissibility.




Figure 1. Factors mediating influenza epidemics [1]

Objective

The purpose of this study is to quantify the role of dry bulb temperature (DBT), relative humidity (RH), absolute humidity (AH), and indoor activity (IA) on the instantaneous reproduction number of COVID-19.

We hypothesize that an interaction between indoor activity and said meteorological variables will be strong predictors of SARS-CoV-2 transmissibility.

Methods

- Daily COVID-19 surveillance data – Johns Hopkins Center for Systems Science and Engineering (JHU CSSE) [2]
- Hourly meteorological measurements – National Centers for Environmental Information (NCEI) [3]
 - AH calculated with humidity R package [4]
- Weekly relative indoor activity (IA) from a study of cell phone records [5]
- R_t for SARS-CoV-2 computed with the Cori et al. method [6]
 - EpiEstim R package
 - Incidence \approx 7 day rolling average of confirmed cases
 - Parametric serial interval with a sliding window of 7 days
 - $\mu = 4.8, \sigma = 2.3$ [7]




Figure 2. Map of regions studied

Results

Equation 1. Multivariate orthogonal binomial model

$$\bar{R} \approx \text{poly}\left(\frac{\text{DBT} \cdot \ln(\text{RH})}{\text{Indoor Activity}}, 2\right)$$

The purpose of this model is to estimate the effect of interactions between DBT, RH, and IA on reproduction number.

Table 1. Summary of model fit and optimal lag*

State	Region	R^2_{adj}	Lag (weeks)
NY	Queens County	0.52	0
	Erie County	0.60	0
VA	Thomas Jefferson Health District	0.28	-1
	Virginia Beach Health District	0.26	-2
TX	Harris County	0.48	0
	Cameron County	0.037	2

*Lags kept at 0 throughout analysis for consistency.

Key Findings

- Significant U-shaped relationship with weekly mean R_t during the "Delta Phase" \rightarrow T, RH, and IA are good predictors




Figure 3. COVID-19 "Alpha Phase" in Queens County

- Weak & non-significant associations during the "Alpha Phase"
- Latitudinal trend \rightarrow stronger further from equator
- Population differences \rightarrow stronger in larger populations




Figure 4. Results by county

Assumptions & Limitations

Table 2. Shapiro-Wilk test of normality for R_t

State	Region	p-value
NY	Queens County	0.0061
	Erie County	0.0831
VA	Thomas Jefferson Health District	0.0578
	Virginia Beach Health District	0.0845
TX	Harris County	0.140
	Cameron County	0.0001

- Weekly mean R_t is mostly normally distributed ($p > 0.05$).
- Heteroskedasticity observed in regression model
- Inconsistencies present in incidence data

Conclusions

DBT, RH, and indoor activity are significant drivers of SARS-CoV-2 transmission in the near-absence of preventative measures.

- Inform predictions of future COVID-19 outbreaks
- Guide public policy planning in affected communities

Future Directions

- Fit model to more counties across the US
- Analyze the Omicron wave
- Control for lags in symptom onset and reporting
- Incorporate demographic parameters (age, risk factors, etc.)

References

- J. Tamerius, M. I. Nelson, S. Z. Zhou, C. Viboud, M. A. Miller, and W. J. Alonso, "Global influenza seasonality: reconciling patterns across temperate and tropical regions," *Environmental health perspectives*, vol. 119, no. 4, pp. 439–445, 2011.
- E. Dong, H. Du, and L. Gardner, "An interactive web-based dashboard to track covid-19 in real time," *The Lancet infectious diseases*, vol. 20, no. 5, pp. 533–534, 2020.
- National Centers for Environmental Information, "U.S. local climatological data," 2021.
- J. Cai, *humidity: Calculate Water Vapor Measures from Temperature and Dew Point*, 2019. R package version 0.1.5.
- S. Susswein, E. C. Rest, and S. Bansal, "Disentangling the rhythms of human activity in the built environment for airborne transmission risk," *medRxiv*, 2022.
- A. Cori, *EpiEstim: Estimate Time Varying Reproduction Numbers from Epidemic Curves*, 2021. R package version 2.2-4.
- H. Nishiura, N. M. Linton, and A. R. Akhmetzhanov, "Serial interval of novel coronavirus (covid-19) infections," *International journal of infectious diseases*, vol. 93, pp. 284–286, 2020.
- R Core Team, *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2022.

Acknowledgments

A very special thank you to Annabel Coyle for her assistance and patience, Linsey Marr for her mentorship, and to David Schmale and Shane Ross for sponsoring this program.

This research was supported in part by the National Science Foundation (NSF) under grant number 1922516.



National
Science
Foundation