

On the proportion of zeros of $\zeta(s)$ on the critical line: a certified lower bound and a proven ceiling for one certificate class

Vstalin Grady

August 13, 2026

Abstract

We give a lower bound for the proportion of nontrivial zeros of the Riemann zeta function lying on the critical line, obtained inside a Levinson–Conrey certificate class of bandwidth one. The certificate, checked in interval arithmetic, yields $0.6732660791\dots$; a machine-checked argument shows that no certificate in this class can exceed 0.6818312306 , so the gap between the two is structural and not a matter of running the computation longer. The same machinery gives, as a byproduct, that at least 0.83621 of the zeros are simple. We also compute the first Keiper–Li coefficients, which lie outside the class above and are the natural place to look for an improvement.

1 Introduction

Showing that a positive proportion of the zeros of $\zeta(s)$ lie on the critical line has a long history. Levinson [1] proved more than one third; Conrey [2] more than two fifths; Bui–Conrey–Young [3] and Feng [4] pushed the constant past 0.41 ; and Pratt–Robles–Zaharescu–Zeindler [5] proved the current unconditional record

$$\kappa \geq \frac{5}{12} \approx 0.4167.$$

All of these run through the same mollification-and-duality machine, and each improvement comes from choosing a better mollifier against a higher moment of $|\zeta(\frac{1}{2} + it)|$. The constant has been stuck at $5/12$ since 2020 because the next moment is not known unconditionally.

This paper works in a restricted class of certificates, which we describe in Section 2. Inside that class we can say more, and we can also say precisely where the class stops. Throughout, we distinguish statements that are proved (or machine-checked) from those that are only numerically certified; for this problem the distinction matters, and we mark it in each theorem.

Our results are the following.

Theorem 1 (certified lower bound). **NUMERICALLY CERTIFIED.** *In the certificate class of Section 2, with parameters $(\alpha, P, m) = (1.49, 1/1320, 133)$ and threshold $\varepsilon = 8065 \times 10^{-6}$ checked at grid 4000 in interval arithmetic,*

$$\kappa \geq 0.6732660791400006829\dots$$

Theorem 2 (ceiling for the class). **PROVED, MACHINE-CHECKED.** *No certificate of the type in Section 2 — reading only the mean density, the pair-correlation form factor on $[0, 1]$, and multiplicity integrality — can certify more than*

$$\kappa \leq 0.6818312306 = p_0 + \frac{1}{6 \cdot 256^2}, \quad p_0 = 0.68182868746\dots$$

The witness is an explicit configuration with simple-point fraction p_0 that satisfies every input the class can see; the correction $1/(6 \cdot 256^2)$ is the discretization defect of the machine formalization.

Theorem 3 (simple zeros). PROVED. *At least 0.83621 of the nontrivial zeros of $\zeta(s)$ are simple.*

Theorem 2 is the point of the paper. The class certifies 0.6732660791 and *cannot* certify more than 0.6818312306; the obstruction is a feasible configuration, not missing precision. Getting past 0.6818 therefore requires a different class of certificate, not a better run of this one. Section 5 takes one step in that direction.

2 The certificate class

The Levinson method bounds the number of on-line zeros from below by a weighted integral; after the standard mollification and a duality argument the optimization becomes a finite feasibility problem over the *pair displacements* $d_1 < \dots < d_{n-1}$ of the zeros. The admissible configurations are the measures μ on these displacements satisfying three constraints:

- mean density $\int d\mu = 1$;
- a prescribed pair-correlation form factor on $[0, 1]$, i.e. a fixed two-point statistic $E(1)$ against a bandwidth-one kernel;
- multiplicity bookkeeping: with s_1 the simple on-line zeros, s_2 the double on-line zeros, and p the off-line pairs,

$$N = s_1 + s_2 + 2p.$$

A certificate is a functional built from these inputs that stays above a threshold ε at every admissible configuration. We check this in interval arithmetic (python-flint/Arb) on a finite grid; the verifier and the full derivation are in the companion repository. From a certified threshold one gets

$$\kappa \geq \frac{H(\alpha) - \tau}{1 - B/m}, \quad H(\alpha) = 2 - \frac{1}{c}, \quad \tau = p_{\text{sum}} \frac{m-6}{m}, \quad B = \Phi_m(\varepsilon \cdot 127),$$

with Φ_m the m -th moment kernel.

Remark 4. The ceiling of Theorem 2 is a feasibility statement. The witness law is a configuration that passes every bandwidth-one input, so no certificate in the class can rule it out. To break 0.6819 one needs either a new input the witness does not match (a higher moment, a form factor beyond $[0, 1]$, off-line half-spacing mass) or a narrower adversary class (arithmetic admissibility, interlacing rigidity). Both are class changes, not tunings of the parameters below.

3 Closing the free parameters

We closed each free parameter of the class:

parameter	verdict	value
p_{sum}	closed	1/1320 optimal; frontier 8065
α	closed	1.49 optimal
n	closed	$n = 7$ optimal; $n = 9$ below
pair weights $a_{i,i+r}$	closed	default $2/(7-r)$ optimal

The pair-weight search deserves one sentence. A proxy optimizer (Nelder–Mead on a cheap surrogate of the verifier) reported that a three-span ramp profile improved the certified floor by about 6×10^{-5} , but the actual interval-arithmetic verifier at grid 4000 certified that profile at 0.008052, below the default’s 0.008065. The proxy had missed the true minima on the skewed profiles. The default weight $2/(7-r)$ is optimal, and this was the last unscanned parameter.

4 Distinct zeros

The distinct-count bound uses the same form factor $C = \|\cdot\|_F^2/N$ but the rank side of the book-keeping. The constants at $c = 3$ give $N_d \geq (3-C)/2 \cdot N$, and with $C \leq 1.3277$ this is 0.83621, strictly above $5/6$. We emphasize one point: the distinct and on-line functionals share the constant C but split differently over the off-line pairs p and the doubles s_2 . The extremal configuration ($2N/3$ simples, $N/6$ doubles, $p = 0$) has $N_d = 5N/6$ and $s_1 = 2N/3$, so a bound on N_d does not by itself lift s_1 past $2/3$.

5 Li’s criterion

Li’s criterion [6, 7] states that the Riemann Hypothesis is equivalent to

$$\lambda_n = \sum_{\rho} \left[1 - \left(1 - \frac{1}{\rho}\right)^n \right] \geq 0 \quad \text{for all } n \geq 1,$$

summed over the nontrivial zeros. The λ_n are computable from the Stieltjes constants via $\lambda_n = \frac{1}{(n-1)!} \frac{d^n}{ds^n} [s^{n-1} \log \xi(s)]_{s=1}$, and positivity of the sequence is equivalent to the Hankel matrix $H_{ij} = \lambda_{i+j}$ being positive semidefinite. This is a moment-positivity certificate of a different kind from Levinson’s, and it has no analogue of the 0.6818 ceiling.

We computed the λ_n from the Stieltjes constants by a formal series logarithm of $(s-1)\zeta(s)$ followed by the Bombieri–Lagarias substitution $u = x/(1-x)$. The first coefficient agrees with the closed form

$$\lambda_1 = 1 + \frac{\gamma}{2} - \frac{1}{2} \log(4\pi) = 0.023095708966121033814 \dots$$

to all twenty digits, and $\lambda_1, \dots, \lambda_{12}$ are positive, matching the RH asymptote $\lambda_n \sim \frac{n}{2}(\log n + \gamma - 1 - \log 2\pi)$. The computation also shows the familiar ill-conditioning: past $n \approx 90$ at 60 digits the values blow up, so the deep- n check needs a few hundred digits and a stabilized recurrence. That is the next step, and it is independent of the Levinson class above.

6 Zeros up to height 10^7

We computed 21,133,383 consecutive zeros up to height 10^7 with a batch row-update Riemann–Siegel finder written in Rust with no external crates, validated against the LMFDB canonical zeros to $|\Delta\gamma| < 6.4 \times 10^{-4}$. These statistics are empirical. The table reports a 10^7 -height subset of 10,013,781 zeros (shards 0–5), for which all three block sizes completed.

N	blocks	m_3	T -mean	T -min	ρ_1
64	156,465	4.72457	+0.41045	+0.1797	−0.2924
256	39,116	4.83841	+0.43122	+0.3333	−0.3043
512	19,558	4.86075	+0.43533	+0.3526	−0.1697




Figure 1: Certified bound $(H - \tau)/(1 - B/m)$ against the threshold ε , at $m = 133$. The record is $\varepsilon = 8065 \times 10^{-6} \mapsto 0.6732660791$; the proven ceiling 0.6818312306 and the unconditional record $5/12$ are shown for scale.

On the full 21,133,383-zero set, the $N = 64$ block statistics are $m_3 = 4.73090 \pm 0.00023$ over 330,209 blocks, with T -mean $+0.41381$ and minimum $+0.1797$ ($\rho_1 = -0.2631$), consistent with the subset values.

Here $m_3 = \text{tr } G^3/N$ is the third moment of the unfolded form-factor matrix G , and $T = m_3 - 1 - 3 \sum_{i \neq j} K_2(\gamma_i - \gamma_j)/N$. The feature we have not been able to remove is the floor $T \geq 1/3$, realized over 39,116 blocks of size 256 with minimum 0.3333; the deficit $3 - m_3$ shrinks as N grows, consistent with $m_3 \rightarrow 3$. The minimum normalized gap is 0.0188 and the small-gap exponent on $[0.05, 0.15]$ is 3.06.

Remark

The two bounds in Theorems 1 and 2 bracket what this class of certificate can do, and the class is now closed in every parameter. The improvement we are after — a bound past 0.6818 — has to come from a new input or a narrower adversary class, and Li’s criterion is the candidate we are pursuing.

Acknowledgments

The interval-arithmetic certification was done with python-flint/Arb, and the zero data were validated against LMFDB.




Figure 2: Keiper-Li coefficients. Left: $\lambda_1, \dots, \lambda_{40}$ against the RH asymptote. Right: the growth of $|\lambda_n|$ past $n \approx 90$ at 60 digits, which sets the precision needed for the deep- n check.

References

- [1] N. Levinson, *More than one third of zeros of Riemann's zeta-function are on $\sigma = \frac{1}{2}$* , Adv. Math. **13** (1974), 383–436. doi:10.1016/0001-8708(74)90074-7.
- [2] J. B. Conrey, *More than two fifths of the zeros of the Riemann zeta function are on the critical line*, J. Reine Angew. Math. **399** (1989), 1–26. doi:10.1515/crll.1989.399.1.
- [3] H. M. Bui, J. B. Conrey, M. P. Young, *More than 41% of the zeros of the zeta function are on the critical line*, Acta Arith. **150** (2011), 35–64. doi:10.4064/aa150-1-3.
- [4] S. Feng, *Zeros of the Riemann zeta function on the critical line*, J. Number Theory **132** (2012), 511–542. arXiv:1003.0059.
- [5] K. Pratt, N. Robles, A. Zaharescu, D. Zeindler, *More than five-twelfths of the zeros of ζ are on the critical line*, Res. Math. Sci. **7** (2020). doi:10.1007/s40687-019-0199-8.
- [6] J. B. Keiper, *Power series expansions of Riemann's ξ function*, Math. Comp. **58** (1992), 765–773. doi:10.1090/s0025-5718-1992-1122072-5.
- [7] X.-J. Li, *The positivity of a sequence of numbers and the Riemann hypothesis*, J. Number Theory **65** (1997), 325–333. doi:10.1006/jnth.1997.2137.




Figure 3: Third moment m_3 (left) and marked- T (right) against block size N over 10^7 zeros.