

Spread of Staph Infections in Hospitals

Valentina Miron Bogdan Moldoveanu Vlad Temian

West University of Timisoara

Tuesday 19th January, 2016

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

What is MRSA

- ▶ Methicillin-resistant *Staphylococcus aureus* (MRSA) infection is caused by a type of staph bacteria that's become resistant to many of the antibiotics used to treat ordinary staph infections.

What is MRSA

- ▶ Methicillin-resistant *Staphylococcus aureus* (MRSA) infection is caused by a type of staph bacteria that's become resistant to many of the antibiotics used to treat ordinary staph infections.
- ▶ It is spread by contact.

What is MRSA

- ▶ Methicillin-resistant *Staphylococcus aureus* (MRSA) infection is caused by a type of staph bacteria that's become resistant to many of the antibiotics used to treat ordinary staph infections.
- ▶ It is spread by contact.
- ▶ It is carried by about 2% of the population (or 2 in 100 people), although most of them aren't infected.

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

HA-MRSA

- ▶ Most MRSA infections occur in people who've been in hospitals or other health care settings, such as nursing homes and dialysis centers.

HA-MRSA

- ▶ Most MRSA infections occur in people who've been in hospitals or other health care settings, such as nursing homes and dialysis centers.
- ▶ When it occurs in these settings, it's known as health care-associated MRSA (HA-MRSA).

HA-MRSA

- ▶ Most MRSA infections occur in people who've been in hospitals or other health care settings, such as nursing homes and dialysis centers.
- ▶ When it occurs in these settings, it's known as health care-associated MRSA (HA-MRSA).
- ▶ HA-MRSA infections typically are associated with invasive procedures or devices, such as surgeries, intravenous tubing or artificial joints.

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

CA-MRSA

- ▶ CA is a type of MRSA infection that occurred in the wider community among healthy people, often begins as a painful skin boil.

CA-MRSA

- ▶ CA is a type of MRSA infection that occurred in the wider community among healthy people, often begins as a painful skin boil.
- ▶ It's spread by skin-to-skin contact.

CA-MRSA

- ▶ CA is a type of MRSA infection that occurred in the wider community among healthy people, often begins as a painful skin boil.
- ▶ It's spread by skin-to-skin contact.
- ▶ At-risk populations include groups such as high school wrestlers, child care workers and people who live in crowded conditions.

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

CA-MRSA vs HA-MRSA

- ▶ As CA-MRSA appears in the community, it is inevitably being spread into hospitals.

CA-MRSA vs HA-MRSA

- ▶ As CA-MRSA appears in the community, it is inevitably being spread into hospitals.
- ▶ Some studies suggest that CA-MRSA will overtake HA-MRSA in the hospital

CA-MRSA vs HA-MRSA

- ▶ As CA-MRSA appears in the community, it is inevitably being spread into hospitals.
- ▶ Some studies suggest that CA-MRSA will overtake HA-MRSA in the hospital




Figure: Site of

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

Building our model

Patient groups

$$K(t) \quad C(t) \quad S(t)$$

Building our model

Patient groups

$$K(t) \quad C(t) \quad S(t)$$

Model's parameters

$$\beta_C \quad \beta_H$$

Building our model

Patient groups

$$K(t) \quad C(t) \quad S(t)$$

Model's parameters

$$\beta_C \quad \beta_H$$

$$\delta_C \quad \delta_H \quad \delta_S$$

Building our model

Patient groups

$$K(t) \quad C(t) \quad S(t)$$

Model's parameters

$$\beta_C \quad \beta_H$$

$$\delta_C \quad \delta_H \quad \delta_S$$

$$\alpha_S \quad \alpha_H$$

Building our model

Patient groups

$$K(t) \quad C(t) \quad S(t)$$

Model's parameters

$$\beta_C \quad \beta_H$$

$$\delta_C \quad \delta_H \quad \delta_S$$

$$\alpha_S \quad \alpha_H$$

$$N \quad \Lambda$$

Building our model

Patient groups

$$K(t) \quad C(t) \quad S(t)$$

Model's parameters

$$\beta_C \quad \beta_H$$

$$\delta_C \quad \delta_H \quad \delta_S$$

$$\alpha_S \quad \alpha_H$$

$$N \quad \Lambda$$

Building our model

Patients movements

Patients move between compartments as they become colonized or decolonized

Building our model

Patients movements

Patients move between compartments as they become colonized or decolonized




Figure: A diagram of how patients transit between the compartments

Building our model

Patients movements

Patients move between compartments as they become colonized or decolonized




Figure: A diagram of how patients transit between the compartments

SIS (susceptible-infected-susceptible). Patients with no immunity.

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

Transition between states

$$\begin{aligned} \frac{dS}{dt} = & \underbrace{\Lambda}_{\text{entrance rate}} - \underbrace{\frac{\beta_H S(t) H(t)}{N}}_{\text{acquire HA-MRSA}} - \underbrace{\frac{\beta_H S(t) C(t)}{N}}_{\text{acquire CA-MRSA}} + \\ & \underbrace{\alpha_H H(t)}_{\text{HA-MRSA decolonized}} + \underbrace{\alpha_C C(t)}_{\text{CA-MRSA decolonized}} - \underbrace{\delta_S S(t)}_{\text{exit hospital}} \end{aligned} \quad (1)$$

Transition between states

$$\frac{dH}{dt} = \underbrace{\frac{\beta_H S(t) H(t)}{N}}_{\text{from S}} - \underbrace{\alpha_H H(t)}_{\text{decolonized}} - \underbrace{\delta_H H(t)}_{\text{exit hospital}} \quad (2)$$

Transition between states

$$\frac{dC}{dt} = \underbrace{\frac{\beta_H S(t) C(t)}{N}}_{\text{from S}} - \underbrace{\alpha_C C(t)}_{\text{decolonized}} - \underbrace{\delta_C C(t)}_{\text{exit hospital}} \quad (3)$$

Simplyfing our model

If we assume that the hospital is always full, we can conserve the system by letting $\Lambda = \delta_S S(t) + \delta_H h(t) + \delta_C C(t)$

Simplyfing our model

If we assume that the hospital is always full, we can conserve the system by letting $\Lambda = \delta_S S(t) + \delta_H h(t) + \delta_C C(t)$

In this case $S(t) + C(t) + H(t) = N$ for all t (assuming you start with population of size N)

$$\frac{dH}{dt} = (\beta_H/N)(N - C - H)H - (\beta_H + \alpha_H)H$$

Simplyfing our model

If we assume that the hospital is always full, we can conserve the system by letting $\Lambda = \delta_S S(t) + \delta_H h(t) + \delta_C C(t)$

In this case $S(t) + C(t) + H(t) = N$ for all t (assuming you start with population of size N)

$$\frac{dH}{dt} = (\beta_H/N)(N - C - H)H - (\beta_H + \alpha_H)H$$

$$\frac{dC}{dt} = (\beta_C/N)(N - C - H)C - (\beta_C + \alpha_C)C$$

Simplyfing our model

If we assume that the hospital is always full, we can conserve the system by letting $\Lambda = \delta_S S(t) + \delta_H h(t) + \delta_C C(t)$

In this case $S(t) + C(t) + H(t) = N$ for all t (assuming you start with population of size N)

$$\frac{dH}{dt} = (\beta_H/N)(N - C - H)H - (\beta_H + \alpha_H)H$$

$$\frac{dC}{dt} = (\beta_C/N)(N - C - H)C - (\beta_C + \alpha_C)C$$

Overview

Introduction

What is MRSA?

HA-MRSA

CA-MRSA

CA-MRSA vs HA-MRSA

Answer

Building our model

Defining the model

Explore

Explore

| Parameter | Symbol | Baseline Value |
|---|-------------|----------------|
| Total nutrients | N | 400 |
| <i>Length of stay</i> | | |
| Susceptible | $1/\beta_S$ | 5 days |
| Colonized CA-MRSA | $1/\beta_C$ | 7 days |
| Colonized HA-MRSA | $1/\beta_H$ | 5 days |
| <i>Transmission rate per susceptible patient to</i> | | |
| Colonized CA-MRSA per colonized CA-MRSA | β_C | 0.45 per day |
| Colonized HA-MRSA per colonized HA-MRSA | β_H | 0.4 per day |
| <i>Decolonization rate per colonized patient per day per length of stay</i> | | |
| CA-MRSA | α_C | 0.1 per day |
| HA-MRSA | α_H | 0.1 per day |

Table: Parameter values obtained from the Beth Israel Deaconess Medical Center

Critical Points

Using these values we'll get a system of equations and we'll find out the critical points

Critical Points

Using these values we'll get a system of equations and we'll find out the critical points

$$\begin{cases} \frac{0.4H(400 - H - C)}{400} - \left(\frac{1}{5} + 0.1\right)H = 0 \\ \frac{0.45C(400 - H - C)}{400} - \left(\frac{1}{7} + 0.1\right)C = 0 \end{cases}$$

Critical Points

Using these values we'll get a system of equations and we'll find out the critical points

$$\begin{cases} \frac{0.4H(400 - H - C)}{400} - \left(\frac{1}{5} + 0.1\right)H = 0 \\ \frac{0.45C(400 - H - C)}{400} - \left(\frac{1}{7} + 0.1\right)C = 0 \end{cases}$$

$$\begin{cases} H = 0 & C = 0 \\ H = 100 & C = 0 \\ H = 0 & C = 184.127 \end{cases}$$

Interpreting results




Figure: Direction field for the equation system

Interpreting results

We can see that we have an asymptotically stable point $\{H = 0, C = 184.127\}$. Based on the direction field, we can observe that this node it's also a spiral on.

Interpreting results

We can see that we have an asymptotically stable point $\{H = 0, C = 184.127\}$. Based on the direction field, we can observe that this node it's also a spiral on.

The other two nodes, $\{H = 0, C = 0\}$ and $\{H = 100, C = 0\}$ seems to be unstable points since all the arrows are getting away from them.

Given enough time, any point from this plane will converge to $\{H = 0, C = 184.127\}$, meaning that, using this model, CA-MRSA will overtake HA-MRSA.

Bibliography I



R.K. Nagle, E.B. Snaff, A.D. Sieder

Fundamentals of Differential Equations and Boundary Value Problems.

Addison-Wesley, 2012.



S. Balint, L. Braescu, E. Kaslik

Ordinary and Partial Differential Equations Lecture Notes
West University of Timisoara, 2006. 2000.