

Lecture 3

Linear Maps: Kernel, Image, Rank–Nullity

Throughout, $\mathbb{F}$ is $\mathbb{R}$ or $\mathbb{C}$, and $\mathcal{L}(V, W)$ denotes the linear maps $V \rightarrow W$. Tags indicate the flavour of each problem; a ★ marks a harder one.

Core tutorial route

Work **3.1, 3.4, 3.8, 3.9, 3.15, 3.19, 3.20, and 3.24** in that order. This route checks linearity and basis-defined maps, a complete kernel/image calculation, an image proof, rank–nullity and dimension obstructions, projection structure, and the finite injective–surjective transfer.

Additional practice: 3.2, 3.5, 3.7, 3.10–3.12, 3.14, 3.17–3.18, 3.21–3.23, 3.26. **Bridge:** 3.3, 3.6, 3.13, 3.16 (calculus applications). **Extension:** 3.25 and 3.27–3.34 (quotients and the First Isomorphism Theorem).

Linear maps

Exercise 3.1 (computational) Which of the following maps $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ (or $\mathbb{R}^2 \rightarrow \mathbb{R}$) are linear?

$$T(x, y) = (x - y, 2x), \quad S(x, y) = (xy, x), \quad R(x, y) = |x| + y.$$

Answer. Only T is linear; S and R are not.

Exercise 3.2 (computational) Define $T \in \mathcal{L}(\mathbb{R}^2, \mathbb{R}^3)$ by $T(1, 0) = (1, 2, 3)$ and $T(0, 1) = (0, 1, -1)$. Find a formula for $T(x, y)$, and determine $\text{rank } T$ and $\dim \ker T$.

Answer. $T(x, y) = (x, 2x + y, 3x - y)$; $\text{rank } T = 2$, $\dim \ker T = 0$.

Exercise 3.3 (computational) On $\mathcal{P}(\mathbb{R})$ let D be differentiation and M multiplication by t . Compute the commutator $[D, M^2] = DM^2 - M^2D$ as an operator.

Answer. $[D, M^2] = 2M$.

Exercise 3.4 (computational) Let $T \in \mathcal{L}(\mathbb{R}^2, \mathbb{R}^2)$ satisfy $T(1, 1) = (3, 1)$ and $T(1, -1) = (1, 3)$. Find a formula for $T(x, y)$.

Answer. $T(x, y) = (2x + y, 2x - y)$.

Exercise 3.5 (computational) Which of the following maps $\mathbb{R}^3 \rightarrow \mathbb{R}^2$ are linear?

$$F(x, y, z) = (x + z, 2y), \quad G(x, y, z) = (x, yz), \quad H(x, y, z) = (x - y, 0).$$

Answer. F and H are linear; G is not.

Exercise 3.6 (computational) On $\mathcal{P}(\mathbb{R})$ let D be differentiation and M multiplication by t . Compute the commutator $[D, M^3] = DM^3 - M^3D$ as an operator.

Answer. $[D, M^3] = 3M^2$.

Exercise 3.7 (proof) Fix $v_0 \in V$. Prove that the evaluation map $E: \mathcal{L}(V, W) \rightarrow W$ defined by $E(T) = Tv_0$ is linear.

Hint. The operations on $\mathcal{L}(V, W)$ are pointwise; write out $(S + T)v_0$ and $(aT)v_0$ and there is nothing left to do.

Kernel and image

Exercise 3.8 (computational) Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 2 & 1 \end{pmatrix}$. Find bases for $\ker T$ and $\operatorname{im} T$, and verify the rank–nullity theorem.

Answer. $\ker T = \operatorname{span}((-1, 1, -1))$ (dim 1); $\operatorname{im} T = \operatorname{span}((1, 0, 1), (1, 1, 2))$ (dim 2); $3 = 2 + 1$.

Exercise 3.9 (proof) Suppose $v_1, \dots, v_n$ is a basis of V and $T \in \mathcal{L}(V, W)$. Prove that $\operatorname{im} T = \operatorname{span}(Tv_1, \dots, Tv_n)$.

Hint. One inclusion is immediate. For the other, expand v in the basis and push T through the sum.

Exercise 3.10 (computational) Let $\varphi: \mathbb{C}^3 \rightarrow \mathbb{C}$ be $\varphi(z_1, z_2, z_3) = z_1 + 2z_2 + 3z_3$. Describe $\ker \varphi$ and its dimension, and decide whether φ is surjective.

Answer. $\dim \ker \varphi = 2$; φ is surjective.

Exercise 3.11 (computational) Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{pmatrix}$. Find bases for $\ker T$ and $\operatorname{im} T$, and verify rank–nullity.

Answer. $\ker T = \operatorname{span}((-1, -1, 1))$ (dim 1); $\operatorname{im} T = \operatorname{span}((1, 0, 1), (2, 1, 1))$ (dim 2); $3 = 2 + 1$.

Exercise 3.12 (computational) Let $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 1 & 3 \\ 1 & 1 & 3 & 4 \end{pmatrix}$. Find bases for $\ker T$ and $\operatorname{im} T$.

Answer. $\ker T = \operatorname{span}((-2, -1, 1, 0), (-1, -3, 0, 1))$ (dim 2); $\operatorname{im} T = \operatorname{span}((1, 0, 1), (0, 1, 1))$ (dim 2).

Exercise 3.13 (computational) Let $T: \mathcal{P}_3(\mathbb{R}) \rightarrow \mathbb{R}^2$ be $Tp = (p(0), p'(0))$. Find $\ker T$ and $\operatorname{rank} T$.

Answer. $\ker T = \operatorname{span}(t^2, t^3)$ (dim 2); $\operatorname{rank} T = 2$ (so T is surjective).

Exercise 3.14 (proof) Let $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$. Prove that $\operatorname{im}(ST) \subseteq \operatorname{im} S$ and $\ker T \subseteq \ker(ST)$.

Hint. Both are one line: $w = (ST)u$ is $S(Tu)$, and $Tv = 0$ makes $(ST)v = S0$.

Rank–nullity

Exercise 3.15 (computational) Use the rank–nullity theorem to fill in each blank. (a) $T \in \mathcal{L}(\mathbb{R}^4, \mathbb{R}^3)$ with $\dim \ker T = 2$: find $\operatorname{rank} T$. (b) $T \in \mathcal{L}(\mathbb{R}^6, \mathbb{R}^2)$ surjective: find $\dim \ker T$. (c) $T \in \mathcal{L}(\mathcal{P}_4(\mathbb{R}), \mathbb{R}^3)$ with $\operatorname{rank} T = 3$: find $\dim \ker T$.

Answer. (a) $\operatorname{rank} T = 2$. (b) $\dim \ker T = 4$. (c) $\dim \ker T = 2$.

Exercise 3.16 (computational) Let D^2 be the second-derivative operator on $\mathcal{P}_4(\mathbb{R})$. Find $\dim \ker(D^2)$ and $\operatorname{rank}(D^2)$.

Answer. $\dim \ker(D^2) = 2$ and $\operatorname{rank}(D^2) = 3$.

Exercise 3.17 (computational) Let $T: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be $Tx = Ax$ with $A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$. Find $\operatorname{rank} T$ and a basis for $\ker T$.

Answer. $\operatorname{rank} T = 2$; $\ker T = \operatorname{span}((-1, 1, 0, 0), (0, 0, -1, 1))$ (dim 2).

Exercise 3.18 (proof) Suppose V is finite-dimensional and $T \in \mathcal{L}(V)$ satisfies $T^2 = 0$. Prove $\operatorname{im} T \subseteq \ker T$, and deduce $\dim \operatorname{im} T \leq \frac{1}{2} \dim V$.

Hint. What does $T^2 = 0$ tell you about Tv for each v ? Combine that containment with rank-nullity.

Exercise 3.19 (proof) Suppose $P \in \mathcal{L}(V)$ satisfies $P^2 = P$ (a *projection*). Prove that $V = \ker P \oplus \operatorname{im} P$.

Hint. Write $v = (v - Pv) + Pv$ and check which summand lies where. For directness, take $w = Pu$ in the intersection and apply P once more.

Exercise 3.20 (★) Prove that every $T \in \mathcal{L}(\mathbb{F}^5, \mathbb{F}^2)$ has $\dim \ker T \geq 3$.

Hint. Rank-nullity, together with the largest the rank can be when the codomain is only $\mathbb{F}^2$.

Exercise 3.21 (proof) Suppose V is finite-dimensional and $S, T \in \mathcal{L}(V)$. Prove that $\operatorname{rank}(S + T) \leq \operatorname{rank} S + \operatorname{rank} T$.

Hint. Show $\operatorname{im}(S + T) \subseteq \operatorname{im} S + \operatorname{im} T$, then compare dimensions using $\dim(U_1 + U_2) \leq \dim U_1 + \dim U_2$.

Operators: injective versus surjective

Exercise 3.22 (computational) Is the operator $T \in \mathcal{L}(\mathbb{R}^3)$ given by $Tx = Ax$ with $A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ invertible?

Answer. Yes; $\ker T = \{0\}$, so T is invertible.

Exercise 3.23 (computational) Is the operator $T \in \mathcal{L}(\mathbb{R}^3)$ given by $Tx = Ax$ with $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 1 & 1 & 1 \end{pmatrix}$ invertible? If not, give $\dim \ker T$.

Answer. Not invertible; $\dim \ker T = 1$.

Exercise 3.24 (proof) Suppose V, W are finite-dimensional with $\dim V = \dim W$, and $T \in \mathcal{L}(V, W)$ is injective. Prove that T is surjective.

Hint. Injectivity fixes $\dim \ker T$; rank-nullity then fixes $\dim \operatorname{im} T$, and a subspace of W of that dimension is all of W .

Exercise 3.25 (★) Suppose V is finite-dimensional and $S, T \in \mathcal{L}(V)$ satisfy $ST = I$. Prove that $TS = I$ as well (so a one-sided inverse of an operator is automatically two-sided).

Hint. First show T is injective: if $Tv = 0$, apply S . In finite dimensions an injective operator is invertible.

Exercise 3.26 (proof) Suppose V is finite-dimensional and $T \in \mathcal{L}(V)$ is surjective. Prove that T is injective.

Hint. Surjectivity fixes $\operatorname{rank} T$; read $\dim \ker T$ off rank-nullity.

Quotient spaces [self-study]

Exercise 3.27 (computational) Let $U = \operatorname{span}((1, 1, 0)) \subseteq \mathbb{R}^3$. Describe the coset of $(2, 0, 1)$ modulo U , and determine $\dim(\mathbb{R}^3/U)$.

Answer. $(2, 0, 1) + U = \{(2 + t, t, 1) : t \in \mathbb{R}\}$; $\dim(\mathbb{R}^3/U) = 2$.

Exercise 3.28 (proof) Prove that the quotient-space operations in the lecture notes are well-defined: if $v + U = v' + U$ and $w + U = w' + U$, show $(v + w) + U = (v' + w') + U$; also show that $v + U = v' + U$ implies $av + U = av' + U$ for every scalar a .

Hint. $v + U = v' + U$ says exactly that $v - v' \in U$. Everything after that is closure of U under addition and scaling.

Exercise 3.29 (computational) Let $T: \mathbb{R}^4 \rightarrow \mathbb{R}^2$ be $T(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_3 - x_4)$. (a) Find $\ker T$. (b) Write the First Isomorphism Theorem for T explicitly: what is the isomorphism $\tilde{T}: \mathbb{R}^4/\ker T \rightarrow \operatorname{im} T$?

Answer. (a) $\dim \ker T = 2$, with basis $\{(-1, 1, 0, 0), (0, 0, 1, 1)\}$. (b) $\operatorname{im} T = \mathbb{R}^2$, hence $\tilde{T}: \mathbb{R}^4/\ker T \xrightarrow{\cong} \mathbb{R}^2$.

Exercise 3.30 (computational) Let $U = \operatorname{span}((1, 1, 1)) \subseteq \mathbb{R}^3$. Describe the coset of $(2, 0, 1)$ modulo U , and determine $\dim(\mathbb{R}^3/U)$.

Answer. $(2, 0, 1) + U = \{(2 + t, t, 1 + t) : t \in \mathbb{R}\}$; $\dim(\mathbb{R}^3/U) = 2$.

Exercise 3.31 (computational) Let $U = \operatorname{span}((1, 0, 1, 0), (0, 1, 0, 1)) \subseteq \mathbb{R}^4$. Do $(1, 1, 1, 1)$ and the zero vector lie in the same coset modulo U ? Find $\dim(\mathbb{R}^4/U)$.

Answer. Yes (since $(1, 1, 1, 1) \in U$); $\dim(\mathbb{R}^4/U) = 2$.

Exercise 3.32 (computational) Let D be differentiation on $\mathcal{P}_3(\mathbb{R})$. Identify $\ker D$ and the isomorphism $\mathcal{P}_3(\mathbb{R})/\ker D \cong \operatorname{im} D$ furnished by the First Isomorphism Theorem.

Answer. $\ker D = \operatorname{span}(1)$ (the constants); $\mathcal{P}_3(\mathbb{R})/\ker D \cong \mathcal{P}_2(\mathbb{R})$, of dimension 3.

Exercise 3.33 (★) Let $T \in \mathcal{L}(V, W)$. Prove that $\tilde{T}: V/\ker T \rightarrow W$, defined by $\tilde{T}(v + \ker T) = Tv$, is well-defined, linear, and injective.

Hint. For well-definedness, replace v by another representative of its coset and check the value is unchanged; for injectivity, see what $\tilde{T}(v + \ker T) = 0$ forces.

Exercise 3.34 (proof) Let U be a subspace of a finite-dimensional space V . Prove that the quotient map $\pi: V \rightarrow V/U$, $\pi(v) = v + U$, is linear and surjective with $\ker \pi = U$, and deduce $\dim(V/U) = \dim V - \dim U$.

Hint. Linearity is the definition of the quotient operations. For the kernel, note $v + U = U$ holds iff $v \in U$; then apply rank-nullity to π itself.