

HSC Mathematics Extension 1: Combinatorics Practice

Vu Hung Nguyen



Download PDF and resources:

<https://github.com/vuhung16au/math-olympiad-ml>

Mathematics may be defined as the economy of counting. There is no problem in the whole of mathematics which cannot be solved by direct counting.

Ernst Mach

Contents

1	Introduction	6
1.1	Project Overview	6
1.2	Target Audience	6
1.3	How to Use This Booklet	6
1.4	Extension 1 Knowledge Needed	6
1.5	Core Formulas and Ideas	7
1.6	Pascal's Triangle	9
1.7	The Binomial Theorem	9
1.8	Binomial Probability	10
1.9	A Brief Note: The Pigeonhole Principle and Remainders	10
1.10	A Brief Extension: Gamma and Beta Functions	11
1.11	A Brief Extension: The Inclusion-Exclusion Principle	11
1.12	A Brief Extension: Stars and Bars	12
1.13	A Brief Extension: Algorithmic Counting and Complexity	12
1.14	Problem-Solving Habits for This Booklet	13
2	Part 1: Worked Problems	13
2.1	Basic	13
	Problem 2.1: Four-digit numbers from 1 to 8	13
	Problem 2.2: Rearranging the letters of COOKED	14
	Problem 2.3: Choosing three coloured balls	14
	Problem 2.4: Rearranging the letters of CONDOBOLIN	15
2.2	Medium	15
	Problem 2.5: Seating around a circle	15
	Problem 2.6: Selections from URGOATED	16
	Problem 2.7: Forming a Dota 2 e-sport team	16
	Problem 2.8: Triangles from points on the sides of a triangle	17
	Problem 2.9: Age-limit check across teams	18
	Problem 2.10: Expected score with replacement	19
	Problem 2.11: Minimum votes needed to win	20
	Problem 2.12: Selecting finalists and awarding places	21
	Problem 2.13: Using the pigeonhole principle with topic choices	21
2.3	Advanced	22
	Problem 2.14: Committee selection	22
	Problem 2.15: Pascal's triangle and a telescoping sum	22
	Problem 2.16: Combining binomial coefficients	24
	Problem 2.17: Equal numbers of men and women	25
	Problem 2.18: A binomial identity and double counting capstone	26
	Problem 2.19: Students and teachers in a circle	28
	Problem 2.20: Guests who refuse to sit together	28
	Problem 2.21: Australian Powerball counting	29
	Problem 2.22: License plates and careful reading	30
	Problem 2.23: A mixed combinatorics review	31
3	Part 2: Practice and Challenge Bank	32
3.1	Warm-Up Drills	32
	Problem 3.1: Expanding a binomial surd expression	32
	Problem 3.2: When is a conjugate surd sum rational?	32
	Problem 3.3: PINs with numbers and letters	33

Problem 3.4: Two coins of the same metal	34
Problem 3.5: Probabilities with the four jacks	35
Problem 3.6: Counting rectangles in a 5×6 grid	35
Problem 3.7: How fast does brute force grow?	36
Problem 3.8: Repeated derivatives of $\frac{1}{x}$	37
Problem 3.9: Using Pascal's triangle to differentiate x^4	38
3.2 Stretch Problems	40
Problem 3.10: Term independent of x	40
Problem 3.11: Five-letter words from Mamungkukumpurangkuntjunya	40
Problem 3.12: Constant term from a product of two expansions	41
Problem 3.13: Adults and children at the cinema	42
Problem 3.14: The alternating row constraint	43
Problem 3.15: Lines determined by points on a line and a circle	45
Problem 3.16: Dota knockout bracket	46
Problem 3.17: Dividing a class into equal groups	47
Problem 3.18: Triangles from points on a circle	48
Problem 3.19: Triangles from a regular octagon and its centre	49
Problem 3.20: Pigeons and pigeonholes	50
3.3 Challenge Corner	51
Problem 3.21: Comparing coefficients in $(1+x)^{m+n}$	51
Problem 3.22: Grid paths and binomial identities	53
Problem 3.23: A coefficient identity for $\binom{4n}{2n}$	55
Problem 3.24: Repeated draws of green counters	57
Problem 3.25: Arrangement formulas for x people	58
Problem 3.26: A pair summing to 21	59
Problem 3.27: Positive integer solutions	59
Problem 3.28: Non-negative integer solutions	60
Problem 3.29: Three identities from $(1+x)^n$	61
Problem 3.30: Making an amount of money with Australian denominations	63
Problem 3.31: Unlocking a phone by trial and error	65
Problem 3.32: Polynomial derivatives and a power-series expansion	67
Problem 3.33: Binary strings and a binomial identity	69
Problem 3.34: Derangements	71
Problem 3.35: A monochromatic triangle on six vertices	74
Problem 3.36: A more general form of the binomial theorem	76
Problem 3.37: Isomers — a possible computing project	78
4 Conclusion	80
A Appendices	81
A.1 Concepts: Injective, Surjective, and Bijective	81

1 Introduction

1.1 Project Overview

This booklet is a growing collection of combinatorics problems for NSW HSC Mathematics Extension 1 students. It now covers a wider range of syllabus ideas including permutations, combinations, Pascal’s triangle, circular arrangements, committee selection, simple probability, expected value, the pigeonhole principle, geometric counting, and proof-style binomial identities.

1.2 Target Audience

This resource is written for students who want extra practice with combinatorics in a clear, classroom-friendly format. The solutions are intentionally brief, but each one points out the key counting idea behind the answer.

1.3 How to Use This Booklet

- Read the short summary of the core formulas first.
- Attempt each problem before reading the solution.
- Pay attention to whether each question is a selection or an arrangement.
- Look for structure: gaps in circles, repeated objects, complementary counting, or balanced groups.
- Use the takeaways to build pattern recognition for future HSC questions.

1.4 Extension 1 Knowledge Needed

Most problems in this booklet can be solved using standard HSC Mathematics Extension 1 ideas rather than advanced university mathematics. The key background knowledge is:

- **Factorials and notation:** understanding $n!$, permutation notation nP_r , and combination notation $\binom{n}{r}$. For a positive integer n ,

$$n! = n(n-1)(n-2) \cdots 2 \cdot 1.$$

Also,

$$0! = 1, \quad 1! = 1, \quad 2! = 2, \quad 3! = 6, \quad 4! = 24, \quad 5! = 120.$$

Here $\binom{n}{r}$ means “the number of ways to choose r objects from n objects”, where order does not matter.

- **The multiplication principle:** breaking a problem into stages and multiplying the number of choices at each stage.
- **Order versus selection:** recognising when order matters (permutations) and when it does not (combinations).
- **Repeated objects:** dividing by repeated factorials when arranging words such as COOKED or CONDOBOLIN.

- **Circular permutations:** using $(n - 1)!$ for arrangements around a circle and handling restrictions by fixing seats, using blocks, or using gaps.
- **Complementary counting:** counting all cases first, then subtracting the unwanted ones.
- **Casework and restrictions:** splitting problems into clean cases such as fixed numbers of girls and boys, men and women, or chosen leaders.
- **Pascal's triangle and binomial coefficients:** using identities such as

$$\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r}$$

and the symmetry rule

$$\binom{n}{r} = \binom{n}{n-r}.$$

- **The binomial theorem:** expanding $(1 + x)^n$ and comparing coefficients when proving identities.
- **Pigeonhole principle reasoning:** proving that some outcome must happen because the total is too large to avoid repetition or overload.
- **Basic probability and expectation:** counting favourable outcomes, dividing by total outcomes, and using expected value as an average score or average outcome.
- **Double counting:** counting the same set of outcomes in two different ways to derive a useful identity.

1.5 Core Formulas and Ideas

Fundamental Counting Principle. If a task is completed in stages with a choices for the first stage, b for the second, and c for the third, then the total number of outcomes is

$$a \times b \times c.$$

Generalized counting principle. Suppose that k experiments are to be performed, with the number of possible outcomes being n_i for the i th experiment. Then there are

$$n_1 \cdot n_2 \cdot \dots \cdot n_k$$

possible outcomes of all k experiments together.

What is a set? A **set** is a collection of distinct objects. In a set, order does not matter, and repeated copies are not listed separately. For example,

$$\{1, 2, 3\} = \{3, 2, 1\},$$

and writing $\{1, 1, 2\}$ does not create a new 3-element set; as a set, it is just $\{1, 2\}$.

In counting problems, this usually means:

- the objects in the set are treated as distinct unless the question says some are equal or repeated
- repetition is *not* automatically allowed just because we are arranging or choosing from a set
- if repetition *is* allowed, the question should say so explicitly.

Permutations versus combinations. A **permutation** is an arrangement, so order matters. A **combination** is a selection, so order does not matter.

Worked example. Suppose 3 students are Alice, Ben, and Chloe.

- If we choose a captain and vice-captain, order matters, so this is a permutation:

$${}^3P_2 = 3 \times 2 = 6.$$

For example, “Alice then Ben” is different from “Ben then Alice”.

- If we simply choose a team of 2, order does not matter, so this is a combination:

$$\binom{3}{2} = 3.$$

The team {Alice, Ben} is the same team as {Ben, Alice}.

Permutations without repetition. The number of ways to arrange r objects chosen from n distinct objects is

$${}^nP_r = \frac{n!}{(n-r)!}.$$

Combinations. The number of ways to choose r objects from n distinct objects is

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}.$$

Multinomial coefficients. If there are n objects in total, divided into groups of sizes $a, b, c, \dots$ with

$$n = a + b + c + \dots,$$

then the number of distinct arrangements is

$$\frac{n!}{a!b!c!\dots}.$$

Multinomial expansion note. In the expansion of

$$(x_1 + x_2 + \dots + x_k)^m,$$

the coefficient of a term

$$x_1^{a_1} x_2^{a_2} \dots x_k^{a_k}$$

with

$$a_1 + a_2 + \dots + a_k = m$$

is the multinomial coefficient

$$\frac{m!}{a_1!a_2!\dots a_k!}.$$

Circular permutations. The number of ways to arrange n distinct objects around a circle is

$$(n-1)!.$$

Pascal’s triangle identity.

$$\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r}.$$


Binomial symmetry.

$$\binom{n}{r} = \binom{n}{n-r}.$$

1.6 Pascal's Triangle

Pascal's triangle is a triangular arrangement of binomial coefficients. For HSC Extension 1, it is useful both for quick coefficient reading and for proving combinatorial identities.

For $n = 0, 1, 2, \dots, 9$ (10 rows), Pascal's triangle can be shown with stacked hexagons:



Three facts to remember:

- **Pascal rule:**

$$\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r}.$$

Each interior entry is the sum of the two entries above it.

- **Symmetry:**

$$\binom{n}{r} = \binom{n}{n-r}.$$

Each row reads the same from left to right and from right to left.

- **Row sum:**

$$\binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{n} = 2^n.$$

The sum of the entries in row n is a power of 2.

1.7 The Binomial Theorem

For a positive integer n ,

$$(x + y)^n = \binom{n}{0}x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \dots + \binom{n}{n}y^n.$$

This theorem tells us how to expand powers such as $(x + y)^n$. The coefficients come directly from row n of Pascal's triangle.

Worked example. Expand $(x + 2)^4$.

Using row 4 of Pascal's triangle, the coefficients are 1, 4, 6, 4, 1:

$$\begin{aligned} (x + 2)^4 &= \binom{4}{0}x^4 + \binom{4}{1}x^3(2) + \binom{4}{2}x^2(2^2) + \binom{4}{3}x(2^3) + \binom{4}{4}(2^4) \\ &= x^4 + 4x^3(2) + 6x^2(4) + 4x(8) + 16 \\ &= x^4 + 8x^3 + 24x^2 + 32x + 16. \end{aligned}$$

This is especially useful later in the booklet when proving identities by comparing coefficients of matching powers of x .

Expected value. If a random variable takes values $x_1, x_2, \dots$ with probabilities $p_1, p_2, \dots$, then

$$E(X) = x_1p_1 + x_2p_2 + \dots$$

1.8 Binomial Probability

Binomial probability is used when the same trial is repeated a fixed number of times and each trial has only two possible outcomes, often called *success* and *failure*.

The main features are:

- there are only two possible outcomes on each trial
- the probability of success is the same on every trial
- the number of trials is fixed
- the trials are independent

If X is the number of successes in n trials, and the probability of success on each trial is p , then

$$P(X = r) = \binom{n}{r} p^r (1 - p)^{n-r}.$$

Here $\binom{n}{r}$ counts the number of ways to arrange the r successes among the n trials.

Example. A green counter is selected with probability 0.2 on each draw, and the draw is repeated 5 times with replacement. The probability of getting exactly 2 green counters is

$$P(X = 2) = \binom{5}{2} (0.2)^2 (0.8)^3.$$

This idea appears in repeated-draw questions later in the booklet.

1.9 A Brief Note: The Pigeonhole Principle and Remainders

Suppose that n objects are placed into d boxes. If we divide n by d , we can write

$$n = qd + r, \quad \text{where } 0 \leq r < d.$$

This helps us read the pigeonhole principle more precisely:

- if $r = 0$, then at least one box contains at least q objects
- if $r > 0$, then at least one box contains at least $q + 1$ objects.

In particular, the smallest number of objects that guarantees at least one box contains at least $q + 1$ objects is

$$qd + 1.$$

This point of view is especially useful in “must happen” questions, where the boxes may be teams, remainders after division, topic choices, or vote totals.

1.10 A Brief Extension: Gamma and Beta Functions

Although this booklet focuses on HSC Mathematics Extension 1, it is worth knowing that factorials and combinations have important extensions in higher mathematics.

Gamma function. The Gamma function is defined for suitable real or complex inputs by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt.$$

It satisfies the recurrence

$$\Gamma(z+1) = z\Gamma(z),$$

which mirrors the factorial rule $n! = n(n-1)!$. In fact,

$$\Gamma(n) = (n-1)!, \quad \text{so} \quad n! = \Gamma(n+1).$$

This means the Gamma function is the natural continuous extension of the factorial function.

Beta function. The Beta function is closely related to Gamma and is defined by

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

Since combinations are built from factorials,

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{\Gamma(n+1)}{\Gamma(r+1)\Gamma(n-r+1)}.$$

So Gamma and Beta functions provide a continuous version of the factorial and binomial-coefficient ideas that appear throughout combinatorics.

Why this matters. These functions form a bridge between discrete counting and continuous mathematics. They are useful in advanced probability, statistics, asymptotics, and areas such as machine learning.

1.11 A Brief Extension: The Inclusion-Exclusion Principle

The inclusion-exclusion principle is used when counting objects that may belong to overlapping sets.

For two sets A and B ,

$$|A \cup B| = |A| + |B| - |A \cap B|.$$


For three sets A , B , and C ,

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|.$$

The idea is simple: when we add the sizes of sets, elements in overlaps are counted more than once, so we must correct for this.

Also note that this is an important topic when you learn probability.

This principle is very useful in restricted counting problems, probability, and Venn diagram questions.



1.12 A Brief Extension: Stars and Bars

Stars and bars is a method for counting integer solutions to equations such as

$$x_1 + x_2 + \cdots + x_k = n.$$

The idea is to represent the total n by n stars and use $k - 1$ bars to divide the stars into k groups.

$$\begin{array}{ccccccc} * & * & | & * & * & * & | & * & * \\ 2 & | & 3 & | & 2 \end{array}$$

If the variables are required to be positive, then each group must contain at least one star, so the number of solutions is

$$\binom{n-1}{k-1}.$$

If the variables are allowed to be zero, then empty groups are allowed, so the number of solutions is

$$\binom{n+k-1}{k-1}.$$

Stars and bars is a standard combinatorial technique for turning algebraic equations into counting problems. It is directly connected to the two Part 2 challenge problems on positive and non-negative integer solutions.

Exam note. Recent HSC Mathematics Extension 1 exams do not usually ask this type of problem directly. However, it is still very good to know, because it is a powerful counting technique to keep as a mathematical “weapon” for olympiad-style problems, extension work, and future university study.

1.13 A Brief Extension: Algorithmic Counting and Complexity

Algorithmic counting is the study of methods for counting objects systematically. Complexity asks how difficult a counting problem becomes as the size of the problem grows.

Some counting problems have simple closed formulas, such as

$$\binom{n}{r}, \quad n!, \quad 2^n.$$

However, many counting problems grow extremely quickly and cannot be solved efficiently by checking every possibility one by one.

For example:

n	$n!$	2^n
5	120	32
10	3628800	1024
20	2432902008176640000	1048576

So even a simple arrangement problem can become enormous very quickly.

This is one reason why combinatorics is important: good counting methods, symmetry arguments, recurrences, and identities help us avoid brute force.

Syllabus note. This topic is outside the HSC Mathematics Extension 1 syllabus. However, it is very useful for higher education, especially in areas such as computer science, discrete mathematics, and algorithm design.

1.14 Problem-Solving Habits for This Booklet

- Ask first: *Does order matter?*
- Check whether repetition is allowed or forbidden.
- In circle problems, decide whether to use a fixed reference point, a block, or a gap method.
- In identity questions, look for Pascal's triangle, symmetry, coefficient matching, or telescoping.
- In probability questions, decide whether the experiment is with replacement or without replacement.
- In “must happen” questions, test whether the pigeonhole principle or a contradiction argument applies.

Remark 1.1

The most common combinatorics mistake is using a permutation when the problem is actually a combination, or vice versa. A good first question is always: *Does order matter?*

Remark 1.2 (How fast counting grows)

A 19×19 Go board has 361 points. If each point is allowed to be black, white, or empty, then a rough count of possible board states is

$$3^{361} \approx 10^{172}.$$

Even after restricting to legal positions, the number is still about 10^{170} , which is unimaginably large. This is one of the best reminders that counting problems can grow far beyond ordinary intuition.

2 Part 1: Worked Problems

2.1 Basic

Problem 2.1: Four-digit numbers from 1 to 8

How many 4-digit numbers can you make using the digits 1 to 8 if no digit is repeated?

Solution 2.1

There are 8 choices for the thousands digit, then 7 choices for the hundreds digit, 6 for the tens digit, and 5 for the units digit.

$$8 \times 7 \times 6 \times 5 = 1680$$

So the number of such 4-digit numbers is 1680.

Takeaways 2.1

- When order matters and repetition is not allowed, use a permutation count.
- The multiplication principle is often the fastest method for straightforward arrangement questions.

Problem 2.2: Rearranging the letters of COOKED

How many different arrangements of the letters in the word COOKED are possible?

Solution 2.2

The word COOKED has 6 letters in total, with:

O repeated 2 times.

If all letters were distinct, there would be $6!$ arrangements. We divide by $2!$ for the repeated O 's:

$$\frac{6!}{2!} = \frac{720}{2} = 360.$$

Hence the number of different arrangements is $\boxed{360}$.

Takeaways 2.2

- For repeated letters, start with the total factorial and divide by the factorial of each repeated group.
- This is the standard multiset permutation formula.

Problem 2.3: Choosing three coloured balls

A bag contains six different coloured balls.

How many different ways can three balls be selected?

Solution 2.3

Since the order of selection does not matter, we use combinations:

$$\binom{6}{3} = \frac{6 \cdot 5 \cdot 4}{3 \cdot 2 \cdot 1} = 20.$$

So the number of different selections is $\boxed{20}$.

Takeaways 2.3

- Use combinations when order does not matter.
- The phrase “select two balls” usually signals a combination, not a permutation.

Problem 2.4: Rearranging the letters of CONDOBOLIN

In how many different ways can all the letters of the word CONDOBOLIN be arranged in a line?

Solution 2.4

The word CONDOBOLIN has 10 letters in total, with:

O repeated 3 times, N repeated 2 times.

If all letters were distinct, there would be $10!$ arrangements. We divide by $3!$ for the repeated O 's and by $2!$ for the repeated N 's:

$$\frac{10!}{3!2!} = \frac{3628800}{12} = 302400.$$

Hence the number of different arrangements is $\boxed{302400}$.

Takeaways 2.4

- This is another multiset permutation problem: start with $n!$ and divide by repeated factorials.
- Always count repeated letters carefully before substituting into the formula.

2.2 Medium

Problem 2.5: Seating around a circle

How many different ways can 9 people be seated in a circle if 2 people have already chosen their seats?

Solution 2.5

Once the two chosen seats are fixed, those two people no longer move. We only need to arrange the remaining 7 people in the remaining 7 seats.

$$7! = 5040$$

Therefore, the number of possible seatings is $\boxed{5040}$.

Takeaways 2.5

- Circular arrangements usually reduce by one factor because rotations are equivalent.
- If some seats are already fixed, the circular symmetry has already been broken.

Problem 2.6: Selections from URGOATED

Five letters are chosen from the letters of the word URGOATED, without ordering them. How many selections are there if the choice:

- a) has no restriction?
- b) must contain all four vowels?
- c) must contain the letter R?
- d) must contain the letter U, but not D?

Solution 2.6

All 8 letters in URGOATED are distinct: U, R, G, O, A, T, E, D .

a) No restriction:

$$\binom{8}{5} = 56$$

b) Must contain all four vowels U, O, A, E : We must choose 1 more letter from the remaining 4 consonants R, G, T, D .

$$\binom{4}{1} = 4$$

c) Must contain R: Fix R, then choose 4 more letters from the remaining 7 letters.

$$\binom{7}{4} = 35$$

d) Must contain U, but not D: Fix U, exclude D, then choose 4 more letters from R, G, O, A, T, E .

$$\binom{6}{4} = 15$$

So the answers are:

56, 4, 35, 15

Takeaways 2.6

- Restriction questions are often easiest if you first force required letters in, or remove forbidden letters.
- After handling the restriction, finish with a simple combination count.

Problem 2.7: Forming a Dota 2 e-sport team

- a) How many different ways can a Dota 2 e-sport team of 5 students be chosen from a class of 30 students?
- b) The team must have 2 girls and 3 boys. If the class contains 12 girls and 18 boys, how many teams are possible?
- c) Suppose Vu and Zachary are both students in the class. Find the probability that both Vu and Zachary are chosen on the team.
- d) Find the probability that neither Vu nor Zachary is chosen.

Solution 2.7

a) Choose any 5 students from 30:

$$\binom{30}{5} = 142506$$

b) Choose 2 girls from 12 and 3 boys from 18:

$$\binom{12}{2} \binom{18}{3} = 66 \times 816 = 53856$$

c) If Vu and Zachary must both be chosen, then choose the remaining 3 team members from the other 28 students:

$$\frac{\binom{28}{3}}{\binom{30}{5}} = \frac{3276}{142506} = \frac{2}{87}$$

So the probability is $\frac{2}{87}$.

d) If neither Vu nor Zachary is chosen, then all 5 team members must come from the other 28 students:

$$\frac{\binom{28}{5}}{\binom{30}{5}} = \frac{98280}{142506} = \frac{460}{667}$$


So the probability is $\frac{460}{667}$.

Takeaways 2.7

- For probabilities in equally likely team selections, count favourable teams over total teams.
- “Both chosen” means fix those people first; “neither chosen” means remove them from the pool.

Problem 2.8: Triangles from points on the sides of a triangle

The diagram shows triangle ABC with points chosen on each of the sides. On side AB , 3 points are chosen. On side AC , 4 points are chosen. On side BC , 5 points are chosen.



How many triangles can be formed using the chosen points as vertices?

Solution 2.8

There are

$$3 + 4 + 5 = 12$$

chosen points in total.

If no three points were collinear, the number of triangles would be

$$\binom{12}{3} = 220.$$

However, any 3 points chosen on the same side do not form a triangle.

On side AB , there are

$$\binom{3}{3} = 1$$

collinear triple.

On side AC , there are

$$\binom{4}{3} = 4$$

collinear triples.

On side BC , there are

$$\binom{5}{3} = 10$$

collinear triples.

So the number of valid triangles is

$$\binom{12}{3} - \binom{3}{3} - \binom{4}{3} - \binom{5}{3} = 220 - 1 - 4 - 10 = 205.$$

Hence the number of triangles is 205.

Takeaways 2.8

- Start by counting all possible triples of points.
- Then subtract the triples that are collinear, since they do not form triangles.
- Geometric counting problems often reduce to combinations plus careful exclusion.

Problem 2.9: Age-limit check across teams

A sports association manages 13 junior teams. It decides to check the age of all players.

Any team that has more than 3 players above the age limit will be penalised.

A total of 41 players are found to be above the age limit.

Will any team be penalised? Justify your answer.

Solution 2.9

Yes, at least one team must be penalised.

Suppose no team were penalised. Then each of the 13 teams would have at most 3 players above the age limit.

So the greatest possible total number of above-limit players would be

$$13 \times 3 = 39.$$

But the actual total is 41, and

$$41 > 39.$$

This is impossible if every team has at most 3 above-limit players.

Therefore, at least one team must have more than 3 players above the age limit, so at least one team will be penalised.

Takeaways 2.9

- This is a pigeonhole-principle style argument: if every group had the maximum allowed amount, the total would still be too small.
- A good way to prove that something *must* happen is to assume it does not, then show the numbers do not work.
- In combinatorics, “justify your answer” often means giving a short contradiction argument like this one.

Problem 2.10: Expected score with replacement

A game consists of randomly selecting 4 balls from a bag. After each ball is selected it is replaced in the bag. The bag contains 3 red balls and 7 green balls.

For each red ball selected, 10 points are earned and for each green ball selected, 5 points are deducted. For instance, if a player picks 3 red balls and 1 green ball, the score will be

$$3 \times 10 - 1 \times 5 = 25$$

points.

What is the expected score in the game?

Solution 2.10

For one draw,

$$P(\text{red}) = \frac{3}{10}, \quad P(\text{green}) = \frac{7}{10}.$$

So the expected score from one draw is

$$\begin{aligned} E(\text{score per draw}) &= 10 \left(\frac{3}{10} \right) + (-5) \left(\frac{7}{10} \right) \\ &= 3 - \frac{35}{10} \\ &= 3 - 3.5 \\ &= -0.5. \end{aligned}$$

Since there are 4 draws with replacement, the expected total score is

$$4 \times (-0.5) = -2.$$

Therefore, the expected score in the game is $\boxed{-2}$ points.

Takeaways 2.10

- With replacement, each draw has the same probabilities.
- The expected value of a total is the sum of the expected values of the individual draws.
- A negative expected value means that, on average, the game favours the house rather than the player.

Problem 2.11: Minimum votes needed to win

The members of a club voted for a new president. There were 15 candidates for the position of president and 3543 members voted. Each member voted for one candidate only.

One candidate received more votes than anyone else and so became the new president. What is the smallest number of votes the new president could have received?

Solution 2.11

To make the winner's vote total as small as possible, we want the other 14 candidates to receive as many votes as possible, while still having fewer votes than the winner.

Suppose the winner received x votes. Then each of the other 14 candidates can receive at most $x - 1$ votes.

So the greatest possible total number of votes is

$$x + 14(x - 1).$$

This must be at least 3543:

$$x + 14(x - 1) \geq 3543$$

$$15x - 14 \geq 3543$$

$$15x \geq 3557$$

$$x \geq \frac{3557}{15} = 237.13\dots$$

So the smallest possible integer value of x is

$$\boxed{238}.$$

This value is achievable, since if the winner gets 238 votes then the other 14 candidates can together receive up to

$$14 \times 237 = 3318$$

votes, and

$$3318 + 238 = 3556 > 3543,$$

so the 3543 votes can be distributed with the winner still strictly ahead.

Takeaways 2.11

- To minimize a maximum, try making all competitors as large as possible without overtaking it.
- This is a classic “worst-case” or extremal counting argument.
- After finding the bound, always check that it is actually achievable.

Problem 2.12: Selecting finalists and awarding places

Out of 10 contestants, six are to be selected for the final round of a competition. Four of those six will be placed 1st, 2nd, 3rd and 4th.

In how many ways can this process be carried out?

Solution 2.12

First choose the 6 finalists from the 10 contestants:

$$\binom{10}{6} = 210.$$

Now choose and order the 4 placed contestants from those 6 finalists:

$${}^6P_4 = 6 \times 5 \times 4 \times 3 = 360.$$

So the total number of ways is

$$\binom{10}{6} \times {}^6P_4 = 210 \times 360 = 75600.$$

Hence the process can be carried out in 75600 ways.

Takeaways 2.12

- Break the process into stages: first select, then arrange.
- Use combinations for choosing the finalists because order does not matter there.
- Use permutations for the placings because 1st, 2nd, 3rd, and 4th are ordered positions.

Problem 2.13: Using the pigeonhole principle with topic choices

To complete a course, a student must choose and pass exactly three topics. There are eight topics from which to choose.

Last year 400 students completed the course.

Explain, using the pigeonhole principle, why at least eight students passed exactly the same three topics.

Solution 2.13

The number of different ways to choose exactly 3 topics from 8 is

$$\binom{8}{3} = 56.$$

So there are only 56 possible topic combinations.

Now distribute the 400 students among these 56 combinations. If at most 7 students had any one particular combination, then the total number of students could be at most

$$56 \times 7 = 392.$$

But in fact there are 400 students, and

$$400 > 392.$$

Therefore, by the pigeonhole principle, at least one of the 56 topic combinations must have been chosen by at least 8 students.

So at least eight students passed exactly the same three topics.

Takeaways 2.13

- Start by counting the number of possible categories, or “pigeonholes”.
- If the total number of objects exceeds the maximum possible without overlap, then one category must contain more than that maximum.
- In these questions, the pigeonhole principle often appears as: “If every group had at most k , the total would be too small.”

2.3 Advanced**Problem 2.14: Committee selection**

A committee containing 4 men and 4 women is to be formed from a group of 10 men and 8 women.

In how many different ways can the committee be formed?

Solution 2.14

Choose the men and women independently:

$$\binom{10}{4} \binom{8}{4} = 210 \times 70 = 14700$$

Hence the number of possible committees is $\boxed{14700}$.

Takeaways 2.14

- When categories are separate, count each category and multiply.
- Committee questions are combination questions because order is irrelevant.

Problem 2.15: Pascal’s triangle and a telescoping sum

i) The Pascal’s triangle relation can be expressed as

$$\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r}.$$

Do not prove this relation. Show that

$$\binom{m}{R} = \binom{m+1}{R+1} - \binom{m}{R+1}.$$

ii) Hence, or otherwise, prove that

$$\binom{2000}{2000} + \binom{2001}{2000} + \binom{2002}{2000} + \cdots + \binom{2050}{2000} = \binom{2051}{2001}.$$

Solution 2.15

i) Apply Pascal's triangle with $n = m + 1$ and $r = R + 1$:

$$\binom{m+1}{R+1} = \binom{m}{R} + \binom{m}{R+1}.$$

Rearranging gives

$$\binom{m}{R} = \binom{m+1}{R+1} - \binom{m}{R+1}.$$

ii) Using part (i) with $R = 2000$,

$$\binom{m}{2000} = \binom{m+1}{2001} - \binom{m}{2001}.$$

Now let $m = 2000, 2001, \dots, 2050$ and add:

$$\begin{aligned} \binom{2000}{2000} + \binom{2001}{2000} + \dots + \binom{2050}{2000} &= \left(\binom{2001}{2001} - \binom{2000}{2001} \right) \\ &\quad + \left(\binom{2002}{2001} - \binom{2001}{2001} \right) \\ &\quad + \dots \\ &\quad + \left(\binom{2051}{2001} - \binom{2050}{2001} \right). \end{aligned}$$

This is a telescoping sum, so all middle terms cancel. Also,

$$\binom{2000}{2001} = 0$$

because we cannot choose 2001 objects from 2000.

Hence the sum becomes

$$\binom{2051}{2001}.$$

Therefore,

$$\binom{2000}{2000} + \binom{2001}{2000} + \binom{2002}{2000} + \dots + \binom{2050}{2000} = \boxed{\binom{2051}{2001}}.$$

Takeaways 2.15

- Pascal's triangle identities are often most useful after rearranging them into a subtraction form.
- A telescoping sum is one in which most terms cancel after expansion.
- In binomial coefficients, $\binom{n}{r} = 0$ when $r > n$, and that can clean up endpoint terms neatly.
- In an HSC proof question, full marks usually come from showing the key substitutions, the cancellation, and the final conclusion clearly, not just writing the final answer.
- Writing the process matters: markers need to see how part (i) is applied in part (ii), so short linking phrases such as “using part (i)” and “this is a telescoping sum” are worth including.

Remark 2.1 (Marking guide note)

For a question like this, students should read the marking guide as a list of *visible ingredients* their solution must contain. In part (i), a correct rearrangement with the substitution $n = m + 1$ and $r = R + 1$ is enough. In part (ii), full marks usually require more than the final identity: show how part (i) is applied to the general term, write out enough of the expanded sum to make the telescoping pattern visible, and then state why the endpoint terms simplify. In short, a neat mathematical answer is good, but an examiner awards marks for a clear mathematical *process*.

Problem 2.16: Combining binomial coefficients

It is known that

$$\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r}$$

for all integers such that $1 \leq r \leq n-1$. Do not prove this.

Find one possible set of values for p and q such that

$$\binom{2025}{86} + \binom{2025}{87} + \binom{2026}{1940} = \binom{p}{q}.$$

Solution 2.16

Using Pascal's triangle identity,

$$\binom{2025}{86} + \binom{2025}{87} = \binom{2026}{87}.$$

Also,

$$\binom{2026}{1940} = \binom{2026}{2026-1940} = \binom{2026}{86}$$

by the symmetry property of binomial coefficients.

So the given expression becomes

$$\binom{2026}{87} + \binom{2026}{86}.$$

Applying Pascal's triangle identity again,

$$\binom{2026}{86} + \binom{2026}{87} = \binom{2027}{87}.$$

Therefore one possible answer is

$$\boxed{p = 2027, q = 87}.$$

Takeaways 2.16

- When two binomial coefficients have the same upper index and consecutive lower indices, try Pascal's triangle immediately.
- The symmetry identity $\binom{n}{r} = \binom{n}{n-r}$ is often the key step before applying Pascal's triangle.
- In HSC multiple-step identities, simplify one pair at a time.

Problem 2.17: Equal numbers of men and women

A club has $2n$ members, with n women and n men.

A group consisting of an even number $(0, 2, 4, \dots, 2n)$ of members is chosen, with the number of men equal to the number of women.

Show, giving reasons, that the number of ways to do this is

$$\binom{2n}{n}.$$

Solution 2.17

Suppose the group contains r women. Since the number of men must equal the number of women, the group must also contain r men.

For a fixed value of r , the number of ways to choose the group is

$$\binom{n}{r} \binom{n}{r}.$$

This is because we choose r women from the n women and r men from the n men.

Now r can take any value from 0 to n . So the total number of possible groups is

$$\sum_{r=0}^n \binom{n}{r}^2.$$

But from the identity

$$\binom{2n}{n} = \binom{n}{0}^2 + \binom{n}{1}^2 + \dots + \binom{n}{n}^2,$$

this sum is equal to

$$\binom{2n}{n}.$$

Therefore, the number of ways to choose such a group is

$$\boxed{\binom{2n}{n}}.$$

Takeaways 2.17

- When a selection has balanced categories, fix the number chosen from one category first.
- A total count can often be written as a sum over all allowable cases.
- This problem is a direct application of the identity $\sum_{r=0}^n \binom{n}{r}^2 = \binom{2n}{n}$.

Problem 2.18: A binomial identity and double counting capstone

- i) Use the identity

$$(1+x)^{2n} = (1+x)^n(1+x)^n$$

to show that

$$\binom{2n}{n} = \binom{n}{0}^2 + \binom{n}{1}^2 + \cdots + \binom{n}{n}^2,$$

where n is a positive integer.

- ii) A club has
- $2n$
- members, with
- n
- women and
- n
- men.

A group consisting of an even number $(0, 2, 4, \dots, 2n)$ of members is chosen, with the number of men equal to the number of women.

Show, giving reasons, that the number of ways to do this is

$$\binom{2n}{n}.$$

- iii) From the group chosen in part (ii), one of the men and one of the women are selected as leaders.

Show, giving reasons, that the number of ways to choose the even number of people and then the leaders is

$$1^2 \binom{n}{1}^2 + 2^2 \binom{n}{2}^2 + \cdots + n^2 \binom{n}{n}^2.$$

- iv) The process is now reversed so that the leaders, one man and one woman, are chosen first. The rest of the group is then selected, still made up of an equal number of women and men.

By considering this reversed process and using part (ii), find a simple expression for the sum in part (iii).

Solution 2.18

i) Compare coefficients of x^n in

$$(1+x)^{2n} = (1+x)^n(1+x)^n.$$

The coefficient of x^n on the left is $\binom{2n}{n}$. On the right it is

$$\sum_{r=0}^n \binom{n}{r} \binom{n}{n-r} = \sum_{r=0}^n \binom{n}{r}^2.$$

Hence

$$\binom{2n}{n} = \binom{n}{0}^2 + \binom{n}{1}^2 + \cdots + \binom{n}{n}^2.$$

ii) If the group contains r women, it must also contain r men, so for each r there are

$$\binom{n}{r} \binom{n}{r} = \binom{n}{r}^2$$

choices. Summing over $r = 0, 1, \dots, n$ gives

$$\sum_{r=0}^n \binom{n}{r}^2 = \binom{2n}{n}$$

by part (i).

iii) For a fixed r , after choosing r women and r men there are r^2 ways to choose one male leader and one female leader. So the total number of outcomes is

$$\sum_{r=1}^n r^2 \binom{n}{r}^2 = 1^2 \binom{n}{1}^2 + 2^2 \binom{n}{2}^2 + \cdots + n^2 \binom{n}{n}^2.$$

iv) Reverse the process. Choose the leaders first in n^2 ways. Then choose the rest of the group from the remaining $n-1$ men and $n-1$ women, with equal numbers of each. By part (ii), this can be done in

$$\binom{2n-2}{n-1}$$

ways. Therefore

$$1^2 \binom{n}{1}^2 + 2^2 \binom{n}{2}^2 + \cdots + n^2 \binom{n}{n}^2 = \boxed{n^2 \binom{2n-2}{n-1}}.$$

Takeaways 2.18

- This is a strong example of counting the same set of outcomes in two different ways.
- In part (iii), the count is grouped by the number of men and women selected.
- In part (iv), choosing the leaders first turns the same problem into a much simpler count.
- Double counting is a powerful way to derive closed forms for complicated sums.

Problem 2.19: Students and teachers in a circle

A group with 5 students and 3 teachers is to be arranged in a circle. In how many ways can this be done if no more than 2 students can sit together?

- A. $4! \times 3!$
- B. $5! \times 3!$
- C. $2! \times 5! \times 3!$
- D. $2! \times 2! \times 2! \times 3!$

Solution 2.19

Arrange the 3 teachers in a circle first. The number of circular arrangements is

$$(3 - 1)! = 2!.$$

These 3 teachers create 3 gaps between them. To ensure that no more than 2 students sit together, the 5 students must be split across these 3 gaps with at most 2 students in each gap.

We need

$$x_1 + x_2 + x_3 = 5$$

with each $x_i \leq 2$.

The only possible distribution is

$$2, 2, 1.$$

There are

$$3$$

ways to choose which gap gets the single student.

Now arrange the 5 distinct students in the chosen student positions:

$$5!$$

ways.

Hence the total number of arrangements is

$$2! \times 3 \times 5! = 2! \times 3! \times 5! = 5! \times 3!.$$

So the correct answer is

$$\boxed{\text{B. } 5! \times 3!}.$$

Takeaways 2.19

- For circular arrangement problems with mixed groups, it is often best to fix one category first and then place the other category into the gaps.
- Restriction questions often become gap-distribution problems.
- After handling the pattern of slots, place the distinct people into those slots at the end.

Problem 2.20: Guests who refuse to sit together

Eight guests are to be seated at a round table. If two of these guests refuse to sit next to each other, how many seating arrangements are possible?

Solution 2.20

First count all circular seating arrangements of 8 distinct guests:

$$(8 - 1)! = 7! = 5040.$$

Now count the arrangements in which the two particular guests do sit next to each other.

Treat those two guests as a single block. Then we have 7 objects around the table: the block plus the other 6 guests. The number of circular arrangements is

$$(7 - 1)! = 6!.$$

Within the block, the two guests can switch places in

$$2!$$

ways.

So the number of arrangements in which they sit together is

$$2 \times 6! = 1440.$$

Therefore, the number of arrangements in which they do *not* sit next to each other is

$$7! - 2 \cdot 6! = 5040 - 1440 = 3600.$$

Hence the required number of seating arrangements is 3600.

Takeaways 2.20

- In circular arrangement questions with a restriction like “not next to each other”, complementary counting is often the cleanest method.
- To count adjacent people, bind them into one block and arrange the block with the remaining people.
- For circular permutations of n distinct objects, use $(n - 1)!$.

Problem 2.21: Australian Powerball counting

In Australian Powerball, a single standard game consists of choosing 7 regular numbers from 35, and then choosing 1 Powerball from 20.

- i) How many different standard Powerball game lines are possible?
- ii) Hence show that the probability of winning Division 1 (all 7 regular numbers and the Powerball) is

$$\frac{1}{134490400}.$$

- iii) For a fixed winning draw, in how many different game lines would a player match exactly 3 regular numbers and the Powerball? Hence show that this probability is about 1 in 188.

Solution 2.21

i) Choose 7 regular numbers from 35, then choose 1 Powerball from 20:

$$\binom{35}{7} \times 20 = 6724520 \times 20 = 134490400.$$

So there are $\boxed{134490400}$ standard game lines.

ii) For a fixed draw, exactly one line wins Division 1, so

$$P(\text{Division 1}) = \frac{1}{134490400}.$$

iii) To match exactly 3 regular numbers, choose 3 of the 7 winning regular numbers and 4 of the 28 non-winning regular numbers. The Powerball must also be correct:

$$\binom{7}{3} \binom{28}{4} = 35 \times 20475 = 716625.$$

Hence

$$P(3 + \text{Powerball}) = \frac{716625}{134490400} \approx 0.005328 \approx \frac{1}{188}.$$

So the probability is about $\boxed{1 \text{ in } 188}$.

Takeaways 2.21

- Lottery questions often split naturally into “choose the regular numbers” and “choose the Powerball”.
- “Exactly k matches” means choose some winning numbers and the rest from the non-winning numbers.
- Published lottery odds are often rounded, so a combinatorial answer may be an approximation rather than a perfect exact fraction.

Problem 2.22: License plates and careful reading

A license plate involves 2 digits, followed by 2 letters, followed by 2 more digits, all chosen at random.

- How many different license plates are possible?
- What is the probability that the last digit is prime?
- Find the probability that the plate says 12AB34.

Solution 2.22

We interpret “all chosen at random” to mean repetition is allowed.

a) There are 10 choices for each digit and 26 choices for each letter:

$$10 \times 10 \times 26 \times 26 \times 10 \times 10 = 10^4 \cdot 26^2 = 6760000$$

So the number of license plates is $\boxed{6760000}$.

b) The last digit can be any of 0 to 9. The prime digits are 2, 3, 5, 7, so there are 4 favourable digits out of 10:

$$\frac{4}{10} = \frac{2}{5}$$

Thus the probability is $\boxed{\frac{2}{5}}$.

c) There is exactly one favourable plate, namely 12AB34, out of all possible license plates. Therefore

$$\frac{1}{6760000}$$

$$\boxed{\frac{1}{6760000}}$$

Takeaways 2.22

- Before counting, check the format carefully.
- If a question asks for one exact plate, the probability is usually $\frac{1}{\text{total number of plates}}$.
- When repetition is not forbidden, assume repeated digits or letters are allowed.

Problem 2.23: A mixed combinatorics review

1. How many different ways can 5 letters be chosen from URGOATED if all 4 vowels must be included?
2. How many different ways could the letters of COOKED be arranged?
3. How many 4-digit numbers can be made from the digits 1 to 8 without repetition?

Explain briefly which parts are combinations and which parts are permutations.

Solution 2.23

1. This is a combination because order does not matter. We must take U, O, A, E and then choose 1 more letter from the remaining 4 consonants:

$$\binom{4}{1} = 4$$

2. This is a permutation with repeated letters:

$$\frac{6!}{2!} = 360$$

3. This is a permutation without repetition:

$$8 \times 7 \times 6 \times 5 = 1680$$

So the answers are:

$$\boxed{4, 360, 1680}$$

Takeaways 2.23

- Combinations are for selections where order does not matter.
- Permutations are for arrangements where order does matter.
- Repeated objects in a permutation require division by factorials of repeated counts.

3 Part 2: Practice and Challenge Bank**3.1 Warm-Up Drills****Problem 3.1: Expanding a binomial surd expression**

Expand

$$(1 - \sqrt{2})^5 = a + b\sqrt{2},$$

where a and b are integers. Find a and b .

Hint: Use the binomial theorem, then simplify powers of $\sqrt{2}$ carefully: $(\sqrt{2})^2 = 2$, $(\sqrt{2})^3 = 2\sqrt{2}$, and so on.

Solution 3.1

Using the binomial theorem,

$$\begin{aligned} (1 - \sqrt{2})^5 &= 1 - 5\sqrt{2} + 10(\sqrt{2})^2 - 10(\sqrt{2})^3 + 5(\sqrt{2})^4 - (\sqrt{2})^5 \\ &= 1 - 5\sqrt{2} + 20 - 20\sqrt{2} + 20 - 4\sqrt{2} \\ &= 41 - 29\sqrt{2}. \end{aligned}$$

Hence

$$\boxed{a = 41, \quad b = -29}.$$

Takeaways 3.1

- After expanding, separate the rational terms from the $\sqrt{2}$ terms.
- Powers of $\sqrt{2}$ alternate between integer multiples and surd multiples.

Problem 3.2: When is a conjugate surd sum rational?For what integer values of n is

$$(1 - \sqrt{2})^n + (1 + \sqrt{2})^n$$

rational?

Hint: Expand both powers using the binomial theorem and compare the terms involving odd and even powers of $\sqrt{2}$. Also think about what happens for negative integers.

Solution 3.2

For a non-negative integer n , expand both terms by the binomial theorem:

$$(1 + \sqrt{2})^n = \binom{n}{0} + \binom{n}{1}\sqrt{2} + \binom{n}{2}(\sqrt{2})^2 + \dots,$$

$$(1 - \sqrt{2})^n = \binom{n}{0} - \binom{n}{1}\sqrt{2} + \binom{n}{2}(\sqrt{2})^2 - \dots.$$

When we add these two expansions, all terms involving odd powers of $\sqrt{2}$ cancel, while the even-power terms remain. Since

$$(\sqrt{2})^{2k} = 2^k,$$

all remaining terms are rational.

Therefore, for every non-negative integer n ,

$$(1 - \sqrt{2})^n + (1 + \sqrt{2})^n$$

is rational.

Now consider negative integers. Since

$$(1 + \sqrt{2})(1 - \sqrt{2}) = 1 - 2 = -1,$$

we have

$$1 - \sqrt{2} = -\frac{1}{1 + \sqrt{2}}.$$

So for any positive integer n ,

$$(1 - \sqrt{2})^{-n} + (1 + \sqrt{2})^{-n} = (-1)^n(1 + \sqrt{2})^n + (-1)^n(1 - \sqrt{2})^n = (-1)^n((1 + \sqrt{2})^n + (1 - \sqrt{2})^n),$$

which is also rational.

Hence the expression is rational for

all integers n .

Takeaways 3.2

- Conjugate surds often become much simpler when added or multiplied.
- In binomial expansions of $(a + b)^n$ and $(a - b)^n$, the odd-power terms cancel when the expansions are added.
- Here that cancellation shows the expression is rational for every integer n .

Problem 3.3: PINs with numbers and letters

The number of possible different PINs with a combination of 4 numbers and 2 letters is:

- A. 4435236
- B. 6760000
- C. 1000000
- D. 10676

Hint: Use the multiplication principle: count the choices for the 4 digits and the 2 letters separately, then multiply.

Solution 3.3

Assume the PIN has 4 digits followed by 2 letters, with repetition allowed.

There are 10 choices for each digit and 26 choices for each letter, so the total number of PINs is

$$10^4 \times 26^2 = 10000 \times 676 = 6760000.$$

So the correct answer is

B. 6760000.

Takeaways 3.3

- If each position can be chosen independently, multiply the number of choices for each position.
- Unless a question says otherwise, repetition is usually allowed in code or PIN questions.

Problem 3.4: Two coins of the same metal

A bag contains n metal coins, where $n \geq 3$. There are k silver coins in the bag and the rest are bronze.

Two coins are drawn at random from the bag, without replacement.

Find an expression for the probability that the two coins drawn are made of the same metal.

Hint: The two coins are the same metal if they are both silver or both bronze.

Solution 3.4

There are two favourable cases:

Both silver:

$$\frac{k}{n} \cdot \frac{k-1}{n-1}.$$

Both bronze: There are $n - k$ bronze coins, so

$$\frac{n-k}{n} \cdot \frac{n-k-1}{n-1}.$$

Hence the required probability is

$$\frac{k(k-1)}{n(n-1)} + \frac{(n-k)(n-k-1)}{n(n-1)}.$$

Takeaways 3.4

- For “same type” probability questions, split into mutually exclusive cases and add.
- Without replacement means the denominator changes on the second draw.

Problem 3.5: Probabilities with the four jacks

In a pack of cards, the 4 jacks are taken out and shuffled.

- What is the probability of picking out the jack of diamonds at random?
- If all the jacks are arranged in order, what is the probability of guessing the correct order?

Hint: For part (a), all four jacks are equally likely. For part (b), count the total number of possible orders of the four jacks.

Solution 3.5

- a) There are 4 jacks, and only one of them is the jack of diamonds. So

$$P(\text{jack of diamonds}) = \frac{1}{4}.$$

- b) The 4 jacks can be arranged in

$$4! = 24$$

different orders. Only one of these is the correct order, so


$$P(\text{guessing the correct order}) = \frac{1}{24}.$$

Takeaways 3.5

- If all outcomes are equally likely, probability is favourable outcomes over total outcomes.
- Ordered arrangements of 4 distinct objects are counted by $4!$.

Problem 3.6: Counting rectangles in a 5×6 grid

How many rectangles can be found in the 5×6 grid below? Remember that squares should also be counted, since every square is a rectangle.



Hint: Choose any 2 of the vertical grid lines and any 2 of the horizontal grid lines.

Solution 3.6

A rectangle is determined by choosing:

- 2 of the 7 vertical grid lines
- 2 of the 6 horizontal grid lines.

So the total number of rectangles is

$$\binom{7}{2} \binom{6}{2} = 21 \times 15 = 315.$$

Hence the number of rectangles is 315.

Takeaways 3.6

- Grid-rectangle questions are usually solved by choosing boundary lines, not by listing shapes one by one.
- Squares are already included, since every square is also a rectangle.

Problem 3.7: How fast does brute force grow?

A password is made from n binary digits, each of which is either 0 or 1.

- How many possible passwords are there when $n = 5$?
- How many possible passwords are there when $n = 10$?
- Explain briefly why checking every possible password becomes difficult when n is large.

Hint: Each position has 2 choices, so use the multiplication principle.

Solution 3.7

- a) For $n = 5$, the number of passwords is

$$2^5 = 32.$$

- b) For $n = 10$, the number of passwords is

$$2^{10} = 1024.$$

- c) The total number of possibilities doubles each time one more digit is added. So the number of passwords grows exponentially, and brute-force checking quickly becomes impractical for large n .

Takeaways 3.7

- Some counting problems grow exponentially, even when the rule looks simple.
- This is one reason efficient counting methods matter.

Problem 3.8: Repeated derivatives of $\frac{1}{x}$

If

$$f(x) = \frac{1}{x},$$

find:

- i) $f'(x)$
- ii) $f''(x)$
- iii) $f^{(5)}(x)$
- iv) $f^{(n)}(x)$.

and differentiate a few times to spot the pattern.

$$f^{-1}(x) = (x)f$$

Hint: Write**Solution 3.8**

Using the power rule,

$$f(x) = x^{-1}.$$

i)

$$f'(x) = -x^{-2} = -\frac{1}{x^2}.$$

ii)

$$f''(x) = 2x^{-3} = \frac{2}{x^3}.$$

iii) Continuing the pattern,

$$f^{(3)}(x) = -\frac{6}{x^4}, \quad f^{(4)}(x) = \frac{24}{x^5},$$

so

$$f^{(5)}(x) = -\frac{120}{x^6}.$$

iv) Each differentiation introduces:

- one extra factor in the numerator
- one extra power of x in the denominator
- alternating signs.

Therefore

$$f^{(n)}(x) = (-1)^n \frac{n!}{x^{n+1}}.$$

Takeaways 3.8

- Repeated differentiation often reveals a pattern after the first few derivatives.
- Here the coefficients are factorials: $1, 1, 2, 6, 24, 120, \dots$
- This is more of a calculus enrichment problem than a core combinatorics problem, but it connects nicely to factorial notation.

Problem 3.9: Using Pascal's triangle to differentiate x^4

i) Use Pascal's triangle to expand

$$(x + h)^4.$$

ii) If

$$f(x) = x^4,$$

simplify

$$f(x + h) - f(x).$$

iii) Hence use the definition

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

to differentiate x^4 .

iv) For the general case

$$f(x) = x^n,$$

write down the expansion of $(x + h)^n$, and hence show that

$$\frac{d}{dx}(x^n) = nx^{n-1}.$$

Hint: Use row 4 of Pascal's triangle for the specific case, then use the binomial theorem for $(x + h)^n$. In both cases, factor out h before taking the limit.

Solution 3.9

i) Using Pascal's triangle or the binomial theorem,

$$(x+h)^4 = x^4 + 4x^3h + 6x^2h^2 + 4xh^3 + h^4.$$

ii) Since $f(x) = x^4$,

$$f(x+h) - f(x) = (x+h)^4 - x^4 = 4x^3h + 6x^2h^2 + 4xh^3 + h^4.$$

iii) Therefore,

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{4x^3h + 6x^2h^2 + 4xh^3 + h^4}{h} \\ &= \lim_{h \rightarrow 0} (4x^3 + 6x^2h + 4xh^2 + h^3) \\ &= 4x^3. \end{aligned}$$

Hence

$$\boxed{f'(x) = 4x^3.}$$

iv) In general, the binomial theorem gives

$$(x+h)^n = x^n + \binom{n}{1}x^{n-1}h + \binom{n}{2}x^{n-2}h^2 + \cdots + h^n.$$

So if

$$f(x) = x^n,$$

then

$$f(x+h) - f(x) = nx^{n-1}h + \binom{n}{2}x^{n-2}h^2 + \cdots + h^n.$$

Therefore,

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \left(nx^{n-1} + \binom{n}{2}x^{n-2}h + \cdots + h^{n-1} \right) \\ &= nx^{n-1}. \end{aligned}$$

Hence

$$\boxed{\frac{d}{dx}(x^n) = nx^{n-1}.}$$

Takeaways 3.9

- Pascal's triangle gives the coefficients in expansions such as $(x+h)^4$.
- In the derivative definition, the key step is to factor out h before taking the limit.
- The same idea works in general: the binomial theorem for $(x+h)^n$ leads to the power rule $\frac{d}{dx}(x^n) = nx^{n-1}$.
- This is a calculus enrichment problem, but it shows one way algebra and Pascal's triangle support differentiation.

3.2 Stretch Problems

Problem 3.10: Term independent of x

Find the term independent of x in the expansion of

$$\left(11x - \frac{4}{x^2}\right)^{15}.$$

Hint: Use the general term in the binomial expansion. Then make the power of x equal to 0.

Solution 3.10

The general term is

$$T_{r+1} = \binom{15}{r} (11x)^{15-r} \left(-\frac{4}{x^2}\right)^r.$$

So the power of x is

$$x^{15-r} x^{-2r} = x^{15-3r}.$$

For the term independent of x , we need

$$15 - 3r = 0 \quad \Rightarrow \quad r = 5.$$

Hence the required term is

$$\begin{aligned} \binom{15}{5} (11x)^{10} \left(-\frac{4}{x^2}\right)^5 &= \binom{15}{5} \cdot 11^{10} \cdot (-4)^5 \\ &= 3003 \cdot 11^{10} \cdot (-1024). \end{aligned}$$

Therefore, the term independent of x is

$$\boxed{-3003 \cdot 1024 \cdot 11^{10}}.$$

Takeaways 3.10

- For “term independent of x ” questions, write the general term first.
- The key step is solving the exponent equation that makes the power of x equal to 0.

Problem 3.11: Five-letter words from Mamungkukumpurangkuntjunya

How many 5-letter “words” can be made from the word Mamungkukumpurangkuntjunya?

Helpful note: To save time, you may use the repeated-letter frequencies

$$u^6, n^4, m^3, a^3, k^3, g^2,$$

and note that the remaining letters each occur once.

Hint: Treat this as an arrangement problem with repeated letters. Split into cases according to the repetition pattern of the 5 letters.

Solution 3.11

The word Mamungkukupurangkuntjunya contains the letter types

$$a, g, j, k, m, n, p, r, t, u, y$$

with frequencies

$$u^6, n^4, m^3, a^3, k^3, g^2,$$

and p, r, t, j, y each occurring once.

We count by repetition pattern.

1. All 5 letters distinct:

$$\binom{11}{5} 5! = 55440.$$

2. One pair and 3 singles:

$$6 \cdot \binom{10}{3} \cdot \frac{5!}{2!} = 43200.$$

3. Two pairs and 1 single:

$$\binom{6}{2} \cdot 9 \cdot \frac{5!}{2!2!} = 4050.$$

4. One triple and 2 singles:

$$5 \cdot \binom{10}{2} \cdot \frac{5!}{3!} = 4500.$$

5. One triple and one pair:

$$5 \cdot 5 \cdot \frac{5!}{3!2!} = 250.$$

6. Four of one letter and one single:

$$2 \cdot 10 \cdot \frac{5!}{4!} = 100.$$

7. Five of one letter:

$$1.$$

Therefore, the total number of 5-letter words is

$$55440 + 43200 + 4050 + 4500 + 250 + 100 + 1 = 107541.$$

Hence the answer is 107541.

Takeaways 3.11

- When letters repeat, count by repetition pattern such as $(2, 1, 1, 1)$ or $(3, 2)$.
- After choosing the letters, use multinomial arrangements such as $\frac{5!}{2!}$ or $\frac{5!}{3!2!}$.

Problem 3.12: Constant term from a product of two expansions

Find the term independent of x in the expansion of

$$\left(x + \frac{1}{x}\right)^5 \left(2x^2 - \frac{3}{x}\right)^6.$$

Hint: Write the general term from each bracket, then find which pair of terms gives a total power of x equal to 0.

Solution 3.12

From

$$\left(x + \frac{1}{x}\right)^5,$$

the general term is

$$\binom{5}{r} x^{5-r} \left(\frac{1}{x}\right)^r = \binom{5}{r} x^{5-2r}.$$

From

$$\left(2x^2 - \frac{3}{x}\right)^6,$$

the general term is

$$\binom{6}{s} (2x^2)^{6-s} \left(-\frac{3}{x}\right)^s = \binom{6}{s} 2^{6-s} (-3)^s x^{12-3s}.$$

So in the product, the power of x is

$$(5 - 2r) + (12 - 3s) = 17 - 2r - 3s.$$

For a constant term, we need

$$17 - 2r - 3s = 0.$$

With $0 \leq r \leq 5$ and $0 \leq s \leq 6$, the only solution is

$$r = 1, \quad s = 5.$$

So the constant term is

$$\begin{aligned} \binom{5}{1} \binom{6}{5} \cdot 1 \cdot 2^1 (-3)^5 &= 5 \cdot 6 \cdot 2 \cdot (-243) \\ &= -14580. \end{aligned}$$

Hence the term independent of x is

$$\boxed{-14580}.$$

Takeaways 3.12

- In products of expansions, add the exponents from the chosen terms.
- The constant term occurs when the total exponent of x is 0.
- After finding the exponent condition, substitute the matching values back into the coefficients.

Problem 3.13: Adults and children at the cinema

Three adults and five children go to the cinema and are seated next to each other in a row of eight seats.

How many ways can these eight people sit so that at least two of the adults sit next to each other?

Hint: Count all seatings first, then subtract the seatings in which no two adults sit together.

Solution 3.13

There are

$$8!$$

ways to seat the 8 distinct people in a row.

Now count the arrangements in which no two adults sit together.

First arrange the 5 children:

$$5!$$

ways.

This creates 6 gaps where adults could be placed:

$$_ C _ C _ C _ C _ C _$$

Choose 3 of these 6 gaps for the adults:

$$\binom{6}{3}$$

ways.

Then arrange the 3 adults in those chosen gaps:

$$3!$$

ways.

So the number of seatings with no two adults together is

$$5! \binom{6}{3} 3! = 120 \cdot 20 \cdot 6 = 14400.$$

Therefore, the number of seatings in which at least two adults sit together is

$$8! - 14400 = 40320 - 14400 = 25920.$$

Hence the required number of ways is

$$\boxed{25920}.$$

Takeaways 3.13

- “At least” often suggests complementary counting.
- To keep certain people apart, arrange the others first and then place people into the gaps.
- Gap methods are especially useful for adjacency restrictions in line arrangements.

Problem 3.14: The alternating row constraint

Four boys, including Bob, and four girls, including Grace, are to stand in a row.

Find the number of distinct arrangements if the boys and girls must alternate positions and Bob must stand immediately next to Grace.

Hint: Count all alternating arrangements first. Then use symmetry: in an alternating row, each specific boy-girl pairing appears equally often.

Solution 3.14

There are two alternating patterns:

$$B G B G B G B G \quad \text{or} \quad G B G B G B G B.$$

For each pattern, the boys can be arranged in $4!$ ways and the girls in $4!$ ways. So the total number of alternating arrangements is

$$2 \cdot 4! \cdot 4! = 1152.$$

Now impose the condition that Bob is next to Grace.

In any alternating row of 8 people, there are 7 adjacent boy-girl pairs. On the other hand, Bob can be paired with any of the 4 girls, and Grace can be paired with any of the 4 boys, giving

$$4 \times 4 = 16$$

possible boy-girl pairings in total.

By symmetry, each specific boy-girl pairing occurs equally often among the alternating arrangements, so the fraction in which Bob and Grace are adjacent is

$$\frac{7}{16}.$$

Hence the required number of arrangements is

$$1152 \cdot \frac{7}{16} = 72 \cdot 7 = 504.$$

Therefore the number of distinct arrangements is

$$\boxed{504}.$$

Takeaways 3.14

- With equal group sizes, alternating rows always begin from exactly two templates.
- Count the full structured sample space first, then apply the extra adjacency restriction.
- This problem is a good example of symmetry simplifying what would otherwise be a messy case split.
- Numbering the positions from 1 to 8. There are 7 adjacent pairs for Bob and Grace: $(1, 2), (2, 3), \dots, (7, 8)$.


Problem 3.15: Lines determined by points on a line and a circle

There are 11 points on the plane.

- 4 of them lie on the line $y = 0$
- 7 of them lie on the circle

$$x^2 + (y - 2)^2 = 1.$$

How many different lines can be formed that pass through at least two of the 11 points?



Hint: Start with all pairs of points, then correct for the overcount caused by the 4 collinear points on $y = 0$.

Solution 3.15

If no three points were collinear, the number of lines would be

$$\binom{11}{2} = 55.$$

However, the 4 points on the line $y = 0$ are collinear. The pairs among these 4 points are

$$\binom{4}{2} = 6,$$

but they produce only *one* line.

So among these 6 counted pairs, we have overcounted by

$$6 - 1 = 5.$$

Therefore, the number of distinct lines is

$$\binom{11}{2} - 5 = 55 - 5 = 50.$$

Hence the required number of lines is

$$\boxed{50}.$$


Takeaways 3.15

- The usual starting point is $\binom{n}{2}$, because each pair of points determines a line.
- If several points are collinear, those pairs all represent the same line and must be adjusted.
- A block of k collinear points contributes $\binom{k}{2}$ pairs but only 1 line.

Problem 3.16: Dota knockout bracket

Suppose a college Dota tournament consists of 64 teams playing head-to-head in a knockout tournament. There are 6 rounds: the round of 64, round of 32, round of 16, round of 8, the final four teams, and the grand final.

Suppose you are filling out a bracket that specifies which team will win each match in each round.



How many possible brackets can you make?

Hint: A completed bracket is determined by choosing the winner of every match. How many matches are played in a 64-team knockout tournament?

Solution 3.16

In a knockout tournament, each match eliminates exactly one team.

Starting with 64 teams, we must eliminate 63 teams to get one champion. So the tournament contains

$$63$$

matches in total.

For each match, there are 2 possible winners. Therefore the total number of possible completed brackets is

$$2^{63}.$$

Hence the number of possible brackets is

$$\boxed{2^{63}}.$$

Takeaways 3.16

- In a knockout tournament with n teams, there are $n - 1$ matches.
- A bracket is determined by the winner of each match.
- Once the number of matches is known, use the multiplication principle.

Problem 3.17: Dividing a class into equal groups

A class has 28 students.

The teacher wants to divide them into 7 groups for an assignment task, with each group containing 4 students.

- How many different ways can this be done if the 7 groups are unlabeled?
- How many different ways can this be done if the groups are labeled Group 1, Group 2, ..., Group 7?

Hint: Start by arranging all 28 students, then correct for the fact that order within each group does not matter. In part (i), the order of the groups also does not matter, but in part (ii) it does.

Solution 3.17

If we arrange all 28 students in a line, there are

$$28!$$

ways.

Now split this line into 7 consecutive blocks of 4.

(i) If the groups are unlabeled, this overcounts in two ways:

- within each group, the 4 students can be arranged in $4!$ ways without changing the group
- the 7 groups themselves can be arranged in $7!$ ways without changing the division.

So the number of different unlabeled groupings is

$$\frac{28!}{(4!)^7 7!}.$$

Hence

$$\boxed{\frac{28!}{(4!)^7 7!}}.$$

(ii) If the groups are labeled, we still divide by $(4!)^7$ because order within each group does not matter, but we do *not* divide by $7!$ because the groups are now distinguishable.

Hence


$$\boxed{\frac{28!}{(4!)^7}}.$$

Takeaways 3.17

- When dividing people into equal-sized groups, first count all arrangements, then divide by repeated internal orders.
- If the groups are unlabeled, also divide by the number of ways to reorder the groups.
- The difference between labeled and unlabeled groups here is exactly a factor of $7!$.

Problem 3.18: Triangles from points on a circle

Seven points $P_1, P_2, \dots, P_7$ are arranged in order around a circle, as shown below.



- How many triangles can be drawn using these points as vertices?
- How many pairs of triangles can be drawn, where the vertices of each triangle are distinct points?

Hint: For part (i), choose any 3 of the 7 points. For part (ii), choose 6 points, then split them into two groups of 3.

Solution 3.18

i) Any 3 of the 7 points form a triangle, so the number of triangles is

$$\binom{7}{3} = 35.$$

ii) First choose the 6 points that will be used:

$$\binom{7}{6} = 7.$$

Now split those 6 points into two groups of 3 to make two triangles:

$$\binom{6}{3} = 20.$$

This counts each pair of triangles twice, because choosing Triangle A first or Triangle B first gives the same pair. So divide by 2:

$$\frac{\binom{7}{6} \binom{6}{3}}{2} = \frac{7 \cdot 20}{2} = 70.$$

Hence the answers are

$$\boxed{35 \text{ and } 70}.$$

Takeaways 3.18


- On a circle, any choice of 3 distinct points gives a triangle.
- When counting pairs of groups, check whether the order of the groups matters.
- If the pair is unordered, divide by 2 after choosing the first and second groups.

Problem 3.19: Triangles from a regular octagon and its centre

On the unit circle with centre O , there are 8 equally spaced vertices $A_1, A_2, \dots, A_8$. How many triangles can be formed using the 9 points

$$A_1, A_2, \dots, A_8, O$$

as vertices?



Hint: Choose any 3 of the 9 points, then subtract the choices that are collinear.

Solution 3.19

If there were no collinear triples, the number of triangles would be

$$\binom{9}{3} = 84.$$

Now identify the collinear triples.

No line can contain three of the points on the circle, because a line meets a circle in at most two points. However, the centre O lies on each diameter, so the following four triples are collinear:

$$(A_1, O, A_5), \quad (A_2, O, A_6), \quad (A_3, O, A_7), \quad (A_4, O, A_8).$$

These 4 triples do not form triangles, so we subtract them:

$$84 - 4 = 80.$$

Hence the number of triangles is

$$\boxed{80}.$$

Takeaways 3.19

- Start with $\binom{n}{3}$ when counting triangles from points, then subtract collinear triples.
- On a circle, the only collinear triples here come from diameters passing through the centre.
- Geometry counting problems often reduce to one small structural observation.

Problem 3.20: Pigeons and pigeonholes

A flock of 41 pigeons fly into 10 different pigeonholes.

- Show that there is at least one pigeonhole with at least 5 pigeons.
- Show that there is at least one pair of pigeonholes in which there are at least 9 pigeons altogether.

Hint: For part (a), divide 41 by 10. For part (b), group the 10 pigeonholes into 5 pairs.

Solution 3.20

a) If every pigeonhole had at most 4 pigeons, then the total number of pigeons would be at most

$$10 \times 4 = 40.$$

But there are actually 41 pigeons, so this is impossible.

Therefore, at least one pigeonhole must contain at least

$$\boxed{5}$$

pigeons.

b) Group the 10 pigeonholes into 5 pairs.

If every pair of pigeonholes had at most 8 pigeons altogether, then the total number of pigeons would be at most

$$5 \times 8 = 40.$$

Again, this is impossible because there are 41 pigeons.

Therefore, at least one pair of pigeonholes must contain at least

$$\boxed{9}$$

pigeons altogether.

Takeaways 3.20

- The pigeonhole principle often begins with: “Suppose every group had at most this many.”
- If the resulting total is too small, then some group must exceed that bound.
- For part (b), the key idea is to create larger pigeonholes by pairing the original ones.

3.3 Challenge Corner**Problem 3.21: Comparing coefficients in $(1+x)^{m+n}$**

Use

$$(1+x)^{m+n} = (1+x)^m(1+x)^n$$

and compare coefficients of x^r to show that

$$\binom{m+n}{r} = \binom{m}{0}\binom{n}{r} + \binom{m}{1}\binom{n}{r-1} + \cdots + \binom{m}{r}\binom{n}{0},$$

where terms with impossible binomial coefficients are taken to be 0.

Hint: Find the coefficient of x^r on the left first. Then, on the right, think about all the ways to obtain x^r by multiplying a term from $(1+x)^m$ with a term from $(1+x)^n$.

Solution 3.21

From the binomial theorem,

$$(1+x)^{m+n} = \sum_{k=0}^{m+n} \binom{m+n}{k} x^k,$$

so the coefficient of x^r on the left is

$$\binom{m+n}{r}.$$

Also,

$$(1+x)^m = \sum_{i=0}^m \binom{m}{i} x^i, \quad (1+x)^n = \sum_{j=0}^n \binom{n}{j} x^j.$$

To obtain an x^r term in the product, we must choose x^i from the first factor and x^{r-i} from the second. Hence the coefficient of x^r on the right is

$$\sum_{i=0}^r \binom{m}{i} \binom{n}{r-i}.$$

Since both expressions are the coefficient of x^r in the same polynomial,

$$\binom{m+n}{r} = \sum_{i=0}^r \binom{m}{i} \binom{n}{r-i}.$$

That is,


$$\boxed{\binom{m+n}{r} = \binom{m}{0} \binom{n}{r} + \binom{m}{1} \binom{n}{r-1} + \cdots + \binom{m}{r} \binom{n}{0}}.$$

Takeaways 3.21

- Expanding the same expression in two ways is a standard way to prove combinatorial identities.
- Comparing coefficients is especially useful when a problem involves powers of $(1+x)$.
- This identity is often called Vandermonde's identity.

Problem 3.22: Grid paths and binomial identities

The diagram shows a 6×8 grid. The aim is to walk from the point A in the top left-hand corner to the point B in the bottom right-hand corner by moving along the grid lines either downwards or to the right. Let C be the grid point that is 3 steps to the right and 2 steps down from A .



a) Find how many different routes are possible:

- i) without restriction
- ii) if you must pass through C .

b) Consider the diagonal of grid points

$$(0, 6), (1, 5), (2, 4), \dots, (6, 0).$$

Explain why every route from A to B passes through exactly one of these points, and hence prove that

$$\binom{14}{6} = \binom{8}{0} \binom{6}{6} + \binom{8}{1} \binom{6}{5} + \dots + \binom{8}{6} \binom{6}{0}.$$

c) Generalise part (b) to prove that

$$\binom{m+n}{n} = \binom{m}{0} \binom{n}{n} + \binom{m}{1} \binom{n}{n-1} + \dots + \binom{m}{n} \binom{n}{0},$$

for integers $m \geq n \geq 0$.

d) By considering an $n \times n$ square grid and the corresponding diagonal, prove that

$$\binom{2n}{n} = \binom{n}{0}^2 + \binom{n}{1}^2 + \dots + \binom{n}{n}^2.$$

Hint: For part (a), count the number of right and down moves. For part (b), split each route at the unique point where it meets the diagonal. Parts (c) and (d) are the corresponding general forms of the same idea.

Solution 3.22

a)(i) A route from A to B uses 8 right moves and 6 down moves, so

$$\binom{14}{6} = \binom{14}{8}.$$

a)(ii) Split the route at C . Then

$$\#(A \rightarrow C) = \binom{5}{2} = 10, \quad \#(C \rightarrow B) = \binom{9}{4} = 126,$$

$$\therefore \#(A \rightarrow C \rightarrow B) = \binom{5}{2} \binom{9}{4} = 1260.$$

b) Every move decreases $x + y$ by 1, so each route meets $x + y = 6$ exactly once, at one of

$$(0, 6), (1, 5), \dots, (6, 0).$$

$$\text{At } (k, 6 - k): \quad \#(A \rightarrow (k, 6 - k)) = \binom{6}{k}, \quad \#((k, 6 - k) \rightarrow B) = \binom{8}{k}.$$

$$\binom{14}{6} = \sum_{k=0}^6 \binom{6}{k} \binom{8}{k} = \sum_{k=0}^6 \binom{8}{k} \binom{6}{6-k}.$$

c) In an $n \times m$ grid with $m \geq n$, the total number of routes is

$$\binom{m+n}{n}.$$

Split routes by the unique point where they meet

$$x + y = n.$$

If this happens after k right moves, then

$$\#(\text{first part}) = \binom{n}{k}, \quad \#(\text{second part}) = \binom{m}{k}.$$

$$\binom{m+n}{n} = \sum_{k=0}^n \binom{m}{k} \binom{n}{n-k}.$$

d) For an $n \times n$ square grid,

$$\binom{2n}{n} = \sum_{k=0}^n \binom{n}{k} \binom{n}{n-k}.$$

Using the symmetry identity

$$\binom{n}{n-k} = \binom{n}{k},$$

we obtain

$$\boxed{\binom{2n}{n} = \binom{n}{0}^2 + \binom{n}{1}^2 + \dots + \binom{n}{n}^2.}$$

Takeaways 3.22

- Grid-path counting gives a very natural combinatorial proof of Vandermonde-type identities.
- Splitting paths at a suitable diagonal point is a standard and powerful technique.
- The square-grid case leads directly to the classic identity

$$\binom{2n}{n} = \sum_{k=0}^n \binom{n}{k}^2.$$

Problem 3.23: A coefficient identity for $\binom{4n}{2n}$

(i) Write down the coefficient of x^{2n} in the binomial expansion of $(1+x)^{4n}$.

(ii) Show that

$$(1+x^2+2x)^{2n} = \sum_{k=0}^{2n} \binom{2n}{k} x^{2n-k} (x+2)^{2n-k}.$$

(iii) It is known that

$$x^{2n-k}(x+2)^{2n-k} = \binom{2n-k}{0} 2^{2n-k} x^{2n-k} + \binom{2n-k}{1} 2^{2n-k-1} x^{2n-k+1} + \dots + \binom{2n-k}{2n-k} 2^0 x^{4n-2k}.$$

Do *not* prove this.

Show that

$$\binom{4n}{2n} = \sum_{k=0}^n 2^{2n-2k} \binom{2n}{k} \binom{2n-k}{k}.$$

Hint: For part (ii), rewrite $1+x(x+2)$ as $1+x^2+2x$ and identify which term in the inner expansion produces that power of x^{2n} on both sides and identify which term in the inner expansion produces that power.

Solution 3.23

(i) From the binomial theorem,

$$(1+x)^{4n} = \sum_{r=0}^{4n} \binom{4n}{r} x^r,$$

so the coefficient of x^{2n} is $\binom{4n}{2n}$.

$$\boxed{\binom{4n}{2n}}.$$

(ii) Since $1 + x^2 + 2x = 1 + x(x+2)$,

$$(1 + x^2 + 2x)^{2n} = (1 + x(x+2))^{2n} = \sum_{k=0}^{2n} \binom{2n}{k} x^{2n-k} (x+2)^{2n-k}.$$

(iii) Also,

$$(1 + x^2 + 2x)^{2n} = ((1+x)^2)^{2n} = (1+x)^{4n},$$

so its coefficient of x^{2n} is again $\binom{4n}{2n}$.

Now in

$$\sum_{k=0}^{2n} \binom{2n}{k} x^{2n-k} (x+2)^{2n-k},$$

a typical term from

$$x^{2n-k} (x+2)^{2n-k}$$

is

$$\binom{2n-k}{r} 2^{2n-k-r} x^{2n-k+r}.$$

For an x^{2n} term we need

$$2n - k + r = 2n, \quad \text{so } r = k.$$

Hence the contribution from the k th summand is

$$\binom{2n}{k} \binom{2n-k}{k} 2^{2n-2k}.$$

This is possible only when $k \leq 2n - k$, i.e. $k \leq n$. Therefore

$$\binom{4n}{2n} = \sum_{k=0}^n 2^{2n-2k} \binom{2n}{k} \binom{2n-k}{k}.$$

Takeaways 3.23

- This is another example of proving an identity by expanding the same expression in two different ways.
- The key step is to compare coefficients of a carefully chosen power of x .
- Bounds matter: the upper limit drops from $2n$ to n because terms with $k > n$ cannot contribute to the coefficient of x^{2n} .

Problem 3.24: Repeated draws of green counters

A bag contains counters, some of which are green.

One hundred trials of an experiment are run. In each trial, one counter is selected from the bag at random and its colour noted. The counter is then returned to the bag.

Let X be the random variable representing the number of times that a green counter is selected.

Given that $E(X) = 20$, identify the distribution of X and find the probability that a green counter is selected exactly 20 times.

Hint: The experiment consists of repeated independent trials with only two outcomes on each trial: green or not green.

Solution 3.24

Since each trial is independent and each draw is with replacement, X follows a binomial distribution:

$$X \sim \text{Bin}(100, p),$$

where p is the probability of selecting a green counter on any one draw.

For a binomial random variable,

$$E(X) = np.$$

Given that $E(X) = 20$, we have

$$100p = 20 \Rightarrow p = 0.2.$$

So

$$X \sim \text{Bin}(100, 0.2).$$

Therefore,

$$P(X = 20) = \binom{100}{20} (0.2)^{20} (0.8)^{80}.$$

Hence the required probability is

$$\boxed{\binom{100}{20} (0.2)^{20} (0.8)^{80}}.$$

Takeaways 3.24

- Repeated draws with replacement lead naturally to a binomial model.
- For $X \sim \text{Bin}(n, p)$, the mean is $E(X) = np$.
- Once p is known, exact probabilities are found using the binomial formula.

Problem 3.25: Arrangement formulas for x people

Find the number of arrangements possible if x people are

- in a straight line
- in a circle
- in a circle with 2 particular people together
- in a straight line with 3 particular people together
- in a circle with 2 particular people not together.

Hint: Use $x!$ for a line and $(x-1)!$ for a circle. For “together” questions, bind people into a block. For “not together”, subtract from the total.

Solution 3.25

a) In a straight line:

$$x!$$

b) In a circle:

$$(x-1)!$$

c) In a circle with 2 particular people together: Treat the 2 people as one block. Then there are $(x-1)$ objects in a circle, giving

$$(x-2)!$$

arrangements, and the pair can switch places in $2!$ ways. So

$$2(x-2)!$$

d) In a straight line with 3 particular people together: Treat the 3 people as one block. Then there are $(x-2)$ objects in a line, giving

$$(x-2)!$$

arrangements, and the 3 people inside the block can be arranged in $3!$ ways. So

$$3!(x-2)! = 6(x-2)!$$

e) In a circle with 2 particular people not together: Start with all circular arrangements:

$$(x-1)!$$

Subtract the arrangements where the two particular people are together:

$$2(x-2)!$$

Hence

$$(x-1)! - 2(x-2)!$$

Takeaways 3.25

- The standard counts are $x!$ for a line and $(x-1)!$ for a circle.
- “Together” usually suggests treating people as a single block.
- “Not together” is often easiest by complementary counting.

Problem 3.26: A pair summing to 21

Eleven numbers are randomly chosen from the set of integers

$$S = \{1, 2, 3, 4, \dots, 20\}.$$

Prove that the sum of two of the eleven numbers selected must equal 21.

Hint: Pair the numbers in the set so that each pair sums to 21.

Solution 3.26

The set S can be split into the following 10 pairs:

$$(1, 20), (2, 19), (3, 18), \dots, (10, 11).$$

Each pair sums to 21.

Now suppose 11 numbers are chosen from the 20 numbers in S .

There are only 10 such pairs, but we have chosen 11 numbers. By the pigeonhole principle, at least one complete pair must be selected.

Therefore, among the 11 chosen numbers, there must be two numbers whose sum is 21.

Takeaways 3.26

- A strong first step in pigeonhole problems is to group objects into meaningful boxes.
- Here, the natural boxes are the pairs that sum to 21.
- Choosing more objects than there are boxes forces one box to contain two chosen objects.

Problem 3.27: Positive integer solutions

Show that there are

$$\binom{n-1}{k-1}$$

distinct positive integer-valued vectors $(x_1, \dots, x_k)$ satisfying

$$x_1 + \dots + x_k = n, \quad x_i > 0 \text{ for all } i = 1, \dots, k.$$

Hint: Think of writing n as a row of n identical objects, then inserting dividers to split them into k positive groups.

Solution 3.27

Represent n as n identical stars:

$$\underbrace{* * * \cdots *}_{n \text{ stars}}.$$

To split these into k positive parts, we must place $k - 1$ dividers into the gaps between consecutive stars. There are

$$n - 1$$

gaps between the n stars, and we choose

$$k - 1$$

of them for the dividers.

Hence the number of positive integer solutions is

$$\boxed{\binom{n-1}{k-1}}.$$

Takeaways 3.27

- This is the standard stars-and-bars count for positive integer solutions.
- Positivity means every group must contain at least one star, so dividers can only go in the internal gaps.

Problem 3.28: Non-negative integer solutions

Show that there are

$$\binom{n+k-1}{k-1}$$

distinct non-negative integer-valued vectors $(x_1, \dots, x_k)$ satisfying

$$x_1 + \cdots + x_k = n, \quad x_i \geq 0 \text{ for all } i = 1, \dots, k.$$

Hint: Use the same stars-and-bars idea, but now allow some groups to be empty.

Solution 3.28

Again represent n as n identical stars. This time we want to split them into k groups, but some groups may be empty.

So we place $k - 1$ dividers among the n stars, allowing dividers to appear at the ends or next to each other.

This is equivalent to arranging

$$n \text{ stars and } k - 1 \text{ dividers}$$

in a row. The total number of symbols is

$$n + k - 1.$$

We simply choose the positions of the $k - 1$ dividers:

$$\boxed{\binom{n+k-1}{k-1}}.$$

Takeaways 3.28

- This is the standard stars-and-bars count for non-negative integer solutions.
- Allowing zero means dividers may be adjacent or may appear at the ends.

Problem 3.29: Three identities from $(1+x)^n$

Consider the binomial expansion

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \cdots + \binom{n}{n}x^n.$$

i) Deduce that

$$\binom{2n}{0} + \binom{2n}{2} + \cdots + \binom{2n}{2n} = \binom{2n}{1} + \binom{2n}{3} + \cdots + \binom{2n}{2n-1}.$$

ii) Integrate both sides with respect to x from 0 to 1, and deduce that

$$\sum_{r=0}^n \frac{1}{r+1} \binom{n}{r} = \frac{2^{n+1} - 1}{n+1}.$$

iii) Differentiate both sides with respect to x , then let $x = 1$, and deduce that

$$\binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \cdots + n\binom{n}{n} = n2^{n-1}.$$

Hint: For each part, start from the same binomial expansion and apply the requested operation directly to both sides.

Solution 3.29

i) Apply the identity

$$(1+x)^{2n} = \sum_{r=0}^{2n} \binom{2n}{r} x^r$$

and let $x = -1$. Then

$$\sum_{r=0}^{2n} \binom{2n}{r} (-1)^r = 0.$$

So

$$\binom{2n}{0} - \binom{2n}{1} + \binom{2n}{2} - \binom{2n}{3} + \cdots + \binom{2n}{2n} = 0.$$

Hence the sum of the even-index terms equals the sum of the odd-index terms:

$$\boxed{\binom{2n}{0} + \binom{2n}{2} + \cdots + \binom{2n}{2n} = \binom{2n}{1} + \binom{2n}{3} + \cdots + \binom{2n}{2n-1}}.$$

ii) Integrating from 0 to 1,

$$\int_0^1 (1+x)^n dx = \int_0^1 \sum_{r=0}^n \binom{n}{r} x^r dx.$$

The left-hand side is

$$\left[\frac{(1+x)^{n+1}}{n+1} \right]_0^1 = \frac{2^{n+1} - 1}{n+1}.$$

The right-hand side is

$$\sum_{r=0}^n \binom{n}{r} \int_0^1 x^r dx = \sum_{r=0}^n \binom{n}{r} \frac{1}{r+1}.$$

Hence

$$\boxed{\sum_{r=0}^n \frac{1}{r+1} \binom{n}{r} = \frac{2^{n+1} - 1}{n+1}}.$$

iii) Differentiating,

$$n(1+x)^{n-1} = \binom{n}{1} + 2\binom{n}{2}x + 3\binom{n}{3}x^2 + \cdots + n\binom{n}{n}x^{n-1}.$$

Now let $x = 1$:

$$n2^{n-1} = \binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \cdots + n\binom{n}{n}.$$

So

$$\boxed{\binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \cdots + n\binom{n}{n} = n2^{n-1}}.$$

Takeaways 3.29

- A single binomial expansion can produce several useful identities by substitution, integration, and differentiation.
- Setting $x = -1$ often produces alternating-sign identities.
- Integrating or differentiating term-by-term can turn coefficients into new weighted sums.

Problem 3.30: Making an amount of money with Australian denominations

Suppose we use the simplified Australian denominations (coins less than \$1 are not used):

$$\text{\$1, \$2, \$5, \$10, \$20, \$50, \$100.}$$

For a fixed positive integer amount M , let $N(M)$ be the number of ways to express $\$M$ using these coins and notes, where order does *not* matter.

- i) Write down a generating function whose coefficient of x^M is $N(M)$.
- ii) Hence explain how the problem of finding $N(M)$ can be turned into a coefficient problem.

Enrichment note: This problem is outside the HSC Mathematics Extension 1 syllabus, but it is a very useful example of how generating functions turn a counting problem into algebra.

Hint: For each denomination d , think about the possible contributions $0, d, 2d, 3d, \dots$

Solution 3.30

For each denomination, we write a generating function for how many of that denomination may be used:

$$1 + x + x^2 + x^3 + \dots \quad \text{for \$1,}$$

$$1 + x^2 + x^4 + x^6 + \dots \quad \text{for \$2,}$$

and similarly for the other notes.

So the full generating function is

$$G(x) = (1 + x + x^2 + \dots)(1 + x^2 + x^4 + \dots)(1 + x^5 + x^{10} + \dots)(1 + x^{10} + x^{20} + \dots) \\ (1 + x^{20} + x^{40} + \dots)(1 + x^{50} + x^{100} + \dots)(1 + x^{100} + x^{200} + \dots).$$

The coefficient of x^M in $G(x)$ is exactly the number of ways to make $\$M$, because each chosen power of x records how much each denomination contributes, and multiplying the choices combines them. So

$$N(M) = [x^M] G(x),$$

where $[x^M] G(x)$ means “the coefficient of x^M in $G(x)$.”

Outline of method. To solve a specific case, such as $M = 20$ or $M = 50$, expand only as far as needed and collect the coefficient of x^M . In practice, this is often done using recurrence methods or computer algebra, because the coefficients grow quickly.

Takeaways 3.30

- A generating function is a power series in which the coefficient of x^M stores the number of ways to make the amount M .
- For a sequence of numbers $a_0, a_1, a_2, \dots$, the generating function is

$$a_0 + a_1x + a_2x^2 + \dots$$

- For the \$1 denomination, the generating function is

$$1 + x + x^2 + x^3 + \dots$$

because we may use 0, 1, 2, 3, ... one-dollar coins, contributing \$0, \$1, \$2, \$3, ... respectively.

- The full generating function is the product of the generating functions for each denomination, because we can choose how many of each denomination to use independently.
- Each other denomination contributes a similar series, and multiplying these series combines the choices from all denominations.
- This is a standard higher-level combinatorics technique and is excellent enrichment beyond the HSC course.

Remark 3.1 (Algorithmic note)

We can also compute $N(M)$ using dynamic programming. This is the algorithmic version of finding the coefficient of x^M in the generating function.

```

1 def calculate_coin_combinations(target_amount, denominations):
2     """
3     Calculates the number of ways to make a target amount using given
4     denominations.
5     This is the algorithmic equivalent of finding the coefficient of  $x^M$ 
6     in our generating function.
7     """
8     # dp is an array where dp[i] will store the number of ways to make
9     # amount i.
10    # We initialize an array of zeros from 0 up to our target_amount.
11    dp = [0] * (target_amount + 1)
12
13    # Base case: There is exactly 1 way to make $0 (by choosing no
14    # coins)
15    dp[0] = 1
16
17    # Outer loop: We process one denomination bracket at a time.
18    # This is exactly like multiplying the next bracket in our
19    # generating function:
20    # (1 + x^2 + x^4...) * (1 + x^5 + x^10...)

```



```

17     for coin in denominations:
18
19         # Inner loop: We update the combinations for every amount
20         # from 'coin' up to the target.
21         for amount in range(coin, target_amount + 1):
22
23             # The number of ways to make the current 'amount'
24             # increases by the
25             # number of ways we could make the amount WITHOUT this
26             # current coin.
27             dp[amount] += dp[amount - coin]
28
29         # The final answer is stored at the index of our target amount
30         return dp[target_amount]
31
32 # --- Example Usage ---
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267
2268
2269
2270
2271
2272
2273
2274
2275
2276
2277
2278
2279
2280
2281
2282
2283
2284
2285
2286
2287
2288
2289
2290
2291
2292
2293
2294
2295
2296
2297
2298
2299
2300
2301
2302
2303
2304
2305
2306
2307
2308
2309
2310
2311
2312
2313
2314
2315
2316
2317
2318
2319
2320
2321
2322
2323
2324
2325
2326
2327
2328
2329
2330
2331
2332
2333
2334
2335
2336
2337
2338
2339
2340
2341
2342
2343
2344
2345
2346
2347
2348
2349
2350
2351
2352
2353
2354
2355
2356
2357
2358
2359
2360
2361
2362
2363
2364
2365
2366
2367
2368
2369
2370
2371
2372
2373
2374
2375
2376
2377
2378
2379
2380
2381
2382
2383
2384
2385
2386
2387
2388
2389
2390
2391
2392
2393
2394
2395
2396
2397
2398
2399
2400
2401
2402
2403
2404
2405
2406
2407
2408
2409
2410
2411
2412
2413
2414
2415
2416
2417
2418
2419
2420
2421
2422
2423
2424
2425
2426
2427
2428
2429
2430
2431
2432
2433
2434
2435
2436
2437
2438
2439
2440
2441
2442
2443
2444
2445
2446
2447
2448
2449
2450
2451
2452
2453
2454
2455
2456
2457
2458
2459
2460
2461
2462
2463
2464
2465
2466
2467
2468
2469
2470
2471
2472
2473
2474
2475
2476
2477
2478
2479
2480
2481
2482
2483
2484
2485
2486
2487
2488
2489
2490
2491
2492
2493
2494
2495
2496
2497
2498
2499
2500
2501
2502
2503
2504
2505
2506
2507
2508
2509
2510
2511
2512
2513
2514
2515
2516
2517
2518
2519
2520
2521
2522
2523
2524
2525
2526
2527
2528
2529
2530
2531
2532
2533
2534
2535
2536
2537
2538
2539
2540
2541
2542
2543
2544
2545
2546
2547
2548
2549
2550
2551
2552
2553
2554
2555
2556
2557
2558
2559
2560
2561
2562
2563
2564
2565
2566
2567
2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582
2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595
2596
2597
2598
2599
2600
2601
2602
2603
2604
2605
2606
2607
2608
2609
2610
2611
2612
2613
2614
2615
2616
2617
2618
2619
2620
2621
2622
2623
2624
2625
2626
2627
2628
2629
2630
2631
2632
2633
2634
2635
2636
2637
2638
2639
2640
2641
2642
2643
2644
2645
2646
2647
2648
2649
2650
2651
2652
2653
2654
2655
2656
2657
2658
2659
2660
2661
2662
266
```

Solution 3.31

The digits 1, 2, 3, 4 can be arranged in

$$4! = 24$$

different ways, so if we test PINs one by one, the correct PIN is equally likely to appear in positions

$$1, 2, 3, \dots, 24.$$

If the correct PIN is found on the m th attempt, then there have been $(m - 1)$ wrong attempts before success.

The waiting times after the wrong attempts are

$$4, 8, 16, \dots, 4 \cdot 2^{m-2}.$$

So the total waiting time before the m th attempt is

$$4 + 8 + 16 + \dots + 4 \cdot 2^{m-2} = 4(2^{m-1} - 1).$$

Therefore the average waiting time is

$$\frac{1}{24} \sum_{m=1}^{24} 4(2^{m-1} - 1).$$

Now

$$\sum_{m=1}^{24} 2^{m-1} = 2^{24} - 1,$$

so

$$\begin{aligned} \frac{1}{24} \sum_{m=1}^{24} 4(2^{m-1} - 1) &= \frac{4}{24} ((2^{24} - 1) - 24) \\ &= \frac{1}{6} (2^{24} - 25) \\ &= \frac{16777191}{6}. \end{aligned}$$

Hence the average time needed is

$$\boxed{\frac{16777191}{6} \text{ seconds}}$$

which is about

$$\boxed{2796198.5 \text{ seconds}}.$$

Takeaways 3.31

- When all possible arrangements are equally likely, the correct one is equally likely to occur in any position of your trial order.
- Exponentially growing waiting times often lead to a geometric series.
- Expected-value questions can often be solved by averaging over all possible positions of success.

Remark 3.2

“Space is big. Really big. You just won’t believe how vastly, hugely, mind-bogglingly big it is. I mean, you may think it’s a long way down the road to the chemist, but that’s just peanuts to space.”

Douglas Adams

Problem 3.32: Polynomial derivatives and a power-series expansion

Show that if

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

is any polynomial, then

$$f^{(k)}(0) = \begin{cases} a_k k!, & \text{for } k = 0, 1, 2, \dots, n, \\ 0, & \text{for } k > n. \end{cases}$$

Hence explain why

$$f(x) = \frac{f(0)}{0!} + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f^{(3)}(0)}{3!}x^3 + \cdots.$$

Enrichment note: This problem is outside the HSC Mathematics Extension 1 syllabus, but it is a very useful first step toward power series and Taylor series.

Hint: Differentiate a general term $a_r x^r$ exactly k times, then substitute $x = 0$. Most terms vanish.

Solution 3.32

Write

$$f(x) = \sum_{r=0}^n a_r x^r.$$

If we differentiate k times, then

$$\frac{d^k}{dx^k}(a_r x^r) = \begin{cases} a_r r(r-1)\cdots(r-k+1)x^{r-k}, & r \geq k, \\ 0, & r < k. \end{cases}$$

Now substitute $x = 0$. Every term with $r > k$ still contains a factor of x and becomes 0. The only surviving term is the one with $r = k$, giving

$$f^{(k)}(0) = a_k k!$$

for $k = 0, 1, 2, \dots, n$.

If $k > n$, every term has already vanished after enough differentiations, so

$$f^{(k)}(0) = 0.$$

Therefore

$$a_k = \frac{f^{(k)}(0)}{k!},$$

so the polynomial can be rewritten as

$$f(x) = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n = \frac{f(0)}{0!} + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \cdots + \frac{f^{(n)}(0)}{n!}x^n.$$

Since f is a polynomial, this series has only finitely many nonzero terms.

Takeaways 3.32

- Repeated derivatives at $x = 0$ recover the coefficients of a polynomial.
- The factor of $k!$ appears because differentiating x^k exactly k times gives $k!$.
- For any function $f(x)$ that can be expressed as a power series, the same idea gives the Taylor series at 0:

$$f(x) = \frac{f(0)}{0!} + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \dots$$

- This is the polynomial version of the idea behind Taylor series, so it is a natural enrichment topic beyond the HSC course.

Problem 3.33: Binary strings and a binomial identity

Computing is based on binary strings, which are sequences of 0s and 1s, such as 00101 and 1001110.

The problem in this question is to find the number N of binary strings that contain exactly a zeros and at most b ones, where a and b are whole numbers. We will count such strings in two different ways, and hence prove an identity.

a) Using nC_r notation, find how many binary strings with exactly a zeros contain:

- i) no 1s
- ii) one 1
- iii) two 1s
- iv) three 1s
- v) r 1s.

b) Hence prove that

$$N = \binom{a}{0} + \binom{a+1}{1} + \binom{a+2}{2} + \cdots + \binom{a+b}{b}.$$

c) Let S be a binary string consisting of a zeros and r ones, where $r \leq b$. Extend S to a longer binary string $S01 \cdots 1$ by adding 0 on the right, and then adding $b-r$ ones. How many 0s and 1s are there in the resulting string, and what is its total length?

d) Describe the inverse process by which a string of $(a+1)$ zeros and b ones can be converted to a string of a zeros and at most b ones, and show that the two processes are one-to-one correspondences.

e) Hence prove that

$$N = \binom{a+b+1}{b}.$$

f) Deduce that

$$\binom{a}{0} + \binom{a+1}{1} + \binom{a+2}{2} + \cdots + \binom{a+b}{b} = \binom{a+b+1}{b}.$$

Hint: For part (a), if a string has exactly a zeros and r ones, then its length is $a+r$, and you only need to choose where the r ones go. For parts (c) and (d), build a bijection between the two families of strings.

Solution 3.33

a) If a binary string has exactly a zeros and exactly r ones, then its total length is

$$a + r.$$

We may choose the positions of the r ones (or equivalently the a zeros), so the number of such strings is

$$\binom{a+r}{r}.$$

Hence:

$$\text{i) } \binom{a}{0}, \quad \text{ii) } \binom{a+1}{1}, \quad \text{iii) } \binom{a+2}{2}, \quad \text{iv) } \binom{a+3}{3}, \quad \text{v) } \binom{a+r}{r}.$$

b) Since the number of ones can be $0, 1, 2, \dots, b$, summing these cases gives

$$N = \binom{a}{0} + \binom{a+1}{1} + \binom{a+2}{2} + \dots + \binom{a+b}{b}.$$

c) Start with a binary string S containing a zeros and r ones, where $r \leq b$. Append one extra 0, then append $(b-r)$ ones. The resulting string has

$$a + 1$$

zeros and

$$r + (b - r) = b$$

ones, so its total length is

$$a + b + 1.$$

d) Conversely, given a binary string with $(a+1)$ zeros and b ones, locate the last zero in the string and delete it together with all the ones to its right.

What remains is a binary string with exactly a zeros and at most b ones.

This process is the inverse of the construction in part (c), so the two sets of strings are in one-to-one correspondence.

e) Therefore, the number N is equal to the number of binary strings with $(a+1)$ zeros and b ones. Such strings have length

$$a + b + 1,$$

and we choose the positions of the b ones, so

$$N = \binom{a+b+1}{b}.$$

f) Equating the two expressions for N gives

$$\boxed{\binom{a}{0} + \binom{a+1}{1} + \binom{a+2}{2} + \dots + \binom{a+b}{b} = \binom{a+b+1}{b}}.$$

Takeaways 3.33

- This is a nice example of proving an identity by counting the same set in two different ways.
- Strings with fixed numbers of zeros and ones are counted by choosing positions.
- Bijections are powerful: if two families of objects can be matched one-to-one, then they have the same size.

Problem 3.34: Derangements

A *derangement* of n distinct letters is a permutation of them so that no letter appears in its original position. For example, DABC is a derangement of ABCD, but DACB is not. Denote the number of derangements of n letters by $D(n)$.

- By listing all the derangements of A, AB, ABC and ABCD, find the values of $D(1)$, $D(2)$, $D(3)$ and $D(4)$.
- Suppose that we have formed a derangement of the five letters ABCDE. Let the last letter in the derangement be X , and exchange X with E. This puts E back into its original position. Either X is now also in its original position so that three letters are away from their original positions, or X is not in its original position so that four letters are away from their original positions. Hence explain why

$$D(5) = 4D(4) + 4D(3).$$

- Use this formula to evaluate $D(5)$. Then apply the corresponding arguments and formulae to evaluate $D(6)$, $D(7)$ and $D(8)$.
- Explain how this can be generalised to

$$D(n) = n! \left(\frac{1}{0!} - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots + \frac{(-1)^n}{n!} \right).$$

- The alternating sum in part (d) approaches $\frac{1}{e}$ as $n \rightarrow \infty$. Explain what this means for the ratio

$$\frac{D(n)}{n!}$$

and give both a combinatorial and a probabilistic interpretation of this result.

Enrichment note: Derangements are a classic higher-level counting topic. They are not part of the standard HSC Mathematics Extension 1 syllabus, but they are an excellent example of recurrence relations in combinatorics.

Hint: For part (a), list small cases directly. For parts (b) and (c), fix where E goes, then split into the two cases described in the question. For part (d), think about inclusion-exclusion: start from all permutations and subtract those with one or more fixed points.

Solution 3.34

a) By direct listing:

$$D(1) = 0, \quad D(2) = 1, \quad D(3) = 2, \quad D(4) = 9.$$

b) In a derangement of ABCDE, the letter E cannot stay in the last position, so it has 4 possible positions. Let the last letter be X , and swap E with X . Either X lands in its own position, giving $D(3)$ possibilities, or it does not, giving $D(4)$ possibilities. Hence

$$D(5) = 4D(4) + 4D(3).$$

c) Using $D(4) = 9$ and $D(3) = 2$,

$$D(5) = 4 \cdot 9 + 4 \cdot 2 = 44.$$

More generally,

$$D(n) = (n-1)(D(n-1) + D(n-2)).$$

Thus

$$D(5) = 44, \quad D(6) = 265, \quad D(7) = 1854, \quad D(8) = 14833.$$

d) Let A_i be the set of permutations in which the i th letter stays fixed. Then

$$|A_i| = (n-1)!, \quad |A_i \cap A_j| = (n-2)!,$$

and in general fixing k specified letters leaves

$$(n-k)!$$

permutations. Hence, by inclusion-exclusion,

$$D(n) = n! - \binom{n}{1}(n-1)! + \binom{n}{2}(n-2)! - \cdots + (-1)^n \binom{n}{n}(n-n)!.$$

Using $\binom{n}{k}(n-k)! = \frac{n!}{k!}$, we can conclude that,

$$D(n) = n! \left(\frac{1}{0!} - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots + \frac{(-1)^n}{n!} \right).$$

e) Dividing by $n!$ gives

$$\frac{D(n)}{n!} = \frac{1}{0!} - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots + \frac{(-1)^n}{n!}.$$

As $n \rightarrow \infty$, the right-hand side approaches $\frac{1}{e}$, so

$$\frac{D(n)}{n!} \rightarrow \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} = \frac{1}{e}.$$

Takeaways 3.34


- *Combinatorially*, about a fraction $\frac{1}{e}$ of all permutations are derangements for large n .
- *Probabilistically*, a random permutation has no fixed points with probability approaching $\frac{1}{e}$.
- A derangement is a permutation with no fixed points.
- Small cases are often best handled by direct listing before a general pattern appears.
- This problem is a good example of a recurrence relation arising from careful case-work.
- Inclusion-exclusion gives a closed formula for $D(n)$ by counting permutations with fixed points.
- The ratio $\frac{D(n)}{n!}$ approaching $\frac{1}{e}$ means derangements make up about $\frac{1}{e}$ of all permutations when n is large.
- The Taylor series for e^x is

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}.$$

Problem 3.35: A monochromatic triangle on six vertices

A regular octahedron has 6 vertices. Label them $A_0, A_1, \dots, A_5$. Each pair of vertices is joined by a rod, and each rod is coloured either yellow or blue.

- Explain why there are 15 rods.
- Each set of three vertices, together with the rods joining them, forms a triangle. Explain why there are 20 such triangles.
- Explain why there must be at least one triangle whose rods all have the same colour.



Enrichment note: This is a small and beautiful example of a Ramsey-type idea. It is beyond the standard HSC Mathematics Extension 1 syllabus, but it is a great illustration of how counting and the pigeonhole principle can force structure.

Hint: For part (c), focus on one vertex, say A_0 , and look at the 5 rods from that vertex. Use the pigeonhole principle on their colours.

Solution 3.35

a) A rod joins a pair of distinct vertices. Since there are 6 vertices, the number of rods is

$$\binom{6}{2} = 15.$$

b) Any choice of 3 vertices determines a triangle. So the number of triangles is

$$\binom{6}{3} = 20.$$

c) Focus on the vertex A_0 . It is joined to the other 5 vertices by 5 rods, and each rod is coloured either yellow or blue.

By the pigeonhole principle, at least 3 of these 5 rods must have the same colour. Without loss of generality, suppose

$$A_0A_1, A_0A_2, A_0A_3$$

are all yellow.

Now consider the triangle formed by A_1, A_2, A_3 .

- If one of the rods A_1A_2 , A_1A_3 , or A_2A_3 is yellow, then together with A_0 it forms a yellow triangle.
- If none of those rods is yellow, then all three are blue, so $A_1A_2A_3$ is a blue triangle.

In either case, there is at least one triangle whose rods all have the same colour.

Takeaways 3.35

- This is a classic example of combining counting with the pigeonhole principle.
- Looking at all edges from one vertex is the key step that forces a repeated colour.
- This is the first nontrivial case of a famous Ramsey-theory fact: in any two-colouring of the edges of K_6 , there is a monochromatic triangle.

Problem 3.36: A more general form of the binomial theorem

The classical binomial theorem expands

$$(1 + x)^n$$

when n is a non-negative integer. In more advanced mathematics, a similar expansion can be used when n is negative, fractional, or any real number, provided we interpret the right-hand side as an infinite power series.

- a) Show that for a non-negative integer n , the binomial expansion can be written as

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$$

- b) Assuming the same pattern continues for more general values of n , generate the power-series expansions of:

i) $\frac{1}{1-x}$

ii) $\frac{1}{(1-x)^2}$

iii) $\frac{1}{(1+x)^2}$

iv) $\sqrt{1+x}$.

- c) Verify, using your expansions in parts (b)(i) and (b)(ii), that

$$\frac{d}{dx} \left(\frac{1}{1-x} \right) = \frac{1}{(1-x)^2},$$

assuming that the power series may be differentiated term-by-term.

Enrichment note: This problem is outside the HSC Mathematics Extension 1 syllabus. It connects the classical binomial theorem to infinite power series, Taylor series, and later to Gamma-function ideas that allow binomial coefficients to be extended beyond whole numbers.

Hint: In part (a), rewrite the usual binomial coefficients in factorial form. In part (b), substitute values such as $n = -1$, $n = -2$, and $n = \frac{1}{2}$ into the general pattern.

Solution 3.36

a) For a non-negative integer n , the usual binomial theorem gives

$$(1+x)^n = \sum_{r=0}^n \binom{n}{r} x^r.$$

Now

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)(n-2)\cdots(n-r+1)}{r!},$$

so the expansion may be written as

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \cdots.$$

b)(i) Take $n = -1$:

$$(1-x)^{-1} = 1 + x + x^2 + x^3 + \cdots.$$

So

$$\boxed{\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots.}$$

b)(ii) Take $n = -2$:

$$(1-x)^{-2} = 1 + 2x + 3x^2 + 4x^3 + \cdots.$$

So

$$\boxed{\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \cdots.}$$

b)(iii) Replace x by $-x$ in part (ii):

$$\boxed{\frac{1}{(1+x)^2} = 1 - 2x + 3x^2 - 4x^3 + \cdots.}$$

b)(iv) Take $n = \frac{1}{2}$:

$$(1+x)^{1/2} = 1 + \frac{1}{2}x + \frac{\frac{1}{2}(-\frac{1}{2})}{2!}x^2 + \frac{\frac{1}{2}(-\frac{1}{2})(-\frac{3}{2})}{3!}x^3 + \cdots.$$

So

$$\boxed{\sqrt{1+x} = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 - \frac{5}{128}x^4 + \cdots.}$$

c) From part (b)(i),

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots.$$

Differentiating term-by-term gives

$$\frac{d}{dx} \left(\frac{1}{1-x} \right) = 1 + 2x + 3x^2 + 4x^3 + \cdots.$$

But this is exactly the expansion found in part (b)(ii), so

$$\boxed{\frac{d}{dx} \left(\frac{1}{1-x} \right) = \frac{1}{(1-x)^2}.}$$

Takeaways 3.36

- The usual finite binomial theorem is the beginning of a much more general power-series expansion.
- For any real number n , and for $|x| < 1$,

$$(1+x)^n = \sum_{r=0}^{\infty} \binom{n}{r} x^r = 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 + \dots$$

- When n is not a non-negative integer, the binomial expansion usually becomes an infinite Taylor-style series rather than a finite polynomial. For any real value of n ,

$$(1+x)^n = \sum_{r=0}^{\infty} \frac{f^{(r)}(0)}{r!} x^r, \quad \text{where } f(x) = (1+x)^n.$$

Refer to the problem “Polynomial derivatives and a power-series expansion” for the idea of finding these coefficients using derivatives at 0.

- The generalized coefficient may be written using Gamma functions:

$$\binom{n}{r} = \frac{\Gamma(n+1)}{\Gamma(r+1)\Gamma(n-r+1)},$$

which extends the factorial formula to non-integer values of n .

- This topic is outside HSC Mathematics Extension 1, but it is excellent enrichment for students interested in university mathematics.

Problem 3.37: Isomers — a possible computing project

Because of branching, it is difficult to count the number of possible hydrocarbons with, say, 20 carbon atoms.

For example, for butane C_4H_{10} there are two carbon chains, and for pentane C_5H_{12} there are three carbon chains.

Investigate the number of possible carbon chains for hydrocarbons with increasing numbers of carbon atoms.

- Explain why this is a counting problem about *connected graphs* with no cycles, where each carbon atom has degree at most 4.
- Find the numbers of possible carbon chains for $n = 1, 2, 3, 4, 5, 6, 7$ carbon atoms.
- If you can write code, design a program to generate all possible carbon chains for a given n and remove duplicates up to relabelling.

Enrichment note: This is a hard problem. Writing code is strongly recommended. It is outside the HSC Mathematics Extension 1 syllabus, but it is an excellent project linking combinatorics, graph theory, chemistry, and computing.

Hint: Model each carbon chain as a tree: vertices are carbon atoms and edges are carbon-carbon bonds. Two drawings represent the same chain if they are the same up to relabelling of the vertices.

Solution 3.37

a) A carbon chain can be modelled by a graph in which:

- each vertex represents a carbon atom
- each edge represents a carbon-carbon bond.

For an alkane chain, the carbon skeleton is connected and has no cycles, so it is a tree. Also, each carbon atom can make at most 4 bonds, so each vertex has degree at most 4.

b) The numbers of possible carbon chains for small n are:

n	1	2	3	4	5	6	7
number of chains	1	1	1	2	3	5	9

These are the counts of non-isomorphic trees on n vertices with maximum degree at most 4. For small n , the degree restriction does not yet exclude any tree.

c) A practical computational approach is:

- generate trees on n vertices by adding one new vertex at a time
- reject any graph with a vertex of degree more than 4
- identify graphs that are the same up to relabelling by converting them to a canonical form
- count the distinct canonical forms.

Remark 3.3 (Python idea)

The following Python snippet generates non-isomorphic trees on n vertices by adding one new vertex at a time and keeping one canonical representative of each isomorphism class. It is suitable for small values such as $n \leq 7$.

```

1 from itertools import permutations
2
3
4 def canonical_form(adj):
5     n = len(adj)
6     best = None
7     for perm in permutations(range(n)):
8         bits = []
9         for i in range(n):
10             for j in range(i + 1, n):
11                 bits.append("1" if perm[j] in adj[perm[i]] else "0")
12             code = "".join(bits)
13             if best is None or code < best:
14                 best = code
15     return best
16
17
18 def generate_trees(n):
19     if n == 1:
20         return [tuple([frozenset()])]
21

```

```
22     previous = generate_trees(n - 1)
23     seen = {}
24
25     for tree in previous:
26         base = [set(neighbours) for neighbours in tree]
27         for v in range(n - 1):
28             if len(base[v]) >= 4:
29                 continue
30
31             graph = [set(neighbours) for neighbours in base] + [set()]
32             graph[v].add(n - 1)
33             graph[n - 1].add(v)
34
35             key = canonical_form(graph)
36             seen[key] = tuple(frozenset(x) for x in graph)
37
38     return list(seen.values())
39
40
41 for n in range(1, 8):
42     print(n, len(generate_trees(n)))
```

Takeaways 3.37

- This problem is better viewed as a graph-theory and computing project than as a routine exam question.
- **Combinatorial explosion** means the number of possible structures grows extremely quickly as n increases, so direct listing soon becomes impractical.
- Because of this explosion, writing code is often the most realistic way to investigate the problem.
- This is outside the HSC Mathematics Extension 1 syllabus, but it is an excellent example of how combinatorics connects to chemistry and computer science.

4 Conclusion

Combinatorics rewards careful reading as much as technical skill. Many HSC questions become much easier once you decide whether the task is an arrangement, a selection, or a probability question built from counting. As this booklet grows, it can become a compact reference for the most common combinatorics patterns students meet in Extension 1.

Contact Information:

LinkedIn: <https://www.linkedin.com/in/nguyenvuhung/>

GitHub: <https://github.com/vuhung16au/>

Repository: <https://github.com/vuhung16au/math-olympiad-ml/tree/main/HSC-Combinatorics>

A Appendices

A.1 Concepts: Injective, Surjective, and Bijective

Understanding these concepts is incredibly important for combinatorics, particularly when answering “how many” or proving that two sets have the same number of elements without actually counting them one by one. In combinatorics, formalizing a counting process as a function provides powerful insights:

- **Injective (One-to-one) tracking:** If you can construct an injective mapping from set X to set Y , you instantly prove that $|X| \leq |Y|$ (the size of X is less than or equal to Y). If you try to map more elements into fewer targets, an injection is impossible—this is the formal basis of the **Pigeonhole Principle**. When counting arrangements, injections describe ways to assign objects to distinct slots without repetition (like nP_r).
- **Surjective (Onto) coverage:** If you can construct a surjective mapping from set X to set Y , you prove that $|X| \geq |Y|$. In counting, surjections are used when we want to count the number of ways to distribute objects into boxes where *no box is left empty*.
- **Bijective (Perfect pairing) counting:** If you can construct a bijection between set X and set Y , you prove that $|X| = |Y|$. This is called **Bijective Proof** and it is one of the most elegant techniques in combinatorics. If set X is very hard to count, but you can link it perfectly to set Y which is easy to count, then you have solved the problem! (For instance, stars and bars is a bijection between dividing items and placing dividers).


Here is the complete breakdown of how these mappings work mathematically.

Remark A.1 (Syllabus Note)

These concepts are outside the HSC Mathematics Extension 1 and 2 syllabus, but they are very good to know for higher education. In combinatorics, they help formalize counting by pairing elements perfectly between sets. In functions, bijections define exactly when an inverse function exists.

Injective / One-to-one

A function is injective if every output is mapped by *at most* one input. In other words, no two different inputs will ever give you the same output.




- **Formal Definition:** If $f(a) = f(b)$, then it must be true that $a = b$.
- **Visualizing it:** Imagine a group of people (domain) and a group of chairs (codomain). An injective mapping means no two people are sitting in the same chair. There might be empty chairs left over, but no chair is double-occupied.

- **Math Example:** $f(x) = 2x$ is injective. Every unique x produces a unique y . However, $f(x) = x^2$ is *not* injective because $f(2) = 4$ and $f(-2) = 4$ (two different inputs hit the same output).

Surjective / Onto


A function is surjective if every possible output in the target set (codomain) is hit by *at least* one input. There are no “leftover” or unreachable outputs.



- **Visualizing it:** Every chair in the room has at least one person sitting in it. (Some chairs might have two people squeezing into them, but no chair is empty).

Bijjective / One-to-one correspondence

A function is bijective if it is **both** injective and surjective. Every input maps to a unique output, and every possible output is mapped to by an input.



- **Visualizing it:** It is a perfect pairing. There is exactly one person for every chair, and exactly one chair for every person. No empty chairs, no double-occupied chairs.
- **Why it matters:** Bijective functions are incredibly important in mathematics because they are the **only** functions that have a perfect **inverse function** (f^{-1}). Because the mapping is a perfect two-way street, you can flawlessly travel backward from the output to the exact input.
- **Math Example:** $f(x) = 2x + 1$ (where x is any real number) is a bijection. You can easily find its inverse: $f^{-1}(x) = \frac{x-1}{2}$.