




HSC Mathematics Advanced

Sequences and Series

Vu Hung Nguyen



Download PDF and resources:

[https://github.com/vuhung16au/
math-olympiad-ml](https://github.com/vuhung16au/math-olympiad-ml)



Read online at Vu's Maths Hub:

<https://vumaths.com/>

This work is licensed under CC BY-NC 4.0 (Attribution-Non-Commercial), see the LICENSE file on the github page for more info.

"Sometimes it is the people no one imagines anything of who do the things that no one can imagine."

Alan Turing

Contents

1	Introduction	6
1.1	Project Overview	6
1.2	Target Audience	6
1.3	How to Use This Booklet	6
1.4	Topics Covered	6
1.5	Where This Leads Next	6
2	Fundamentals Review	7
2.1	How Sequences Are Specified	7
2.2	Arithmetic Sequences (AP)	7
	Problem 2.1: Recognizing an AP	7
2.3	Arithmetic Series	7
	Problem 2.2: AP Sum	7
2.4	Geometric Sequences (GP)	8
	Problem 2.3: Recognizing a GP	8
2.5	Geometric Series	8
	Problem 2.4: Finite GP Sum	8
2.6	Limiting Sum of a Geometric Series	8
	Problem 2.5: Infinite GP	9
2.7	Recurring Decimals and Geometric Series	9
2.8	Sequence Convergence Theorems	9
2.9	Sigma and Pi Notation Quick Rules	9
3	Part 1: Problems and Solutions (Detailed)	11
3.1	Basic Sequence Problems	11
	Problem 3.1: Arithmetic nth term	11
	Problem 3.2: Find the common ratio	11
	Problem 3.3: Sum of an AP	11
	Problem 3.4: Recurring decimal as a fraction	11
	Problem 3.5: Chebyshev recurrence: first kind	12
3.2	Medium Sequence Problems	12
	Problem 3.6: Mixed AP constraints	12
	Problem 3.7: Finite GP with unknown index	13
	Problem 3.8: Financial sequences and geometric sums	14
	Problem 3.9: Telescoping sums and inductive proof	16
	Problem 3.10: Binomial coefficients and partial sums	18
	Problem 3.11: The Recursive Geometry of Chebyshev Sequences	20
3.3	Advanced Sequence Problems	22
	Problem 3.12: Recursive sequence and limiting value	22
	Problem 3.13: The Dynamics of Iteration	23
	Problem 3.14: The Geometry of Harmonic Means	25
	Problem 3.15: Chebyshev composition	28
	Problem 3.16: Bounding a sigma sum	29
4	Part 2: Problems with Hints and Solutions (Concise)	31
4.1	Basic Sequence Problems	31
	Problem 4.1: Quick AP term	31
	Problem 4.2: Quick GP sum	31
	Problem 4.3: Quick Chebyshev value	31

Problem 4.4: Arithmetic pricing model	32
Problem 4.5: Simple interest and AP	32
Problem 4.6: Recurring decimal to fraction	33
Problem 4.7: Nested shaded squares	33
Problem 4.8: Fraction of a recurring decimal	33
4.2 Medium Sequence Problems	34
Problem 4.9: Indexing with sigma notation	34
Problem 4.10: Infinite geometric interpretation	34
Problem 4.11: Second-kind Chebyshev recurrence	35
Problem 4.12: Alternative telescoping patterns	35
Problem 4.13: Surd telescoping	36
Problem 4.14: Central symmetry in a GP	36
Problem 4.15: Fibonacci and Lucas	37
Problem 4.16: Linear combinations of two GPs	37
Problem 4.17: Partial fractions and telescoping series	38
4.3 Advanced Sequence Problems	40
Problem 4.18: Convergence condition	40
Problem 4.19: AP and GP crossover	40
Problem 4.20: Chebyshev roots	41
Problem 4.21: Architecture of GP sums	41
Problem 4.22: AP space and difference operator	42
Problem 4.23: Basis and projection in AP space	42
Problem 4.24: Countable and uncountable sets	43
Problem 4.25: Prime denominators and cycle length	43
Problem 4.26: Pythagorean means	44
Problem 4.27: Recurring decimals and modular structure	44
Problem 4.28: Sissa and arithmetico-geometric sums	45
Problem 4.29: Discrete calculus and sums of powers	46
Problem 4.30: The Fibonacci Generating Function	47
5 Conclusion	49
6 Appendices	50
6.1 Appendix A: Formula Sheet	50
6.2 Appendix B: Common Sequence Traps	51
6.3 Appendix C: Rigorous Definition of the Limit of a Sequence	51
6.4 Appendix D: Limit Superior and Limit Inferior	53
6.5 Appendix E: Discrete Calculus and the Difference Operator	54

1 Introduction

1.1 Project Overview

This booklet is a focused companion for NSW HSC Mathematics Extension 1 students studying sequences and series. It is designed to connect core sequence skills to exam-style problem solving with clear methods and structured solutions.

1.2 Target Audience

This resource is written for Year 12 students in Extension 1, and for tutors and teachers who want a compact booklet with consistent notation and worked examples.

1.3 How to Use This Booklet

- Read the fundamentals review first, then attempt the detailed problems in Part 1.
- Use Part 2 for timed practice and fluency, using hints only when needed.
- Keep a formula sheet while you work, but always justify each step from first principles.
- Track common slips, especially sign errors in geometric sums and indexing mistakes in sigma notation.

1.4 Topics Covered

This booklet develops the following Extension 1 ideas:

- defining sequences by term formula, recursion, and context
- arithmetic sequences and arithmetic series
- geometric sequences and geometric series
- finite sum formulas and sigma notation
- limiting sum of a geometric series when $|r| < 1$
- links between recurring decimals and geometric series
- mixed AP/GP problem solving in HSC-style questions

1.5 Where This Leads Next

Sequence fluency supports later work in calculus, proof, and Extension 2 style reasoning. In particular, convergence arguments and recursive definitions form a bridge to harder topics such as iterative methods and infinite processes.

2 Fundamentals Review

This section is a compact reference for notation, formulas, and common reasoning patterns used throughout the booklet.

2.1 How Sequences Are Specified

A sequence is an ordered list of numbers, usually written as $(a_n)_{n \geq 1}$. Common ways to define a sequence are:

- explicit formula: a_n written directly in terms of n
- recursive formula: each term written using previous terms
- contextual rule: terms generated from a worded process

Examples.

$$a_n = 3n - 1, \quad b_1 = 2, \quad b_{n+1} = b_n + 5, \quad c_n = (-1)^n.$$

2.2 Arithmetic Sequences (AP)

An arithmetic sequence has constant difference d :

$$a_{n+1} - a_n = d.$$

If the first term is a , then

$$a_n = a + (n - 1)d.$$

Problem 2.1: Recognizing an AP

Given $a_1 = 7$ and $a_{n+1} = a_n - 3$, find a_{10} .

Solution 2.1

Here $d = -3$, so

$$a_{10} = 7 + (10 - 1)(-3) = 7 - 27 = -20.$$

2.3 Arithmetic Series

The sum of the first n terms of an AP is:

$$S_n = \frac{n}{2}(2a + (n - 1)d) = \frac{n}{2}(a + l),$$

where l is the n th term.

Problem 2.2: AP Sum

Find $\sum_{k=1}^{20} (5 + 2(k - 1))$.

Solution 2.2

This is an AP with $a = 5$, $d = 2$, and $n = 20$. The 20th term is

$$l = 5 + 19 \cdot 2 = 43.$$

So

$$S_{20} = \frac{20}{2}(5 + 43) = 10 \cdot 48 = 480.$$

2.4 Geometric Sequences (GP)

A geometric sequence has constant ratio r :

$$\frac{a_{n+1}}{a_n} = r \quad (a_n \neq 0).$$

If the first term is a , then

$$a_n = ar^{n-1}.$$

Problem 2.3: Recognizing a GP

If $a_1 = 81$ and $r = \frac{1}{3}$, find a_5 .

Solution 2.3

$$a_5 = 81 \left(\frac{1}{3}\right)^4 = 81 \cdot \frac{1}{81} = 1.$$

2.5 Geometric Series

For $r \neq 1$, the sum of the first n terms is

$$S_n = a \frac{1 - r^n}{1 - r} = a \frac{r^n - 1}{r - 1}.$$

Use the form that keeps signs clean for your r value.

Problem 2.4: Finite GP Sum

Find $\sum_{k=0}^6 3 \cdot 2^k$.

Solution 2.4

This is $a = 3$, $r = 2$, $n = 7$ terms (from $k = 0$ to 6):

$$S_7 = 3 \frac{2^7 - 1}{2 - 1} = 3(128 - 1) = 381.$$

2.6 Limiting Sum of a Geometric Series

If $|r| < 1$, then $r^n \rightarrow 0$ as $n \rightarrow \infty$, and the infinite sum exists:

$$S_\infty = \frac{a}{1 - r}.$$

If $|r| \geq 1$, the geometric series does not converge to a finite limiting sum.

Problem 2.5: Infinite GP

Evaluate $2 + \frac{2}{3} + \frac{2}{9} + \frac{2}{27} + \dots$.

Solution 2.5

Here $a = 2$, $r = \frac{1}{3}$, and $|r| < 1$, so

$$S_{\infty} = \frac{2}{1 - \frac{1}{3}} = \frac{2}{\frac{2}{3}} = 3.$$

2.7 Recurring Decimals and Geometric Series

Recurring decimals can be written as geometric series. For example,

$$0.\overline{7} = 0.7 + 0.07 + 0.007 + \dots$$

is geometric with $a = 0.7$ and $r = 0.1$. So

$$0.\overline{7} = \frac{0.7}{1 - 0.1} = \frac{0.7}{0.9} = \frac{7}{9}.$$

The “Cyclic” Magic of $\frac{1}{7}$

The fraction $\frac{1}{7}$ produces the most famous repeating pattern in elementary number theory.

The Pattern:

$$\frac{1}{7} = 0.\dot{1}4285\dot{7} \text{ (a 6-digit period).}$$

2.8 Sequence Convergence Theorems

Monotone Convergence Theorem (MCT) The MCT provides a “shortcut” for proving a sequence converges by looking at its overall behavior rather than its specific terms.

- **The Rule:** If a sequence is monotone (always increasing or always decreasing) and it is bounded (it never goes past a certain value), then it must converge.

The Squeeze Theorem (Sandwich Theorem) The Squeeze Theorem is used when a sequence is “messy” (like one containing a sine or cosine function) but is trapped between two “cleaner” sequences that share the same limit.

- **The Rule:** Suppose we have three sequences (a_n) , (b_n) , and (c_n) such that for all n beyond a certain point:

$$a_n \leq b_n \leq c_n$$

- **The Result:** If $\lim_{n \rightarrow \infty} a_n = L$ and $\lim_{n \rightarrow \infty} c_n = L$, then the “squeezed” sequence in the middle must also have that limit: $\lim_{n \rightarrow \infty} b_n = L$.

2.9 Sigma and Pi Notation Quick Rules

Summation (Σ):

$$\sum_{k=1}^n (u_k + v_k) = \sum_{k=1}^n u_k + \sum_{k=1}^n v_k, \quad \sum_{k=1}^n c u_k = c \sum_{k=1}^n u_k.$$

Index shifts must preserve both the expression and bounds consistently.

Product (II):

$$\prod_{k=1}^n (u_k v_k) = \left(\prod_{k=1}^n u_k \right) \left(\prod_{k=1}^n v_k \right), \quad \prod_{k=1}^n c u_k = c^n \prod_{k=1}^n u_k.$$

Note that a constant c is raised to the power of the number of terms when pulled outside a product, unlike sums where it acts as a simple multiplier.

Takeaways 2.1

- Always identify whether the question asks for a term a_n , a finite sum S_n , or a limit.
- AP questions depend on constant difference; GP questions depend on constant ratio.
- For infinite geometric sums, check $|r| < 1$ before using $S_\infty = \frac{a}{1-r}$.
- Track indexing carefully in sigma notation to avoid off-by-one errors.
- Chebyshev polynomial questions usually depend on either the recurrence relation or the trigonometric identities.

3 Part 1: Problems and Solutions (Detailed)

Part 1 contains full solutions with method explanations.

3.1 Basic Sequence Problems

Problem 3.1: Arithmetic nth term

An arithmetic sequence has first term 4 and common difference 5. Find a_{30} .

Solution 3.1

Use $a_n = a + (n - 1)d$:

$$a_{30} = 4 + 29 \cdot 5 = 149.$$

Problem 3.2: Find the common ratio

In a geometric sequence, $a_2 = 12$ and $a_5 = 324$. Find the common ratio r and the first term a_1 .

Solution 3.2

From $a_n = ar^{n-1}$,

$$a_2 = ar = 12, \quad a_5 = ar^4 = 324.$$

Divide to remove a :

$$\frac{a_5}{a_2} = r^3 = \frac{324}{12} = 27 \implies r = 3.$$

Then $a_1 = a = \frac{12}{3} = 4$.

Problem 3.3: Sum of an AP

Find the sum of the first 40 terms of the arithmetic sequence

$$9, 13, 17, \dots$$

Solution 3.3

Here $a = 9$, $d = 4$, $n = 40$. The 40th term is

$$l = 9 + 39 \cdot 4 = 165.$$

So

$$S_{40} = \frac{40}{2}(9 + 165) = 20 \cdot 174 = 3480.$$

Problem 3.4: Recurring decimal as a fraction

Express $0.\overline{23}$ as a rational number.

Solution 3.4

$$0.\overline{23} = \frac{23}{99}.$$

A geometric view is

$$0.23 + 0.0023 + 0.000023 + \dots$$

with first term $\frac{23}{100}$ and ratio $\frac{1}{100}$.

Problem 3.5: Chebyshev recurrence: first kind

The Chebyshev polynomials of the first kind satisfy

$$T_0(x) = 1, \quad T_1(x) = x, \quad T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x).$$

Use the recurrence to find $T_2(x)$ and $T_3(x)$, then evaluate $T_3\left(\frac{1}{2}\right)$.

Hint: Build the polynomials one at a time: first use $T_2 = 2xT_1 - T_0$, then use $T_3 = 2xT_2 - T_1$.

Solution 3.5

$$T_2(x) = 2x^2 - 1.$$

Then

$$T_3(x) = 2x(2x^2 - 1) - x = 4x^3 - 3x.$$

So

$$T_3\left(\frac{1}{2}\right) = 4\left(\frac{1}{8}\right) - 3\left(\frac{1}{2}\right) = \frac{1}{2} - \frac{3}{2} = -1.$$

Takeaways 3.1

- Chebyshev polynomials can be generated recursively from two starting values.
- Work in order; each new polynomial depends on the previous two.
- Substitution is easiest after simplifying the polynomial.

3.2 Medium Sequence Problems**Problem 3.6: Mixed AP constraints**

An arithmetic sequence has $a_3 = 11$ and $a_{12} = 47$. Find a_1 and d , then determine S_{25} .

Solution 3.6

Use $a_n = a + (n - 1)d$:

$$a + 2d = 11, \quad a + 11d = 47.$$

Subtract:

$$9d = 36 \implies d = 4,$$

then $a = 11 - 8 = 3$. Now

$$a_{25} = 3 + 24 \cdot 4 = 99,$$

so

$$S_{25} = \frac{25}{2}(3 + 99) = \frac{25}{2} \cdot 102 = 1275.$$

Problem 3.7: Finite GP with unknown index

For the geometric sequence with first term 5 and ratio 2, find n such that

$$S_n = 2555.$$

Solution 3.7

$$S_n = 5 \frac{2^n - 1}{2 - 1} = 5(2^n - 1) = 2555.$$

So

$$2^n - 1 = 511 \implies 2^n = 512 = 2^9.$$

Hence $n = 9$.

Problem 3.8: Financial sequences and geometric sums

A loan of $\$L$ is taken out at an interest rate of r per period. At the end of each period, a constant repayment of $\$P$ is made after interest has been added to the balance. Let M_n be the balance owing after n periods.

- (i) Write down expressions for M_1 and M_2 in terms of L, r , and P .
- (ii) Show that the balance after n periods can be written as

$$M_n = L(1+r)^n - P[1 + (1+r) + (1+r)^2 + \cdots + (1+r)^{n-1}].$$

- (iii) Using the formula for the sum of a geometric progression, prove that

$$M_n = L(1+r)^n - \frac{P((1+r)^n - 1)}{r}.$$

- (iv) A car loan of $\$20\,000$ is taken at 1% interest per month. Calculate the monthly repayment P required to reduce the balance to zero at the end of 24 months.

Hint:

- **For (i):** Add interest first, then subtract the repayment.
- **For (ii):** Expand the first few balances and look for the pattern.
- **For (iii):** The bracket is a finite GP with first term 1 and ratio $1+r$.
- **For (iv):** Set $M_{24} = 0$, with $L = 20000$ and $r = 0.01$.

Solution 3.8

(i) After one period,

$$M_1 = L(1 + r) - P.$$

After two periods,

$$M_2 = (L(1 + r) - P)(1 + r) - P = L(1 + r)^2 - P(1 + r) - P.$$

(ii) Continuing this pattern gives

$$M_n = L(1 + r)^n - P[(1 + r)^{n-1} + (1 + r)^{n-2} + \cdots + (1 + r) + 1].$$

Reordering the bracket,

$$M_n = L(1 + r)^n - P[1 + (1 + r) + (1 + r)^2 + \cdots + (1 + r)^{n-1}].$$

(iii) The bracket is a finite GP with first term 1, ratio $1 + r$, and n terms:

$$1 + (1 + r) + \cdots + (1 + r)^{n-1} = \frac{(1 + r)^n - 1}{(1 + r) - 1} = \frac{(1 + r)^n - 1}{r}.$$

Hence

$$M_n = L(1 + r)^n - \frac{P((1 + r)^n - 1)}{r}.$$

(iv) Here $L = 20000$, $r = 0.01$, and $n = 24$. Set $M_{24} = 0$:

$$0 = 20000(1.01)^{24} - \frac{P((1.01)^{24} - 1)}{0.01}.$$

Thus

$$P = \frac{20000(1.01)^{24}(0.01)}{(1.01)^{24} - 1}.$$

Using $(1.01)^{24} \approx 1.269735$,

$$P \approx \$941.47.$$

Takeaways 3.2

- Loan balances can be modelled by applying interest and repayment recursively.
- Repeated repayments form a finite geometric sum after compounding.
- Setting the final balance to zero gives the repayment required to clear a loan.

Problem 3.9: Telescoping sums and inductive proof

Consider the sequence

$$a_k = \frac{2}{k(k+1)(k+2)} \quad (k \geq 1).$$

Define the partial sums

$$S_n = \sum_{k=1}^n a_k.$$

(i) Show, using partial fractions, that

$$\frac{2}{k(k+1)(k+2)} = \frac{1}{k(k+1)} - \frac{1}{(k+1)(k+2)}.$$

(ii) Demonstrate that the sequence of partial sums telescopes. Hence find a simplified expression for S_n in terms of n .

(iii) Prove the expression for S_n found in part (ii) by mathematical induction for all $n \geq 1$.

(iv) Determine the limiting sum of the series as $n \rightarrow \infty$.

Hint:

- **For (i):** Put the two fractions over the common denominator $k(k+1)(k+2)$.
- **For (ii):** Write out the first few terms of S_n and look for cancellation.
- **For (iii):** Use $S_{m+1} = S_m + a_{m+1}$ in the induction step.
- **For (iv):** Take the limit of the closed form for S_n .

Solution 3.9

(i)

$$\frac{1}{k(k+1)} - \frac{1}{(k+1)(k+2)} = \frac{k+2-k}{k(k+1)(k+2)} = \frac{2}{k(k+1)(k+2)}.$$

(ii) Therefore

$$a_k = \frac{1}{k(k+1)} - \frac{1}{(k+1)(k+2)}.$$

So

$$\begin{aligned} S_n &= \left(\frac{1}{1 \cdot 2} - \frac{1}{2 \cdot 3} \right) + \left(\frac{1}{2 \cdot 3} - \frac{1}{3 \cdot 4} \right) + \cdots + \left(\frac{1}{n(n+1)} - \frac{1}{(n+1)(n+2)} \right) \\ &= \frac{1}{2} - \frac{1}{(n+1)(n+2)}. \end{aligned}$$

(iii) We prove

$$S_n = \frac{1}{2} - \frac{1}{(n+1)(n+2)}.$$

For $n = 1$,

$$S_1 = \frac{2}{1 \cdot 2 \cdot 3} = \frac{1}{3}$$

and

$$\frac{1}{2} - \frac{1}{2 \cdot 3} = \frac{1}{2} - \frac{1}{6} = \frac{1}{3}.$$

Now assume the formula holds for $n = m$. Then

$$\begin{aligned} S_{m+1} &= S_m + \frac{2}{(m+1)(m+2)(m+3)} \\ &= \frac{1}{2} - \frac{1}{(m+1)(m+2)} + \left(\frac{1}{(m+1)(m+2)} - \frac{1}{(m+2)(m+3)} \right) \\ &= \frac{1}{2} - \frac{1}{(m+2)(m+3)}. \end{aligned}$$

This is the required formula with $n = m + 1$, so the result holds for all $n \geq 1$.

(iv)

$$\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \left(\frac{1}{2} - \frac{1}{(n+1)(n+2)} \right) = \frac{1}{2}.$$

Takeaways 3.3

- Telescoping sums work by rewriting terms so that most neighbouring parts cancel.
- Induction is a useful way to verify a closed form found by pattern spotting.
- A finite partial-sum formula can make the limiting sum immediate.

Problem 3.10: Binomial coefficients and partial sums

Let

$$S_n = \sum_{k=0}^n \binom{n}{k}$$

be the sum of the binomial coefficients for a fixed integer $n \geq 1$.

(i) Use the binomial theorem expansion of $(1+x)^n$ to prove that $S_n = 2^n$.

(ii) Use Pascal's identity

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$$

to prove the recurrence relation $S_n = 2S_{n-1}$.

(iii) Evaluate the alternating partial sum

$$A_n = \sum_{k=0}^n (-1)^k \binom{n}{k}$$

and explain why the sum of even-indexed coefficients equals the sum of odd-indexed coefficients.

(iv) By differentiating the expansion of $(1+x)^n$ with respect to x , show that

$$\sum_{k=1}^n k \binom{n}{k} = n2^{n-1}.$$

Hint:

- **For (i):** Differentiate first, then substitute $x = 1$.
- **For (iii):** Substitute $x = -1$ into the binomial theorem.
- **For (ii):** Sum Pascal's identity over the appropriate range of k .
- **For (i):** Substitute $x = 1$ into the binomial theorem.

Solution 3.10

(i) By the binomial theorem,

$$(1+x)^n = \sum_{k=0}^n \binom{n}{k} x^k.$$

Putting $x = 1$ gives

$$2^n = \sum_{k=0}^n \binom{n}{k} = S_n.$$

(ii) Using Pascal's identity,

$$\begin{aligned} S_n &= \sum_{k=0}^n \binom{n}{k} \\ &= \sum_{k=0}^n \binom{n-1}{k} + \sum_{k=0}^n \binom{n-1}{k-1}. \end{aligned}$$

The out-of-range terms are zero, so each sum equals S_{n-1} . Hence

$$S_n = 2S_{n-1}.$$

(iii) By the binomial theorem again,

$$A_n = \sum_{k=0}^n (-1)^k \binom{n}{k} = (1-1)^n = 0.$$

Therefore

$$\sum_{\substack{0 \leq k \leq n \\ k \text{ even}}} \binom{n}{k} - \sum_{\substack{0 \leq k \leq n \\ k \text{ odd}}} \binom{n}{k} = 0,$$

so the even-indexed and odd-indexed sums are equal.

(iv) Differentiate

$$(1+x)^n = \sum_{k=0}^n \binom{n}{k} x^k$$

to get

$$n(1+x)^{n-1} = \sum_{k=1}^n k \binom{n}{k} x^{k-1}.$$

Putting $x = 1$ gives

$$\sum_{k=1}^n k \binom{n}{k} = n2^{n-1}.$$

Takeaways 3.4

- Substituting special values into the binomial theorem turns an expansion into a sum identity.
- Pascal's identity explains why the total sum of coefficients doubles from one row to the next.
- Differentiating a generating expansion is a powerful way to evaluate weighted sums.

Problem 3.11: The Recursive Geometry of Chebyshev Sequences

Let a sequence of functions $(T_n(x))$ be defined for $n \geq 0$ by the recurrence relation

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x),$$

with initial terms $T_0(x) = 1$ and $T_1(x) = x$.

- (i) Calculate the explicit polynomial expressions for $T_2(x)$ and $T_3(x)$.
- (ii) Using the identity

$$\cos(A + B) + \cos(A - B) = 2 \cos A \cos B,$$

prove by mathematical induction that for all $n \geq 0$,

$$T_n(\cos \theta) = \cos(n\theta).$$

- (iii) Let x_0 be a root of $T_n(x)$. Show that $|x_0| \leq 1$. Hence, determine the exact values of the n distinct roots of $T_n(x)$ in terms of n .
- (iv) Consider the sequence of values $v_n = T_n(1.5)$. Show that (v_n) satisfies a linear recurrence relation with constant coefficients. Find a closed-form expression for v_n and determine the value of the limit

$$\lim_{n \rightarrow \infty} \frac{v_{n+1}}{v_n}.$$

Hint:

- **For (i):** Solve the characteristic equation for the recurrence $v_{n+1} = 3v_n - v_{n-1}$.
- **For (iii):** Solve $\cos(n\theta) = 0$ for θ , then use $x = \cos \theta$.
- **For (ii):** Use the sum-to-product identity with $A = k\theta$ and $B = \theta$.
- **For (i):** Substitute the previous two polynomials into the recurrence formula.

Solution 3.11

(i) Directly from the recurrence,

$$T_2(x) = 2x^2 - 1, \quad T_3(x) = 2x(2x^2 - 1) - x = 4x^3 - 3x.$$

(ii) The cases $n = 0, 1$ give $1 = \cos 0$ and $T_1(\cos \theta) = \cos \theta$. Suppose

$$T_k(\cos \theta) = \cos(k\theta), \quad T_{k-1}(\cos \theta) = \cos((k-1)\theta).$$

Then

$$\begin{aligned} T_{k+1}(\cos \theta) &= 2 \cos \theta T_k(\cos \theta) - T_{k-1}(\cos \theta) \\ &= 2 \cos \theta \cos(k\theta) - \cos((k-1)\theta) \\ &= \cos((k+1)\theta), \end{aligned}$$

using $\cos((k+1)\theta) + \cos((k-1)\theta) = 2 \cos(k\theta) \cos \theta$. Hence $T_n(\cos \theta) = \cos(n\theta)$ for all $n \geq 0$.

(iii) If $x > 1$, write $x = \cosh u$; then $T_n(x) = \cosh(nu) > 0$. If $x < -1$, write $x = -\cosh u$; then $T_n(x) = (-1)^n \cosh(nu) \neq 0$. Thus every real root satisfies $|x_0| \leq 1$.

Now write $x = \cos \theta$. The roots satisfy

$$T_n(\cos \theta) = 0 \iff \cos(n\theta) = 0 \iff n\theta = \frac{(2k-1)\pi}{2}.$$

So the n distinct roots are

$$x_k = \cos\left(\frac{(2k-1)\pi}{2n}\right), \quad k = 1, 2, \dots, n.$$

(iv) With $v_n = T_n(1.5)$,

$$v_{n+1} = 3v_n - v_{n-1}, \quad v_0 = 1, \quad v_1 = \frac{3}{2}.$$

The characteristic equation is $\lambda^2 - 3\lambda + 1 = 0$, with roots

$$\alpha = \frac{3 + \sqrt{5}}{2}, \quad \beta = \frac{3 - \sqrt{5}}{2}.$$

Hence $v_n = A\alpha^n + B\beta^n$. From $v_0 = 1$ and $v_1 = \frac{3}{2}$, $A = B = \frac{1}{2}$, so

$$v_n = \frac{1}{2}\alpha^n + \frac{1}{2}\beta^n.$$

Since $|\beta| < \alpha$, the dominant term is α^n , and therefore

$$\lim_{n \rightarrow \infty} \frac{v_{n+1}}{v_n} = \alpha = \frac{3 + \sqrt{5}}{2}.$$

Takeaways 3.5

- The recurrence produces the Chebyshev polynomials one term at a time.
- The identity $T_n(\cos \theta) = \cos(n\theta)$ turns polynomial roots into trigonometric roots.
- Evaluating a polynomial recurrence at a fixed value gives an ordinary numerical recurrence.

3.3 Advanced Sequence Problems

Problem 3.12: Recursive sequence and limiting value

Let (x_n) be defined by

$$x_1 = 16, \quad x_{n+1} = 2 + \frac{1}{x_n} \quad (n \geq 1).$$

- (i) Find exact values of x_2, x_3, x_4 .
- (ii) Assuming $x_n \rightarrow L > 0$, show that $L = 1 + \sqrt{2}$.
- (iii) Prove that

$$|x_{n+1} - L| = \frac{|x_n - L|}{Lx_n},$$

and deduce that for $n \geq 2$,

$$|x_{n+1} - L| < \frac{1}{4}|x_n - L|.$$

- (iv) Hence show $x_n \rightarrow 1 + \sqrt{2}$.

Solution 3.12

(i)

$$x_2 = 2 + \frac{1}{16} = \frac{33}{16}, \quad x_3 = 2 + \frac{1}{33} = \frac{82}{33}, \quad x_4 = 2 + \frac{1}{82} = \frac{197}{82}.$$

For $n \geq 2$, we have $x_n > 2$ because each term is $2 + \frac{1}{x_{n-1}}$ with positive second part.

(ii) If $x_n \rightarrow L$, then taking limits in the recurrence gives

$$L = 2 + \frac{1}{L} \implies L^2 - 2L - 1 = 0.$$

So $L = 1 \pm \sqrt{2}$. Since terms are positive and > 2 from $n \geq 2$, we take

$$L = 1 + \sqrt{2}.$$

(iii)

$$x_{n+1} - L = \left(2 + \frac{1}{x_n}\right) - \left(2 + \frac{1}{L}\right) = \frac{L - x_n}{Lx_n},$$

thus

$$|x_{n+1} - L| = \frac{|x_n - L|}{Lx_n}.$$

Now $L = 1 + \sqrt{2} > 2$ and $x_n > 2$ for $n \geq 2$, so $Lx_n > 4$. Hence

$$|x_{n+1} - L| < \frac{1}{4}|x_n - L|, \quad n \geq 2.$$

(iv) Repeatedly applying the inequality gives geometric decay of the error:

$$|x_n - L| \leq \left(\frac{1}{4}\right)^{n-2} |x_2 - L| \rightarrow 0.$$

Therefore $x_n \rightarrow L = 1 + \sqrt{2}$.

Takeaways 3.6

- For recursive limits, first solve the fixed-point equation.
- A contraction inequality gives a rigorous convergence proof.
- The same method works for many rational recursions.
- The sequence x_n is related to the continued fraction of $1 + \sqrt{2}$. It can be expressed as

$$x_{n+1} = 2 + \frac{1}{x_n} = 2 + \frac{1}{2 + \frac{1}{x_{n-1}}} = \cdots = 2 + \frac{1}{2 + \frac{1}{2 + \cdots + \frac{1}{x_1}}}$$

Problem 3.13: The Dynamics of Iteration

Consider the function

$$g(x) = \sqrt{x+2}.$$

We define an iterative sequence by

$$x_{n+1} = g(x_n), \quad x_1 = 1.$$

- (i) Calculate x_2, x_3 , and x_4 to three decimal places. By solving

$$L = \sqrt{L+2},$$

determine the exact positive limit L that the sequence appears to be approaching.

- (ii) Show that

$$0 < g'(x) < \frac{1}{2}$$

for all $x > 0$.

- (iii) Prove that

$$|x_{n+1} - L| < \frac{1}{2}|x_n - L|.$$

Hence deduce that the sequence converges to L for any initial value $x_1 > 0$.

- (iv) Define a function $f(x)$ such that Newton's method would converge to the same limit L found in part (i), and write down the specific iterative formula for this $f(x)$.

Hint:

- **For (i):** Substitute repeatedly into $x_{n+1} = \sqrt{x_n + 2}$, then solve the fixed-point equation.
- **For (ii):** Differentiate $g(x) = \sqrt{x + 2}$ and note where the derivative is largest for $x > 0$.
- **For (iii):** Use either the mean value theorem or rationalise $\sqrt{x_n + 2} - \sqrt{L + 2}$.
- **For (iv):** Rearrange $L = \sqrt{L + 2}$ into a polynomial equation.

Solution 3.13

(i)

$$x_2 = \sqrt{3} \approx 1.732, \quad x_3 = \sqrt{2 + \sqrt{3}} \approx 1.932, \quad x_4 \approx 1.983.$$

If $x_n \rightarrow L > 0$, then

$$L = \sqrt{L + 2} \implies L^2 - L - 2 = 0 \implies (L - 2)(L + 1) = 0.$$

Thus the positive limit is

$$L = 2.$$

(ii)

$$g'(x) = \frac{1}{2\sqrt{x + 2}}.$$

For $x > 0$, this is positive and

$$g'(x) < \frac{1}{2\sqrt{2}} < \frac{1}{2}.$$

(iii) Since $L = 2$ and $g(L) = L$,

$$\begin{aligned} |x_{n+1} - L| &= \left| \sqrt{x_n + 2} - \sqrt{L + 2} \right| \\ &= \frac{|x_n - L|}{\sqrt{x_n + 2} + \sqrt{L + 2}} \\ &< \frac{1}{2} |x_n - L|, \end{aligned}$$

because $x_n > 0$ and $\sqrt{L + 2} = 2$. Repeating this inequality gives

$$|x_n - L| < \left(\frac{1}{2}\right)^{n-1} |x_1 - L| \rightarrow 0,$$

so $x_n \rightarrow L = 2$ for any $x_1 > 0$.

(iv) From the fixed-point equation,

$$L^2 - L - 2 = 0.$$

So take

$$f(x) = x^2 - x - 2.$$

Newton's method gives

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^2 - x_n - 2}{2x_n - 1}.$$

Takeaways 3.7

- Fixed point is the value of x such that $f(x) = x$.
- Fixed points often reveal the likely limit of an iterative sequence.
- A contraction inequality proves convergence by forcing errors to shrink geometrically.
- Newton's method can target the same limit by solving the fixed-point equation as a root problem.

Problem 3.14: The Geometry of Harmonic Means

A sequence (h_n) is in harmonic progression (HP) if the sequence of reciprocals

$$x_n = \frac{1}{h_n}$$

forms an arithmetic progression (AP).

- Given that the third term of an HP is 1 and the sixth term is $\frac{1}{2}$, find the first term h_1 and the common difference d of the underlying AP.
- The harmonic mean H of two positive numbers a and b is defined so that a, H, b is an HP. Show algebraically that

$$H = \frac{2ab}{a+b}.$$

- Let

$$A = \frac{a+b}{2}, \quad G = \sqrt{ab}$$

be the arithmetic and geometric means of a and b . Prove that $G^2 = AH$. Hence prove that, for distinct positive real numbers a and b ,

$$H < G < A.$$

- Consider the infinite harmonic series

$$\sum_{n=1}^{\infty} \frac{1}{n}.$$

By comparing the sum of the first 2^k terms to blocks of constant lower bounds, prove that this harmonic series does not have a finite limiting sum.

Hint:

- **For (i):** Convert the HP terms into AP terms x_3 and x_6 , then use $x_n = a + (n-1)d$.
- **For (ii):** If a, H, b is an HP, then $\frac{1}{a}, \frac{1}{H}, \frac{1}{b}$ is an AP.
- **For (iii):** Multiply A and H . For the inequality, use $(a-b)^2 > 0$ or $(\sqrt{a} - \sqrt{b})^2 > 0$.
- **For (iv):** Group terms as $\left(\frac{3}{1} + \frac{1}{1}\right)$, then $\left(\frac{5}{1} + \frac{1}{1} + \frac{1}{8}\right)$, and so on.

Solution 3.14

(i) Since $x_n = \frac{1}{h_n}$,

$$x_3 = 1, \quad x_6 = 2.$$

For the underlying AP,

$$x_6 - x_3 = 3d \implies 2 - 1 = 3d \implies d = \frac{1}{3}.$$

Also

$$x_1 = x_3 - 2d = 1 - \frac{2}{3} = \frac{1}{3},$$

so

$$h_1 = \frac{1}{x_1} = 3.$$

(ii) Since $\frac{1}{a}, \frac{1}{H}, \frac{1}{b}$ is an AP, the middle term is the average of the outer terms:

$$\frac{1}{H} = \frac{1}{2} \left(\frac{1}{a} + \frac{1}{b} \right) = \frac{a+b}{2ab}.$$

Therefore

$$H = \frac{2ab}{a+b}.$$

(iii) Using the formulas for A and H ,

$$AH = \frac{a+b}{2} \cdot \frac{2ab}{a+b} = ab = G^2.$$

For $a \neq b$,

$$(\sqrt{a} - \sqrt{b})^2 > 0 \implies a + b > 2\sqrt{ab} \implies A > G.$$

Since $G^2 = AH$ and all quantities are positive, $A > G$ gives

$$H = \frac{G^2}{A} < G.$$

Hence

$$H < G < A.$$

(iv) Group the harmonic series into dyadic blocks:

$$1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4} \right) + \left(\frac{1}{5} + \cdots + \frac{1}{8} \right) + \cdots.$$

Each block after the first has sum greater than or equal to $\frac{1}{2}$:

$$\frac{1}{3} + \frac{1}{4} > \frac{1}{4} + \frac{1}{4} = \frac{1}{2}, \quad \frac{1}{5} + \cdots + \frac{1}{8} > 4 \cdot \frac{1}{8} = \frac{1}{2},$$

and similarly

$$\frac{1}{2^{j-1}+1} + \cdots + \frac{1}{2^j} > 2^{j-1} \cdot \frac{1}{2^j} = \frac{1}{2}.$$

Thus the partial sums exceed

$$1 + \frac{1}{2} + \underbrace{\frac{1}{2} + \frac{1}{2} + \cdots + \frac{1}{2}}_{\text{arbitrarily many terms}},$$

so they grow without bound. Therefore the harmonic series has no finite limiting sum.

Takeaways 3.8

- Harmonic progressions are arithmetic progressions viewed through reciprocals.
- For positive distinct numbers, the classical means satisfy $H < G < A$.
- A sequence can have terms tending to zero while its infinite series still diverges.

Problem 3.15: Chebyshev composition

The first-kind Chebyshev polynomials $T_n(x)$ are defined by the recurrence

$$T_0(x) = 1, \quad T_1(x) = x, \quad T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x).$$

It is given that $T_n(\cos \theta) = \cos(n\theta)$ for all $n \geq 0$ and $\theta \in \mathbb{R}$. (Do NOT prove this fact.) For first-kind Chebyshev polynomials, prove that

$$T_m(T_n(x)) = T_{mn}(x)$$

for all x of the form $x = \cos \theta$. Then use this result to find

$$T_2(T_3(x)).$$

Hint: Use $T^n(\cos \theta) = \cos(n\theta)$. For the explicit polynomial, use $T_6(x)$.

Solution 3.15

Let $x = \cos \theta$. Then

$$T_m(T_n(x)) = T_m(\cos(n\theta)) = \cos(mn\theta) = T_{mn}(\cos \theta) = T_{mn}(x).$$

Therefore

$$T_2(T_3(x)) = T_6(x).$$

Using the recurrence after $T_5(x) = 16x^5 - 20x^3 + 5x$,

$$T_6(x) = 2xT_5(x) - T_4(x) = 32x^6 - 48x^4 + 18x^2 - 1.$$

Takeaways 3.9

- The trigonometric definition makes Chebyshev composition simple.
- Composition of first-kind Chebyshev polynomials multiplies the indices.
- Hard-looking polynomial composition can often be avoided by using structure.

Problem 3.16: Bounding a sigma sum

Let the sequence u_n be defined by

$$u_n = \frac{1^2}{n^3 + 1} + \frac{2^2}{n^3 + 2} + \cdots + \frac{n^2}{n^3 + n}$$

for integers $n \geq 1$. You may assume the standard result

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}. (\text{Do NOT prove this.})$$

(i) Explain why, for all integers k such that $1 \leq k \leq n$,

$$\frac{k^2}{n^3 + n} \leq \frac{k^2}{n^3 + k} \leq \frac{k^2}{n^3 + 1}.$$

(ii) Hence, evaluate $\lim_{n \rightarrow \infty} u_n$.

- Hint:**
- **Part (i):** Since k is positive, compare the sizes of $n^3 + 1$, $n^3 + k$, and $n^3 + n$.
 - **Part (ii):** Sum the inequality from part (i) over $k = 1$ to n . Factor out denominators independent of k , substitute the sum-of-squares formula, and apply the Squeeze Theorem.

Solution 3.16

(i) For $1 \leq k \leq n$, the denominators satisfy

$$n^3 + 1 \leq n^3 + k \leq n^3 + n.$$

Taking reciprocals reverses the inequality (all terms are positive):

$$\frac{1}{n^3 + n} \leq \frac{1}{n^3 + k} \leq \frac{1}{n^3 + 1}.$$

Multiplying through by the positive term k^2 gives

$$\frac{k^2}{n^3 + n} \leq \frac{k^2}{n^3 + k} \leq \frac{k^2}{n^3 + 1}.$$

(ii) Summing from $k = 1$ to n ,

$$\sum_{k=1}^n \frac{k^2}{n^3 + n} \leq u_n \leq \sum_{k=1}^n \frac{k^2}{n^3 + 1}.$$

Factoring out the k -independent denominators,

$$\frac{1}{n^3 + n} \sum_{k=1}^n k^2 \leq u_n \leq \frac{1}{n^3 + 1} \sum_{k=1}^n k^2.$$

Substituting the given formula,

$$\frac{n(n+1)(2n+1)}{6(n^3+n)} \leq u_n \leq \frac{n(n+1)(2n+1)}{6(n^3+1)}.$$

As $n \rightarrow \infty$, the highest power in numerator and denominator is n^3 , so both bounds tend to

$$\frac{2}{6} = \frac{1}{3}.$$

By the Squeeze Theorem,

$$\lim_{n \rightarrow \infty} u_n = \frac{1}{3}.$$

Takeaways 3.10

- Bounding by manipulating denominators is a core technique for rigorous sequence limits.
- Denominators independent of the summation index k can be factored out of sigma notation.
- This style of question links algebraic series manipulation with limit evaluation via the Squeeze Theorem.

4 Part 2: Problems with Hints and Solutions (Concise)

Part 2 gives extra practice with upside-down hints and concise solutions.

4.1 Basic Sequence Problems

Problem 4.1: Quick AP term

Find a_{18} for the arithmetic sequence $2, 7, 12, \dots$

Hint: Use $a_n = a + (n-1)d$ with $a = 2$ and $d = 5$.

Solution 4.1

$$a_{18} = 2 + 17 \cdot 5 = 87.$$

Problem 4.2: Quick GP sum

Find

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{128}.$$

Hint: This is a finite GP with $a = 1$, $r = \frac{1}{2}$, and 8 terms.

Solution 4.2

$$S_8 = \frac{1 - (\frac{1}{2})^8}{1 - \frac{1}{2}} = 2 \left(1 - \frac{1}{256} \right) = \frac{255}{128}.$$

Problem 4.3: Quick Chebyshev value

For the second-kind Chebyshev polynomial

$$U_2(x) = 4x^2 - 1,$$

find $U_2\left(\frac{1}{2}\right)$.

Hint: Substitute $x = \frac{1}{2}$ carefully.

Solution 4.3

$$U_2\left(\frac{1}{2}\right) = 4\left(\frac{1}{2}\right)^2 - 1 = 1 - 1 = 0.$$

Takeaways 4.1

- For a given Chebyshev polynomial, direct substitution is often enough.
- Square fractions before multiplying by the coefficient.

Problem 4.4: Arithmetic pricing model

An installer charges \$400 for the first window and \$100 for each additional window.

- Write the total cost for 1, 2, 3, and 4 windows.
- Find a formula for the total cost A_n for n windows.
- Find A_{15} .
- If the budget is strictly less than \$10,000, what is the maximum number of windows?

Hint: Model A_n as an arithmetic sequence with first term 400 and difference 100.

Solution 4.4

$$A_n = 400 + (n - 1)100 = 100n + 300.$$

So $A_{15} = 1800$. For the budget:

$$100n + 300 < 10000 \implies n < 97,$$

hence the maximum integer is $n = 96$.

Problem 4.5: Simple interest and AP

A principal of \$2,000,000 is invested at 5.7% p.a. simple interest. Let A_n be the total amount after n years.

- Find A_1, A_2, A_3, A_4 .
- Find a formula for A_n and evaluate A_{12} .
- How many whole years before the amount exceeds \$6,000,000?

Hint: Simple interest adds a fixed amount each year, so (A_n) is arithmetic.

Solution 4.5

Yearly interest is $0.057 \times 2,000,000 = 114,000$. Hence

$$A_n = 2,000,000 + 114,000n.$$

So

$$A_1 = 2,114,000, \quad A_2 = 2,228,000, \quad A_3 = 2,342,000, \quad A_4 = 2,456,000,$$

and

$$A_{12} = 3,368,000.$$

For exceeding \$6,000,000:

$$2,000,000 + 114,000n > 6,000,000 \implies n > 35.087 \dots$$

so $n = 36$ years.

Problem 4.6: Recurring decimal to fraction

Convert each to a fraction:

$$0.2, \quad 0.\dot{2}, \quad 0.1\dot{2}.$$

Hint: Use place value for terminating decimals and GP ideas for recurring decimals.

Solution 4.6

$$0.2 = \frac{1}{5}, \quad 0.\dot{2} = \frac{2}{9}, \quad 0.1\dot{2} = \frac{11}{90}.$$

Problem 4.7: Nested shaded squares

In a unit square, shade the bottom-left quarter. Then repeatedly subdivide the top-right quarter into four equal squares and shade its bottom-left quarter. Find the total shaded fraction.

Hint: Shaded areas form an infinite GP with first term $\frac{1}{4}$ and ratio $\frac{1}{4}$.

Solution 4.7

$$S_\infty = \frac{\frac{1}{4}}{1 - \frac{1}{4}} = \frac{1}{3}.$$

So the shaded fraction is $\frac{1}{3}$.

Problem 4.8: Fraction of a recurring decimal

Find the fraction for $0.1\dot{2}\dot{3}$?

Hint: Use $a = \frac{1000}{123}$ and $r = \frac{1}{1000}$ in your S_∞ formula.

Solution 4.8

$$S_{\infty} = \frac{\frac{123}{1000}}{1 - \frac{1}{1000}} = \frac{123}{999} = \frac{41}{333}.$$

Takeaways 4.2

- Every recurring decimal is a geometric series sum S_{∞} .

4.2 Medium Sequence Problems**Problem 4.9: Indexing with sigma notation**

Given

$$\sum_{k=1}^n (3k - 1) = 119,$$

find n .

Hint: Split the sum into $3 \sum k - \sum 1$.

Solution 4.9

$$\sum_{k=1}^n (3k - 1) = 3 \cdot \frac{n(n+1)}{2} - n = \frac{3n^2 + n}{2}.$$

Set equal to 119:

$$\frac{3n^2 + n}{2} = 119 \implies 3n^2 + n - 238 = 0.$$

$$\Delta = 1 + 2856 = 2857.$$

This is not a perfect square, so there is no integer n with exact value 119.

Problem 4.10: Infinite geometric interpretation

Evaluate

$$0.4\overline{2} = 0.42222 \dots$$

as a fraction.

Hint: Write as $0.4 + 0.02222 \dots$ and convert the recurring part using a GP.

Solution 4.10

$$0.02222 \dots = \frac{2}{90} = \frac{1}{45}.$$

So

$$0.4\overline{2} = \frac{2}{5} + \frac{1}{45} = \frac{18}{45} + \frac{1}{45} = \frac{19}{45}.$$

Problem 4.11: Second-kind Chebyshev recurrence

The second-kind Chebyshev polynomials satisfy

$$U_0(x) = 1, \quad U_1(x) = 2x, \quad U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x).$$

Use this recurrence to find $U_3(x)$ and $U_4(x)$.

Hint: First find $U_2(x)$, then continue to $U_3(x)$ and $U_4(x)$.

Solution 4.11

$$U_2(x) = 2x(2x) - 1 = 4x^2 - 1.$$

$$U_3(x) = 2x(4x^2 - 1) - 2x = 8x^3 - 4x.$$

$$U_4(x) = 2x(8x^3 - 4x) - (4x^2 - 1) = 16x^4 - 12x^2 + 1.$$

Takeaways 4.3

- The first-kind and second-kind Chebyshev polynomials use the same recurrence form.
- The starting values determine which family is produced.
- Organize recurrence work line by line to avoid algebra slips.

Problem 4.12: Alternative telescoping patterns

(i) Evaluate $\sum_{n=0}^4 (-1)^{n+1}(n+5)$.

(ii) Show $\frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}$.

(iii) Hence find

$$S_N = \sum_{n=1}^N \left(\frac{1}{n} - \frac{1}{n+1} \right)$$

and $\lim_{N \rightarrow \infty} S_N$.

Hint: Part (iii) is a direct telescoping sum after part (ii).

Solution 4.12

$$\sum_{n=0}^4 (-1)^{n+1} (n+5) = -7.$$

Also

$$\frac{1}{n(n+1)} = \frac{(n+1) - n}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}.$$

So

$$S_N = 1 - \frac{1}{N+1}, \quad \lim_{N \rightarrow \infty} S_N = 1.$$

Problem 4.13: Surd telescoping

Let

$$a_k = \frac{1}{\sqrt{k+1} + \sqrt{k}}, \quad S_n = \sum_{k=1}^n a_k.$$

- (i) Show $a_k = \sqrt{k+1} - \sqrt{k}$.
- (ii) Find S_n .
- (iii) Evaluate S_{99} .
- (iv) Find the smallest integer N such that $S_N > 10$.

Hint: Rationalize with the conjugate and then telescope.

Solution 4.13

$$a_k = \frac{\sqrt{k+1} - \sqrt{k}}{(k+1) - k} = \sqrt{k+1} - \sqrt{k}.$$

Hence

$$S_n = (\sqrt{2} - \sqrt{1}) + \cdots + (\sqrt{n+1} - \sqrt{n}) = \sqrt{n+1} - 1.$$

So $S_{99} = 10 - 1 = 9$. For $S_N > 10$:

$$\sqrt{N+1} - 1 > 10 \implies N > 120,$$

smallest such integer is $N = 121$.

Problem 4.14: Central symmetry in a GP

In a GP, define $P_n = T_1 T_2 \cdots T_n$.

- (i) Show $P_n = a^n r^{\frac{n(n-1)}{2}}$.
- (ii) Show $T_k T_{n-k+1} = T_1 T_n$.
- (iii) If $n = 11$ and middle term $T_6 = 5$, find P_{11} .

Hint: Use $T_k = ar^{k-1}$ and pair symmetric terms.

Solution 4.14

$$P_n = a(ar) \cdots (ar^{n-1}) = a^n r^{0+1+\cdots+(n-1)} = a^n r^{\frac{n(n-1)}{2}}.$$

Also

$$T_k T_{n-k+1} = ar^{k-1} \cdot ar^{n-k} = a^2 r^{n-1} = T_1 T_n.$$

For odd $n = 11$, the product is middle-term power:

$$P_{11} = T_6^{11} = 5^{11}.$$

Problem 4.15: Fibonacci and Lucas

Define

$$F_1 = 1, F_2 = 1, F_n = F_{n-1} + F_{n-2},$$

$$L_1 = 1, L_2 = 3, L_n = L_{n-1} + L_{n-2}.$$

Show that

$$L_n = F_{n-1} + F_{n+1}.$$

Hint: Set $A_n = L_n + F_n$ and $B_n = L_n - F_n$, then compare with Fibonacci terms.

Solution 4.15

From initial terms and recurrence,

$$A_n = L_n + F_n = 2F_{n+1}, \quad B_n = L_n - F_n = 2F_{n-1}.$$

Adding gives

$$2L_n = 2F_{n+1} + 2F_{n-1},$$

so

$$L_n = F_{n-1} + F_{n+1}.$$

Problem 4.16: Linear combinations of two GPs

Let $T_n = ar^{n-1}$ and $U_n = AR^{n-1}$. Define $W_n = T_n + U_n$.

(i) Show (W_n) satisfies

$$W_{n+2} - (r + R)W_{n+1} + rRW_n = 0.$$

(ii) Solve

$$X_{n+2} - 5X_{n+1} + 6X_n = 0, \quad X_1 = 5, \quad X_2 = 13.$$

Hint: For (ii), the characteristic roots are 2 and 3.

Solution 4.16

Substitute $W_n = ar^{n-1} + AR^{n-1}$ directly to verify (i). For (ii), write

$$X_n = \alpha 2^{n-1} + \beta 3^{n-1}.$$

Using $X_1 = 5$, $X_2 = 13$:

$$\alpha + \beta = 5, \quad 2\alpha + 3\beta = 13$$

gives $\alpha = 2$, $\beta = 3$. Hence

$$X_n = 2^n + 3^n.$$

Problem 4.17: Partial fractions and telescoping series

Let

$$U_n = \frac{1}{n(n+1)(n+2)}$$

where n is a positive integer.

(i) Express the general term U_n in partial fractions by decomposing it into the form:

$$\frac{1}{n(n+1)(n+2)} = \frac{A}{n} + \frac{B}{n+1} + \frac{C}{n+2}$$

where A , B , and C are constants to be determined.

(ii) By grouping the resulting fractions to reveal the telescoping nature of the sequence, find an expression for the sum of the first N terms,

$$S_N = \sum_{n=1}^N U_n = \sum_{n=1}^N \frac{1}{n(n+1)(n+2)}$$

.

Hint:

- **Part (i):** Multiply both sides of the equation by the common denominator $n(n+1)(n+2)$. To quickly find A , B , and C , substitute values for n that make the bracketed terms equal to zero (e.g., $n = 0, -1, -2$).
- **Part (ii) - Grouping:** You will find that $B = -1$. It is helpful to factor out $\frac{1}{2}$ from the entire expression. Then, split the middle -2 term into two separate -1 fractions: $-\frac{1}{n+1} - \frac{1}{n+2}$. Group one with the $\frac{1}{n}$ term and the other with the $\frac{1}{n+2}$ term.
- **Part (ii) - Summing:** Write out the grouped terms for $n = 1, n = 2$, and $n = N$ explicitly. Look for the diagonal cancellation pattern to see which terms at the beginning and end survive.

Solution 4.17**(i) Decompose the general term:**

Multiply by the denominator:

$$1 = A(n+1)(n+2) + B(n)(n+2) + C(n)(n+1)$$

Substitute values to solve for the constants:

- Let $n = 0 \implies 1 = A(1)(2) \implies A = \frac{1}{2}$
- Let $n = -1 \implies 1 = B(-1)(1) \implies B = -1$
- Let $n = -2 \implies 1 = C(-2)(-1) \implies C = \frac{1}{2}$

Therefore:

$$U_n = \frac{1/2}{n} - \frac{1}{n+1} + \frac{1/2}{n+2} = \frac{1}{2} \left(\frac{1}{n} - \frac{2}{n+1} + \frac{1}{n+2} \right)$$

(ii) Group and sum the sequence:

Group the fractions inside the parenthesis to form differences:

$$U_n = \frac{1}{2} \left[\left(\frac{1}{n} - \frac{1}{n+1} \right) - \left(\frac{1}{n+1} - \frac{1}{n+2} \right) \right]$$

Expand S_N to observe the cascading cancellations:

$$\begin{aligned} n=1 &: \frac{1}{2} \left[\left(\frac{1}{1} - \frac{1}{2} \right) - \left(\frac{1}{2} - \frac{1}{3} \right) \right] \\ n=2 &: \frac{1}{2} \left[\left(\frac{1}{2} - \frac{1}{3} \right) - \left(\frac{1}{3} - \frac{1}{4} \right) \right] \\ &\dots \\ n=N &: \frac{1}{2} \left[\left(\frac{1}{N} - \frac{1}{N+1} \right) - \left(\frac{1}{N+1} - \frac{1}{N+2} \right) \right] \end{aligned}$$

Summing vertically, all intermediate brackets cancel, leaving only the first half of the $n=1$ term and the second half of the $n=N$ term:

$$S_N = \frac{1}{2} \left[\left(1 - \frac{1}{2} \right) - \left(\frac{1}{N+1} - \frac{1}{N+2} \right) \right] = \frac{1}{2} \left(\frac{1}{2} - \frac{1}{N+1} + \frac{1}{N+2} \right)$$

Takeaways 4.4

- **The Method of Cover-up:** When decomposing partial fractions with distinct linear factors, substituting the roots of the denominator is the fastest way to find the numerator constants.
- **Structuring Telescoping Series:** A 3-term partial fraction expansion (like $\frac{1}{2}, -1, \frac{1}{2}$) can often be factored to reveal a $(1, -2, 1)$ pattern, which hides a two-part telescoping series. Splitting the middle coefficient transforms it into a recognizable $V_n - V_{n+1}$ pattern.
- **Write it Out:** Always manually expand the first few and last few terms of a telescoping sum. Relying solely on mental math for the cancellations frequently leads to dropping surviving terms.

4.3 Advanced Sequence Problems

Problem 4.18: Convergence condition

Consider the geometric series

$$\sum_{n=0}^{\infty} 7 \left(\frac{m-1}{3} \right)^n.$$

Find all real values of m for which the series converges, and then state its sum.

Hint: Use the condition $|r| < 1$ for convergence of an infinite GP.

Solution 4.18

Here

$$r = \frac{m-1}{3}.$$

Convergence requires

$$\left| \frac{m-1}{3} \right| < 1 \implies |m-1| < 3 \implies -2 < m < 4.$$

For these values,

$$S_{\infty} = \frac{7}{1 - \frac{m-1}{3}} = \frac{21}{4-m}.$$

Problem 4.19: AP and GP crossover

A sequence is both arithmetic and geometric. Given $a_1 = 6$ and $a_3 = 6$, find all possible sequences.

Hint: Let AP difference be d and GP ratio be r . Use both models on a_3 .

Solution 4.19

AP gives

$$a_3 = a_1 + 2d = 6 + 2d = 6 \implies d = 0,$$

so AP is constant: $a_n = 6$. For GP,

$$a_3 = a_1 r^2 = 6r^2 = 6 \implies r^2 = 1 \implies r = \pm 1.$$

If $r = -1$, terms alternate $6, -6, 6, \dots$ and are not arithmetic. Hence the only sequence that is both AP and GP is

$$a_n = 6 \quad \text{for all } n,$$

with $d = 0$ and $r = 1$.

Problem 4.20: Chebyshev roots

The first-kind Chebyshev polynomial of degree 5 is

$$T_5(x) = 16x^5 - 20x^3 + 5x.$$

Use the identity $T_5(\cos \theta) = \cos(5\theta)$ to find all roots of $T_5(x)$ in the interval $[-1, 1]$.

Hint: Set $\cos(5\theta) = 0$, then convert the resulting angles back to $x = \cos \theta$.

Solution 4.20

Let $x = \cos \theta$. We need

$$\cos(5\theta) = 0.$$

Thus

$$5\theta = \frac{(2k+1)\pi}{2}, \quad k = 0, 1, 2, 3, 4.$$

So

$$\theta = \frac{(2k+1)\pi}{10},$$

and the five roots are

$$x = \cos \frac{\pi}{10}, \cos \frac{3\pi}{10}, \cos \frac{5\pi}{10}, \cos \frac{7\pi}{10}, \cos \frac{9\pi}{10}.$$

Takeaways 4.5

- Roots of $T_n(x)$ on $[-1, 1]$ come from zeros of $\cos(n\theta)$.
- The substitution $x = \cos \theta$ is natural because $\cos \theta$ covers $[-1, 1]$.
- Degree 5 gives five roots, which is a useful check.

Problem 4.21: Architecture of GP sums

Let a GP have first term $a \neq 0$, ratio $r \neq 0, \pm 1$, and partial sums S_n .

- Prove $\frac{S_{2n}}{S_n} = r^n + 1$.
- Given $S_{12} = 65S_6$ and second term $ar = 12$, find possible (a, r) .
- If Σ_n is the partial-sum sequence of a GP with first term a and ratio r^2 , show

$$\frac{\Sigma_n}{S_n} = \frac{r^n + 1}{r + 1}.$$

Hint: Use $S_n = \frac{a(1-r^n)}{1-r}$ and difference-of-squares factorization.

Solution 4.21

$$\frac{S_{2n}}{S_n} = \frac{r^{2n} - 1}{r^n - 1} = r^n + 1.$$

So $r^6 + 1 = 65 \implies r^6 = 64 \implies r = \pm 2$. Since $ar = 12$, possible pairs are $(a, r) = (6, 2)$ or $(-6, -2)$. Also

$$\Sigma_n = \frac{a(r^{2n} - 1)}{r^2 - 1},$$

thus

$$\frac{\Sigma_n}{S_n} = \frac{r^{2n} - 1}{(r + 1)(r^n - 1)} = \frac{r^n + 1}{r + 1}.$$

Problem 4.22: AP space and difference operator

Let $\mathcal{A}(a, d)$ denote an AP. Define

$$\mathcal{I} = \mathcal{A}(1, 0), \quad \mathcal{J} = \mathcal{A}(0, 1).$$

- (i) Show any AP is uniquely $a\mathcal{I} + d\mathcal{J}$.
- (ii) Let $\Delta x_n = x_{n+1} - x_n$. Show $\Delta\mathcal{I} = 0$ and $\Delta\mathcal{J} = \mathcal{I}$.
- (iii) Deduce $\Delta(a\mathcal{I} + d\mathcal{J}) = d\mathcal{I}$.

Hint: Compare first term and common difference.

Solution 4.22

An AP has n th term $a + (n - 1)d = a \cdot 1 + d(n - 1)$, so $X = a\mathcal{I} + d\mathcal{J}$, unique by matching (a, d) . Also $\mathcal{I} = (1, 1, 1, \dots)$ so $\Delta\mathcal{I} = 0$, and $\mathcal{J} = (0, 1, 2, \dots)$ so $\Delta\mathcal{J} = (1, 1, 1, \dots) = \mathcal{I}$. Hence

$$\Delta(a\mathcal{I} + d\mathcal{J}) = a\Delta\mathcal{I} + d\Delta\mathcal{J} = d\mathcal{I}.$$

Problem 4.23: Basis and projection in AP space

Let $\mathcal{E}_1 = \mathcal{A}(1, 1)$ and $\mathcal{E}_2 = \mathcal{A}(1, -1)$.

- (i) Express any AP $\mathcal{A}(a, d)$ as $\lambda\mathcal{E}_1 + \mu\mathcal{E}_2$.
- (ii) Decompose $P = (5, 8, 11, 14, \dots)$ in this basis.

Hint: Match first terms and differences: $\lambda + \mu = a$, $\lambda - \mu = d$.

Solution 4.23

Solving

$$\lambda + \mu = a, \quad \lambda - \mu = d$$

gives

$$\lambda = \frac{a+d}{2}, \quad \mu = \frac{a-d}{2}.$$

For P , $a = 5$, $d = 3$, so $\lambda = 4$, $\mu = 1$.**Problem 4.24: Countable and uncountable sets**

- (i) Explain why positive rationals are countable by diagonal listing.
- (ii) Use Cantor's diagonal argument to show reals in $[0, 1)$ are uncountable.

Hint: Construct a new decimal differing from the n th listed decimal at position n .

Solution 4.24

Rationals can be arranged in an infinite grid $\frac{p}{q}$ and traversed by diagonals, skipping duplicates after simplification; this yields a sequence listing all positive rationals. For reals in $[0, 1)$, assume a list $T_1, T_2, \dots$ exists. Build x so its n th digit differs from the n th digit of T_n . Then $x \neq T_n$ for every n , contradicting completeness of the list. Hence uncountable.

Problem 4.25: Prime denominators and cycle length

Let p be prime, $p \neq 2, 5$, and L be period length of $\frac{1}{p}$.

- (i) Show L is the least positive integer with $10^L \equiv 1 \pmod{p}$.
- (ii) Deduce $L \mid (p-1)$.
- (iii) Find periods for $\frac{1}{37}$ and $\frac{1}{41}$.

Hint: Use Fermat's theorem and first occurrence of $p \mid (10^n - 1)$.

Solution 4.25

Recurring block theory gives $(10^L - 1)/p \in \mathbb{Z}$, hence $10^L \equiv 1 \pmod{p}$; minimality gives least such L . By Fermat, $10^{p-1} \equiv 1 \pmod{p}$, so the order L divides $p-1$. Since $37 \mid (10^3 - 1)$ and not $10^1 - 1, 10^2 - 1$, period is 3. Since $41 \mid (10^5 - 1)$ and not $10^1 - 1$, period is 5.

Problem 4.26: Pythagorean means

For $0 < a < b$, define

$$A = \frac{a+b}{2}, \quad G = \sqrt{ab}, \quad H = \frac{2ab}{a+b}.$$

- (i) Prove $H < G < A$.
- (ii) Find $\frac{b}{a}$ such that a, G, A is an AP.
- (iii) Show H, G, A cannot be an AP unless $a = b$.

Hint: Use $(\sqrt{a} - \sqrt{b})^2 > 0$ and $0 < G$ and $AH = G^2$.

Solution 4.26

From $(\sqrt{a} - \sqrt{b})^2 > 0$, get $A > G$. Also

$$AH = \frac{a+b}{2} \cdot \frac{2ab}{a+b} = ab = G^2,$$

so H, G, A form a GP and therefore $H < G < A$. For a, G, A in AP:

$$2G = a + A \implies 4\sqrt{ab} = 3a + b.$$

Let $x = b/a$; then $(x-9)(x-1) = 0$, and $x > 1$ gives $x = 9$. If H, G, A were AP, then $2G = H + A = G^2/A + A$, forcing $A = G$, contradiction for $a \neq b$.

Problem 4.27: Recurring decimals and modular structure

Let p be prime ($p \neq 2, 5$), and let n be the period of $\frac{1}{p}$.

- (i) Show $n \mid (p-1)$.
- (ii) If $n = 2k$, prove $10^k \equiv -1 \pmod{p}$ and the two half-blocks of the repetend sum to $10^k - 1$.

Hint: Factor $10^{2k} - 1 = (10^k - 1)(10^k + 1)$ and use minimality of n .

Solution 4.27

From order arguments, n is the multiplicative order of $10 \pmod{p}$, so $n \mid (p-1)$. If $n = 2k$, then $p \mid (10^{2k} - 1)$. Since $2k$ is minimal, $p \nmid (10^k - 1)$, hence $p \mid (10^k + 1)$, so $10^k \equiv -1 \pmod{p}$. Writing repetend as two k -digit blocks A, B gives Midy's theorem:

$$A + B = 10^k - 1.$$

Problem 4.28: Sissa and arithmetico-geometric sums

(i) Evaluate $S_{64} = 1 + 2 + 4 + \cdots + 2^{63}$.

(ii) Let

$$T = \sum_{n=1}^{64} n2^{n-1}.$$

Use shift-and-subtract to find a closed form.

(iii) Evaluate

$$\sum_{n=1}^{\infty} \frac{n}{2^n}.$$

Hint: Use $T - 2T$ for the finite AGP, and similarly $S - \frac{1}{2}S$ for the infinite one.

Solution 4.28

$$S_{64} = 2^{64} - 1.$$

Also

$$T - 2T = \sum_{n=1}^{64} 2^{n-1} - 64 \cdot 2^{64} = (2^{64} - 1) - 64 \cdot 2^{64},$$

so

$$T = 63 \cdot 2^{64} + 1.$$

For $S = \sum_{n=1}^{\infty} \frac{n}{2^n}$:

$$S - \frac{1}{2}S = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots = 1,$$

hence $S = 2$.

Problem 4.29: Discrete calculus and sums of powers

∇ is the backward difference operator, defined as $\nabla f(n) = f(n) - f(n-1)$ for discrete functions $f(n)$. In this problem, you will explore the properties of the backward difference operator and its applications to sums of powers.

For example, The first order of ∇ is

$$\nabla(n^2) = n^2 - (n-1)^2 = 2n - 1$$

The second order of ∇ is

$$\nabla^2 f(n) = \nabla(\nabla f(n)) = \nabla(f(n) - f(n-1)) = f(n) - 2f(n-1) + f(n-2)$$

and the third order of ∇ is

$$\nabla^3 f(n) = \nabla(\nabla^2 f(n)) = \nabla(f(n) - 2f(n-1) + f(n-2)) = f(n) - 3f(n-1) + 3f(n-2) - f(n-3)$$

In this problem, we use the ∇ operator to derive formulas for sums of powers.

(i) Show $n^3 - (n-1)^3 = 3n^2 - 3n + 1$.

(ii) Sum this identity from 1 to N to derive

$$\sum_{n=1}^N n^2 = \frac{N(N+1)(2N+1)}{6}.$$

(iii) Evaluate $\nabla^3(n^3)$

(iv) State the leading term of $\sum_{n=1}^N n^k$.

Hint: Use telescoping on the left and known formulas for $\sum n$.

Solution 4.29

Expansion gives part (i). Summing:

$$N^3 = 3 \sum_{n=1}^N n^2 - 3 \sum_{n=1}^N n + N,$$

which simplifies to

$$\sum_{n=1}^N n^2 = \frac{N(N+1)(2N+1)}{6}.$$

Also $\nabla^3(n^3) = 6$. In general,

$$\sum_{n=1}^N n^k = \frac{1}{k+1} N^{k+1} + (\text{lower powers of } N).$$

Problem 4.30: The Fibonacci Generating Function

Let F_n be the n -th Fibonacci number ($F_1 = 1, F_2 = 1, F_n = F_{n-1} + F_{n-2}$). We define the power series $S(x)$ as:

$$S(x) = \sum_{n=1}^{\infty} F_n x^{n+1}$$

- (i) By expanding $S(x) - xS(x) - x^2S(x)$ and using the recurrence property of Fibonacci numbers, prove that:

$$S(x) = \frac{x^2}{1 - x - x^2}$$

You may assume that the sum converges without proof.

- (ii) Given the growth rate of Fibonacci numbers, explain why this series converges for $x = 0.1$.
- (iii) Show that $S(0.1) = \frac{1}{89}$.
- (iv) The decimal expansion of $\frac{1}{89}$ is $0.011235955\dots$. Explain why the digit in the 10^{-7} place is 9 rather than the expected Fibonacci number $F_6 = 8$.

Hint: Consider the definition of $S(x)$ term by term for (i). For (iv), look closely at how the next Fibonacci number $F_7 = 13$ interacts with the decimal places.

Solution 4.30

(i) Substitution into $S(x)(1 - x - x^2)$ cancels out all terms except the first, leaving $F_1 x^2 = x^2$. Hence $S(x) = \frac{x^2}{1 - x - x^2}$.

(ii) Fibonacci numbers grow roughly geometrically by the golden ratio $\phi \approx 1.618$. Because $1.618 \dots \times 0.1 < 1$, the series converges.

(iii) Substitute $x = 0.1$:


$$S(0.1) = \frac{0.1^2}{1 - 0.1 - 0.1^2} = \frac{0.01}{1 - 0.1 - 0.01} = \frac{0.01}{0.89} = \frac{1}{89}.$$

(iv) The required sum gives $F_6 \times (0.1)^7 + F_7 \times (0.1)^8$. $F_6 = 8$ and $F_7 = 13$, so the sum overlapping the 10^{-7} place includes the carry-over from F_7 : $8 + 1 = 9$.

Takeaways 4.6

- Power series generating functions can be explicitly derived using recurrence relations.
- The decimal expansion of fractions naturally encodes the base-10 carry-over of power series evaluations.
- Fibonacci recursion can also be visualised geometrically via a quarter-turn spiral whose radii follow consecutive Fibonacci numbers.

$$F_1 = 1, F_2 = 1, F_n = F_{n-1} + F_{n-2}$$



5 Conclusion

Strong sequence work in Extension 1 is not just memorizing formulas. It is about choosing the right model, controlling notation, and checking whether each step is logically justified.

The key habits are:

- define terms clearly, including first term and common difference or ratio
- check indexing before and after every sigma manipulation
- identify whether a question is about terms, partial sums, or limits
- test answers for reasonableness by checking growth and sign behavior

With steady practice, sequences become one of the most reliable scoring areas in Extension 1.

Contact Information:

LinkedIn: <https://www.linkedin.com/in/nguyenvuhung/>

GitHub: <https://github.com/vuhung16au/>

Repository: <https://github.com/vuhung16au/math-olympiad-ml/tree/main/HSC-Sequences>

6 Appendices

6.1 Appendix A: Formula Sheet

- Arithmetic n th term:

$$a_n = a + (n - 1)d.$$

- Arithmetic sum of first n terms:

$$S_n = \frac{n}{2}(2a + (n - 1)d) = \frac{n}{2}(a + l).$$

- Geometric n th term:

$$a_n = ar^{n-1}.$$

- Geometric sum of first n terms ($r \neq 1$):

$$S_n = a \frac{1 - r^n}{1 - r} = a \frac{r^n - 1}{r - 1}.$$

- Infinite geometric sum ($|r| < 1$):

$$S_\infty = \frac{a}{1 - r}.$$

- Sums of powers for the first n positive integers:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2},$$

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6},$$

$$\sum_{k=1}^n k^3 = \left(\frac{n(n+1)}{2} \right)^2.$$

Chebyshev Polynomials

The Chebyshev polynomials of the first kind, denoted $T_n(x)$, are defined by

$$T_0(x) = 1, \quad T_1(x) = x, \quad T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x) \quad (n \geq 1).$$

Equivalently,

$$T_n(\cos \theta) = \cos(n\theta).$$

The Chebyshev polynomials of the second kind, denoted $U_n(x)$, are defined by

$$U_0(x) = 1, \quad U_1(x) = 2x, \quad U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x) \quad (n \geq 1).$$

Equivalently,

$$U_n(\cos \theta) = \frac{\sin((n+1)\theta)}{\sin \theta} \quad (\sin \theta \neq 0).$$

Degree	First kind $T_n(x)$	Second kind $U_n(x)$
1	x	$2x$
2	$2x^2 - 1$	$4x^2 - 1$
3	$4x^3 - 3x$	$8x^3 - 4x$
4	$8x^4 - 8x^2 + 1$	$16x^4 - 12x^2 + 1$
5	$16x^5 - 20x^3 + 5x$	$32x^5 - 32x^3 + 6x$

6.2 Appendix B: Common Sequence Traps

- Mixing up term formulas and sum formulas: a_n and S_n answer different questions.
- Losing track of index starts: check whether the sequence starts at $n = 0$ or $n = 1$.
- Using $S_\infty = \frac{a}{1-r}$ when $|r| \geq 1$: this formula only applies for $|r| < 1$.
- Sign errors in GP sums: keep the exact formula form and substitute carefully.
- Assuming recursion converges without proof: identify monotonicity or contraction before claiming limits.
- Treating recurring decimals as finite truncations: they are infinite geometric expansions.

6.3 Appendix C: Rigorous Definition of the Limit of a Sequence

Definition 1 (ϵ - N Definition of a Limit). Let (a_n) be a sequence of real numbers. We say that (a_n) **converges to a limit** L , written

$$\lim_{n \rightarrow \infty} a_n = L,$$

if for every real number $\epsilon > 0$ there exists a positive integer N such that

$$n > N \implies |a_n - L| < \epsilon.$$

In formal logical notation:

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ such that } n > N \implies |a_n - L| < \epsilon.$$

If no such L exists, the sequence is said to **diverge**.

Unpacking the definition.

- $\epsilon > 0$ is the “margin of error.” The challenger picks any strictly positive number, no matter how small. A smaller ϵ places a tighter band around L .
- N is the “cutoff index.” The responder must produce a concrete N that works for the chosen ϵ . Typically, smaller ϵ forces a larger N .
- Only the *tail* matters. The condition $n > N$ means the first N terms may behave arbitrarily; only the infinite tail of the sequence is constrained.
- $|a_n - L| < \epsilon$ is the “closeness” condition. Every term beyond index N must lie strictly inside the interval $(L - \epsilon, L + \epsilon)$.

Visual interpretation (the “ ϵ -tube”).

Plot the sequence on a graph with the index n on the horizontal axis and a_n on the vertical axis. Draw a horizontal dashed line at $y = L$. Choosing $\epsilon > 0$ creates a horizontal band $(L - \epsilon, L + \epsilon)$ of total width 2ϵ around the limit. The definition states that for every such band there is a vertical line $x = N$ such that *every point of the sequence to the right of that line lies inside the band*.

Example: Proving $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$.

Proof. Let $\epsilon > 0$ be given.

Scratchwork. We require

$$\left| \frac{1}{n} - 0 \right| < \epsilon.$$

Since $n \in \mathbb{N}$ is positive, this simplifies to $\frac{1}{n} < \epsilon$, i.e. $n > \frac{1}{\epsilon}$.

Choice of N . Let N be any positive integer satisfying $N > \frac{1}{\epsilon}$. (Such an integer exists by the Archimedean property of $\mathbb{R}$.)

Verification. Suppose $n > N$. Then

$$n > N > \frac{1}{\epsilon} \implies \frac{1}{n} < \epsilon \implies \left| \frac{1}{n} - 0 \right| = \frac{1}{n} < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, the definition is satisfied and $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$. □

Concrete illustration.

For $\epsilon = 0.01$ we need $N > 100$, so we may take $N = 101$. Every term $a_n = \frac{1}{n}$ with $n \geq 102$ satisfies $\frac{1}{102} < 0.01$, placing all those terms inside the band $(-0.01, 0.01)$.

Note: The following advanced theorems are outside the scope of the regular HSC syllabus, but they are extremely good-to-know fundamentals for higher education and university mathematics.

1. Bolzano-Weierstrass Theorem

The Bolzano-Weierstrass Theorem is a “guarantee of stability.” It tells us that as long as a sequence doesn’t head off to infinity, it must contain at least one “hidden” convergent behavior.

- **The Intuition:** Imagine a sequence as a set of points trapped inside a finite box (bounded). Even if the points bounce around randomly and never settle on one spot as a whole (like $a_n = (-1)^n$), the points must eventually start “clustering” somewhere because they have nowhere else to go.
- **Subsequences:** A subsequence is formed by picking infinitely many terms from the original sequence in their original order.
 - *Example:* Take $a_n = (-1)^n \rightarrow \{-1, 1, -1, 1, \dots\}$. The sequence itself oscillates and diverges. However, if we pick only the even terms, we get $\{1, 1, 1, \dots\}$, which converges to 1.
- **The Utility:** In proofs, if you know a sequence is bounded, you can immediately “extract” a convergent subsequence to use for further logical steps.

2. Cauchy Criterion for Convergence

The Cauchy Criterion is the ultimate “internal” test for convergence.

- **The Definition:** In a standard limit, we say terms get close to a **limit** L . In a Cauchy sequence, we say the terms get **close to each other**.

- **Formal Statement:** For every $\epsilon > 0$, there exists an N such that for all $m, n > N$:

$$|a_n - a_m| < \epsilon$$

- **Why it Matters:** Usually, to prove a sequence converges, you first have to *guess* what the limit L is. The Cauchy Criterion removes this requirement. If you can prove that the terms of the sequence eventually “huddle together” and the gaps between them shrink to zero, the sequence **must** converge to some real number.
- **The “Completeness” of Real Numbers:** This theorem works because the Real Number line has no “holes.” If the terms are huddling together, they are guaranteed to be huddling around a real number. (This is not true in the Rational numbers, where a sequence of fractions could huddle around an “irrational hole” like $\sqrt{2}$).

6.4 Appendix D: Limit Superior and Limit Inferior

Note on Syllabus

The concepts of limit superior ($\limsup$) and limit inferior ($\liminf$) are strictly **outside the scope** of the HSC Mathematics Extension 1 and Extension 2 syllabuses. However, they provide an excellent bridge into higher mathematics (such as university-level real analysis) and serve as useful tools to “estimate a boundary” for the limits of sequences and functions.

Estimating Boundaries with Limsup and Liminf

In standard mathematics, the limit of a sequence a_n as $n \rightarrow \infty$ describes the exact value the sequence settles on. However, many sequences do not settle on a single value; they might oscillate or jump around (for example, $a_n = (-1)^n$).

Instead of asking “What is the exact limit?”, we can ask “What are the ultimate upper and lower boundaries of this sequence?”

- **Limit Superior** ($\limsup_{n \rightarrow \infty} a_n$): The eventual upper bound of the sequence. It describes the maximum value that the sequence continues to reach or approach as n becomes arbitrarily large.
- **Limit Inferior** ($\liminf_{n \rightarrow \infty} a_n$): The eventual lower bound of the sequence. It describes the minimum value that the sequence continues to reach or approach as n becomes arbitrarily large.

A sequence (a_n) converges to a limit L if and only if both its limit superior and limit inferior are equal to that same expected limit L . That is:

$$\lim_{n \rightarrow \infty} a_n = L \iff \limsup_{n \rightarrow \infty} a_n = \liminf_{n \rightarrow \infty} a_n = L$$

Simple Examples

Example 1: A Converging Sequence

Consider the sequence $a_n = \frac{1}{n}$. The terms are $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$. As $n \rightarrow \infty$, the sequence approaches 0. Here, the eventual upper boundary and lower boundary are the same.

$$\limsup_{n \rightarrow \infty} \frac{1}{n} = 0 \quad \text{and} \quad \liminf_{n \rightarrow \infty} \frac{1}{n} = 0$$

Since they are equal, the standard limit exists and $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$.

Example 2: An Oscillating Sequence

Consider the sequence $a_n = (-1)^n$. The terms are $-1, 1, -1, 1, -1, 1, \dots$. This sequence never settles on a single converging value, so the standard limit does not exist. However, the sequence is clearly bounded.

- The highest value the sequence keeps hitting as $n \rightarrow \infty$ is 1. Thus, $\limsup_{n \rightarrow \infty} (-1)^n = 1$.
- The lowest value the sequence keeps hitting as $n \rightarrow \infty$ is -1 . Thus, $\liminf_{n \rightarrow \infty} (-1)^n = -1$.

Example 3: A Sequence with Growth and Oscillation

Consider $a_n = 5 + \frac{(-1)^n}{n}$. The sequence oscillates slightly above and below 5, getting closer and closer to 5.

$$\limsup_{n \rightarrow \infty} \left(5 + \frac{(-1)^n}{n} \right) = 5 \quad \text{and} \quad \liminf_{n \rightarrow \infty} \left(5 + \frac{(-1)^n}{n} \right) = 5$$

Both estimates boundary conditions pinch together at 5, confirming the exact limit is 5.

6.5 Appendix E: Discrete Calculus and the Difference Operator

Note on Syllabus

The concept of the backward difference operator (∇) belongs to the field of *Discrete Calculus*. While the mechanics of telescoping sums are implicitly part of Extension 1 and Extension 2, formalizing them into discrete derivative operators is **outside the scope** of the formal HSC syllabus. Nevertheless, it deeply connects the study of sequences to continuous calculus that is taught at the university level.

The Difference Operator: The "Derivative" for Sequences

In continuous calculus, the derivative measures the instantaneous rate of change of a function $f(x)$ with respect to x :

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h}$$

When we study sequences, the domain is restricted to discrete integers. This means the smallest possible change or "step" in the variable n is exactly 1 (we cannot let $h \rightarrow 0$). Therefore, the discrete counterpart to the derivative is the **backward difference operator**, denoted by ∇ (nabla):

$$\nabla f(n) = \frac{f(n) - f(n-1)}{1} = f(n) - f(n-1)$$

(Note: There is also a forward difference operator $\Delta f(n) = f(n+1) - f(n)$, which functions very similarly).

Just as we can take multiple derivatives of functions like $f''(x)$, we can repeatedly apply difference operators to sequences, evaluating things like $\nabla^3(n^3)$.

Why is it Useful?

In calculus, the Fundamental Theorem of Calculus links derivatives to integrals:

$$\int_a^b f'(x) dx = f(b) - f(a)$$

In discrete mathematics involving sequences, the operation equivalent to continuous integration is **summation**. Summing a difference operator perfectly mirrors the Fundamental Theorem of Calculus through **telescoping sums**:

$$\sum_{n=1}^N \nabla f(n) = \sum_{n=1}^N (f(n) - f(n-1)) = f(N) - f(0)$$

This elegant connection turns difficult summations into basic boundary evaluations. If you want to compute the sum of a sequence $g(n)$, and you can guess or find a discrete "anti-derivative" $f(n)$ such that $\nabla f(n) = g(n)$, the sum is trivially computed.

Application to Sums of Powers

Consider the continuous derivative of a polynomial power: $\frac{d}{dx}(x^3) \approx 3x^2$. In the discrete world, the difference operator applied to n^3 yields:

$$\nabla(n^3) = n^3 - (n-1)^3 = 3n^2 - 3n + 1$$

By summing this identity, we mimic the concept of integrating a derivative:

$$\sum_{n=1}^N \nabla(n^3) = \sum_{n=1}^N (3n^2 - 3n + 1)$$

Because the left hand side is just a sum of differences, it telescopes and collapses, leaving only the bounds:

$$N^3 - 0^3 = 3 \sum_{n=1}^N n^2 - 3 \sum_{n=1}^N n + \sum_{n=1}^N 1$$

This equation is immensely powerful: it provides a rapid mechanism to find the closed-form formula for $\sum_{n=1}^N n^2$. Furthermore, it intuitively explains why the continuous integral $\int x^2 dx = \frac{x^3}{3}$ corresponds directly to the discrete sum having a leading polynomial term of $\frac{N^3}{3}$.