HSC Mathematics Extension 1: Sequences and Series

Vu Hung Nguyen

Download PDF and resources:
`https://github.com/vuhung16au/math-olympiad-ml`

This work is licensed under CC BY 4.0, see the LICENSE file on the github page for more info.

"Sometimes it is the people no one imagines anything of who do the things that no one can imagine."

Alan Turing

Contents

1 Introduction **5**
1.1 Project Overview . . . 5
1.2 Target Audience . . . 5
1.3 How to Use This Booklet . . . 5
1.4 Topics Covered . . . 5
1.5 Where This Leads Next . . . 5

2 Fundamentals Review **6**
2.1 How Sequences Are Specified . . . 6
2.2 Arithmetic Sequences (AP) . . . 6
2.3 Arithmetic Series . . . 6
2.4 Geometric Sequences (GP) . . . 7
2.5 Geometric Series . . . 7
2.6 Limiting Sum of a Geometric Series . . . 8
2.7 Recurring Decimals and Geometric Series . . . 8
2.8 Sigma Notation Quick Rules . . . 8

3 Part 1: Problems and Solutions (Detailed) **9**
3.1 Basic Sequence Problems . . . 9
3.2 Medium Sequence Problems . . . 10
3.3 Advanced Sequence Problems . . . 21

4 Part 2: Problems with Hints and Solutions (Concise) **28**
4.1 Basic Sequence Problems . . . 28
4.2 Medium Sequence Problems . . . 31
4.3 Advanced Sequence Problems . . . 35

5 Conclusion **42**

6 Appendices **43**
6.1 Appendix A: Formula Sheet . . . 43
6.2 Appendix B: Common Sequence Traps . . . 44
6.3 Appendix C: Rigorous Definition of the Limit of a Sequence . . . 44

 Vu Hung Nguyen · CC BY 4.0

1 Introduction

1.1 Project Overview

This booklet is a focused companion for NSW HSC Mathematics Extension 1 students studying sequences and series. It is designed to connect core sequence skills to exam-style problem solving with clear methods and structured solutions.

1.2 Target Audience

This resource is written for Year 12 students in Extension 1, and for tutors and teachers who want a compact booklet with consistent notation and worked examples.

1.3 How to Use This Booklet

- Read the fundamentals review first, then attempt the detailed problems in Part 1.
- Use Part 2 for timed practice and fluency, using hints only when needed.
- Keep a formula sheet while you work, but always justify each step from first principles.
- Track common slips, especially sign errors in geometric sums and indexing mistakes in sigma notation.

1.4 Topics Covered

This booklet develops the following Extension 1 ideas:

- defining sequences by term formula, recursion, and context
- arithmetic sequences and arithmetic series
- geometric sequences and geometric series
- finite sum formulas and sigma notation
- limiting sum of a geometric series when $|r| < 1$
- links between recurring decimals and geometric series
- mixed AP/GP problem solving in HSC-style questions

1.5 Where This Leads Next

Sequence fluency supports later work in calculus, proof, and Extension 2 style reasoning. In particular, convergence arguments and recursive definitions form a bridge to harder topics such as iterative methods and infinite processes.

 Vu Hung Nguyen · CC BY 4.0

2 Fundamentals Review

This section is a compact reference for notation, formulas, and common reasoning patterns used throughout the booklet.

2.1 How Sequences Are Specified

A sequence is an ordered list of numbers, usually written as $(a_n)_{n\geq 1}$. Common ways to define a sequence are:

- explicit formula: a_n written directly in terms of n
- recursive formula: each term written using previous terms
- contextual rule: terms generated from a worded process

Examples.

$$a_n = 3n - 1, \quad b_1 = 2,\ b_{n+1} = b_n + 5, \quad c_n = (-1)^n.$$

2.2 Arithmetic Sequences (AP)

An arithmetic sequence has constant difference d:

$$a_{n+1} - a_n = d.$$

If the first term is a, then

$$a_n = a + (n-1)d.$$

Problem 2.1: Recognizing an AP

Given $a_1 = 7$ and $a_{n+1} = a_n - 3$, find a_{10}.

Solution 2.1

Here $d = -3$, so

$$a_{10} = 7 + (10-1)(-3) = 7 - 27 = -20.$$

2.3 Arithmetic Series

The sum of the first n terms of an AP is:

$$S_n = \frac{n}{2}\big(2a + (n-1)d\big) = \frac{n}{2}(a + l),$$

where l is the nth term.

Problem 2.2: AP Sum

Find $\sum_{k=1}^{20}(5 + 2(k-1))$.

 Vu Hung Nguyen · CC BY 4.0

Solution 2.2

This is an AP with $a = 5$, $d = 2$, and $n = 20$. The 20th term is

$$l = 5 + 19 \cdot 2 = 43.$$

So

$$S_{20} = \frac{20}{2}(5 + 43) = 10 \cdot 48 = 480.$$

2.4 Geometric Sequences (GP)

A geometric sequence has constant ratio r:

$$\frac{a_{n+1}}{a_n} = r \quad (a_n \neq 0).$$

If the first term is a, then

$$a_n = ar^{n-1}.$$

Problem 2.3: Recognizing a GP

If $a_1 = 81$ and $r = \frac{1}{3}$, find a_5.

Solution 2.3

$$a_5 = 81\left(\frac{1}{3}\right)^4 = 81 \cdot \frac{1}{81} = 1.$$

2.5 Geometric Series

For $r \neq 1$, the sum of the first n terms is

$$S_n = a\frac{1 - r^n}{1 - r} = a\frac{r^n - 1}{r - 1}.$$

Use the form that keeps signs clean for your r value.

Problem 2.4: Finite GP Sum

Find $\sum_{k=0}^{6} 3 \cdot 2^k$.

Solution 2.4

This is $a = 3$, $r = 2$, $n = 7$ terms (from $k = 0$ to 6):

$$S_7 = 3\frac{2^7 - 1}{2 - 1} = 3(128 - 1) = 381.$$

 Vu Hung Nguyen · CC BY 4.0

2.6 Limiting Sum of a Geometric Series

If $|r| < 1$, then $r^n \to 0$ as $n \to \infty$, and the infinite sum exists:

$$S_\infty = \frac{a}{1-r}.$$

If $|r| \geq 1$, the geometric series does not converge to a finite limiting sum.

Problem 2.5: Infinite GP

Evaluate $2 + \frac{2}{3} + \frac{2}{9} + \frac{2}{27} + \cdots$.

Solution 2.5

Here $a = 2$, $r = \frac{1}{3}$, and $|r| < 1$, so

$$S_\infty = \frac{2}{1-\frac{1}{3}} = \frac{2}{\frac{2}{3}} = 3.$$

2.7 Recurring Decimals and Geometric Series

Recurring decimals can be written as geometric series. For example,

$$0.\overline{7} = 0.7 + 0.07 + 0.007 + \cdots$$

is geometric with $a = 0.7$ and $r = 0.1$. So

$$0.\overline{7} = \frac{0.7}{1-0.1} = \frac{0.7}{0.9} = \frac{7}{9}.$$

2.8 Sigma Notation Quick Rules

$$\sum_{k=1}^{n}(u_k + v_k) = \sum_{k=1}^{n} u_k + \sum_{k=1}^{n} v_k, \quad \sum_{k=1}^{n} cu_k = c\sum_{k=1}^{n} u_k.$$

Index shifts must preserve both the expression and bounds consistently.

Takeaways 2.1

- Always identify whether the question asks for a term a_n, a finite sum S_n, or a limit.
- AP questions depend on constant difference; GP questions depend on constant ratio.
- For infinite geometric sums, check $|r| < 1$ before using $S_\infty = \frac{a}{1-r}$.
- Track indexing carefully in sigma notation to avoid off-by-one errors.
- Chebyshev polynomial questions usually depend on either the recurrence relation or the trigonometric identities.

 Vu Hung Nguyen · CC BY 4.0

3 Part 1: Problems and Solutions (Detailed)

Part 1 contains full solutions with method explanations.

3.1 Basic Sequence Problems

Problem 3.1: Arithmetic nth term

An arithmetic sequence has first term 4 and common difference 5. Find a_{30}.

Solution 3.1

Use $a_n = a + (n-1)d$:

$$a_{30} = 4 + 29 \cdot 5 = 149.$$

Problem 3.2: Find the common ratio

In a geometric sequence, $a_2 = 12$ and $a_5 = 324$. Find the common ratio r and the first term a_1.

Solution 3.2

From $a_n = ar^{n-1}$,

$$a_2 = ar = 12, \qquad a_5 = ar^4 = 324.$$

Divide to remove a:

$$\frac{a_5}{a_2} = r^3 = \frac{324}{12} = 27 \implies r = 3.$$

Then $a_1 = a = \frac{12}{3} = 4$.

Problem 3.3: Sum of an AP

Find the sum of the first 40 terms of the arithmetic sequence

$$9,\ 13,\ 17,\ \ldots$$

Solution 3.3

Here $a = 9$, $d = 4$, $n = 40$. The 40th term is

$$l = 9 + 39 \cdot 4 = 165.$$

So

$$S_{40} = \frac{40}{2}(9 + 165) = 20 \cdot 174 = 3480.$$

 Vu Hung Nguyen · CC BY 4.0

Problem 3.4: Recurring decimal as a fraction

Express $0.\overline{23}$ as a rational number.

Solution 3.4

$$0.\overline{23} = \frac{23}{99}.$$

A geometric view is

$$0.23 + 0.0023 + 0.000023 + \cdots$$

with first term $\frac{23}{100}$ and ratio $\frac{1}{100}$.

Problem 3.5: Chebyshev recurrence: first kind

The Chebyshev polynomials of the first kind satisfy

$$T_0(x) = 1, \qquad T_1(x) = x, \qquad T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x).$$

Use the recurrence to find $T_2(x)$ and $T_3(x)$, then evaluate $T_3\left(\frac{1}{2}\right)$.

Hint: Build the polynomials one at a time: first use $T_2 = 2xT_1 - T_0$, then use $T_3 = 2xT_2 - T_1$.

Solution 3.5

$$T_2(x) = 2x^2 - 1.$$

Then

$$T_3(x) = 2x(2x^2 - 1) - x = 4x^3 - 3x.$$

So

$$T_3\left(\frac{1}{2}\right) = 4\left(\frac{1}{8}\right) - 3\left(\frac{1}{2}\right) = \frac{1}{2} - \frac{3}{2} = -1.$$

Takeaways 3.1

- Chebyshev polynomials can be generated recursively from two starting values.
- Work in order; each new polynomial depends on the previous two.
- Substitution is easiest after simplifying the polynomial.

3.2 Medium Sequence Problems

Problem 3.6: Mixed AP constraints

An arithmetic sequence has $a_3 = 11$ and $a_{12} = 47$. Find a_1 and d, then determine S_{25}.

 Vu Hung Nguyen · CC BY 4.0

Solution 3.6

Use $a_n = a + (n-1)d$:

$$a + 2d = 11, \qquad a + 11d = 47.$$

Subtract:

$$9d = 36 \implies d = 4,$$

then $a = 11 - 8 = 3$. Now

$$a_{25} = 3 + 24 \cdot 4 = 99,$$

so

$$S_{25} = \frac{25}{2}(3 + 99) = \frac{25}{2} \cdot 102 = 1275.$$

Problem 3.7: Finite GP with unknown index

For the geometric sequence with first term 5 and ratio 2, find n such that

$$S_n = 2555.$$

Solution 3.7

$$S_n = 5\frac{2^n - 1}{2 - 1} = 5(2^n - 1) = 2555.$$

So

$$2^n - 1 = 511 \implies 2^n = 512 = 2^9.$$

Hence $n = 9$.

Problem 3.8: Consecutive cubes and induction

Prove by induction that

$$(n-1)^3 + n^3 + (n+1)^3$$

is divisible by 3 for every integer n.

Hint: First prove the result for $n \geq 0$. In the inductive step, compare the expression at $n = k + 1$ with the expression at $n = k$.

Solution 3.8

Let

$$P(n) = (n-1)^3 + n^3 + (n+1)^3.$$

We first prove that $P(n)$ is divisible by 3 for all integers $n \geq 0$.
Base case. For $n = 0$,

$$P(0) = (-1)^3 + 0^3 + 1^3 = 0,$$

which is divisible by 3.

 Vu Hung Nguyen · CC BY 4.0

Inductive step. Suppose $P(k)$ is divisible by 3 for some integer $k \geq 0$. Then

$$\begin{aligned} P(k+1) - P(k) &= k^3 + (k+1)^3 + (k+2)^3 - \left((k-1)^3 + k^3 + (k+1)^3\right) \\ &= (k+2)^3 - (k-1)^3 \\ &= 9k^2 + 9k + 9 \\ &= 9(k^2 + k + 1), \end{aligned}$$

which is divisible by 3. Since $P(k)$ is divisible by 3, it follows that $P(k+1)$ is also divisible by 3.
Thus $P(n)$ is divisible by 3 for all integers $n \geq 0$. For a negative integer $n = -m$, where $m > 0$,

$$P(-m) = (-m-1)^3 + (-m)^3 + (-m+1)^3 = -P(m),$$

so $P(-m)$ is also divisible by 3. Therefore $P(n)$ is divisible by 3 for every integer n.

Takeaways 3.2

- Induction can prove divisibility by comparing consecutive cases.
- After proving the non-negative cases, symmetry can extend the result to negative integers.
- Direct expansion gives $P(n) = 3n^3 + 6n$, which is a useful check on the induction proof.

 Vu Hung Nguyen · CC BY 4.0

Problem 3.9: Financial sequences and geometric sums

A loan of $\$L$ is taken out at an interest rate of r per period. At the end of each period, a constant repayment of $\$P$ is made after interest has been added to the balance. Let M_n be the balance owing after n periods.

(i) Write down expressions for M_1 and M_2 in terms of L, r, and P.

(ii) Show that the balance after n periods can be written as

$$M_n = L(1+r)^n - P\left[1 + (1+r) + (1+r)^2 + \cdots + (1+r)^{n-1}\right].$$

(iii) Using the formula for the sum of a geometric progression, prove that

$$M_n = L(1+r)^n - \frac{P((1+r)^n - 1)}{r}.$$

(iv) A car loan of \$20 000 is taken at 1% interest per month. Calculate the monthly repayment P required to reduce the balance to zero at the end of 24 months.

Hint:

- **For (i):** Add interest first, then subtract the repayment.
- **For (ii):** Expand the first few balances and look for the pattern.
- **For (iii):** The bracket is a finite GP with first term 1 and ratio $1 + r$.
- **For (iv):** Set $M_{24} = 0$, with $L = 20000$ and $r = 0.01$.

Solution 3.9

(i) After one period,

$$M_1 = L(1+r) - P.$$

After two periods,

$$M_2 = (L(1+r) - P)(1+r) - P = L(1+r)^2 - P(1+r) - P.$$

(ii) Continuing this pattern gives

$$M_n = L(1+r)^n - P\left[(1+r)^{n-1} + (1+r)^{n-2} + \cdots + (1+r) + 1\right].$$

Reordering the bracket,

$$M_n = L(1+r)^n - P\left[1 + (1+r) + (1+r)^2 + \cdots + (1+r)^{n-1}\right].$$

(iii) The bracket is a finite GP with first term 1, ratio $1 + r$, and n terms:

$$1 + (1+r) + \cdots + (1+r)^{n-1} = \frac{(1+r)^n - 1}{(1+r) - 1} = \frac{(1+r)^n - 1}{r}.$$

 Vu Hung Nguyen · CC BY 4.0

Hence

$$M_n = L(1+r)^n - \frac{P((1+r)^n - 1)}{r}.$$

(iv) Here $L = 20000$, $r = 0.01$, and $n = 24$. Set $M_{24} = 0$:

$$0 = 20000(1.01)^{24} - \frac{P((1.01)^{24} - 1)}{0.01}.$$

Thus

$$P = \frac{20000(1.01)^{24}(0.01)}{(1.01)^{24} - 1}.$$

Using $(1.01)^{24} \approx 1.269735$,

$$P \approx \$941.47.$$

Takeaways 3.3

- Loan balances can be modelled by applying interest and repayment recursively.
- Repeated repayments form a finite geometric sum after compounding.
- Setting the final balance to zero gives the repayment required to clear a loan.

 Vu Hung Nguyen · CC BY 4.0

Problem 3.10: Telescoping sums and inductive proof

Consider the sequence

$$a_k = \frac{2}{k(k+1)(k+2)} \qquad (k \geq 1).$$

Define the partial sums

$$S_n = \sum_{k=1}^{n} a_k.$$

(i) Show, using partial fractions, that

$$\frac{2}{k(k+1)(k+2)} = \frac{1}{k(k+1)} - \frac{1}{(k+1)(k+2)}.$$

(ii) Demonstrate that the sequence of partial sums telescopes. Hence find a simplified expression for S_n in terms of n.

(iii) Prove the expression for S_n found in part (ii) by mathematical induction for all $n \geq 1$.

(iv) Determine the limiting sum of the series as $n \to \infty$.

Hint:

- **For (i):** Put the two fractions over the common denominator $k(k+1)(k+2)$.
- **For (ii):** Write out the first few terms of S_n and look for cancellation.
- **For (iii):** Use $S_{m+1} = S_m + a_{m+1}$ in the induction step.
- **For (iv):** Take the limit of the closed form for S_n.

Solution 3.10

(i)

$$\frac{1}{k(k+1)} - \frac{1}{(k+1)(k+2)} = \frac{k+2-k}{k(k+1)(k+2)} = \frac{2}{k(k+1)(k+2)}.$$

(ii) Therefore

$$a_k = \frac{1}{k(k+1)} - \frac{1}{(k+1)(k+2)}.$$

So

$$\begin{aligned} S_n &= \left(\frac{1}{1\cdot 2} - \frac{1}{2\cdot 3}\right) + \left(\frac{1}{2\cdot 3} - \frac{1}{3\cdot 4}\right) + \cdots + \left(\frac{1}{n(n+1)} - \frac{1}{(n+1)(n+2)}\right) \\ &= \frac{1}{2} - \frac{1}{(n+1)(n+2)}. \end{aligned}$$

(iii) We prove

$$S_n = \frac{1}{2} - \frac{1}{(n+1)(n+2)}.$$

 Vu Hung Nguyen · CC BY 4.0

For $n = 1$,

$$S_1 = \frac{2}{1 \cdot 2 \cdot 3} = \frac{1}{3}$$

and

$$\frac{1}{2} - \frac{1}{2 \cdot 3} = \frac{1}{2} - \frac{1}{6} = \frac{1}{3}.$$

Now assume the formula holds for $n = m$. Then

$$\begin{aligned} S_{m+1} &= S_m + \frac{2}{(m+1)(m+2)(m+3)} \\ &= \frac{1}{2} - \frac{1}{(m+1)(m+2)} + \left(\frac{1}{(m+1)(m+2)} - \frac{1}{(m+2)(m+3)} \right) \\ &= \frac{1}{2} - \frac{1}{(m+2)(m+3)}. \end{aligned}$$

This is the required formula with $n = m + 1$, so the result holds for all $n \geq 1$.

(iv)

$$\lim_{n \to \infty} S_n = \lim_{n \to \infty} \left(\frac{1}{2} - \frac{1}{(n+1)(n+2)} \right) = \frac{1}{2}.$$

Takeaways 3.4

- Telescoping sums work by rewriting terms so that most neighbouring parts cancel.
- Induction is a useful way to verify a closed form found by pattern spotting.
- A finite partial-sum formula can make the limiting sum immediate.

Problem 3.11: Binomial coefficients and partial sums

Let

$$S_n = \sum_{k=0}^{n} \binom{n}{k}$$

be the sum of the binomial coefficients for a fixed integer $n \geq 1$.

(i) Use the binomial theorem expansion of $(1+x)^n$ to prove that $S_n = 2^n$.

(ii) Use Pascal's identity

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$$

to prove the recurrence relation $S_n = 2S_{n-1}$.

(iii) Evaluate the alternating partial sum

$$A_n = \sum_{k=0}^{n} (-1)^k \binom{n}{k}$$

and explain why the sum of even-indexed coefficients equals the sum of odd-indexed coefficients.

(iv) By differentiating the expansion of $(1+x)^n$ with respect to x, show that

$$\sum_{k=1}^{n} k \binom{n}{k} = n2^{n-1}.$$

Hint:

- **For (i):** Substitute $x = 1$ into the binomial theorem.
- **For (ii):** Sum Pascal's identity over the appropriate range of k.
- **For (iii):** Substitute $x = -1$ into the binomial theorem.
- **For (iv):** Differentiate first, then substitute $x = 1$.

Solution 3.11

(i) By the binomial theorem,

$$(1+x)^n = \sum_{k=0}^{n} \binom{n}{k} x^k.$$

Putting $x = 1$ gives

$$2^n = \sum_{k=0}^{n} \binom{n}{k} = S_n.$$

 Vu Hung Nguyen · CC BY 4.0

(ii) Using Pascal's identity,

$$\begin{aligned} S_n &= \sum_{k=0}^{n} \binom{n}{k} \\ &= \sum_{k=0}^{n} \binom{n-1}{k} + \sum_{k=0}^{n} \binom{n-1}{k-1}. \end{aligned}$$

The out-of-range terms are zero, so each sum equals S_{n-1}. Hence

$$S_n = 2S_{n-1}.$$

(iii) By the binomial theorem again,

$$A_n = \sum_{k=0}^{n} (-1)^k \binom{n}{k} = (1-1)^n = 0.$$

Therefore

$$\sum_{\substack{0 \le k \le n \\ k \text{ even}}} \binom{n}{k} - \sum_{\substack{0 \le k \le n \\ k \text{ odd}}} \binom{n}{k} = 0,$$

so the even-indexed and odd-indexed sums are equal.
(iv) Differentiate

$$(1+x)^n = \sum_{k=0}^{n} \binom{n}{k} x^k$$

to get

$$n(1+x)^{n-1} = \sum_{k=1}^{n} k \binom{n}{k} x^{k-1}.$$

Putting $x = 1$ gives

$$\sum_{k=1}^{n} k \binom{n}{k} = n2^{n-1}.$$

Takeaways 3.5

- Substituting special values into the binomial theorem turns an expansion into a sum identity.
- Pascal's identity explains why the total sum of coefficients doubles from one row to the next.
- Differentiating a generating expansion is a powerful way to evaluate weighted sums.

 Vu Hung Nguyen · CC BY 4.0

Problem 3.12: The Recursive Geometry of Chebyshev Sequences

Let a sequence of functions $(T_n(x))$ be defined for $n \geq 0$ by the recurrence relation

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x),$$

with initial terms $T_0(x) = 1$ and $T_1(x) = x$.

(i) Calculate the explicit polynomial expressions for $T_2(x)$ and $T_3(x)$.

(ii) Using the identity

$$\cos(A+B) + \cos(A-B) = 2\cos A \cos B,$$

prove by mathematical induction that for all $n \geq 0$,

$$T_n(\cos\theta) = \cos(n\theta).$$

(iii) Let x_0 be a root of $T_n(x)$. Show that $|x_0| \leq 1$. Hence, determine the exact values of the n distinct roots of $T_n(x)$ in terms of n.

(iv) Consider the sequence of values $v_n = T_n(1.5)$. Show that (v_n) satisfies a linear recurrence relation with constant coefficients. Find a closed-form expression for v_n and determine the value of the limit

$$\lim_{n\to\infty} \frac{v_{n+1}}{v_n}.$$

Hint:

- **For (i):** Substitute the previous two polynomials into the recurrence formula.
- **For (ii):** Use the sum-to-product identity with $A = k\theta$ and $B = \theta$.
- **For (iii):** Solve $\cos(n\theta) = 0$ for θ, then use $x = \cos\theta$.
- **For (iv):** Solve the characteristic equation for the recurrence $v_{n+1} = 3v_n - v_{n-1}$.

 Vu Hung Nguyen · CC BY 4.0

Solution 3.12

(i) Directly from the recurrence,

$$T_2(x) = 2x^2 - 1, \qquad T_3(x) = 2x(2x^2 - 1) - x = 4x^3 - 3x.$$

(ii) The cases $n = 0, 1$ give $1 = \cos 0$ and $T_1(\cos\theta) = \cos\theta$. Suppose

$$T_k(\cos\theta) = \cos(k\theta), \qquad T_{k-1}(\cos\theta) = \cos((k-1)\theta).$$

Then

$$\begin{aligned} T_{k+1}(\cos\theta) &= 2\cos\theta\, T_k(\cos\theta) - T_{k-1}(\cos\theta) \\ &= 2\cos\theta\cos(k\theta) - \cos((k-1)\theta) \\ &= \cos((k+1)\theta), \end{aligned}$$

using $\cos((k+1)\theta) + \cos((k-1)\theta) = 2\cos(k\theta)\cos\theta$. Hence $T_n(\cos\theta) = \cos(n\theta)$ for all $n \geq 0$.

(iii) If $x > 1$, write $x = \cosh u$; then $T_n(x) = \cosh(nu) > 0$. If $x < -1$, write $x = -\cosh u$; then $T_n(x) = (-1)^n \cosh(nu) \neq 0$. Thus every real root satisfies $|x_0| \leq 1$.
Now write $x = \cos\theta$. The roots satisfy

$$T_n(\cos\theta) = 0 \iff \cos(n\theta) = 0 \iff n\theta = \frac{(2k-1)\pi}{2}.$$

So the n distinct roots are

$$x_k = \cos\left(\frac{(2k-1)\pi}{2n}\right), \qquad k = 1, 2, \ldots, n.$$

(iv) With $v_n = T_n(1.5)$,

$$v_{n+1} = 3v_n - v_{n-1}, \qquad v_0 = 1, \quad v_1 = \frac{3}{2}.$$

The characteristic equation is $\lambda^2 - 3\lambda + 1 = 0$, with roots

$$\alpha = \frac{3+\sqrt{5}}{2}, \qquad \beta = \frac{3-\sqrt{5}}{2}.$$

Hence $v_n = A\alpha^n + B\beta^n$. From $v_0 = 1$ and $v_1 = \frac{3}{2}$, $A = B = \frac{1}{2}$, so

$$v_n = \frac{1}{2}\alpha^n + \frac{1}{2}\beta^n.$$

Since $|\beta| < \alpha$, the dominant term is α^n, and therefore

$$\lim_{n\to\infty} \frac{v_{n+1}}{v_n} = \alpha = \frac{3+\sqrt{5}}{2}.$$

 Vu Hung Nguyen · CC BY 4.0

Takeaways 3.6

- The recurrence produces the Chebyshev polynomials one term at a time.
- The identity $T_n(\cos\theta) = \cos(n\theta)$ turns polynomial roots into trigonometric roots.
- Evaluating a polynomial recurrence at a fixed value gives an ordinary numerical recurrence.

3.3 Advanced Sequence Problems

Problem 3.13: Recursive sequence and limiting value

Let (x_n) be defined by

$$x_1 = 16, \qquad x_{n+1} = 2 + \frac{1}{x_n} \quad (n \geq 1).$$

(i) Find exact values of x_2, x_3, x_4.

(ii) Assuming $x_n \to L > 0$, show that $L = 1 + \sqrt{2}$.

(iii) Prove that

$$|x_{n+1} - L| = \frac{|x_n - L|}{Lx_n},$$

and deduce that for $n \geq 2$,

$$|x_{n+1} - L| < \frac{1}{4}|x_n - L|.$$

(iv) Hence show $x_n \to 1 + \sqrt{2}$.

Solution 3.13

(i)

$$x_2 = 2 + \frac{1}{16} = \frac{33}{16}, \quad x_3 = 2 + \frac{16}{33} = \frac{82}{33}, \quad x_4 = 2 + \frac{33}{82} = \frac{197}{82}.$$

For $n \geq 2$, we have $x_n > 2$ because each term is $2 + \frac{1}{x_{n-1}}$ with positive second part.

(ii) If $x_n \to L$, then taking limits in the recurrence gives

$$L = 2 + \frac{1}{L} \implies L^2 - 2L - 1 = 0.$$

So $L = 1 \pm \sqrt{2}$. Since terms are positive and > 2 from $n \geq 2$, we take

$$L = 1 + \sqrt{2}.$$

(iii)

$$x_{n+1} - L = \left(2 + \frac{1}{x_n}\right) - \left(2 + \frac{1}{L}\right) = \frac{L - x_n}{Lx_n},$$

thus

$$|x_{n+1} - L| = \frac{|x_n - L|}{Lx_n}.$$

 Vu Hung Nguyen · CC BY 4.0

Now $L = 1 + \sqrt{2} > 2$ and $x_n > 2$ for $n \geq 2$, so $Lx_n > 4$. Hence

$$|x_{n+1} - L| < \frac{1}{4}|x_n - L|, \quad n \geq 2.$$

(iv) Repeatedly applying the inequality gives geometric decay of the error:

$$|x_n - L| \leq \left(\frac{1}{4}\right)^{n-2} |x_2 - L| \to 0.$$

Therefore $x_n \to L = 1 + \sqrt{2}$.

Takeaways 3.7

- For recursive limits, first solve the fixed-point equation.
- A contraction inequality gives a rigorous convergence proof.
- The same method works for many rational recursions.

Problem 3.14: The Dynamics of Iteration

Consider the function

$$g(x) = \sqrt{x+2}.$$

We define an iterative sequence by

$$x_{n+1} = g(x_n), \qquad x_1 = 1.$$

(i) Calculate x_2, x_3, and x_4 to three decimal places. By solving

$$L = \sqrt{L+2},$$

determine the exact positive limit L that the sequence appears to be approaching.

(ii) Show that

$$0 < g'(x) < \frac{1}{2}$$

for all $x > 0$.

(iii) Prove that

$$|x_{n+1} - L| < \frac{1}{2}|x_n - L|.$$

Hence deduce that the sequence converges to L for any initial value $x_1 > 0$.

(iv) Define a function $f(x)$ such that Newton's method would converge to the same limit L found in part (i), and write down the specific iterative formula for this $f(x)$.

 Vu Hung Nguyen · CC BY 4.0

Hint:

- **For (i):** Substitute repeatedly into $x_{n+1} = \sqrt{x_n + 2}$, then solve the fixed-point equation.
- **For (ii):** Differentiate $g(x) = \sqrt{x+2}$ and note where the derivative is largest for $x > 0$.
- **For (iii):** Use either the mean value theorem or rationalise $\sqrt{x_n + 2} - \sqrt{L+2}$.
- **For (iv):** Rearrange $L = \sqrt{L+2}$ into a polynomial equation.

Solution 3.14

(i)

$$x_2 = \sqrt{3} \approx 1.732, \quad x_3 = \sqrt{2+\sqrt{3}} \approx 1.932, \quad x_4 \approx 1.983.$$

If $x_n \to L > 0$, then

$$L = \sqrt{L+2} \implies L^2 - L - 2 = 0 \implies (L-2)(L+1) = 0.$$

Thus the positive limit is

$$L = 2.$$

(ii)

$$g'(x) = \frac{1}{2\sqrt{x+2}}.$$

For $x > 0$, this is positive and

$$g'(x) < \frac{1}{2\sqrt{2}} < \frac{1}{2}.$$

(iii) Since $L = 2$ and $g(L) = L$,

$$\begin{aligned} |x_{n+1} - L| &= \left|\sqrt{x_n+2} - \sqrt{L+2}\right| \\ &= \frac{|x_n - L|}{\sqrt{x_n+2} + \sqrt{L+2}} \\ &< \frac{1}{2}|x_n - L|, \end{aligned}$$

because $x_n > 0$ and $\sqrt{L+2} = 2$. Repeating this inequality gives

$$|x_n - L| < \left(\frac{1}{2}\right)^{n-1} |x_1 - L| \to 0,$$

so $x_n \to L = 2$ for any $x_1 > 0$.

(iv) From the fixed-point equation,

$$L^2 - L - 2 = 0.$$

So take

$$f(x) = x^2 - x - 2.$$

Newton's method gives

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^2 - x_n - 2}{2x_n - 1}.$$

 Vu Hung Nguyen · CC BY 4.0

Takeaways 3.8

- Fixed points often reveal the likely limit of an iterative sequence.
- A contraction inequality proves convergence by forcing errors to shrink geometrically.
- Newton's method can target the same limit by solving the fixed-point equation as a root problem.

Problem 3.15: The Geometry of Harmonic Means

A sequence (h_n) is in harmonic progression (HP) if the sequence of reciprocals

$$x_n = \frac{1}{h_n}$$

forms an arithmetic progression (AP).

(i) Given that the third term of an HP is 1 and the sixth term is $\frac{1}{2}$, find the first term h_1 and the common difference d of the underlying AP.

(ii) The harmonic mean H of two positive numbers a and b is defined so that a, H, b is an HP. Show algebraically that

$$H = \frac{2ab}{a+b}.$$

(iii) Let

$$A = \frac{a+b}{2}, \qquad G = \sqrt{ab}$$

be the arithmetic and geometric means of a and b. Prove that $G^2 = AH$. Hence prove that, for distinct positive real numbers a and b,

$$H < G < A.$$

(iv) Consider the infinite harmonic series

$$\sum_{n=1}^{\infty} \frac{1}{n}.$$

By comparing the sum of the first 2^k terms to blocks of constant lower bounds, prove that this harmonic series does not have a finite limiting sum.

 Vu Hung Nguyen · CC BY 4.0

Hint:

- **For (i):** Convert the HP terms into AP terms x_3 and x_6, then use $x_n = a + (n-1)d$.
- **For (ii):** If a, H, b is an HP, then $\frac{1}{a}, \frac{1}{H}, \frac{1}{b}$ is an AP.
- **For (iii):** Multiply A and H. For the inequality, use $(a-b)^2 > 0$ or $(\sqrt{a}-\sqrt{b})^2 > 0$.
- **For (iv):** Group terms as $\left(\frac{1}{3}+\frac{1}{4}\right)$, then $\left(\frac{1}{5}+\cdots+\frac{1}{8}\right)$, and so on.

Solution 3.15

(i) Since $x_n = \frac{1}{h_n}$,

$$x_3 = 1, \qquad x_6 = 2.$$

For the underlying AP,

$$x_6 - x_3 = 3d \implies 2 - 1 = 3d \implies d = \frac{1}{3}.$$

Also

$$x_1 = x_3 - 2d = 1 - \frac{2}{3} = \frac{1}{3},$$

so

$$h_1 = \frac{1}{x_1} = 3.$$

(ii) Since $\frac{1}{a}, \frac{1}{H}, \frac{1}{b}$ is an AP, the middle term is the average of the outer terms:

$$\frac{1}{H} = \frac{1}{2}\left(\frac{1}{a} + \frac{1}{b}\right) = \frac{a+b}{2ab}.$$

Therefore

$$H = \frac{2ab}{a+b}.$$

(iii) Using the formulas for A and H,

$$AH = \frac{a+b}{2} \cdot \frac{2ab}{a+b} = ab = G^2.$$

For $a \neq b$,

$$(\sqrt{a} - \sqrt{b})^2 > 0 \implies a + b > 2\sqrt{ab} \implies A > G.$$

Since $G^2 = AH$ and all quantities are positive, $A > G$ gives

$$H = \frac{G^2}{A} < G.$$

Hence

$$H < G < A.$$

(iv) Group the harmonic series into dyadic blocks:

$$1+\frac{1}{2}+\left(\frac{1}{3}+\frac{1}{4}\right)+\left(\frac{1}{5}+\cdots+\frac{1}{8}\right)+\cdots.$$

Each block after the first has sum greater than or equal to $\frac{1}{2}$:

$$\frac{1}{3}+\frac{1}{4}>\frac{1}{4}+\frac{1}{4}=\frac{1}{2}, \qquad \frac{1}{5}+\cdots+\frac{1}{8}>4\cdot\frac{1}{8}=\frac{1}{2},$$

and similarly

$$\frac{1}{2^{j-1}+1}+\cdots+\frac{1}{2^j}>2^{j-1}\cdot\frac{1}{2^j}=\frac{1}{2}.$$

Thus the partial sums exceed

$$1+\frac{1}{2}+\underbrace{\frac{1}{2}+\frac{1}{2}+\cdots+\frac{1}{2}}_{\text{arbitrarily many terms}},$$

so they grow without bound. Therefore the harmonic series has no finite limiting sum.

Takeaways 3.9

- Harmonic progressions are arithmetic progressions viewed through reciprocals.
- For positive distinct numbers, the classical means satisfy $H < G < A$.
- A sequence can have terms tending to zero while its infinite series still diverges.

Problem 3.16: Chebyshev composition

The first-kind Chebyshev polynomials $T_n(x)$ are defined by the recurrence

$$T_0(x)=1, \quad T_1(x)=x, \quad T_{n+1}(x)=2xT_n(x)-T_{n-1}(x).$$

It is given that $T_n(\cos\theta)=\cos(n\theta)$ for all $n\geq 0$ and $\theta\in\mathbb{R}$. (Do NOT prove this fact.) For first-kind Chebyshev polynomials, prove that

$$T_m(T_n(x))=T_{mn}(x)$$

for all x of the form $x=\cos\theta$. Then use this result to find

$$T_2(T_3(x)).$$

Hint: Use $T_n(\cos\theta)=\cos(n\theta)$. For the explicit polynomial, use $T_6(x)$.

 Vu Hung Nguyen · CC BY 4.0

Solution 3.16

Let $x = \cos\theta$. Then

$$T_m(T_n(x)) = T_m(\cos(n\theta)) = \cos(mn\theta) = T_{mn}(\cos\theta) = T_{mn}(x).$$

Therefore

$$T_2(T_3(x)) = T_6(x).$$

Using the recurrence after $T_5(x) = 16x^5 - 20x^3 + 5x$,

$$T_6(x) = 2xT_5(x) - T_4(x) = 32x^6 - 48x^4 + 18x^2 - 1.$$

Takeaways 3.10

- The trigonometric definition makes Chebyshev composition simple.
- Composition of first-kind Chebyshev polynomials multiplies the indices.
- Hard-looking polynomial composition can often be avoided by using structure.

 Vu Hung Nguyen · CC BY 4.0

4 Part 2: Problems with Hints and Solutions (Concise)

Part 2 gives extra practice with upside-down hints and concise solutions.

4.1 Basic Sequence Problems

Problem 4.1: Quick AP term

Find a_{18} for the arithmetic sequence $2, 7, 12, \ldots$.

Hint: Use $a_n = a + (n-1)d$ with $a = 2$ and $d = 5$.

Solution 4.1

$$a_{18} = 2 + 17 \cdot 5 = 87.$$

Problem 4.2: Quick GP sum

Find

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots + \frac{1}{128}.$$

Hint: This is a finite GP with $a = 1$, $r = \frac{1}{2}$, and 8 terms.

Solution 4.2

$$S_8 = \frac{1 - (\frac{1}{2})^8}{1 - \frac{1}{2}} = 2\left(1 - \frac{1}{256}\right) = \frac{255}{128}.$$

Problem 4.3: Quick Chebyshev value

For the second-kind Chebyshev polynomial

$$U_2(x) = 4x^2 - 1,$$

find $U_2\left(\frac{1}{2}\right)$.

Hint: Substitute $x = \frac{1}{2}$ carefully.

Solution 4.3

$$U_2\left(\frac{1}{2}\right) = 4\left(\frac{1}{4}\right) - 1 = 1 - 1 = 0.$$

Takeaways 4.1

- For a given Chebyshev polynomial, direct substitution is often enough.
- Square fractions before multiplying by the coefficient.

Problem 4.4: Arithmetic pricing model

An installer charges \$400 for the first window and \$100 for each additional window.

(i) Write the total cost for 1, 2, 3, and 4 windows.

(ii) Find a formula for the total cost A_n for n windows.

(iii) Find A_{15}.

(iv) If the budget is strictly less than \$10,000, what is the maximum number of windows?

Hint: Model A_n as an arithmetic sequence with first term 400 and difference 100.

Solution 4.4

$$A_n = 400 + (n-1)100 = 100n + 300.$$

So $A_{15} = 1800$. For the budget:

$$100n + 300 < 10000 \implies n < 97,$$

hence the maximum integer is $n = 96$.

Problem 4.5: Simple interest and AP

A principal of \$2,000,000 is invested at 5.7% p.a. simple interest. Let A_n be the total amount after n years.

(i) Find A_1, A_2, A_3, A_4.

(ii) Find a formula for A_n and evaluate A_{12}.

(iii) How many whole years before the amount exceeds \$6,000,000?

Hint: Simple interest adds a fixed amount each year, so (A_n) is arithmetic.

 Vu Hung Nguyen · CC BY 4.0

Solution 4.5

Yearly interest is $0.057 \times 2{,}000{,}000 = 114{,}000$. Hence

$$A_n = 2{,}000{,}000 + 114{,}000n.$$

So

$$A_1 = 2{,}114{,}000,\ A_2 = 2{,}228{,}000,\ A_3 = 2{,}342{,}000,\ A_4 = 2{,}456{,}000,$$

and

$$A_{12} = 3{,}368{,}000.$$

For exceeding \$6,000,000:

$$2{,}000{,}000 + 114{,}000n > 6{,}000{,}000 \implies n > 35.087\ldots$$

so $n = 36$ years.

Problem 4.6: Recurring decimal to fraction

Convert each to a fraction:

$$0.2, \qquad 0.\dot{2}, \qquad 0.1\dot{2}.$$

Hint: Use place value for terminating decimals and GP ideas for recurring decimals.

Solution 4.6

$$0.2 = \frac{1}{5}, \qquad 0.\dot{2} = \frac{2}{9}, \qquad 0.1\dot{2} = \frac{11}{90}.$$

Problem 4.7: Nested shaded squares

In a unit square, shade the bottom-left quarter. Then repeatedly subdivide the top-right quarter into four equal squares and shade its bottom-left quarter. Find the total shaded fraction.

Hint: Shaded areas form an infinite GP with first term $\frac{1}{4}$ and ratio $\frac{1}{4}$.

Solution 4.7

$$S_\infty = \frac{\frac{1}{4}}{1 - \frac{1}{4}} = \frac{1}{3}.$$

So the shaded fraction is $\frac{1}{3}$.

 Vu Hung Nguyen · CC BY 4.0

4.2 Medium Sequence Problems

Problem 4.8: Indexing with sigma notation

Given

$$\sum_{k=1}^{n}(3k-1) = 119,$$

find n.

Hint: Split the sum into $3\sum k - \sum 1$.

Solution 4.8

$$\sum_{k=1}^{n}(3k-1) = 3 \cdot \frac{n(n+1)}{2} - n = \frac{3n^2+n}{2}.$$

Set equal to 119:

$$\frac{3n^2+n}{2} = 119 \implies 3n^2+n-238 = 0.$$

$$\Delta = 1 + 2856 = 2857.$$

This is not a perfect square, so there is no integer n with exact value 119.

Problem 4.9: Infinite geometric interpretation

Evaluate

$$0.4\overline{2} = 0.42222\ldots$$

as a fraction.

Hint: Write as $0.4 + 0.02222\ldots$ and convert the recurring part using a GP.

Solution 4.9

$$0.02222\ldots = \frac{2}{90} = \frac{1}{45}.$$

So

$$0.4\overline{2} = \frac{2}{5} + \frac{1}{45} = \frac{18}{45} + \frac{1}{45} = \frac{19}{45}.$$

Problem 4.10: Second-kind Chebyshev recurrence

The second-kind Chebyshev polynomials satisfy

$$U_0(x) = 1, \qquad U_1(x) = 2x, \qquad U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x).$$

Use this recurrence to find $U_3(x)$ and $U_4(x)$.

 Vu Hung Nguyen · CC BY 4.0

Hint: First find $U_2(x)$, then continue to $U_3(x)$ and $U_4(x)$.

Solution 4.10

$$U_2(x) = 2x(2x) - 1 = 4x^2 - 1.$$
$$U_3(x) = 2x(4x^2 - 1) - 2x = 8x^3 - 4x.$$
$$U_4(x) = 2x(8x^3 - 4x) - (4x^2 - 1) = 16x^4 - 12x^2 + 1.$$

Takeaways 4.2

- The first-kind and second-kind Chebyshev polynomials use the same recurrence form.
- The starting values determine which family is produced.
- Organize recurrence work line by line to avoid algebra slips.

Problem 4.11: Alternative telescoping patterns

(i) Evaluate $\sum_{n=0}^{4}(-1)^{n+1}(n+5)$.

(ii) Show $\frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}$.

(iii) Hence find

$$S_N = \sum_{n=1}^{N}\left(\frac{1}{n} - \frac{1}{n+1}\right)$$

and $\lim_{N\to\infty} S_N$.

Hint: Part (iii) is a direct telescoping sum after part (ii).

Solution 4.11

$$\sum_{n=0}^{4}(-1)^{n+1}(n+5) = -7.$$

Also

$$\frac{1}{n(n+1)} = \frac{(n+1)-n}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}.$$

So

$$S_N = 1 - \frac{1}{N+1}, \qquad \lim_{N\to\infty} S_N = 1.$$

 Vu Hung Nguyen · CC BY 4.0

Problem 4.12: Surd telescoping

Let

$$a_k = \frac{1}{\sqrt{k+1}+\sqrt{k}}, \qquad S_n = \sum_{k=1}^{n} a_k.$$

(i) Show $a_k = \sqrt{k+1} - \sqrt{k}$.

(ii) Find S_n.

(iii) Evaluate S_{99}.

(iv) Find the smallest integer N such that $S_N > 10$.

Hint: Rationalize with the conjugate and then telescope.

Solution 4.12

$$a_k = \frac{\sqrt{k+1}-\sqrt{k}}{(k+1)-k} = \sqrt{k+1}-\sqrt{k}.$$

Hence

$$S_n = (\sqrt{2}-\sqrt{1}) + \cdots + (\sqrt{n+1}-\sqrt{n}) = \sqrt{n+1}-1.$$

So $S_{99} = 10 - 1 = 9$. For $S_N > 10$:

$$\sqrt{N+1} - 1 > 10 \implies N > 120,$$

smallest such integer is $N = 121$.

Problem 4.13: Central symmetry in a GP

In a GP, define $P_n = T_1 T_2 \cdots T_n$.

(i) Show $P_n = a^n r^{\frac{n(n-1)}{2}}$.

(ii) Show $T_k T_{n-k+1} = T_1 T_n$.

(iii) If $n = 11$ and middle term $T_6 = 5$, find P_{11}.

Hint: Use $T_k = ar^{k-1}$ and pair symmetric terms.

Solution 4.13

$$P_n = a(ar)\cdots(ar^{n-1}) = a^n r^{0+1+\cdots+(n-1)} = a^n r^{\frac{n(n-1)}{2}}.$$

Also

$$T_k T_{n-k+1} = ar^{k-1} \cdot ar^{n-k} = a^2 r^{n-1} = T_1 T_n.$$

 Vu Hung Nguyen · CC BY 4.0

For odd $n = 11$, the product is middle-term power:

$$P_{11} = T_6^{11} = 5^{11}.$$

Problem 4.14: Fibonacci and Lucas

Define

$$F_1 = 1,\ F_2 = 1,\ F_n = F_{n-1} + F_{n-2},$$
$$L_1 = 1,\ L_2 = 3,\ L_n = L_{n-1} + L_{n-2}.$$

Show that

$$L_n = F_{n-1} + F_{n+1}.$$

Hint: Set $A_n = L_n + F_n$ and $B_n = L_n - F_n$, then compare with Fibonacci terms.

Solution 4.14

From initial terms and recurrence,

$$A_n = L_n + F_n = 2F_{n+1}, \qquad B_n = L_n - F_n = 2F_{n-1}.$$

Adding gives

$$2L_n = 2F_{n+1} + 2F_{n-1},$$

so

$$L_n = F_{n-1} + F_{n+1}.$$

Problem 4.15: Linear combinations of two GPs

Let $T_n = ar^{n-1}$ and $U_n = AR^{n-1}$. Define $W_n = T_n + U_n$.

(i) Show (W_n) satisfies

$$W_{n+2} - (r + R)W_{n+1} + rRW_n = 0.$$

(ii) Solve

$$X_{n+2} - 5X_{n+1} + 6X_n = 0, \quad X_1 = 5,\ X_2 = 13.$$

Hint: For (ii), the characteristic roots are 2 and 3.

Solution 4.15

Substitute $W_n = ar^{n-1} + AR^{n-1}$ directly to verify (i). For (ii), write

$$X_n = \alpha 2^{n-1} + \beta 3^{n-1}.$$

 Vu Hung Nguyen · CC BY 4.0

Using $X_1 = 5,\ X_2 = 13$:

$$\alpha + \beta = 5, \quad 2\alpha + 3\beta = 13$$

gives $\alpha = 2,\ \beta = 3$. Hence

$$X_n = 2^n + 3^n.$$

4.3 Advanced Sequence Problems

Problem 4.16: Convergence condition

Consider the geometric series

$$\sum_{n=0}^{\infty} 7\left(\frac{m-1}{3}\right)^n.$$

Find all real values of m for which the series converges, and then state its sum.

Hint: Use the condition $|r| < 1$ for convergence of an infinite GP.

Solution 4.16

Here

$$r = \frac{m-1}{3}.$$

Convergence requires

$$\left|\frac{m-1}{3}\right| < 1 \implies |m-1| < 3 \implies -2 < m < 4.$$

For these values,

$$S_\infty = \frac{7}{1 - \frac{m-1}{3}} = \frac{21}{4-m}.$$

Problem 4.17: AP and GP crossover

A sequence is both arithmetic and geometric. Given $a_1 = 6$ and $a_3 = 6$, find all possible sequences.

Hint: Let AP difference be d and GP ratio be r. Use both models on a_3.

Solution 4.17

AP gives

$$a_3 = a_1 + 2d = 6 + 2d = 6 \implies d = 0,$$

so AP is constant: $a_n = 6$. For GP,

$$a_3 = a_1 r^2 = 6r^2 = 6 \implies r^2 = 1 \implies r = \pm 1.$$

 Vu Hung Nguyen · CC BY 4.0

If $r = -1$, terms alternate $6, -6, 6, \ldots$ and are not arithmetic. Hence the only sequence that is both AP and GP is

$$a_n = 6 \quad \text{for all } n,$$

with $d = 0$ and $r = 1$.

Problem 4.18: Chebyshev roots

The first-kind Chebyshev polynomial of degree 5 is

$$T_5(x) = 16x^5 - 20x^3 + 5x.$$

Use the identity $T_5(\cos\theta) = \cos(5\theta)$ to find all roots of $T_5(x)$ in the interval $[-1, 1]$.

Hint: Set $\cos(5\theta) = 0$, then convert the resulting angles back to $x = \cos\theta$.

Solution 4.18

Let $x = \cos\theta$. We need

$$\cos(5\theta) = 0.$$

Thus

$$5\theta = \frac{(2k+1)\pi}{2}, \qquad k = 0, 1, 2, 3, 4.$$

So

$$\theta = \frac{(2k+1)\pi}{10},$$

and the five roots are

$$x = \cos\frac{\pi}{10},\ \cos\frac{3\pi}{10},\ \cos\frac{5\pi}{10},\ \cos\frac{7\pi}{10},\ \cos\frac{9\pi}{10}.$$

Takeaways 4.3

- Roots of $T_n(x)$ on $[-1, 1]$ come from zeros of $\cos(n\theta)$.
- The substitution $x = \cos\theta$ is natural because $\cos\theta$ covers $[-1, 1]$.
- Degree 5 gives five roots, which is a useful check.

 Vu Hung Nguyen · CC BY 4.0

Problem 4.19: Architecture of GP sums

Let a GP have first term $a \neq 0$, ratio $r \neq 0, \pm 1$, and partial sums S_n.

(i) Prove $\dfrac{S_{2n}}{S_n} = r^n + 1$.

(ii) Given $S_{12} = 65S_6$ and second term $ar = 12$, find possible (a, r).

(iii) If Σ_n is the partial-sum sequence of a GP with first term a and ratio r^2, show

$$\frac{\Sigma_n}{S_n} = \frac{r^n + 1}{r + 1}.$$

Hint: Use $S_n = \dfrac{a(r^n - 1)}{r - 1}$ and difference-of-squares factorization.

Solution 4.19

$$\frac{S_{2n}}{S_n} = \frac{r^{2n} - 1}{r^n - 1} = r^n + 1.$$

So $r^6 + 1 = 65 \implies r^6 = 64 \implies r = \pm 2$. Since $ar = 12$, possible pairs are $(a, r) = (6, 2)$ or $(-6, -2)$. Also

$$\Sigma_n = \frac{a(r^{2n} - 1)}{r^2 - 1},$$

thus

$$\frac{\Sigma_n}{S_n} = \frac{r^{2n} - 1}{(r + 1)(r^n - 1)} = \frac{r^n + 1}{r + 1}.$$

Problem 4.20: AP space and difference operator

Let $\mathcal{A}(a, d)$ denote an AP. Define

$$\mathcal{I} = \mathcal{A}(1, 0), \qquad \mathcal{J} = \mathcal{A}(0, 1).$$

(i) Show any AP is uniquely $a\mathcal{I} + d\mathcal{J}$.

(ii) Let $\Delta x_n = x_{n+1} - x_n$. Show $\Delta\mathcal{I} = 0$ and $\Delta\mathcal{J} = \mathcal{I}$.

(iii) Deduce $\Delta(a\mathcal{I} + d\mathcal{J}) = d\mathcal{I}$.

Hint: Compare first term and common difference.

Solution 4.20

An AP has nth term $a + (n-1)d = a \cdot 1 + d(n-1)$, so $X = a\mathcal{I} + d\mathcal{J}$, unique by matching (a, d). Also $\mathcal{I} = (1, 1, 1, \dots)$ so $\Delta\mathcal{I} = 0$, and $\mathcal{J} = (0, 1, 2, \dots)$ so $\Delta\mathcal{J} = (1, 1, 1, \dots) = \mathcal{I}$.

 Vu Hung Nguyen · CC BY 4.0

Hence

$$\Delta(a\mathcal{I} + d\mathcal{J}) = a\Delta\mathcal{I} + d\Delta\mathcal{J} = d\mathcal{I}.$$

Problem 4.21: Basis and projection in AP space

Let $\mathcal{E}_1 = \mathcal{A}(1,1)$ and $\mathcal{E}_2 = \mathcal{A}(1,-1)$.

(i) Express any AP $\mathcal{A}(a,d)$ as $\lambda\mathcal{E}_1 + \mu\mathcal{E}_2$.

(ii) Decompose $P = (5, 8, 11, 14, \dots)$ in this basis.

Hint: Match first terms and differences: $\lambda + \mu = a$, $\lambda - \mu = d$.

Solution 4.21

Solving

$$\lambda + \mu = a, \qquad \lambda - \mu = d$$

gives

$$\lambda = \frac{a+d}{2}, \qquad \mu = \frac{a-d}{2}.$$

For P, $a = 5,\ d = 3$, so $\lambda = 4,\ \mu = 1$.

Problem 4.22: Countable and uncountable sets

(i) Explain why positive rationals are countable by diagonal listing.

(ii) Use Cantor's diagonal argument to show reals in $[0,1)$ are uncountable.

Hint: Construct a new decimal differing from the nth listed decimal at position n.

Solution 4.22

Rationals can be arranged in an infinite grid $\frac{p}{q}$ and traversed by diagonals, skipping duplicates after simplification; this yields a sequence listing all positive rationals. For reals in $[0,1)$, assume a list $T_1, T_2, \dots$ exists. Build x so its nth digit differs from the nth digit of T_n. Then $x \neq T_n$ for every n, contradicting completeness of the list. Hence uncountable.

 Vu Hung Nguyen · CC BY 4.0

Problem 4.23: Prime denominators and cycle length

Let p be prime, $p \neq 2, 5$, and L be period length of $\frac{1}{p}$.

(i) Show L is the least positive integer with $10^L \equiv 1 \pmod{p}$.

(ii) Deduce $L \mid (p-1)$.

(iii) Find periods for $\frac{1}{37}$ and $\frac{1}{41}$.

Hint: Use Fermat's theorem and first occurrence of $p \mid (10^n - 1)$.

Solution 4.23

Recurring block theory gives $(10^L - 1)/p \in \mathbb{Z}$, hence $10^L \equiv 1 \pmod{p}$; minimality gives least such L. By Fermat, $10^{p-1} \equiv 1 \pmod{p}$, so the order L divides $p-1$. Since $37 \mid (10^3 - 1)$ and not $10^1 - 1, 10^2 - 1$, period is 3. Since $41 \mid (10^5 - 1)$ and not $10^1 - 1$, period is 5.

Problem 4.24: Pythagorean means

For $0 < a < b$, define

$$A = \frac{a+b}{2}, \quad G = \sqrt{ab}, \quad H = \frac{2ab}{a+b}.$$

(i) Prove $H < G < A$.

(ii) Find $\frac{b}{a}$ such that a, G, A is an AP.

(iii) Show H, G, A cannot be an AP unless $a = b$.

Hint: Use $(\sqrt{a} - \sqrt{b})^2 > 0$ and $G^2 = AH$.

Solution 4.24

From $(\sqrt{a} - \sqrt{b})^2 > 0$, get $A > G$. Also

$$AH = \frac{a+b}{2} \cdot \frac{2ab}{a+b} = ab = G^2,$$

so H, G, A form a GP and therefore $H < G < A$. For a, G, A in AP:

$$2G = a + A \implies 4\sqrt{ab} = 3a + b.$$

Let $x = b/a$; then $(x-9)(x-1) = 0$, and $x > 1$ gives $x = 9$. If H, G, A were AP, then $2G = H + A = G^2/A + A$, forcing $A = G$, contradiction for $a \neq b$.

 Vu Hung Nguyen · CC BY 4.0

Problem 4.25: Recurring decimals and modular structure

Let p be prime ($p \neq 2, 5$), and let n be the period of $\frac{1}{p}$.

(i) Show $n \mid (p-1)$.

(ii) If $n = 2k$, prove $10^k \equiv -1 \pmod{p}$ and the two half-blocks of the repetend sum to $10^k - 1$.

Hint: Factor $10^{2k} - 1 = (10^k - 1)(10^k + 1)$ and use minimality of n.

Solution 4.25

From order arguments, n is the multiplicative order of 10 mod p, so $n \mid (p-1)$. If $n = 2k$, then $p \mid (10^{2k} - 1)$. Since $2k$ is minimal, $p \nmid (10^k - 1)$, hence $p \mid (10^k + 1)$, so $10^k \equiv -1 \pmod{p}$. Writing repetend as two k-digit blocks A, B gives Midy's theorem:

$$A + B = 10^k - 1.$$

Problem 4.26: Sissa and arithmetico-geometric sums

(i) Evaluate $S_{64} = 1 + 2 + 4 + \cdots + 2^{63}$.

(ii) Let

$$T = \sum_{n=1}^{64} n 2^{n-1}.$$

Use shift-and-subtract to find a closed form.

(iii) Evaluate

$$\sum_{n=1}^{\infty} \frac{n}{2^n}.$$

Hint: Use $T - 2T$ for the finite AGP, and similarly $S - \frac{1}{2}S$ for the infinite one.

Solution 4.26

$$S_{64} = 2^{64} - 1.$$

Also

$$T - 2T = \sum_{n=1}^{64} 2^{n-1} - 64 \cdot 2^{64} = (2^{64} - 1) - 64 \cdot 2^{64},$$

so

$$T = 63 \cdot 2^{64} + 1.$$

 Vu Hung Nguyen · CC BY 4.0

For $S = \sum_{n=1}^{\infty} \frac{n}{2^n}$:

$$S - \frac{1}{2}S = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots = 1,$$

hence $S = 2$.

Problem 4.27: Discrete calculus and sums of powers

(i) Show $n^3 - (n-1)^3 = 3n^2 - 3n + 1$.

(ii) Sum this identity from 1 to N to derive

$$\sum_{n=1}^{N} n^2 = \frac{N(N+1)(2N+1)}{6}.$$

(iii) Evaluate $\nabla^3(n^3)$ for $\nabla f(n) = f(n) - f(n-1)$.

(iv) State the leading term of $\sum_{n=1}^{N} n^k$.

Hint: Use telescoping on the left and known formulas for $\sum n$.

Solution 4.27

Expansion gives part (i). Summing:

$$N^3 = 3\sum_{n=1}^{N} n^2 - 3\sum_{n=1}^{N} n + N,$$

which simplifies to

$$\sum_{n=1}^{N} n^2 = \frac{N(N+1)(2N+1)}{6}.$$

Also $\nabla^3(n^3) = 6$. In general,

$$\sum_{n=1}^{N} n^k = \frac{1}{k+1} N^{k+1} + (\text{lower powers of } N).$$

 Vu Hung Nguyen · CC BY 4.0

5 Conclusion

Strong sequence work in Extension 1 is not just memorizing formulas. It is about choosing the right model, controlling notation, and checking whether each step is logically justified.

The key habits are:

- define terms clearly, including first term and common difference or ratio
- check indexing before and after every sigma manipulation
- identify whether a question is about terms, partial sums, or limits
- test answers for reasonableness by checking growth and sign behavior

With steady practice, sequences become one of the most reliable scoring areas in Extension 1.

Contact Information:

LinkedIn: https://www.linkedin.com/in/nguyenvuhung/

GitHub: https://github.com/vuhung16au/

Repository: https://github.com/vuhung16au/math-olympiad-ml/tree/main/HSC-Sequences

 Vu Hung Nguyen · CC BY 4.0

6 Appendices

6.1 Appendix A: Formula Sheet

- Arithmetic nth term:
$$a_n = a + (n-1)d.$$
- Arithmetic sum of first n terms:
$$S_n = \frac{n}{2}(2a + (n-1)d) = \frac{n}{2}(a + l).$$
- Geometric nth term:
$$a_n = ar^{n-1}.$$
- Geometric sum of first n terms ($r \neq 1$):
$$S_n = a\frac{1-r^n}{1-r} = a\frac{r^n - 1}{r-1}.$$
- Infinite geometric sum ($|r| < 1$):
$$S_\infty = \frac{a}{1-r}.$$
- Sums of powers for the first n positive integers:
$$\sum_{k=1}^{n} k = \frac{n(n+1)}{2},$$
$$\sum_{k=1}^{n} k^2 = \frac{n(n+1)(2n+1)}{6},$$
$$\sum_{k=1}^{n} k^3 = \left(\frac{n(n+1)}{2}\right)^2.$$

Chebyshev Polynomials

The Chebyshev polynomials of the first kind, denoted $T_n(x)$, are defined by
$$T_0(x) = 1, \qquad T_1(x) = x, \qquad T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x) \quad (n \geq 1).$$
Equivalently,
$$T_n(\cos\theta) = \cos(n\theta).$$
The Chebyshev polynomials of the second kind, denoted $U_n(x)$, are defined by
$$U_0(x) = 1, \qquad U_1(x) = 2x, \qquad U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x) \quad (n \geq 1).$$
Equivalently,
$$U_n(\cos\theta) = \frac{\sin((n+1)\theta)}{\sin\theta} \quad (\sin\theta \neq 0).$$

Degree	First kind $T_n(x)$	Second kind $U_n(x)$
1	x	$2x$
2	$2x^2 - 1$	$4x^2 - 1$
3	$4x^3 - 3x$	$8x^3 - 4x$
4	$8x^4 - 8x^2 + 1$	$16x^4 - 12x^2 + 1$
5	$16x^5 - 20x^3 + 5x$	$32x^5 - 32x^3 + 6x$

 Vu Hung Nguyen · CC BY 4.0

6.2 Appendix B: Common Sequence Traps

- Mixing up term formulas and sum formulas: a_n and S_n answer different questions.
- Losing track of index starts: check whether the sequence starts at $n = 0$ or $n = 1$.
- Using $S_\infty = \frac{a}{1-r}$ when $|r| \geq 1$: this formula only applies for $|r| < 1$.
- Sign errors in GP sums: keep the exact formula form and substitute carefully.
- Assuming recursion converges without proof: identify monotonicity or contraction before claiming limits.
- Treating recurring decimals as finite truncations: they are infinite geometric expansions.

6.3 Appendix C: Rigorous Definition of the Limit of a Sequence

Definition 1 (ϵ-N Definition of a Limit)**.** Let (a_n) be a sequence of real numbers. We say that (a_n) **converges to a limit** L, written

$$\lim_{n\to\infty} a_n = L,$$

if for every real number $\epsilon > 0$ there exists a positive integer N such that

$$n > N \implies |a_n - L| < \epsilon.$$

In formal logical notation:

$$\forall\, \epsilon > 0,\ \exists\, N \in \mathbb{N} \text{ such that } n > N \implies |a_n - L| < \epsilon.$$

If no such L exists, the sequence is said to **diverge**.

Unpacking the definition.

- $\epsilon > 0$ **is the "margin of error."** The challenger picks any strictly positive number, no matter how small. A smaller ϵ places a tighter band around L.
- N **is the "cutoff index."** The responder must produce a concrete N that works for the chosen ϵ. Typically, smaller ϵ forces a larger N.
- **Only the *tail* matters.** The condition $n > N$ means the first N terms may behave arbitrarily; only the infinite tail of the sequence is constrained.
- $|a_n - L| < \epsilon$ **is the "closeness" condition.** Every term beyond index N must lie strictly inside the interval $(L - \epsilon,\ L + \epsilon)$.

Visual interpretation (the "ϵ-tube").

Plot the sequence on a graph with the index n on the horizontal axis and a_n on the vertical axis. Draw a horizontal dashed line at $y = L$. Choosing $\epsilon > 0$ creates a horizontal band $\big(L-\epsilon,\ L+\epsilon\big)$ of total width 2ϵ around the limit. The definition states that for every such band there is a vertical line $x = N$ such that *every point of the sequence to the right of that line lies inside the band.*

Example: Proving $\lim_{n\to\infty} \dfrac{1}{n} = 0$**.**

Proof. Let $\epsilon > 0$ be given.

Scratchwork. We require

$$\left|\frac{1}{n} - 0\right| < \epsilon.$$

Since $n \in \mathbb{N}$ is positive, this simplifies to $\dfrac{1}{n} < \epsilon$, i.e. $n > \dfrac{1}{\epsilon}$.

Choice of N. Let N be any positive integer satisfying $N > \dfrac{1}{\epsilon}$. (Such an integer exists by the Archimedean property of $\mathbb{R}$.)

Verification. Suppose $n > N$. Then

$$n > N > \frac{1}{\epsilon} \implies \frac{1}{n} < \epsilon \implies \left|\frac{1}{n} - 0\right| = \frac{1}{n} < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, the definition is satisfied and $\lim\limits_{n\to\infty} \dfrac{1}{n} = 0$. □

Concrete illustration.

For $\epsilon = 0.01$ we need $N > 100$, so we may take $N = 101$. Every term $a_n = \frac{1}{n}$ with $n \geq 102$ satisfies $\frac{1}{102} < 0.01$, placing all those terms inside the band $(-0.01, 0.01)$.

 Vu Hung Nguyen · CC BY 4.0