

An abstract fractal pattern, resembling a complex, self-similar geometric design, is centered on the page. It features a central point from which numerous lines radiate outwards, creating a star-like or flower-like appearance. The pattern is composed of many small, repeating geometric shapes, giving it a textured, almost crystalline look. The colors are primarily dark blue and black, with some lighter blue and white highlights that create a sense of depth and movement. The overall effect is one of intricate, mathematical beauty.

HSC Mathematics Extension 1

Vectors Mastery

Vu Hung Nguyen



Download PDF and resources:

[https://github.com/vuhung16au/
math-olympiad-ml](https://github.com/vuhung16au/math-olympiad-ml)



Read online at Vu's Maths Hub:

<https://vumaths.com/>

"A Vector is a quantity represented by an arrow with both direction and magnitude."

- Mr. Vector (Despicable Me)

Contents

1	Introduction	8
1.1	Project Overview	8
1.2	Target Audience	8
1.3	How to Use This Booklet	8
1.3.1	Notations	8
1.3.2	Choosing Vector Notation	10
1.4	Vectors Primer	10
1.4.1	Coordinates in 3D	10
1.4.2	Vectors in Three Dimensions	10
1.4.3	The Dot Product	10
1.4.4	Applications of the Dot Product	10
1.4.5	Vector Proofs in Geometry	11
1.4.6	Vector Equation of a Line	11
1.4.7	Vector Equations of Circles, Spheres and Planes	11
1.4.8	Planes in HSC Vectors (Short Note)	11
1.4.9	The Dot Product (Scalar Product)	12
1.4.10	Vector Equation of a Line	13
1.4.11	Spheres & Planes	14
1.4.12	Geometric Proofs	14
1.4.13	The Cross Product (Optional Enrichment, Not in the NESA Syllabus)	14
1.5	Conic Sections Reference (Enrichment)	15
2	Part 1: Problems and Solutions (Detailed)	15
2.1	Part 1 Basic Problems (Easy)	16
	P 2.1: Vector Projection Formula	16
	S 2.1	16
	P 2.2: Shortest Distance from Point to Line	17
	S 2.2	17
	P 2.3: Parallelogram Area via Dot-Product Identity	18
	S 2.3	18
	P 2.4: Cosine Difference Formula Proof	18
	S 2.4	18
	P 2.5: Perpendicular Vectors Condition	19
	S 2.5	19
2.2	Part 1 Medium Problems	20
	P 2.6: Line Tangent to Sphere	20
	S 2.6	20
	P 2.7: Perpendicular Lines and Plane Equation	21
	S 2.7	21
	P 2.8: Three Conditions on Vectors	22
	S 2.8	22
	P 2.9: Projectile Motion Vector Proof	23
	S 2.9	23
	P 2.10: Distance from Point to Line and Sphere	24
	S 2.10	24
	P 2.11: Elegant Vector Proof of Concurrent Medians	24
	S 2.11	25
2.3	Part 1 Advanced Problems (Hard)	26

P 2.12: Position Vector Ratios and Parallelogram Division	26
S 2.12	26
P 2.13: Triangle Inequality and Cauchy-Schwarz on Sphere	27
S 2.13	27
P 2.14: Tetrahedron Bimedians Equality	28
S 2.14	28
P 2.15: Regular Tetrahedron, Common Perpendicular, and Circumsphere	29
S 2.15	30
P 2.16: The Projection Paradox	31
S 2.16	32
P 2.17: Point-to-Line Distance and Sphere Intersection	33
S 2.17	33
P 2.18: 3D Helix Projection	34
S 2.18	34
P 2.19: Parametric Equation of a Helix	35
S 2.19	35
P 2.20: Curve on a Cylinder and Arc Length	36
S 2.20	37
P 2.21: Sketching 3D Parametric Curves	38
S 2.21	38
P 2.22: Spiral Curve on a Paraboloid	39
S 2.22	39

3 Part 2: Problems and Solutions (Concise + Hints) 41

3.1 Part 2 Basic Problems	41
P 3.1: Vector to Cartesian Line Equation	41
S 3.1: Sketch	41
P 3.2: Sketch 3D Helix	41
S 3.2: Sketch	41
P 3.3: Angle Between 3D Vectors	42
S 3.3: Sketch	42
P 3.4: Unit Vector in Direction	42
S 3.4: Sketch	42
P 3.5: Distance to Plane and Axis	42
S 3.5: Sketch	43
P 3.6: Unit Vector Perpendicular to Two Vectors	43
S 3.6: Sketch	43
P 3.7: Point on Line	43
S 3.7: Sketch	44
P 3.8: Equal Magnitude Perpendicular Vectors	44
S 3.8: Sketch	44
P 3.9: Direction Cosines	45
S 3.9: Sketch	45
P 3.10: Line Intersection by Components	46
S 3.10: Sketch	46
P 3.11: Multiple Choice: Cartesian Equation	46
S 3.11: Sketch	46
P 3.12: Closest Point on Line to Origin	46
S 3.12: Sketch	47
P 3.13: Unit Vector Perpendicular to Two	47

	S 3.13: Sketch	47
	P 3.14: Perpendicular Dot Product Proof	47
	S 3.14: Sketch	48
3.2	Part 2 Medium Problems	48
	P 3.15: Line Intersection in 3D	48
	S 3.15: Sketch	48
	P 3.16: Triangular Pyramid — Find Vertex Coordinates	48
	S 3.16: Sketch	49
	P 3.17: Perpendicular Intersecting Lines	49
	S 3.17: Sketch	50
	P 3.18: Tetrahedron Collinearity	51
	S 3.18: Sketch	51
	P 3.19: Varignon's Theorem - Midpoint Parallelogram	52
	S 3.19: Sketch	53
	P 3.20: Force Vector Analysis	54
	S 3.20: Sketch	54
	P 3.21: Double Angle with Vectors	55
	S 3.21: Sketch	55
	P 3.22: Vector Projection	55
	S 3.22: Sketch	55
	P 3.23: Perpendicular Vectors Condition	55
	S 3.23: Sketch	56
	P 3.24: Parallel and Perpendicular Lines	56
	S 3.24: Sketch	56
	P 3.25: Angle BCD Using Dot Product	56
	S 3.25: Sketch	57
	P 3.26: Direction Cosines — Compute and Verify	57
	S 3.26: Sketch	57
	P 3.27: Section Formula Proof	57
	S 3.27: Sketch	58
	P 3.28: Skew or Intersecting Lines	58
	S 3.28: Sketch	58
	P 3.29: Line Through Points, Intersection Check	58
	S 3.29: Sketch	59
	P 3.30: Line-Sphere Intersection Angle	59
	S 3.30: Sketch	59
	P 3.31: Linear Combination of Vectors	60
	S 3.31: Sketch	60
	P 3.32: Parallelogram Fourth Vertex	62
	S 3.32: Sketch	63
3.3	Part 2 Advanced Problems	63
	P 3.33: Triangle Inequality and Cauchy-Schwarz	63
	S 3.33: Sketch	64
	P 3.34: Bimedians of Tetrahedron	64
	S 3.34: Sketch	65
	P 3.35: Concurrent Medians of a Tetrahedron	67
	S 3.35: Sketch	67
	P 3.36: Triangle Intersection Ratios	68
	S 3.36: Sketch	68
	P 3.37: Circle Intersection of Sets	69

S 3.37: Sketch	69
P 3.38: Complex Numbers and Centroid	70
S 3.38: Sketch	70
P 3.39: Pyramid Centroid	71
S 3.39	72
P 3.40: Line-Sphere Intersection Points	72
S 3.40: Sketch	72
P 3.41: Regular Octagon Vector Sum	72
S 3.41	74
4 Conclusion	74
A Appendices	75
A.1 Appendix 1: 2D Parametric Curves	75
A.2 Appendix 2: 3D Parametric Curves	76
A.3 Appendix 3: Projections of a 3D Parametric Curve	77
A.4 Appendix 4: Polar Form of Two-Dimensional Vectors	80
A.5 Appendix 5: The Cross Product of Two Vectors	82
A.6 Appendix 6: Equations of Planes in Three Dimensions	86
A.7 Appendix 7: Linear Dependence and Independence	88
A.8 Appendix 8: Vector-Valued Motion in Three Dimensions	90

1 Introduction

1.1 Project Overview

This booklet compiles high quality vector problems curated specifically for the HSC Mathematics Extension 2 syllabus. The collection covers core 3D vector topics including geometric proofs, line equations, dot products, spheres, planes, projections, and distance calculations. Each problem is selected to challenge students while building toward exam readiness.

1.2 Target Audience

These problems are designed for Extension 2 students who want to master vectors through systematic practice. Part 1 provides detailed solutions showing complete algebraic steps, geometric intuition, and key takeaways. Part 2 offers hints and concise solutions to encourage independent problem-solving.

1.3 How to Use This Booklet

- Read the vectors primer below to review fundamental concepts and notation.
- Attempt problems in Part 1 independently; study the detailed solutions to understand model approaches.
- Use the upside-down hints in Part 2 only after making a genuine attempt.
- Focus on understanding geometric meaning alongside algebraic manipulation.
- Revisit challenging problems after a few days to reinforce techniques.

1.3.1 Notations

Throughout this booklet, we use the following standard mathematical notations:

Syllabus Note. In the current NESA Mathematics Extension 1 and Extension 2 syllabuses, the cross product is *not* prescribed content. When it appears later in this booklet, it is clearly marked as optional enrichment only. It can still be useful background knowledge for later university study in areas such as mathematics, physics, and engineering.

Vectors:

- $\mathbf{v}, \mathbf{a}, \mathbf{b}, \mathbf{u}$ — vectors in bold font
- $\overrightarrow{AB}$ — directed line segment from point A to point B
- $\vec{AB}$ — alternative notation for directed line segment
- $\mathbf{i}, \mathbf{j}, \mathbf{k}$ — standard unit vectors along x, y, z axes
- $|\mathbf{v}|$ — magnitude (length) of vector $\mathbf{v}$
- $\hat{\mathbf{u}}$ — unit vector in the direction of $\mathbf{u}$
- $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$ — column vector notation
- (x, y, z) — coordinate/component notation

Vector Operations:

- $\mathbf{a} \cdot \mathbf{b}$ — dot product (scalar product) of vectors $\mathbf{a}$ and $\mathbf{b}$
- $\mathbf{a} \perp \mathbf{b}$ — vectors $\mathbf{a}$ and $\mathbf{b}$ are perpendicular
- $\mathbf{a} \parallel \mathbf{b}$ — vectors $\mathbf{a}$ and $\mathbf{b}$ are parallel
- $\text{proj}_{\mathbf{b}} \mathbf{a}$ — projection of vector $\mathbf{a}$ onto vector $\mathbf{b}$

Lines, Planes, and Spheres:

- $\mathbf{r}$ — position vector of a general point
- $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$ — parametric equation of a line through $\mathbf{a}$ with direction $\mathbf{b}$
- λ, μ, t — scalar parameters
- $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$ — vector equation of a plane through $\mathbf{a}$ with normal $\mathbf{n}$
- $|\mathbf{r} - \mathbf{c}| = R$ — vector equation of a sphere with center $\mathbf{c}$ and radius R
- $ax + by + cz = d$ — Cartesian equation of a plane
- $(x - h)^2 + (y - k)^2 + (z - \ell)^2 = R^2$ — Cartesian equation of a sphere

Angles and Geometry:

- θ — angle between vectors or geometric objects
- $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|}$ — formula for angle between vectors
- $\angle ABC$ — angle at vertex B formed by rays BA and BC
- $\triangle ABC$ — triangle with vertices A, B, C
- $ABCD$ — quadrilateral with vertices listed in order

Other Symbols:

- $\mathbb{R}$ — the set of real numbers
- $\in$ — belongs to (element of)
- $\Leftrightarrow$ — if and only if (equivalence)
- $\Rightarrow$ — implies
- $\equiv$ — identically equal to
- $\approx$ — approximately equal to

1.3.2 Choosing Vector Notation

The NSW Extension 1 and Extension 2 syllabuses accept several equivalent notations. Read all forms fluently; when writing, match the medium (handwritten vs. typed) and follow the notation used in each question.

Notation	Example	When to use	How to write
Boldface	a, i, j, k	Typed exams, textbooks, formal print	<code>\mathbf{a}</code>
Arrow	$\overrightarrow{AB}, \overrightarrow{OP}$	Handwritten work; vectors from one point to another	<code>\overrightarrow{AB}</code>
Tilde	$\underset{\sim}{a}$	Handwriting only—impractical to typeset	draw a tilde under the letter
Component	(x, y, z)	Coordinate algebra; ordered triples	plain coordinates
Column	$\begin{pmatrix} x \\ y \\ z \end{pmatrix}$	When the question uses column form	<code>\pmatrix</code> environment

This booklet uses boldface and $\overrightarrow{AB}$ throughout. In your own handwritten working, prefer arrow notation; reserve tildes for quick pen-and-paper work if your teacher uses them.

1.4 Vectors Primer

1.4.1 Coordinates in 3D

In three dimensions, every point is described by an ordered triple (x, y, z) , representing its position along the x , y , and z axes. This extension from 2D to 3D allows us to model space more realistically and solve geometric problems involving depth.

1.4.2 Vectors in Three Dimensions

A vector in 3D is an object with both magnitude and direction, written as $\mathbf{v} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ or $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$. The standard unit vectors $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ point along the x , y , and z axes, respectively, and form a basis for all vectors in space.

1.4.3 The Dot Product

The dot product (or scalar product) combines two vectors to produce a real number. It measures how much one vector extends in the direction of another and is fundamental for finding angles, projections, and testing perpendicularity in 3D geometry.

1.4.4 Applications of the Dot Product

The dot product is used to calculate angles between vectors, determine orthogonality, find projections, and solve geometric problems such as distances from points to planes or lines. It is also essential in physics for work and energy calculations.

1.4.5 Vector Proofs in Geometry

Vector methods provide elegant, coordinate-free proofs for geometric properties of shapes like triangles, parallelograms, and tetrahedrons. By expressing points and relationships with vectors, many results can be derived more simply than with traditional coordinate geometry.

1.4.6 Vector Equation of a Line

In 3D, the equation $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$ describes a line through point $\mathbf{a}$ in the direction of vector $\mathbf{b}$. This form is more flexible than $y = mx + b$ and is well-suited for problems involving intersections and distances in space.


1.4.7 Vector Equations of Circles, Spheres and Planes

Circles and spheres can be described using vector equations such as $|\mathbf{r} - \mathbf{c}| = R$, where $\mathbf{c}$ is the center and R the radius. Planes are described by $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$, where $\mathbf{n}$ is a normal vector. These forms unify many geometric objects under a common vector framework.


1.4.8 Planes in HSC Vectors (Short Note)

In HSC Extension 2, students should understand the *idea* of planes, not the full university algebra of general plane equations. Focus on the standard coordinate planes xy , xz , and yz , where one coordinate is zero (for example, $z = 0$ for the xy -plane), and simple translated planes such as $z = 5$. These are central when interpreting 3D line intersections and projections.

In proof questions, coplanarity is usually shown through vector relationships (such as linear combinations and dependence), rather than advanced techniques. The priority in this course is strong spatial reasoning and clean vector logic, not formal linear algebra machinery.



Standard unit vectors: $\mathbf{i}$ (red), $\mathbf{j}$ (blue), and $\mathbf{k}$ (green) are mutually perpendicular unit vectors that form the basis of 3D coordinate space. Any vector can be written as a combination of these three directions.



Left: A vector $\mathbf{v}$ in 3D space decomposed into components $x\mathbf{i}$, $y\mathbf{j}$, and $z\mathbf{k}$. *Right:* The magnitude formula derived using Pythagoras' theorem twice.

1.4.9 The Dot Product (Scalar Product)

The dot product is the central algebraic tool for vectors in Extension 2.

- **Definition:** $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$ and algebraically: $\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$
- **Perpendicular Vectors:** $\mathbf{a} \cdot \mathbf{b} = 0 \Leftrightarrow \mathbf{a} \perp \mathbf{b}$
- **Angle Between Vectors:** $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$
- **Scalar Projection:** $\text{proj}_{\mathbf{b}} \mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{b}|}$
- **Vector Projection:** $\text{proj}_{\mathbf{b}} \mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{b} \cdot \mathbf{b}} \mathbf{b}$

Remark (Useful Identity for Vector Proofs):

For any vectors $\mathbf{a}$ and $\mathbf{b}$, the squared length of the component of $\mathbf{a}$ perpendicular to $\mathbf{b}$ is

$$|\mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a}|^2 = |\mathbf{a}|^2 - \frac{(\mathbf{a} \cdot \mathbf{b})^2}{|\mathbf{b}|^2}.$$

Derivation: Let $\mathbf{p} = \text{proj}_{\mathbf{b}} \mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{b}|^2} \mathbf{b}$. Then

$$\begin{aligned} |\mathbf{a} - \mathbf{p}|^2 &= (\mathbf{a} - \mathbf{p}) \cdot (\mathbf{a} - \mathbf{p}) \\ &= |\mathbf{a}|^2 - 2\mathbf{a} \cdot \mathbf{p} + |\mathbf{p}|^2 \\ &= |\mathbf{a}|^2 - 2\frac{(\mathbf{a} \cdot \mathbf{b})^2}{|\mathbf{b}|^2} + \frac{(\mathbf{a} \cdot \mathbf{b})^2}{|\mathbf{b}|^2} \\ &= |\mathbf{a}|^2 - \frac{(\mathbf{a} \cdot \mathbf{b})^2}{|\mathbf{b}|^2}. \end{aligned}$$

Connection to Cauchy-Schwarz: Since the left-hand side is always non-negative,


$$|\mathbf{a}|^2 - \frac{(\mathbf{a} \cdot \mathbf{b})^2}{|\mathbf{b}|^2} \geq 0$$

which rearranges to

$$(\mathbf{a} \cdot \mathbf{b})^2 \leq |\mathbf{a}|^2 |\mathbf{b}|^2.$$

Equality holds exactly when $\mathbf{a}$ is parallel to $\mathbf{b}$, because then the perpendicular component is zero.

Applications: This identity is useful in geometric proofs involving projections, perpendicular distances, and shortest-distance arguments without needing any out-of-syllabus techniques.




Left: The dot product $\mathbf{a} \cdot \mathbf{b}$ equals $|\mathbf{a}|$ times the projection of $\mathbf{b}$ onto $\mathbf{a}$ (or vice versa). *Right:* When vectors are perpendicular ($\theta = 90^\circ$), the dot product is zero.

1.4.10 Vector Equation of a Line

Extension 2 uses vector form rather than $y = mx + b$.


- **Parametric Form:** $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$ where $\mathbf{a}$ is a point on the line, $\mathbf{b}$ is the direction vector, and λ is a parameter
- **Parallel Lines:** Direction vectors are scalar multiples
- **Skew Lines:** In 3D, lines can be non-parallel and non-intersecting



Left: A line in 3D space defined by $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$, where $\mathbf{a}$ is a fixed point and $\mathbf{b}$ is the direction vector. *Right:* Different values of λ generate different points along the line; $\lambda = 0$ gives point $\mathbf{a}$, positive λ extends forward, negative λ extends backward.

1.4.11 Spheres & Planes

- **Sphere:** Vector form $|\mathbf{r} - \mathbf{c}| = R$ or Cartesian $(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = R^2$
- **Plane (Point-Normal):** $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$ where $\mathbf{n}$ is normal to the plane
- **Plane (Cartesian):** $ax + by + cz = d$ where (a, b, c) is the normal vector



Left: A sphere centered at $\mathbf{c}$ with radius R . Any point $\mathbf{r}$ on the sphere satisfies $|\mathbf{r} - \mathbf{c}| = R$. *Right:* A plane containing point $\mathbf{a}$ with normal vector $\mathbf{n}$. Any point $\mathbf{r}$ on the plane has $(\mathbf{r} - \mathbf{a}) \perp \mathbf{n}$, so their dot product is zero.

1.4.12 Geometric Proofs

Use pure vector logic (without coordinates) to prove theorems about parallelograms, triangles, centroids, medians, and 3D shapes like tetrahedrons.

1.4.13 The Cross Product (Optional Enrichment, Not in the NESA Syllabus)

The **cross product** $\mathbf{a} \times \mathbf{b}$ of two 3D vectors produces a new vector that is perpendicular to both $\mathbf{a}$ and $\mathbf{b}$. Its magnitude equals the area of the parallelogram they form:





$$|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta.$$

The cross product is **not** prescribed in the current NESA Mathematics Extension 1 or Extension 2 syllabus, so none of the core worked methods in this booklet depend on it. It is covered in detail in **Appendix A.5** as part of the enrichment material drawn from Victorian and Queensland Specialist Mathematics.

1.5 Conic Sections Reference (Enrichment)

Reference Table: Key Conic Sections (Enrichment & Further Study)

Note on the 2026 Syllabus Update: Conic sections have been removed from the new NSW HSC Mathematics Extension 2 syllabus. However, this section is retained as enrichment material. A strong grasp of these parametric forms is highly recommended for students intending to pursue university-level engineering, physics, or computer science. Furthermore, representing points on conics parametrically as position vectors allows instant geometric computations (such as triangle areas and tangent vectors) via the vector cross product (see Appendix A.5) and matrix determinants (see our companion booklet Matrix Practice).

Shape	Parametric Equations	Cartesian Equation	Sketch
Circle (General)	$x = r \cos t$ $y = r \sin t$ $t \in [0, 2\pi)$	$x^2 + y^2 = r^2$	
Ellipse (Horizontal)	$x = a \cos t$ $y = b \sin t$ $t \in [0, 2\pi)$ (where $a > b > 0$)	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$	
Parabola (Vertical)	$x = 2at$ $y = at^2$ $t \in \mathbb{R}$ (standard form)	$x^2 = 4ay$	
Rectangular Hyperbola	$x = \frac{ct}{t}$ $y = \frac{c}{t}$ $t \in \mathbb{R}, t \neq 0$	$y = \frac{c^2}{x}$ or $xy = c^2$	

2 Part 1: Problems and Solutions (Detailed)

Part 1 contains 18 carefully selected problems covering all 10 topic areas. Each problem includes complete solutions with all algebraic steps, geometric diagrams where helpful, and key takeaways. Solutions use compact notation to fit on one A4 page while maintaining clarity.

2.1 Part 1 Basic Problems (Easy)

Problem 2.1: Vector Projection Formula

The vector $\mathbf{a}$ is $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and the vector $\mathbf{b}$ is $\begin{pmatrix} 2 \\ 0 \\ -4 \end{pmatrix}$.

- (i) Find $\frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{b} \cdot \mathbf{b}} \mathbf{b}$.
- (ii) Show that $\mathbf{a} - \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{b} \cdot \mathbf{b}} \mathbf{b}$ is perpendicular to $\mathbf{b}$.

Solution 2.1

Given: $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 2 \\ 0 \\ -4 \end{pmatrix}$

- (i) Calculate the dot products:

$$\mathbf{a} \cdot \mathbf{b} = (1)(2) + (2)(0) + (3)(-4) = 2 - 12 = -10$$

$$\mathbf{b} \cdot \mathbf{b} = 4 + 0 + 16 = 20$$

Thus:

$$\frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{b} \cdot \mathbf{b}} \mathbf{b} = \frac{-10}{20} \begin{pmatrix} 2 \\ 0 \\ -4 \end{pmatrix} = -\frac{1}{2} \begin{pmatrix} 2 \\ 0 \\ -4 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ 2 \end{pmatrix}$$

- (ii) Let $\mathbf{v} = \mathbf{a} - \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{b} \cdot \mathbf{b}} \mathbf{b}$. Then:

$$\mathbf{v} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} - \begin{pmatrix} -1 \\ 0 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$$

Check perpendicularity: $\mathbf{v} \cdot \mathbf{b} = (2)(2) + (2)(0) + (1)(-4) = 4 - 4 = 0 \quad \therefore$ perpendicular.

Takeaways 2.1

The expression $\text{proj}_{\mathbf{b}} \mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{b} \cdot \mathbf{b}} \mathbf{b}$ is the **vector projection** of $\mathbf{a}$ onto $\mathbf{b}$. The remainder $\mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a}$ is the component perpendicular to $\mathbf{b}$. This orthogonal decomposition is fundamental: any vector can be split into parallel and perpendicular components relative to another vector. Always verify perpendicularity by checking that the dot product equals zero.

Problem 2.2: Shortest Distance from Point to Line

$\mathbf{r}_1$ and $\mathbf{r}_2$ are two lines with vector equations:

$$\mathbf{r}_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \text{ and } \mathbf{r}_2 = \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}, \lambda, \mu \in \mathbb{R}$$

- (i) Show that these two lines intersect.
- (ii) Find the angle between the lines.
- (iii) Find the shortest distance from point $P(1, 2, 0)$ to line $\mathbf{r}_1$.

Solution 2.2

(i) Equate components: $1 = 2 + \mu$ gives $\mu = -1$. Then $\lambda = 3\mu = -3$. Check z : $1 + \lambda = 1 - 3 = -2$ and $2\mu = -2$. Consistent. Lines intersect at $(1, -3, -2)$.

(ii) Direction vectors: $\mathbf{d}_1 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$, $\mathbf{d}_2 = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}$

$$\mathbf{d}_1 \cdot \mathbf{d}_2 = 0 + 3 + 2 = 5$$

$$|\mathbf{d}_1| = \sqrt{2}, \quad |\mathbf{d}_2| = \sqrt{1 + 9 + 4} = \sqrt{14}$$

$$\cos \theta = \frac{5}{\sqrt{2}\sqrt{14}} = \frac{5}{2\sqrt{7}} \implies \theta \approx 19.1^\circ$$

(iii) Let $A(1, 0, 1)$ be on line $\mathbf{r}_1$, direction $\mathbf{d} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$, and

$$\vec{AP} = \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}.$$

Project $\vec{AP}$ onto $\mathbf{d}$:

$$\text{proj}_{\mathbf{d}} \vec{AP} = \frac{\vec{AP} \cdot \mathbf{d}}{\mathbf{d} \cdot \mathbf{d}} \mathbf{d} = \frac{1}{2} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}.$$

So the perpendicular component is

$$\vec{AP}_{\perp} = \vec{AP} - \text{proj}_{\mathbf{d}} \vec{AP} = \begin{pmatrix} 0 \\ \frac{3}{2} \\ -\frac{3}{2} \end{pmatrix}.$$

Hence

$$D = |\vec{AP}_{\perp}| = \sqrt{\left(\frac{3}{2}\right)^2 + \left(-\frac{3}{2}\right)^2} = \sqrt{\frac{9}{2}} = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2} \text{ units.}$$

Takeaways 2.2

To find line intersection in 3D: equate components and solve the system (3 equations, 2 unknowns). Consistent solution means intersection; inconsistent means skew lines. For point-to-line distance, use projection: subtract the component of $\vec{AP}$ parallel to the direction vector, then take the magnitude of the perpendicular remainder. This approach is fully syllabus-aligned and reinforces the idea of resolving a vector into parallel and perpendicular parts.

Problem 2.3: Parallelogram Area via Dot-Product Identity

The adjacent sides of a parallelogram are represented by vectors $\mathbf{a} = 4\mathbf{i} + 3\mathbf{j} - \mathbf{k}$ and $\mathbf{b} = 2\mathbf{i} - \mathbf{j} + 3\mathbf{k}$. Show that the area of the parallelogram is $6\sqrt{10}$ square units.

Solution 2.3

If θ is the angle between $\mathbf{a}$ and $\mathbf{b}$, then the area of the parallelogram is

$$|\mathbf{a}| |\mathbf{b}| \sin \theta.$$

Using $\sin^2 \theta = 1 - \cos^2 \theta$ and

$$\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|},$$

we get

$$\text{Area}^2 = |\mathbf{a}|^2 |\mathbf{b}|^2 - (\mathbf{a} \cdot \mathbf{b})^2.$$

Now compute:

$$|\mathbf{a}|^2 = 4^2 + 3^2 + (-1)^2 = 26,$$

$$|\mathbf{b}|^2 = 2^2 + (-1)^2 + 3^2 = 14,$$

$$\mathbf{a} \cdot \mathbf{b} = (4)(2) + (3)(-1) + (-1)(3) = 2.$$

Therefore

$$\text{Area}^2 = (26)(14) - 2^2 = 360,$$

so

$$\text{Area} = \sqrt{360} = 6\sqrt{10}.$$

Takeaways 2.3

For vectors $\mathbf{a}$ and $\mathbf{b}$, the area formula

$$\text{Area}^2 = |\mathbf{a}|^2 |\mathbf{b}|^2 - (\mathbf{a} \cdot \mathbf{b})^2$$

comes directly from the dot-product definition of angle. It is a convenient in-syllabus way to find 3D areas without introducing any extra vector operation.

Problem 2.4: Cosine Difference Formula Proof

Let $\mathbf{a}$ and $\mathbf{b}$ be 2-dimensional unit vectors, inclined to the x -axis at angles α and β respectively. You may assume $\mathbf{a} = \cos \alpha \mathbf{i} + \sin \alpha \mathbf{j}$ and $\mathbf{b} = \cos \beta \mathbf{i} + \sin \beta \mathbf{j}$. Prove that $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$.

Solution 2.4

Method 1: Geometric dot product

The angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \alpha - \beta$. Since both are unit vectors:

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta = (1)(1) \cos(\alpha - \beta) = \cos(\alpha - \beta)$$

Method 2: Algebraic dot product

$$\mathbf{a} \cdot \mathbf{b} = (\cos \alpha)(\cos \beta) + (\sin \alpha)(\sin \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

Equating the two expressions: $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$ $\square$

Takeaways 2.4

This elegant proof demonstrates the power of vectors in deriving trigonometric identities. The key insight: the dot product has both a **geometric definition** ($|\mathbf{a}||\mathbf{b}| \cos \theta$) and an **algebraic definition** (sum of component products). Equating these yields the cosine difference formula. Similar approaches can derive $\cos(\alpha + \beta)$, $\sin(\alpha \pm \beta)$, and other compound angle formulas. This vector method is often cleaner than traditional geometric proofs.

Problem 2.5: Perpendicular Vectors Condition

Consider two vectors $\mathbf{u} = -2\mathbf{i} - \mathbf{j} + 3\mathbf{k}$ and $\mathbf{v} = p\mathbf{i} + \mathbf{j} + 2\mathbf{k}$. For what values of p are $\mathbf{u} - \mathbf{v}$ and $\mathbf{u} + \mathbf{v}$ perpendicular?

Solution 2.5

First compute $\mathbf{u} - \mathbf{v}$ and $\mathbf{u} + \mathbf{v}$:

$$\mathbf{u} - \mathbf{v} = (-2 - p)\mathbf{i} + (-1 - 1)\mathbf{j} + (3 - 2)\mathbf{k} = (-2 - p)\mathbf{i} - 2\mathbf{j} + \mathbf{k}$$

$$\mathbf{u} + \mathbf{v} = (-2 + p)\mathbf{i} + (-1 + 1)\mathbf{j} + (3 + 2)\mathbf{k} = (-2 + p)\mathbf{i} + 0\mathbf{j} + 5\mathbf{k}$$

For perpendicularity, their dot product must equal zero:

$$\begin{aligned} (\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v}) &= 0 \\ (-2 - p)(-2 + p) + (-2)(0) + (1)(5) &= 0 \\ 4 - p^2 + 5 &= 0 \\ 9 - p^2 &= 0 \\ p^2 &= 9 \\ p &= \pm 3 \end{aligned}$$

Takeaways 2.5

Perpendicularity problems always reduce to setting the dot product equal to zero. This problem illustrates the algebraic identity: $(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v}) = |\mathbf{u}|^2 - |\mathbf{v}|^2$ (vector form of difference of squares). The general principle: two vectors are perpendicular if and only if the sum of products of corresponding components equals zero. When the problem involves a parameter, solve the resulting equation carefully—don't forget both positive and negative solutions for squared terms.

2.2 Part 1 Medium Problems

Problem 2.6: Line Tangent to Sphere

A sphere has centre at $(3, -3, 4)$ and radius 6 units.

A line has equation $\mathbf{r} = \begin{pmatrix} 1 \\ 5 \\ 4 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix}$.

- Write down the vector equation of the sphere.
- Determine whether the line is a tangent to the sphere, clearly justifying your conclusion.

Solution 2.6

(i) Let centre $\mathbf{c} = \begin{pmatrix} 3 \\ -3 \\ 4 \end{pmatrix}$ and radius $R = 6$. Vector equation:

$$\left| \mathbf{r} - \begin{pmatrix} 3 \\ -3 \\ 4 \end{pmatrix} \right| = 6$$

(ii) Substitute line into sphere. Cartesian form: $(x - 3)^2 + (y + 3)^2 + (z - 4)^2 = 36$. Parametric line: $x = 1 + 2\lambda$, $y = 5 - \lambda$, $z = 4 - \lambda$.

$$(1 + 2\lambda - 3)^2 + (5 - \lambda + 3)^2 + (4 - \lambda - 4)^2 = 36$$

$$(2\lambda - 2)^2 + (8 - \lambda)^2 + (-\lambda)^2 = 36$$

$$4\lambda^2 - 8\lambda + 4 + 64 - 16\lambda + \lambda^2 + \lambda^2 = 36$$

$$6\lambda^2 - 24\lambda + 68 = 36$$

$$6\lambda^2 - 24\lambda + 32 = 0$$

$$3\lambda^2 - 12\lambda + 16 = 0$$

Discriminant: $\Delta = (-12)^2 - 4(3)(16) = 144 - 192 = -48 < 0$

No real solutions $\implies$ **line does NOT intersect sphere** (not a tangent).

Takeaways 2.6

For line-sphere intersection: substitute parametric line equations into sphere equation to get a quadratic in λ . The discriminant determines the nature: $\Delta > 0$ (2 intersections, secant), $\Delta = 0$ (1 intersection, tangent), $\Delta < 0$ (no intersection). A tangent touches the sphere at exactly one point, requiring distance from centre to line equals radius. Always check discriminant rather than just counting solutions.

Problem 2.7: Perpendicular Lines and Plane Equation

Consider lines $L_1 : \mathbf{r} = \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$ and $L_2 : \mathbf{r} = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$

- (i) Show that L_1 and L_2 intersect and are perpendicular, stating the point of intersection.
- (ii) Deduce that the plane containing both lines has equation $y + z = 1$.
- (iii) Find the perpendicular distance from the origin to this plane.

Solution 2.7

(i) Directions: $\mathbf{d}_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$, $\mathbf{d}_2 = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$. Perpendicular: $\mathbf{d}_1 \cdot \mathbf{d}_2 = 2 - 1 - 1 = 0 \checkmark$

Intersection: Equate $(3 + 2\lambda, 2 - \lambda, -1 + \lambda) = (-1 + \mu, 1 + \mu, -\mu)$

$$x : 3 + 2\lambda = -1 + \mu \implies \mu - 2\lambda = 4 \quad (1)$$

$$y : 2 - \lambda = 1 + \mu \implies \mu + \lambda = 1 \quad (2)$$

$$z : -1 + \lambda = -\mu \implies \mu + \lambda = 1 \quad (3)$$

(1) - (2): $-3\lambda = 3 \implies \lambda = -1$, so $\mu = 2$. Point: $(3 - 2, 2 + 1, -1 - 1) = (1, 3, -2) \checkmark$

(ii) Plane through $(1, 3, -2)$ spanned by $\mathbf{d}_1, \mathbf{d}_2$. Let a normal vector be

$$\mathbf{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}.$$

Since $\mathbf{n}$ is perpendicular to both direction vectors,

$$\mathbf{n} \cdot \mathbf{d}_1 = 0 \implies 2a - b + c = 0,$$

$$\mathbf{n} \cdot \mathbf{d}_2 = 0 \implies a + b - c = 0.$$

From the second equation, $a = -b + c$. Substituting into the first gives

$$2(-b + c) - b + c = 0 \implies -3b + 3c = 0 \implies b = c.$$

Hence $a = 0$, so we may take

$$\mathbf{n} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}.$$

The plane has form $0x + y + z = k$. Using $(1, 3, -2)$: $3 + (-2) = 1 \implies k = 1$. Thus $y + z = 1$.

(iii) Distance from origin $(0, 0, 0)$ to plane $y + z - 1 = 0$:

$$D = \frac{|0+0-1|}{\sqrt{0^2+1^2+1^2}} = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2} \text{ units}$$

Takeaways 2.7

When two lines intersect, they determine a unique plane. To find the plane's Cartesian equation: (1) verify intersection, (2) let the normal be (a, b, c) and use perpendicularity with the two direction vectors, (3) use point-normal form. For distance from point (x_0, y_0, z_0) to plane $Ax + By + Cz + D = 0$: $d = \frac{|Ax_0 + By_0 + Cz_0 + D|}{\sqrt{A^2 + B^2 + C^2}}$. Memorize this formula.

Problem 2.8: Three Conditions on Vectors

Which of the following is a true statement about the lines

$$\ell_1 = \begin{pmatrix} -1 \\ 2 \\ 5 \end{pmatrix} + \lambda \begin{pmatrix} -1 \\ 3 \\ 1 \end{pmatrix} \text{ and } \ell_2 = \begin{pmatrix} 3 \\ -10 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -3 \\ -1 \end{pmatrix}?$$

- A. ℓ_1 and ℓ_2 are the same line.
- B. ℓ_1 and ℓ_2 are not parallel and they intersect.
- C. ℓ_1 and ℓ_2 are parallel and they do not intersect.
- D. ℓ_1 and ℓ_2 are not parallel and they do not intersect.

Solution 2.8

Direction vectors: $\mathbf{d}_1 = \begin{pmatrix} -1 \\ 3 \\ 1 \end{pmatrix}$, $\mathbf{d}_2 = \begin{pmatrix} 1 \\ -3 \\ -1 \end{pmatrix}$

Step 1: Check parallelism

$$\mathbf{d}_2 = -1 \cdot \mathbf{d}_1$$

Lines are **parallel** (eliminates B and D).

Step 2: Check if coincident Test if point $P_1 = (-1, 2, 5)$ from ℓ_1 lies on ℓ_2 :

$$\begin{pmatrix} -1 \\ 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 3 \\ -10 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -3 \\ -1 \end{pmatrix}$$

Component equations:

$$\begin{aligned} x : -1 &= 3 + \mu \implies \mu = -4 \\ y : 2 &= -10 - 3\mu \implies \mu = -4 \\ z : 5 &= 1 - \mu \implies \mu = -4 \end{aligned}$$

Consistent value $\mu = -4$ for all components $\implies$ point lies on ℓ_2 .

Answer: A (same line).

Takeaways 2.8

For two lines to be identical, two conditions must hold: (1) direction vectors are scalar multiples (parallel), and (2) they share at least one common point. If parallel but don't share a point, they're distinct parallel lines. In 3D, non-parallel lines can be skew (no intersection). The systematic approach: first check parallelism via direction vectors, then test a point from one line on the other. This problem reinforces the distinction between "parallel" and "coincident."

Problem 2.9: Projectile Motion Vector Proof

A particle is projected from the origin with initial velocity u m/s at angle θ to the horizontal. The acceleration vector is $\mathbf{a} = \begin{pmatrix} 0 \\ -g \end{pmatrix}$.

- (i) Show that the position vector is $\mathbf{r}(t) = \begin{pmatrix} ut \cos \theta \\ ut \sin \theta - \frac{1}{2}gt^2 \end{pmatrix}$.
- (ii) Show that the Cartesian equation of the path is $y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta}$.
- (iii) Given $u^2 > gR$, prove there are two distinct values of θ for which the particle lands at $x = R$.

Solution 2.9

- (i) Initial velocity: $\mathbf{v}_0 = u \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$. Integrate acceleration:

$$\mathbf{v}(t) = \mathbf{v}_0 + \int \mathbf{a} dt = \begin{pmatrix} u \cos \theta \\ u \sin \theta - gt \end{pmatrix}$$

Integrate velocity (starting from origin):

$$\mathbf{r}(t) = \int \mathbf{v}(t) dt = \begin{pmatrix} ut \cos \theta \\ ut \sin \theta - \frac{1}{2}gt^2 \end{pmatrix}$$

- (ii) From $x = ut \cos \theta$, we get $t = \frac{x}{u \cos \theta}$. Substitute into y :

$$\begin{aligned} y &= u \sin \theta \cdot \frac{x}{u \cos \theta} - \frac{1}{2}g \left(\frac{x}{u \cos \theta} \right)^2 \\ &= x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta} \end{aligned}$$

- (iii) At landing, $y = 0$ and $x = R$:

$$0 = R \tan \theta - \frac{gR^2}{2u^2 \cos^2 \theta}$$

Multiply by $\cos^2 \theta$: $0 = R \sin \theta \cos \theta - \frac{gR^2}{2u^2}$

Using $\sin(2\theta) = 2 \sin \theta \cos \theta$:

$$\sin(2\theta) = \frac{gR}{u^2}$$

Since $u^2 > gR$, we have $\frac{gR}{u^2} < 1$. Thus 2θ has two solutions in $[0, 180^\circ]$: one acute, one obtuse. This gives two distinct values of θ in $[0, 90^\circ]$.

Takeaways 2.9

Projectile motion problems integrate naturally with vectors: acceleration $\rightarrow$ velocity $\rightarrow$ position. The key insight for part (iii): for a given range R , there are two launch angles (complementary angles to 45°) that achieve the same horizontal distance, provided the initial speed is sufficient. The condition $u^2 > gR$ ensures the target is within reach. The identity $\sin(2\theta) = 2 \sin \theta \cos \theta$ is crucial for converting between different forms.

Problem 2.10: Distance from Point to Line and Sphere

Consider point B with position vector $\mathbf{b}$ and line $\ell : \mathbf{a} + \lambda\mathbf{d}$, where $|\mathbf{d}| = 1$ and λ is a parameter. Let $f(\lambda)$ be the distance between a point on ℓ and point B .

- (i) Find λ_0 , the value of λ that minimises f , in terms of $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{d}$.
- (ii) Let P be the point with position vector $\mathbf{a} + \lambda_0\mathbf{d}$. Show that PB is perpendicular to the direction of ℓ .
- (iii) Hence find the shortest distance between line ℓ and the sphere of radius 1 centred at origin O , in terms of $\mathbf{d}$ and $\mathbf{a}$.

Solution 2.10

- (i) Minimize $S(\lambda) = f(\lambda)^2 = |(\mathbf{a} + \lambda\mathbf{d}) - \mathbf{b}|^2$:

$$S(\lambda) = |\mathbf{a} - \mathbf{b}|^2 + 2\lambda(\mathbf{a} - \mathbf{b}) \cdot \mathbf{d} + \lambda^2|\mathbf{d}|^2$$

$$\frac{dS}{d\lambda} = 2(\mathbf{a} - \mathbf{b}) \cdot \mathbf{d} + 2\lambda = 0$$

$$\lambda_0 = -(\mathbf{a} - \mathbf{b}) \cdot \mathbf{d} = (\mathbf{b} - \mathbf{a}) \cdot \mathbf{d}$$
 - (ii) $\vec{PB} = \mathbf{b} - (\mathbf{a} + \lambda_0\mathbf{d}) = (\mathbf{b} - \mathbf{a}) - \lambda_0\mathbf{d}$ Check: $\vec{PB} \cdot \mathbf{d} = (\mathbf{b} - \mathbf{a}) \cdot \mathbf{d} - \lambda_0(\mathbf{d} \cdot \mathbf{d}) = \lambda_0 - \lambda_0(1) = 0 \checkmark$
 - (iii) For origin (set $\mathbf{b} = \mathbf{0}$): $\lambda_{min} = -\mathbf{a} \cdot \mathbf{d}$. Point on line closest to O : $\mathbf{p} = \mathbf{a} - (\mathbf{a} \cdot \mathbf{d})\mathbf{d}$

$$|\mathbf{p}|^2 = |\mathbf{a}|^2 - 2(\mathbf{a} \cdot \mathbf{d})^2 + (\mathbf{a} \cdot \mathbf{d})^2 = |\mathbf{a}|^2 - (\mathbf{a} \cdot \mathbf{d})^2$$
- Shortest distance to sphere: $\sqrt{|\mathbf{a}|^2 - (\mathbf{a} \cdot \mathbf{d})^2} - 1$.

Takeaways 2.10

Point-to-line distance problems use calculus (minimize squared distance) or projection (subtract parallel component). The perpendicularity in part (ii) confirms the minimum: the shortest path is always perpendicular to the line. For sphere-line distance, first find line-to-centre distance, then subtract the radius. The identity $|\mathbf{a} - (\mathbf{a} \cdot \mathbf{d})\mathbf{d}|^2 = |\mathbf{a}|^2 - (\mathbf{a} \cdot \mathbf{d})^2$ (when $|\mathbf{d}| = 1$) is the key algebraic shortcut.

Problem 2.11: Elegant Vector Proof of Concurrent Medians

Let the vertices of $\triangle ABC$ have position vectors $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ relative to an origin O . Let D , E , and F be the midpoints of sides BC , AC , and AB respectively.

- (i) Show that

$$\mathbf{d} = \frac{\mathbf{b} + \mathbf{c}}{2}, \quad \mathbf{e} = \frac{\mathbf{a} + \mathbf{c}}{2}, \quad \mathbf{f} = \frac{\mathbf{a} + \mathbf{b}}{2}.$$

- (ii) Define

$$\mathbf{g} = \frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3}.$$

Prove that G lies on all three medians AD , BE , and CF .

- (iii) Hence show that the medians of a triangle are concurrent and that the common point divides each median in the ratio 2 : 1.

Solution 2.11

(i) By the midpoint formula for position vectors:

$$\mathbf{d} = \frac{\mathbf{b} + \mathbf{c}}{2}, \quad \mathbf{e} = \frac{\mathbf{a} + \mathbf{c}}{2}, \quad \mathbf{f} = \frac{\mathbf{a} + \mathbf{b}}{2}.$$

(ii) Start from

$$\mathbf{g} = \frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3} = \frac{\mathbf{a} + (\mathbf{b} + \mathbf{c})}{3}.$$

Since $\mathbf{b} + \mathbf{c} = 2\mathbf{d}$, we get

$$\mathbf{g} = \frac{\mathbf{a} + 2\mathbf{d}}{3} = \frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{d}.$$

So G lies on line segment AD . By the section formula, $AG : GD = 2 : 1$.

Now use symmetry of $\mathbf{g}$:

$$\mathbf{g} = \frac{\mathbf{b} + (\mathbf{a} + \mathbf{c})}{3} = \frac{\mathbf{b} + 2\mathbf{e}}{3} = \frac{1}{3}\mathbf{b} + \frac{2}{3}\mathbf{e},$$

so G lies on BE and divides it in ratio $2 : 1$.

Similarly,

$$\mathbf{g} = \frac{\mathbf{c} + (\mathbf{a} + \mathbf{b})}{3} = \frac{\mathbf{c} + 2\mathbf{f}}{3} = \frac{1}{3}\mathbf{c} + \frac{2}{3}\mathbf{f},$$

so G lies on CF and divides it in ratio $2 : 1$.

(iii) The same point G lies on all three medians AD, BE, CF , so the medians are concurrent. The coefficient form in each case gives the same internal division ratio:

$$AG : GD = BG : GE = CG : GF = 2 : 1.$$

Takeaways 2.11

This proof is a model vector-geometry argument: define a symmetric candidate point $\mathbf{g} = \frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3}$, then rewrite it in section-form along each median. Symmetry avoids repeating heavy calculations and guarantees all three medians meet at one point (the centroid). The coefficient pattern $\frac{1}{3}$ and $\frac{2}{3}$ immediately encodes the $2 : 1$ division property.

2.3 Part 1 Advanced Problems (Hard)

Problem 2.12: Position Vector Ratios and Parallelogram Division

Point C divides interval AB so that $\frac{CB}{AC} = \frac{m}{n}$. Position vectors of A and B are $\mathbf{a}, \mathbf{b}$.


(i) Show that $\vec{AC} = \frac{n}{m+n}(\mathbf{b} - \mathbf{a})$.

(ii) Prove that $\vec{OC} = \frac{m}{m+n}\mathbf{a} + \frac{n}{m+n}\mathbf{b}$.

Let $OPQR$ be a parallelogram with $\vec{OP} = \mathbf{p}$, $\vec{OR} = \mathbf{r}$. S is the midpoint of QR , T is the intersection of PR and OS .

(iii) Show that $\vec{OT} = \frac{2}{3}\mathbf{r} + \frac{1}{3}\mathbf{p}$.

(iv) Prove that T divides PR in the ratio $2 : 1$.



Parallelogram $OPQR$ with $\vec{OP} = \mathbf{p}$, $\vec{OR} = \mathbf{r}$. Point S is the midpoint of QR , and T is where diagonal PR intersects OS , dividing PR in ratio $2 : 1$.

Solution 2.12

(i) Given $\frac{CB}{AC} = \frac{m}{n}$, let $\vec{AC} = k(\mathbf{b} - \mathbf{a})$ for some k . Then $C = A + k(\mathbf{b} - \mathbf{a})$. Since $\vec{CB} = \mathbf{b} - \mathbf{c} = (1 - k)(\mathbf{b} - \mathbf{a})$ and $\vec{AC} = k(\mathbf{b} - \mathbf{a})$:

$$\frac{CB}{AC} = \frac{1-k}{k} = \frac{m}{n} \implies n(1-k) = mk \implies n = k(m+n) \implies k = \frac{n}{m+n}$$

Thus $\vec{AC} = \frac{n}{m+n}(\mathbf{b} - \mathbf{a})$.

(ii) $\vec{OC} = \vec{OA} + \vec{AC} = \mathbf{a} + \frac{n}{m+n}(\mathbf{b} - \mathbf{a}) = \mathbf{a} \left(1 - \frac{n}{m+n}\right) + \frac{n}{m+n}\mathbf{b} = \frac{m}{m+n}\mathbf{a} + \frac{n}{m+n}\mathbf{b}$

(iii) In parallelogram $OPQR$: $\vec{OQ} = \mathbf{p} + \mathbf{r}$. S is midpoint of QR :

$$\vec{OS} = \frac{1}{2}(\vec{OQ} + \vec{OR}) = \frac{1}{2}(\mathbf{p} + \mathbf{r} + \mathbf{r}) = \frac{1}{2}\mathbf{p} + \mathbf{r}$$

T lies on OS : $\vec{OT} = s(\frac{1}{2}\mathbf{p} + \mathbf{r})$ for some s . T also lies on PR . Since $\vec{OP} = \mathbf{p}$ and $\vec{OR} = \mathbf{r}$:

$$\vec{OT} = (1-t)\mathbf{p} + t\mathbf{r}$$

for some $t \in [0, 1]$. Equating: $s \cdot \frac{1}{2}\mathbf{p} + s\mathbf{r} = (1-t)\mathbf{p} + t\mathbf{r}$

Since $\mathbf{p}, \mathbf{r}$ are independent: $\frac{s}{2} = 1-t$ and $s = t$. From second: $s = t$. Substitute into first: $\frac{s}{2} = 1-s \implies \frac{3s}{2} = 1 \implies s = \frac{2}{3}$. Thus: $\vec{OT} = \frac{2}{3}(\frac{1}{2}\mathbf{p} + \mathbf{r}) = \frac{1}{3}\mathbf{p} + \frac{2}{3}\mathbf{r}$ (or equivalently $\frac{2}{3}\mathbf{r} + \frac{1}{3}\mathbf{p}$).

(iv) From part (iii), $t = \frac{2}{3}$ means T divides PR such that $\vec{PT} = \frac{2}{3}\vec{PR}$, so $PT : TR = 2 : 1$.

Takeaways 2.12

The **section formula** $\vec{OC} = \frac{m\mathbf{a}+n\mathbf{b}}{m+n}$ (when C divides AB in ratio $n : m$ from A) is fundamental for position vectors. For geometric proofs: (1) express target point in terms of base vectors using two different paths, (2) equate and solve using **linear independence** (two non-zero vectors are linearly independent when neither is a scalar multiple of the other—see Appendix A.7 for a formal treatment). Parts (iii)-(iv) show how medians and diagonals in parallelograms create consistent ratios—here T is the centroid-like point dividing both segments in ratio $2 : 1$.

Problem 2.13: Triangle Inequality and Cauchy-Schwarz on Sphere

- (i) Point $P(x, y, z)$ lies on the unit sphere centred at origin O . Using the triangle inequality, show that $|x| + |y| + |z| \geq 1$.
- (ii) Given vectors $\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$, show that $|a_1b_1 + a_2b_2 + a_3b_3| \leq \sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}$.
- (iii) As in part (i), point $P(x, y, z)$ lies on the unit sphere. Using part (ii), show that $|x| + |y| + |z| \leq \sqrt{3}$.

Solution 2.13

(i) Position vector: $\vec{OP} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ with $|\vec{OP}| = 1$.

Triangle inequality: $|\mathbf{u} + \mathbf{v} + \mathbf{w}| \leq |\mathbf{u}| + |\mathbf{v}| + |\mathbf{w}|$

$$1 = |x\mathbf{i} + y\mathbf{j} + z\mathbf{k}| \leq |x\mathbf{i}| + |y\mathbf{j}| + |z\mathbf{k}| = |x| + |y| + |z|$$

(ii) Cauchy-Schwarz from dot product: $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}|\cos\theta$ Taking absolute value: $|\mathbf{a} \cdot \mathbf{b}| = |\mathbf{a}||\mathbf{b}|\cos\theta| \leq |\mathbf{a}||\mathbf{b}|$ (since $|\cos\theta| \leq 1$). In components: $|a_1b_1 + a_2b_2 + a_3b_3| \leq \sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}$

(iii) Let $\mathbf{a} = \begin{pmatrix} |x| \\ |y| \\ |z| \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$. Apply Cauchy-Schwarz:

$$|x| + |y| + |z| = |x|(1) + |y|(1) + |z|(1) \leq \sqrt{x^2 + y^2 + z^2} \sqrt{1^2 + 1^2 + 1^2}$$

Since P is on unit sphere: $x^2 + y^2 + z^2 = 1$. Thus:

$$|x| + |y| + |z| \leq \sqrt{1} \cdot \sqrt{3} = \sqrt{3}$$

Takeaways 2.13


This problem demonstrates two fundamental inequalities. The **triangle inequality** $|\mathbf{u} + \mathbf{v}| \leq |\mathbf{u}| + |\mathbf{v}|$ gives lower bounds. The **Cauchy-Schwarz inequality** $|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}||\mathbf{b}|$ gives upper bounds. Both arise from the dot product definition. Part (iii) uses a clever choice of vectors to convert a sum constraint into a dot product. Combined: $1 \leq |x| + |y| + |z| \leq \sqrt{3}$ for points on the unit sphere—tight bounds achieved at corners of inscribed cube.

Problem 2.14: Tetrahedron Bimedians Equality

On triangular pyramid $ABCD$, L, M, N, P, Q, R are midpoints of edges AB, AC, AD, CD, BD, BC respectively. Let $\mathbf{b} = \vec{AB}$, $\mathbf{c} = \vec{AC}$, $\mathbf{d} = \vec{AD}$.

(i) Show that $\vec{LP} = \frac{1}{2}(-\mathbf{b} + \mathbf{c} + \mathbf{d})$.

(ii) Given that $\vec{MQ} = \frac{1}{2}(\mathbf{b} - \mathbf{c} + \mathbf{d})$ and $\vec{NR} = \frac{1}{2}(\mathbf{b} + \mathbf{c} - \mathbf{d})$, prove that $|\vec{AB}|^2 + |\vec{AC}|^2 + |\vec{AD}|^2 + |\vec{BC}|^2 + |\vec{BD}|^2 + |\vec{CD}|^2 = 4(|\vec{LP}|^2 + |\vec{MQ}|^2 + |\vec{NR}|^2)$.



Tetrahedron $ABCD$ with midpoints: L (of AB), M (of AC), N (of AD), P (of CD), Q (of BD), R (of BC). The three bimedians LP, MQ, NR connect midpoints of opposite edges.

Solution 2.14

(i) L is midpoint of AB : $\vec{AL} = \frac{1}{2}\mathbf{b}$. P is midpoint of CD : $\vec{AP} = \frac{1}{2}(\vec{AC} + \vec{AD}) = \frac{1}{2}(\mathbf{c} + \mathbf{d})$.

$$\vec{LP} = \vec{AP} - \vec{AL} = \frac{1}{2}(\mathbf{c} + \mathbf{d}) - \frac{1}{2}\mathbf{b} = \frac{1}{2}(-\mathbf{b} + \mathbf{c} + \mathbf{d})$$

(ii) **LHS:** Sum of squared edge lengths.

$$\begin{aligned} \text{LHS} &= |\mathbf{b}|^2 + |\mathbf{c}|^2 + |\mathbf{d}|^2 + |\mathbf{c} - \mathbf{b}|^2 + |\mathbf{d} - \mathbf{b}|^2 + |\mathbf{d} - \mathbf{c}|^2 \\ &= b^2 + c^2 + d^2 + (c^2 + b^2 - 2\mathbf{b} \cdot \mathbf{c}) + (d^2 + b^2 - 2\mathbf{b} \cdot \mathbf{d}) \\ &\quad + (d^2 + c^2 - 2\mathbf{c} \cdot \mathbf{d}) \\ &= 3(b^2 + c^2 + d^2) - 2(\mathbf{b} \cdot \mathbf{c} + \mathbf{b} \cdot \mathbf{d} + \mathbf{c} \cdot \mathbf{d}) \end{aligned}$$

RHS: Calculate $4|\vec{LP}|^2$:

$$\begin{aligned} |\vec{LP}|^2 &= \frac{1}{4}(-\mathbf{b} + \mathbf{c} + \mathbf{d}) \cdot (-\mathbf{b} + \mathbf{c} + \mathbf{d}) \\ &= \frac{1}{4}(b^2 + c^2 + d^2 - 2\mathbf{b} \cdot \mathbf{c} - 2\mathbf{b} \cdot \mathbf{d} + 2\mathbf{c} \cdot \mathbf{d}) \\ 4|\vec{LP}|^2 &= b^2 + c^2 + d^2 - 2\mathbf{b} \cdot \mathbf{c} - 2\mathbf{b} \cdot \mathbf{d} + 2\mathbf{c} \cdot \mathbf{d} \end{aligned}$$

Similarly:

$$\begin{aligned} 4|\vec{MQ}|^2 &= b^2 + c^2 + d^2 - 2\mathbf{b} \cdot \mathbf{c} + 2\mathbf{b} \cdot \mathbf{d} - 2\mathbf{c} \cdot \mathbf{d} \\ 4|\vec{NR}|^2 &= b^2 + c^2 + d^2 + 2\mathbf{b} \cdot \mathbf{c} - 2\mathbf{b} \cdot \mathbf{d} - 2\mathbf{c} \cdot \mathbf{d} \end{aligned}$$

Sum:

$$\begin{aligned} \text{RHS} &= 4(|\vec{LP}|^2 + |\vec{MQ}|^2 + |\vec{NR}|^2) \\ &= 3(b^2 + c^2 + d^2) + (-2 - 2 + 2)\mathbf{b} \cdot \mathbf{c} + (-2 + 2 - 2)\mathbf{b} \cdot \mathbf{d} \\ &\quad + (2 - 2 - 2)\mathbf{c} \cdot \mathbf{d} \\ &= 3(b^2 + c^2 + d^2) - 2(\mathbf{b} \cdot \mathbf{c} + \mathbf{b} \cdot \mathbf{d} + \mathbf{c} \cdot \mathbf{d}) = \text{LHS} \quad \square \end{aligned}$$

Takeaways 2.14

Bimedians (segments joining midpoints of opposite edges) in a tetrahedron have remarkable properties. This identity relates all six edge lengths to the three bimedian lengths—a 3D analogue of the parallelogram law. The result is known as **Commandino's theorem** (16th century): the sum of squares of all edges equals four times the sum of squares of the three bimedians.

The proof strategy: express everything in terms of base vectors $\mathbf{b}, \mathbf{c}, \mathbf{d}$, expand dot products carefully, and verify algebraic cancellation. Note the symmetry in coefficients $(+2, -2, -2)$ cycling through the three cross terms.

Special case: When $D = A$ (degenerate tetrahedron), $ABCD$ collapses to triangle ABC . Then $\mathbf{d} = \mathbf{0}$, and the identity reduces to: $|AB|^2 + |AC|^2 + |BC|^2 = 4(|LP|^2 + |MQ|^2 + |NR|^2)$, which becomes the triangle median formula: the sum of squares of the sides equals $\frac{4}{3}$ times the sum of squares of the medians. Such identities are useful in crystallography and structural analysis.

Problem 2.15: Regular Tetrahedron, Common Perpendicular, and Circumsphere

The points $A(1, 0, 0)$, $B(0, 1, 0)$, and $C(0, 0, 1)$ form an equilateral triangle in 3D space. A fourth point $D(x, y, z)$ is placed so that $ABCD$ forms a regular tetrahedron.

- (i) By equating the magnitudes of $\vec{DA}$, $\vec{DB}$, and $\vec{DC}$, prove that the position vector of D lies on the line

$$\mathbf{r} = \lambda(\mathbf{i} + \mathbf{j} + \mathbf{k}).$$

- (ii) Hence find the two possible coordinates of D .
- (iii) Let D_1 be the point from part (ii) lying in the first octant. Let M be the midpoint of AB , and N the midpoint of the opposite edge CD_1 . By finding $\vec{AB}$, $\vec{CD_1}$, and $\vec{MN}$, show that $\vec{MN}$ is the common perpendicular to the skew lines AB and CD_1 .
- (iv) A sphere passes through A, B, C , and D_1 . Given that the centre of the circumscribed sphere of a regular tetrahedron is the centroid of its four vertices, find the vector equation of the sphere in the form $|\mathbf{r} - \mathbf{c}| = R$.

Solution 2.15

(i)

$$\vec{DA} = (1-x, -y, -z), \quad \vec{DB} = (-x, 1-y, -z), \quad \vec{DC} = (-x, -y, 1-z).$$

$$|\vec{DA}|^2 = (1-x)^2 + y^2 + z^2, \quad |\vec{DB}|^2 = x^2 + (1-y)^2 + z^2, \quad |\vec{DC}|^2 = x^2 + y^2 + (1-z)^2.$$

Hence $x = y$ and $x = z$, so $x = y = z = \lambda$. Thus

$$\vec{OD} = (\lambda, \lambda, \lambda) = \lambda(\mathbf{i} + \mathbf{j} + \mathbf{k}),$$

so D lies on $\mathbf{r} = \lambda(\mathbf{i} + \mathbf{j} + \mathbf{k})$.

(ii) Let $D = (a, a, a)$. Then

$$|\vec{AB}|^2 = 2, \quad |\vec{DA}|^2 = 1 - 2a + 3a^2 \implies 3a^2 - 2a - 1 = 0 \implies a = 1 \text{ or } -\frac{1}{3},$$

$$\therefore D = (1, 1, 1) \text{ or } \left(-\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}\right).$$

(iii) $D_1 = (1, 1, 1)$,

$$M = (1/2, 1/2, 0), \quad N = (1/2, 1/2, 1),$$

$$\vec{MN} = (0, 0, 1), \quad \vec{AB} = (-1, 1, 0), \quad \vec{CD}_1 = (1, 1, 0).$$

$$\vec{MN} \cdot \vec{AB} = 0, \quad \vec{MN} \cdot \vec{CD}_1 = 0,$$

hence $\vec{MN}$ is the common perpendicular to AB and CD_1 .

(iv) The circumsphere centre is the centroid:

$$\mathbf{c} = \frac{1}{4}(\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{d}_1) = \frac{1}{4}(2, 2, 2) = (1/2, 1/2, 1/2).$$

$$R = |(1, 0, 0) - (1/2, 1/2, 1/2)| = |(1/2, -1/2, -1/2)| = \frac{\sqrt{3}}{2}.$$

Hence

$$|\mathbf{r} - (1/2, 1/2, 1/2)| = \frac{\sqrt{3}}{2}.$$

Takeaways 2.15

This problem blends several major Extension 2 vector skills into one structure: symmetry arguments, equal-edge conditions in 3D, skew-line geometry, and the vector equation of a sphere. Part (i) shows a classic proof technique: equate squared lengths to force coordinates into a symmetric pattern. Part (iii) is a model dot-product proof, since a common perpendicular must be perpendicular to both direction vectors. Part (iv) reinforces the syllabus form $|\mathbf{r} - \mathbf{c}| = R$, which is usually the cleanest way to present a sphere in vector questions.

Problem 2.16: The Projection Paradox

Two straight lines L_1 and L_2 pass through the origin in 3D space, with vector equations

$$L_1 : \mathbf{r} = \lambda \begin{pmatrix} 1 \\ a \\ b \end{pmatrix}, \quad L_2 : \mathbf{r} = \mu \begin{pmatrix} 1 \\ c \\ d \end{pmatrix},$$

where a, b, c, d are non-zero real constants.

- (i) State the Cartesian equations of the projections of L_1 and L_2 onto the xy -plane and the xz -plane.
- (ii) A student observes that the projections onto the xy -plane are perpendicular, and the projections onto the xz -plane are also perpendicular. Using 2D gradient properties, write down two algebraic equations connecting a, b, c , and d .
- (iii) By evaluating the dot product of the 3D direction vectors, prove that the actual lines L_1 and L_2 can never be perpendicular in 3D space.
- (iv) Let θ be the acute angle between the two 3D lines. Show that

$$\cos \theta = \frac{1}{\sqrt{3 + a^2 + \frac{1}{a^2} + b^2 + \frac{1}{b^2} + \frac{a^2}{b^2} + \frac{b^2}{a^2}}}.$$

- (v) Hence, using the AM-GM inequality, find the minimum possible acute angle between L_1 and L_2 , correct to the nearest degree.

Solution 2.16

(i) Projections:

$$y = ax, \quad y = cx, \quad z = bx, \quad z = dx.$$

(ii) Perpendicular projections give

$$ac = -1, \quad bd = -1,$$

so

$$c = -\frac{1}{a}, \quad d = -\frac{1}{b}.$$

(iii) Let

$$\mathbf{u} = \begin{pmatrix} 1 \\ a \\ b \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} 1 \\ c \\ d \end{pmatrix}.$$

Then

$$\mathbf{u} \cdot \mathbf{v} = 1 + ac + bd = 1 - 1 - 1 = -1 \neq 0,$$

so L_1 and L_2 are never perpendicular in 3D.

(iv) Since θ is acute,

$$\cos \theta = \frac{|\mathbf{u} \cdot \mathbf{v}|}{|\mathbf{u}||\mathbf{v}|} = \frac{1}{\sqrt{(1+a^2+b^2)(1+c^2+d^2)}}.$$

Substitute $c = -\frac{1}{a}$, $d = -\frac{1}{b}$:

$$\begin{aligned} (1+a^2+b^2)(1+c^2+d^2) &= 1 + \frac{1}{a^2} + \frac{1}{b^2} + a^2 + 1 + \frac{a^2}{b^2} + b^2 + \frac{b^2}{a^2} + 1 \\ &= 3 + a^2 + \frac{1}{a^2} + b^2 + \frac{1}{b^2} + \frac{a^2}{b^2} + \frac{b^2}{a^2}. \end{aligned}$$

Hence

$$\cos \theta = \frac{1}{\sqrt{3 + a^2 + \frac{1}{a^2} + b^2 + \frac{1}{b^2} + \frac{a^2}{b^2} + \frac{b^2}{a^2}}}.$$

(v) By AM-GM,

$$a^2 + \frac{1}{a^2} \geq 2, \quad b^2 + \frac{1}{b^2} \geq 2, \quad \frac{a^2}{b^2} + \frac{b^2}{a^2} \geq 2.$$

So

$$\cos \theta \leq \frac{1}{\sqrt{3+2+2+2}} = \frac{1}{3}.$$

Thus

$$\theta_{\min} = \cos^{-1} \left(\frac{1}{3} \right) \approx 70.5^\circ \approx \boxed{71^\circ}.$$

Takeaways 2.16

This problem is a great reminder that 2D projections can be deceptive: two lines may look perpendicular from multiple viewpoints without being perpendicular in 3D. It also shows how Extension 2 problems often combine topics naturally—vector geometry, gradients, dot products, and inequalities all work together here. Once a complicated expression appears in a denominator, inequalities such as AM-GM become a natural tool for optimising the angle.

Problem 2.17: Point-to-Line Distance and Sphere Intersection

Consider the line ℓ with equation $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$.

- (i) Find the perpendicular distance from point $P(1, 2, 0)$ to the line ℓ .
- (ii) Find the shortest distance from the origin to the line ℓ .
- (iii) Determine the points where line ℓ intersects the sphere $x^2 + y^2 + z^2 = 4$.

Solution 2.17

(i) Let $A(1, 0, 1)$ be on ℓ , direction $\mathbf{d} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$.

$$\vec{AP} = \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}.$$

Project $\vec{AP}$ onto $\mathbf{d}$:

$$\text{proj}_{\mathbf{d}} \vec{AP} = \frac{\vec{AP} \cdot \mathbf{d}}{|\mathbf{d}|^2} \mathbf{d} = \frac{1}{2} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}.$$

So the perpendicular component is

$$\vec{AP}_{\perp} = \vec{AP} - \text{proj}_{\mathbf{d}} \vec{AP} = \begin{pmatrix} 0 \\ \frac{3}{2} \\ -\frac{3}{2} \end{pmatrix},$$

and therefore

$$d = |\vec{AP}_{\perp}| = \sqrt{\frac{9}{4} + \frac{9}{4}} = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2} \text{ units.}$$

(ii) Use projection method. Let $\vec{OA} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$. Projection of $\vec{OA}$ onto $\mathbf{d}$: $\text{proj}_{\mathbf{d}} \vec{OA} = \frac{\vec{OA} \cdot \mathbf{d}}{|\mathbf{d}|^2} \mathbf{d} = \frac{1}{2} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$

Perpendicular component: $\vec{OA}_{\perp} = \vec{OA} - \text{proj}_{\mathbf{d}} \vec{OA} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \begin{pmatrix} 0 \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix} = \begin{pmatrix} 1 \\ -\frac{1}{2} \\ \frac{1}{2} \end{pmatrix}$

Distance: $|\vec{OA}_{\perp}| = \sqrt{1 + \frac{1}{4} + \frac{1}{4}} = \sqrt{\frac{3}{2}} = \frac{\sqrt{6}}{2}$ units.

(iii) Substitute line into sphere: $x = 1, y = \lambda, z = 1 + \lambda$.

$$1 + \lambda^2 + (1 + \lambda)^2 = 4$$

$$1 + \lambda^2 + 1 + 2\lambda + \lambda^2 = 4$$

$$2\lambda^2 + 2\lambda - 2 = 0 \implies \lambda^2 + \lambda - 1 = 0$$

$$\lambda = \frac{-1 \pm \sqrt{1+4}}{2} = \frac{-1 \pm \sqrt{5}}{2}$$

Points: For $\lambda_1 = \frac{-1+\sqrt{5}}{2}$: $(1, \frac{-1+\sqrt{5}}{2}, \frac{1+\sqrt{5}}{2})$

For $\lambda_2 = \frac{-1-\sqrt{5}}{2}$: $(1, \frac{-1-\sqrt{5}}{2}, \frac{1-\sqrt{5}}{2})$

Takeaways 2.17


Three key distance techniques demonstrated: (1) **projection** for point-to-line from a given point, (2) **projection** again for the distance from the origin, and (3) **substitution** for line-sphere intersection. Both distance questions rely on the same idea: resolve the relevant vector into components parallel and perpendicular to the line direction. For intersections, substitute parametric equations into the surface equation to get a quadratic; the number of real solutions indicates the geometric relationship. The golden ratio $\phi = \frac{1+\sqrt{5}}{2}$ appearing in λ is a nice coincidence!

Problem 2.18: 3D Helix Projection

Each point P on a helix has position vector:

$$\mathbf{r} = 3 \cos \theta \mathbf{i} + 3 \sin \theta \mathbf{j} + \frac{\theta}{\pi} \mathbf{k} \quad \text{for } \theta \geq 0.$$

- (a) The vector $\mathbf{r}$ is projected into the xz -plane to obtain $\mathbf{s} = \text{proj}_{\mathbf{i}} \mathbf{r} + \text{proj}_{\mathbf{k}} \mathbf{r}$, the position vector of point S . Write $\mathbf{s}$ in terms of the parameter θ .
 (b) Determine the Cartesian equation of the curve traced out by S for $0 \leq \theta \leq 2\pi$, specifying its domain.



Hint:(a) Projecting a 3D vector onto standard basis vectors simply isolates that specific component. What happens to the $\mathbf{j}$ component when projecting onto the xz -plane?
 (b) Write out the parametric equations for x and z based on your answer from (a). Isolate θ in the simpler equation and substitute it into the other to eliminate the parameter. Don't forget to convert the domain of θ into a domain for your new independent variable.

Solution 2.18

- (a) The projection of $\mathbf{r}$ onto $\mathbf{i}$ is simply the x -component, and the projection onto $\mathbf{k}$ is the z -component.

$$\mathbf{s} = 3 \cos \theta \mathbf{i} + \frac{\theta}{\pi} \mathbf{k}$$

- (b) From the position vector $\mathbf{s}$, we extract the parametric equations: $x = 3 \cos \theta$, $z = \frac{\theta}{\pi}$.
 From the z equation, we isolate θ : $\theta = \pi z$. Substitute this into the x equation:

$$x = 3 \cos(\pi z)$$

Now, determine the domain. We are given $0 \leq \theta \leq 2\pi$.

Substitute $\theta = \pi z$ into this inequality:

$$0 \leq \pi z \leq 2\pi \implies 0 \leq z \leq 2$$

The Cartesian equation is $x = 3 \cos(\pi z)$ for $0 \leq z \leq 2$.


Takeaways 2.18

- **The "Shadow" Concept:** Vector projection in 3D geometry is exactly the math used in rendering pipelines to cast shadows. By projecting a 3D path ($\mathbf{r}$) onto a 2D plane ($\mathbf{s}$), we are mathematically defining the "shadow" the helix casts on the wall (the xz -plane) when a light is shone straight down the y -axis. The result is a simple 2D cosine wave!
- **Domain Conversion:** A common student error in parametric elimination is forgetting to translate the domain of the parameter (θ) into the domain of the Cartesian variables (x or z).

Problem 2.19: Parametric Equation of a Helix

A circular helix spirals around the y -axis as shown in the diagram. Every point on the helix is 5 units from the y -axis. The helix completes exactly one full rotation for every 2π units it travels along the positive y -axis, starting from the point $(5, 0, 0)$.

Write down the position vector, $\mathbf{r}$, for this helix in terms of a parameter θ .



- **The Axis of Propagation:** Since the helix moves constantly along the y -axis, the y -component of your vector will be linear (e.g., just the parameter θ).
- **The Circular Cross-Section:** If you look straight down the y -axis, the shadow of the helix on the xz -plane is just a circle of radius 5. Use sine and cosine for the x and z components.
- **Starting Position:** Plug $\theta = 0$ into your proposed sine and cosine components to make sure they yield the correct starting coordinate of $x = 5$ and $z = 0$.

Hint:

Solution 2.19

Let the parameter be θ .

Linear Component (y -axis): The helix travels along the y -axis. Since it completes one rotation (2π radians) over a distance of 2π units, the y -component is directly proportional to θ . $y = \theta$.

Circular Components (x and z axes): The distance from the y -axis is 5, meaning $x^2 + z^2 = 5^2$. This is modeled using $5 \cos \theta$ and $5 \sin \theta$. At $y = 0$ (meaning $\theta = 0$), the position is $(5, 0, 0)$.

For x : we need a function where $f(0) = 5$. Since $5 \cos(0) = 5$, we set $x = 5 \cos \theta$.

For z : we need a function where $f(0) = 0$. Since $5 \sin(0) = 0$, we set $z = 5 \sin \theta$.

Final Position Vector:

$$\mathbf{r} = 5 \cos \theta \mathbf{i} + \theta \mathbf{j} + 5 \sin \theta \mathbf{k} \quad \text{for } \theta \geq 0$$

Takeaways 2.19

- **Component Separation:** When building 3D parametric equations, it helps to isolate the "movement" vector from the "shape" vectors. The axis with the solitary variable ($y = \theta$) acts as the spine of the shape, while the trigonometric axes wrap around it.
- **Initial Value Testing:** The most common mistake is swapping sine and cosine. Always plug 0 into your parameter to verify it matches the given starting coordinate.


Problem 2.20: Curve on a Cylinder and Arc Length

A graph of $h = 2 \sin 2\theta$ for $0 \leq \theta \leq 2\pi$ is drawn on a rectangular sheet of paper. The ends are then glued together to form a cylinder with only the curve showing. It is found that the cylinder has a radius of 3 cm. The xy -plane corresponds with $h = 0$. The point $(3, 0, 0)$ corresponds with $\theta = 0$ and $(0, 3, 0)$ corresponds with $\theta = \frac{\pi}{2}$ on the graph paper.

(a) Suggest a possible parametric vector equation $\mathbf{r}(\theta)$ for this curve.

(b) Re-write the equation of the curve in terms of the new parameter ℓ that is the arc length around the base circle $x^2 + y^2 = 9$, measured anti-clockwise from $(3, 0, 0)$.

(c) A different design uses the graph $h = 3 \cos \theta$. This curve is projected onto the yz -plane to obtain $\mathbf{s}(\theta) = \text{proj}_{\mathbf{j}} \mathbf{r} + \text{proj}_{\mathbf{k}} \mathbf{r}$. Find the Cartesian equation of the shape seen in the yz -plane.



Hint:

- **Part (a):** The x and y coordinates form a standard circle of radius 3. The z -coordinate represents the height h directly.
- **Part (b):** Recall the formula for the arc length of a circle: $\ell = r\theta$. Express θ in terms of ℓ and substitute it into your vector equation.
- **Part (c):** Projecting onto the yz -plane means you only keep the $\mathbf{j}$ and $\mathbf{k}$ components. Write out the parametric equations for y and z , then use a trigonometric identity to eliminate θ .

Solution 2.20

(a) The base of the cylinder is a circle of radius 3 in the xy -plane.

Given $(3, 0, 0)$ is at $\theta = 0$ and $(0, 3, 0)$ is at $\theta = \frac{\pi}{2}$, the standard circular parameterisation applies: $x = 3 \cos \theta$ and $y = 3 \sin \theta$.

The height z is given by $h = 2 \sin 2\theta$.

$$\mathbf{r}(\theta) = 3 \cos \theta \mathbf{i} + 3 \sin \theta \mathbf{j} + 2 \sin 2\theta \mathbf{k}$$

(b) The arc length around a circle is $\ell = r\theta$.

Since the radius $r = 3$, we have $\ell = 3\theta$, which rearranging gives $\theta = \frac{\ell}{3}$.

Substitute this into the vector equation:

$$\mathbf{r}(\ell) = 3 \cos \left(\frac{\ell}{3} \right) \mathbf{i} + 3 \sin \left(\frac{\ell}{3} \right) \mathbf{j} + 2 \sin \left(\frac{2\ell}{3} \right) \mathbf{k}$$

(c) The new curve has the equation:

$$\mathbf{r}(\theta) = 3 \cos \theta \mathbf{i} + 3 \sin \theta \mathbf{j} + 3 \cos \theta \mathbf{k}$$

Projecting onto the yz -plane isolates the y and z components:

$$\mathbf{s}(\theta) = 3 \sin \theta \mathbf{j} + 3 \cos \theta \mathbf{k}$$

This gives parametric equations $y = 3 \sin \theta$ and $z = 3 \cos \theta$.

Squaring both equations gives $y^2 = 9 \sin^2 \theta$ and $z^2 = 9 \cos^2 \theta$.

Adding them together:

$$y^2 + z^2 = 9(\sin^2 \theta + \cos^2 \theta) = 9$$

The Cartesian equation is $y^2 + z^2 = 9$, which is a circle of radius 3.

Takeaways 2.20


- **Read the question carefully:** This problem ask you to write "a possible parametric vector equation", not all.
- **2D to 3D Mappings:** Wrapping a 2D graph around a cylinder is simply assigning the domain (the horizontal axis) to wrap around a standard parametric circle, and the range (the vertical axis) to the z -axis.
- **Natural Parameterisation:** Using arc length (ℓ) as a parameter instead of an angle (θ) is a crucial concept in advanced vector calculus. It means the curve is parameterised by the actual physical distance travelled along it. See also Appendix A.8 for the formal framework of velocity, acceleration, and speed in three-dimensional vector calculus.
- **Orthogonal Projections:** A complex 3D curve (like the tilted ellipse in part c) can look like a perfect circle when viewed exactly side-on.

Problem 2.21: Sketching 3D Parametric Curves

Sketch the curves defined parametrically by the following vector equations. Ensure you clearly label the axes, starting points, and endpoints.

(a) $\mathbf{r}(t) = (2 \sin t)\mathbf{i} + (2 \cos t)\mathbf{j} + \frac{t}{\pi}\mathbf{k}$ for $0 \leq t \leq 4\pi$

(b) $\mathbf{r}(t) = t\mathbf{i} + t\mathbf{j} + (4 - t^2)\mathbf{k}$ for $-2 \leq t \leq 2$



Hint:

- **Part (a):** Look at the $\mathbf{i}$ and $\mathbf{j}$ components. It traces a circle of radius 2, but pay attention to where it *starts* when $t = 0$.
- **Part (b):** Notice that the x and y components are identical ($x = t, y = t$). This means the entire curve is trapped flat inside the diagonal plane $y = x$. What 2D shape does the z equation make?


Solution 2.21

(a) Circular Helix:

Base Shape: $x^2 + y^2 = 4 \sin^2 t + 4 \cos^2 t = 4$. This is a circular helix of radius 2.

Start/End: At $t = 0$, $\mathbf{r}(0) = (0, 2, 0)$. At $t = 4\pi$, $\mathbf{r}(4\pi) = (0, 2, 4)$.

Sketch Requirement: A helix starting on the y -axis, spiralling upwards around the z -axis, completing exactly two full turns, ending at height $z = 4$.




(b) Parabola on a Plane:

Base Shape: Since $x = t$ and $y = t$, the curve lies completely on the plane $y = x$.

Vertical Profile: $z = 4 - t^2 = 4 - x^2$. This is an inverted parabola.

Start/End: At $t = -2$, position is $(-2, -2, 0)$. At $t = 2$, position is $(2, 2, 0)$. The maximum turning point is at $t = 0 \Rightarrow (0, 0, 4)$.

Sketch Requirement: A parabola starting on the xy -plane in the 3rd quadrant, passing through the z -axis at height 4, and dropping back to the xy -plane in the 1st quadrant.




Takeaways 2.21

- **Reading Traces:** When sketching 3D curves, always look for relationships between pairs of variables. If $x = t$ and $y = t$, the 3D curve is just a 2D curve drawn on a diagonal sheet of glass (a plane).
- Think of the projections of the curve onto the coordinate planes as "shadows".

Problem 2.22: Spiral Curve on a Paraboloid

A curve spirals three times from the origin to the point $(3, 0, 9)$ ascending around the surface of the paraboloid $z = x^2 + y^2$, as shown in the diagram. Determine a possible parametric vector equation for this curve, and state the exact restriction on the parameter.



Hint: Let your parameter θ represent the angle of rotation. If it spirals exactly three times, what is your domain? Find a linear equation for the radius r in terms of θ so that $r = 3$ at the very end of the domain. Finally, substitute x and y into the paraboloid equation to get z .

Solution 2.22

Let the parameter be θ , representing the angle of rotation.

- Domain:** Three full spirals means θ must range from 0 to $3 \times 2\pi$. Restriction: $0 \leq \theta \leq 6\pi$.
- Radius $r(\theta)$:** The curve starts at the origin ($r = 0$) and ends at $(3, 0, 9)$ where $r = 3$. Assuming constant outward growth, r is directly proportional to θ .

$$r(\theta) = \frac{\text{Final Radius}}{\text{Final Angle}} \theta = \frac{3}{6\pi} \theta = \frac{\theta}{2\pi}$$

- x and y Components:**

$$x = r \cos \theta = \frac{\theta}{2\pi} \cos \theta, \quad y = r \sin \theta = \frac{\theta}{2\pi} \sin \theta$$

- z Component:** The curve lies on $z = x^2 + y^2 = r^2$.

$$z = \left(\frac{\theta}{2\pi} \right)^2 = \frac{\theta^2}{4\pi^2}$$

Vector Equation:

$$\mathbf{r}(\theta) = \left(\frac{\theta}{2\pi} \cos \theta \right) \mathbf{i} + \left(\frac{\theta}{2\pi} \sin \theta \right) \mathbf{j} + \left(\frac{\theta^2}{4\pi^2} \right) \mathbf{k} \quad \text{for } 0 \leq \theta \leq 6\pi$$

Takeaways 2.22


- Cylindrical Coordinates:** This problem introduces the concept of cylindrical coordinates (r, θ, z) . By writing x, y , and z all in terms of r and θ first, building complex spiral vector equations becomes a trivial step-by-step process.

Remark 2.1: Concluding: The Challenge of 3D Parametrics

If you find 3D parametric equations conceptually difficult, you are not alone. They are universally considered one of the most challenging topics in the Extension 2 syllabus.


Why are they so hard? Our brains are wired to interpret mathematics through 2D planar relationships, and parametrics force us to break that habit. Instead of a direct geometric relationship between x , y , and z , you juggle three coordinates tied to an “invisible” fourth variable (the parameter t). Mastering this topic requires you to mentally dismantle a 3D shape into its 2D planar “shadows”—its projections onto the coordinate planes—to understand the core geometry and behaviour of the curve.

By this point, you should recognise the fundamental difference between a curve that wraps around a 3D volume, and a 3D curve that is actually trapped on a flat, diagonal slice of space.



The Spiral (Helix)

$$x = \cos t, \quad y = \sin t, \quad z = t$$



The Planar Parabola

$$x = t, \quad y = 2t, \quad z = t^2$$

3 Part 2: Problems and Solutions (Concise + Hints)

Part 2 contains 46 additional problems distributed across three difficulty levels. Each problem includes an upside-down hint and a concise solution sketch to encourage independent thinking.

3.1 Part 2 Basic Problems

Problem 3.1: Vector to Cartesian Line Equation

What is the Cartesian equation of the line $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 4 \end{pmatrix}$?

Hint: Eliminate the parameter λ by expressing it from one equation and substituting into the other.

Solution 3.1: Sketch

Write parametric equations: $x = 1 + 2\lambda$, $y = 3 + 4\lambda$. From the first equation, $\lambda = \frac{x-1}{2}$. Substitute into second: $y = 3 + 4 \cdot \frac{x-1}{2} = 3 + 2(x-1) = 2x + 1$. Therefore $y = 2x + 1$ or $2x - y + 1 = 0$.


Problem 3.2: Sketch 3D Helix

Sketch the curve described by $\mathbf{r}(t) = -5 \cos t \mathbf{i} + 5 \sin t \mathbf{j} + t \mathbf{k}$ for $0 \leq t \leq 4\pi$.

Hint: The x and y components trace a circle while z increases linearly with t .

Solution 3.2: Sketch

In the xy -plane, $x^2 + y^2 = 25 \cos^2 t + 25 \sin^2 t = 25$, which is a circle of radius 5. As t increases from 0 to 4π , the point moves around the circle twice while rising from $z = 0$ to $z = 4\pi$. This creates a helix spiraling upward around the z -axis.



Problem 3.3: Angle Between 3D Vectors

Find the angle between vectors $\mathbf{a} = \begin{pmatrix} 2 \\ 0 \\ 4 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} -3 \\ 1 \\ 2 \end{pmatrix}$ to the nearest degree.

$$\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|}$$

Solution 3.3: Sketch

$\mathbf{a} \cdot \mathbf{b} = (2)(-3) + (0)(1) + (4)(2) = -6 + 8 = 2$. $|\mathbf{a}| = \sqrt{4 + 16} = \sqrt{20} = 2\sqrt{5}$, $|\mathbf{b}| = \sqrt{9 + 1 + 4} = \sqrt{14}$.
Therefore $\cos \theta = \frac{2}{2\sqrt{5}\sqrt{14}} = \frac{1}{\sqrt{70}}$, so $\theta = \cos^{-1}(1/\sqrt{70}) \approx 83^\circ$.

Problem 3.4: Unit Vector in Direction

Given $\mathbf{u} = 2\mathbf{i} - 2\mathbf{j} - 5\mathbf{k}$, find a vector parallel to $\mathbf{u}$ with length 5 units.

Hint: Find the unit vector in the direction of $\mathbf{u}$, then scale by 5.

Solution 3.4: Sketch

$|\mathbf{u}| = \sqrt{4 + 4 + 25} = \sqrt{33}$. Unit vector: $\hat{\mathbf{u}} = \frac{1}{\sqrt{33}}(2\mathbf{i} - 2\mathbf{j} - 5\mathbf{k})$. Required vector: $5\hat{\mathbf{u}} = \frac{5}{\sqrt{33}}(2\mathbf{i} - 2\mathbf{j} - 5\mathbf{k})$.

Problem 3.5: Distance to Plane and Axis

Find the distance from point $P(1, 4, 3)$ to: (a) the xz -plane, (b) the y -axis.

Hint: (a) Distance to xz -plane is the y -coordinate. (b) Distance to y -axis uses x and z coordinates.

Solution 3.5: Sketch

Part (a): Distance from $P(1, 4, 3)$ to the xz -plane

The xz -plane consists of all points where $y = 0$. The perpendicular distance from point $P(1, 4, 3)$ to this plane is simply the absolute value of the y -coordinate:

$$\text{Distance} = |y| = |4| = 4 \text{ units}$$

The closest point on the xz -plane to P is $(1, 0, 3)$, which we can verify: $\sqrt{(1-1)^2 + (4-0)^2 + (3-3)^2} = 4$.

Part (b): Distance from $P(1, 4, 3)$ to the y -axis

The y -axis consists of all points of the form $(0, y, 0)$ where $y \in \mathbb{R}$. To find the closest point on the y -axis to $P(1, 4, 3)$, we note that this point has the same y -coordinate as P . Therefore, the closest point is $(0, 4, 0)$.

The distance from $P(1, 4, 3)$ to $(0, 4, 0)$ is:

$$\text{Distance} = \sqrt{(1-0)^2 + (4-4)^2 + (3-0)^2} = \sqrt{1+0+9} = \sqrt{10} \text{ units}$$

Geometric interpretation: The distance to the y -axis is the radius of the circle in a plane perpendicular to the y -axis passing through P . This radius involves only the x and z coordinates: $\sqrt{x^2 + z^2} = \sqrt{1^2 + 3^2} = \sqrt{10}$.

Problem 3.6: Unit Vector Perpendicular to Two Vectors

Find a unit vector perpendicular to both $\mathbf{u} = \mathbf{i} + \mathbf{j}$ and $\mathbf{v} = \mathbf{i} + \mathbf{k}$, where $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ are the standard unit vectors along the x -, y -, and z -axes respectively.

Hint: Let the required vector be (x, y, z) and use perpendicularity with both given vectors.

Solution 3.6: Sketch

Let $\mathbf{w} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$. Since $\mathbf{w}$ is perpendicular to both vectors,

$$\mathbf{w} \cdot \mathbf{u} = x + y = 0, \quad \mathbf{w} \cdot \mathbf{v} = x + z = 0.$$

So $y = -x$ and $z = -x$, giving $\mathbf{w} = x(\mathbf{i} - \mathbf{j} - \mathbf{k})$. Choosing $x = 1$ gives a convenient perpendicular vector $\mathbf{i} - \mathbf{j} - \mathbf{k}$, whose magnitude is $\sqrt{3}$. Hence a unit vector is

$$\frac{1}{\sqrt{3}}(\mathbf{i} - \mathbf{j} - \mathbf{k}).$$

Problem 3.7: Point on Line

Consider the line passing through points $A(0, 2, 1)$ and $B(2, 7, 4)$. Find the values of a and b such that the point $P(a, 1, b)$ also lies on this line.

Hint: Find parametric equations using direction vector $(2, 5, 3)$, then solve for the parameter when $y = 1$.

Solution 3.7: Sketch

Step 1: Find the direction vector

The direction vector of the line is:

$$\vec{AB} = B - A = (2, 7, 4) - (0, 2, 1) = (2, 5, 3)$$

Step 2: Write parametric equations

Using point $A(0, 2, 1)$ and direction vector $(2, 5, 3)$, the parametric equations of the line are:

$$x = 0 + 2t = 2t$$

$$y = 2 + 5t$$

$$z = 1 + 3t$$

where t is the parameter.

Step 3: Find the parameter when $y = 1$

Since point P has y -coordinate equal to 1, we substitute into the y equation:

$$1 = 2 + 5t \Rightarrow 5t = -1 \Rightarrow t = -\frac{1}{5}$$

Step 4: Find a and b

Substituting $t = -\frac{1}{5}$ into the x and z equations:

$$a = x = 2 \left(-\frac{1}{5} \right) = -\frac{2}{5}$$

$$b = z = 1 + 3 \left(-\frac{1}{5} \right) = 1 - \frac{3}{5} = \frac{2}{5}$$

Therefore, $a = -\frac{2}{5}$ and $b = \frac{2}{5}$, giving the point $P \left(-\frac{2}{5}, 1, \frac{2}{5} \right)$.

Verification: We can verify by checking that this point satisfies the line equation with $t = -\frac{1}{5}$.

Problem 3.8: Equal Magnitude Perpendicular Vectors

Prove that if $\mathbf{u}$ and $\mathbf{v}$ are non-zero vectors of equal magnitude, then $\mathbf{u} - \mathbf{v}$ is perpendicular to $\mathbf{u} + \mathbf{v}$.

Hint: Show that $(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v}) = 0$.

Solution 3.8: Sketch

To prove perpendicularity, we need to show that the dot product equals zero.

Expand the dot product using distributivity:

$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v}) = \mathbf{u} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u} - \mathbf{v} \cdot \mathbf{v}$$

Since the dot product is commutative ($\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$), the middle terms cancel:

$$= \mathbf{u} \cdot \mathbf{u} - \mathbf{v} \cdot \mathbf{v}$$

$$= |\mathbf{u}|^2 - |\mathbf{v}|^2$$

Given that $|\mathbf{u}| = |\mathbf{v}|$, we have:

$$|\mathbf{u}|^2 - |\mathbf{v}|^2 = 0$$

Therefore, $(\mathbf{u} - \mathbf{v}) \perp (\mathbf{u} + \mathbf{v})$.

Takeaways 3.1

- This result has a beautiful geometric interpretation: if vectors $\mathbf{u}$ and $\mathbf{v}$ have equal magnitude, then the vectors $\mathbf{u} + \mathbf{v}$ (the diagonal) and $\mathbf{u} - \mathbf{v}$ (the other diagonal) form perpendicular diagonals.
- **Rectangle interpretation:** The vectors $\mathbf{u}$, $\mathbf{v}$, $\mathbf{u} + \mathbf{v}$, and $\mathbf{u} - \mathbf{v}$ form a *rhombus* (not a rectangle in general), since $|\mathbf{u}| = |\mathbf{v}|$ makes all four sides equal length. The diagonals of this rhombus are perpendicular, which is what we just proved.
- Special case: If $\mathbf{u} \perp \mathbf{v}$ and $|\mathbf{u}| = |\mathbf{v}|$, then the rhombus becomes a *square*, and the diagonals are both perpendicular and equal in length.
- This identity is useful in many geometric proofs and demonstrates the power of algebraic vector manipulation to prove geometric relationships.

Problem 3.9: Direction Cosines

Prove that $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$ for a position vector $\mathbf{r} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ making angles α, β, γ with the positive x -, y -, and z -axes respectively. (The quantities $\cos \alpha$, $\cos \beta$, and $\cos \gamma$ are called the *direction cosines* of the vector $\mathbf{r}$.)

Hint: Use dot products with unit vectors $\mathbf{i}, \mathbf{j}, \mathbf{k}$ to find the direction cosines.

Solution 3.9: Sketch

Here α , β , and γ are the angles between $\mathbf{r}$ and the *positive* x -, y -, and z -axes. Their cosines are the **direction cosines** of $\mathbf{r}$.

Step 1: Use the dot product with a unit axis vector

For any two vectors $\mathbf{u}$ and $\mathbf{v}$, $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}||\mathbf{v}| \cos \theta$, where θ is the angle between them. The axis unit vectors have length $|\mathbf{i}| = |\mathbf{j}| = |\mathbf{k}| = 1$.

Step 2: Find each direction cosine

Dot $\mathbf{r} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ with $\mathbf{i}$:

$$\mathbf{r} \cdot \mathbf{i} = a \quad \text{and} \quad \mathbf{r} \cdot \mathbf{i} = |\mathbf{r}||\mathbf{i}| \cos \alpha = |\mathbf{r}| \cos \alpha.$$

So $\cos \alpha = \frac{a}{|\mathbf{r}|}$. The same idea with $\mathbf{j}$ and $\mathbf{k}$ gives

$$\cos \beta = \frac{b}{|\mathbf{r}|}, \quad \cos \gamma = \frac{c}{|\mathbf{r}|}.$$

Step 3: Square and add

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \frac{a^2}{|\mathbf{r}|^2} + \frac{b^2}{|\mathbf{r}|^2} + \frac{c^2}{|\mathbf{r}|^2} = \frac{a^2 + b^2 + c^2}{|\mathbf{r}|^2}.$$

Step 4: Replace with the magnitude

By definition, $|\mathbf{r}| = \sqrt{a^2 + b^2 + c^2}$, so $|\mathbf{r}|^2 = a^2 + b^2 + c^2$. Hence

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \frac{|\mathbf{r}|^2}{|\mathbf{r}|^2} = 1. \quad \square$$

Geometric meaning: the direction cosines tell you how much of $\mathbf{r}$ points along each axis; squaring and adding recovers the full length of the vector.

Problem 3.10: Line Intersection by Components

Find the intersection point of lines $\mathbf{r} = \begin{pmatrix} 3 \\ -1 \\ 7 \end{pmatrix} + \lambda_1 \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ and $\mathbf{r} = \begin{pmatrix} 3 \\ -6 \\ 2 \end{pmatrix} + \lambda_2 \begin{pmatrix} -2 \\ 1 \\ 3 \end{pmatrix}$.

Hint: Equate components and solve the system for λ_1 and λ_2 .

Solution 3.10: Sketch

Equating: $3 + \lambda_1 = 3 - 2\lambda_2$, $-1 + 2\lambda_1 = -6 + \lambda_2$, $7 + \lambda_1 = 2 + 3\lambda_2$. From first equation: $\lambda_1 = -2\lambda_2$. Substitute into second: $-1 - 4\lambda_2 = -6 + \lambda_2 \Rightarrow \lambda_2 = 1$, so $\lambda_1 = -2$. Check third equation: $7 - 2 = 5 = 2 + 3$ ✓. Intersection point: $(1, -5, 5)$.

Problem 3.11: Multiple Choice: Cartesian Equation

What is the Cartesian equation of $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} -2 \\ 4 \end{pmatrix}$?

Options:

- A. $2y + x = 7$,
- B. $y - 2x = -5$,
- C. $y + 2x = 5$,
- D. $2y - x = -1$.

Hint: Eliminate λ from $x = 1 - 2\lambda$ and $y = 3 + 4\lambda$.

Solution 3.11: Sketch

From $x = 1 - 2\lambda$, get $\lambda = \frac{1-x}{2}$. Substitute: $y = 3 + 4 \cdot \frac{1-x}{2} = 3 + 2(1-x) = 5 - 2x$, giving $y + 2x = 5$. Answer: C.

Problem 3.12: Closest Point on Line to Origin

Find the point on line $\mathbf{r} = \begin{pmatrix} 1 \\ 4 \\ 6 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix}$ closest to the origin. That is, find the value of λ such that the distance from the point $\mathbf{r}(\lambda)$ to the origin $O(0, 0, 0)$ is minimized.

Hint: The closest point occurs when $\mathbf{r}$ is perpendicular to the direction vector.

Solution 3.12: Sketch

Let $\mathbf{r} = \begin{pmatrix} 1+2\lambda \\ 4+\lambda \\ 6+\lambda \end{pmatrix}$. For closest point: $\mathbf{r} \cdot \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} = 0$. This gives $2(1+2\lambda) + (4+\lambda) + (6+\lambda) = 0 \Rightarrow 6\lambda + 12 = 0 \Rightarrow \lambda = -2$. Point: $(-3, 2, 4)$.

Problem 3.13: Unit Vector Perpendicular to Two

Find a unit vector perpendicular to $\begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 4 \\ -1 \end{bmatrix}$.

Hint: Let the required vector be (x, y, z) and solve two perpendicularity equations.

Solution 3.13: Sketch

Let $\mathbf{w} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$. Then

$$\mathbf{w} \cdot \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix} = 0 \quad \text{and} \quad \mathbf{w} \cdot \begin{bmatrix} 1 \\ 4 \\ -1 \end{bmatrix} = 0.$$

So

$$3x - 2y + z = 0, \quad x + 4y - z = 0.$$

Adding gives $4x + 2y = 0$, so $y = -2x$. Substituting into $x + 4y - z = 0$ gives

$$x - 8x - z = 0 \Rightarrow z = -7x.$$

Thus $\mathbf{w} = x \begin{bmatrix} 1 \\ -2 \\ -7 \end{bmatrix}$. Taking $x = 1$, a convenient perpendicular vector is $\begin{bmatrix} 1 \\ -2 \\ -7 \end{bmatrix}$, whose magnitude is

$$\sqrt{1 + 4 + 49} = 3\sqrt{6}.$$

Hence a unit vector is

$$\frac{1}{3\sqrt{6}} \begin{bmatrix} 1 \\ -2 \\ -7 \end{bmatrix}.$$

Problem 3.14: Perpendicular Dot Product Proof

Prove that for non-zero vectors $\mathbf{a}, \mathbf{b}$, the equation $(\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) = |\mathbf{a}|^2 + |\mathbf{b}|^2$ holds only if $\mathbf{a} \perp \mathbf{b}$.

Hint: Expand the left side and compare with the right side.

Solution 3.14: Sketch

Expanding: $(\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) = |\mathbf{a}|^2 + 2\mathbf{a} \cdot \mathbf{b} + |\mathbf{b}|^2$. Comparing with RHS: $|\mathbf{a}|^2 + 2\mathbf{a} \cdot \mathbf{b} + |\mathbf{b}|^2 = |\mathbf{a}|^2 + |\mathbf{b}|^2 \Rightarrow 2\mathbf{a} \cdot \mathbf{b} = 0 \Rightarrow \mathbf{a} \cdot \mathbf{b} = 0$, which means $\mathbf{a} \perp \mathbf{b}$.

3.2 Part 2 Medium Problems

Problem 3.15: Line Intersection in 3D

Find the intersection point of lines $\mathbf{r} = \begin{pmatrix} 3 \\ -1 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ and $\mathbf{r} = \begin{pmatrix} 1 \\ -6 \\ 2 \end{pmatrix} + \mu \begin{pmatrix} -2 \\ 1 \\ 3 \end{pmatrix}$.


Hint: Set up three equations by equating components, then solve for parameters.

Solution 3.15: Sketch

Equating components: $3 + \lambda = 1 - 2\mu$, $-1 + 2\lambda = -6 + \mu$, $7 + \lambda = 2 + 3\mu$. From equations 1 and 3: $\lambda = -2\mu$ and $\lambda = -5 + 3\mu$. Solving: $\mu = 1$, $\lambda = -2$. Verify in equation 2. Intersection point: $(1, -5, 5)$.

Problem 3.16: Triangular Pyramid — Find Vertex Coordinates

The diagram below shows a triangular pyramid with vertices $O(0, 0, 0)$, $P(4, 3, 0)$, $Q(3, 4, 0)$ and $R(x, y, z)$, where x, y, z are positive real numbers. Given that $\angle RPO = \angle RQO = \frac{\pi}{3}$ and $|\vec{PR}| = |\vec{QR}| = 5$, find the coordinates of R .



- Calculate $|OP|$ and $|OQ|$ to recognise any special triangle.
- Use the fact that equal sides with included 60° imply equilateral properties.
- Set up three sphere-distance equations for $R(x, y, z)$ from O, P, Q and subtract to get linear relations for x, y .

Hint:

Solution 3.16: Sketch

Compute $|OP| = \sqrt{4^2 + 3^2} = 5$ and $|OQ| = 5$. Since $|PR| = 5$ and $\angle RPO = 60^\circ$, triangle RPO is equilateral so $|OR| = 5$. Thus

$$\begin{aligned}x^2 + y^2 + z^2 &= 25, \\(x - 4)^2 + (y - 3)^2 + z^2 &= 25, \\(x - 3)^2 + (y - 4)^2 + z^2 &= 25.\end{aligned}$$

Subtracting the first from the second and third gives the linear system

$$\begin{aligned}8x + 6y &= 25, \\6x + 8y &= 25,\end{aligned}$$

so by symmetry $x = y = \frac{25}{14}$. Substituting into $x^2 + y^2 + z^2 = 25$ yields $z = \frac{5\sqrt{47}}{7}$.

Takeaways

- Intersection of spheres often reduces 3D location problems to solving linear systems by subtraction.
- Symmetry in the base points can force $x = y$ and simplify algebra.

Problem 3.17: Perpendicular Intersecting Lines

Consider the line $L_1 : \mathbf{r} = \begin{pmatrix} 3 \\ -1 \\ 7 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ and the line $L_2 : \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ p \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ q \\ -1 \end{pmatrix}$.

Given that these two lines intersect and are perpendicular to each other, find the values of p and q .

(Note: Two conditions must be satisfied: the lines must intersect at some point, and their direction vectors must be perpendicular.)

Hint: Use perpendicularity condition (dot product = 0) and intersection condition (solve system).

Solution 3.17: Sketch

Step 1: Find q from perpendicularity. Direction vectors: $\mathbf{d}_1 = (1, 2, 1)$ and $\mathbf{d}_2 = (2, q, -1)$. Since $\mathbf{d}_1 \cdot \mathbf{d}_2 = 0$: $(1)(2) + (2)(q) + (1)(-1) = 0 \Rightarrow 2q = -1 \Rightarrow q = -\frac{1}{2}$.

Step 2: Set up intersection condition. The lines intersect when $\begin{pmatrix} 3 \\ -1 \\ 7 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ p \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1/2 \\ -1 \end{pmatrix}$. This gives: $3 + \mu = 1 + 2\lambda$, $-1 + 2\mu = -\frac{1}{2}\lambda$, and $7 + \mu = p - \lambda$.

Step 3: Solve for μ and λ . From first equation: $\mu = 2\lambda - 2$. Substituting into second: $-1 + 2(2\lambda - 2) = -\frac{1}{2}\lambda \Rightarrow \frac{9}{2}\lambda = 5 \Rightarrow \lambda = \frac{10}{9}$. Thus $\mu = \frac{2}{9}$.

Step 4: Find p . From third equation: $p = 7 + \frac{2}{9} + \frac{10}{9} = 7 + \frac{4}{3} = \frac{25}{3}$.

Answer: $p = \frac{25}{3}$ and $q = -\frac{1}{2}$.

Problem 3.18: Tetrahedron Collinearity

- (i) The two non-parallel vectors $\vec{u}$ and $\vec{v}$ satisfy $\lambda\vec{u} + \mu\vec{v} = \vec{0}$ for some real numbers λ and μ .

Show that $\lambda = \mu = 0$.


- (ii) The two non-parallel vectors $\vec{u}$ and $\vec{v}$ satisfy $\lambda_1\vec{u} + \mu_1\vec{v} = \lambda_2\vec{u} + \mu_2\vec{v}$ for some real numbers $\lambda_1, \lambda_2, \mu_1$ and μ_2 .

Using part (i), or otherwise, show that $\lambda_1 = \lambda_2$ and $\mu_1 = \mu_2$.

The diagram below shows a tetrahedron with vertices A, B, C and S .

The point K is defined by $S\vec{K} = \frac{1}{4}S\vec{B} + \frac{1}{3}S\vec{C}$, as shown in the diagram.

The point L is the point of intersection of the straight lines SK and BC .



- (iii) Using part (ii), or otherwise, determine the position of L by showing that $\vec{BL} = \frac{4}{7}\vec{BC}$.

- (iv) The point P is defined by $\vec{AP} = -6\vec{AB} - 8\vec{AC}$.

Does P lie on the line AL ? Justify your answer.


Hint: (i) Use proof by contradiction: if $\lambda \neq 0$, then $\vec{u} = -\frac{\mu}{\lambda}\vec{v}$, contradicting non-parallel assumption. (ii) Rearrange to get $(\lambda_1 - \lambda_2)\vec{u} + (\mu_1 - \mu_2)\vec{v} = \vec{0}$ and apply part (i). (iii) Express $S\vec{L}$ in two ways: as $mS\vec{K}$ and as $(1-k)S\vec{B} + kS\vec{C}$, then equate coefficients. (iv) Express $\vec{AL}$ in terms of $\vec{AB}$ and $\vec{AC}$, then check if $\vec{AP} = c\vec{AL}$ for some scalar c .

Solution 3.18: Sketch


- (i) Assume $\lambda \neq 0$. Then $\vec{u} = -\frac{\mu}{\lambda}\vec{v}$, implying $\vec{u} \parallel \vec{v}$, contradiction. Thus $\lambda = 0$, which gives $\mu = 0$.
 (ii) Rearranging: $(\lambda_1 - \lambda_2)\vec{u} + (\mu_1 - \mu_2)\vec{v} = \vec{0}$. By part (i), both coefficients equal zero.
 (iii) Since S, K, L are collinear: $S\vec{L} = mS\vec{K} = \frac{m}{4}S\vec{B} + \frac{m}{3}S\vec{C}$. Since L is on BC : $S\vec{L} = (1-k)S\vec{B} + kS\vec{C}$ where $\vec{BL} = k\vec{BC}$. Equating coefficients: $\frac{m}{4} = 1-k$ and $\frac{m}{3} = k$. Solving: $m = \frac{12}{7}$ and $k = \frac{4}{7}$.
 (iv) $\vec{AL} = \vec{AB} + \frac{4}{7}\vec{BC} = \vec{AB} + \frac{4}{7}(\vec{AC} - \vec{AB}) = \frac{3}{7}\vec{AB} + \frac{4}{7}\vec{AC}$. Check: $\vec{AP} = -6\vec{AB} - 8\vec{AC} = -14(\frac{3}{7}\vec{AB} + \frac{4}{7}\vec{AC}) = -14\vec{AL}$. Yes, P lies on line AL (extended backwards through A).

Problem 3.19: Varignon's Theorem - Midpoint Parallelogram

Let $ABCD$ be any quadrilateral in three-dimensional space (not necessarily planar). Let P , Q , R , and S be the midpoints of sides AB , BC , CD , and DA respectively.



3D Non-Planar Quadrilateral:



- (i) Express the position vectors of P , Q , R , and S in terms of the position vectors $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$, and $\mathbf{d}$ of points A , B , C , and D respectively.
- (ii) Show that $\overrightarrow{PQ} = \overrightarrow{SR}$.
- (iii) Hence, prove that $PQRS$ is a parallelogram.
- (iv) If $ABCD$ is a parallelogram, what special type of quadrilateral is $PQRS$? Justify your answer.

Note: This result is known as Varignon's Theorem (1731). It states that the midpoints of any quadrilateral always form a parallelogram, regardless of the shape of the original quadrilateral.

Hint:(i) Use the midpoint formula: the position vector of the midpoint of two points is the average of their position vectors. (ii) Express both $\overrightarrow{PQ}$ and $\overrightarrow{SR}$ in terms of $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$, and $\mathbf{d}$, then show they are equal. (iii) Use the result from (ii) - if one pair of opposite sides is equal and parallel, then $PQRS$ is a parallelogram. (iv) When $ABCD$ is a parallelogram, use $\overrightarrow{DC} = \overrightarrow{AB}$ and investigate the relationship between adjacent sides of $PQRS$.

Solution 3.19: Sketch

(i) Using the midpoint formula:

$$\begin{aligned} \mathbf{p} &= \frac{\mathbf{a} + \mathbf{b}}{2} \\ \mathbf{q} &= \frac{\mathbf{b} + \mathbf{c}}{2} \\ \mathbf{r} &= \frac{\mathbf{c} + \mathbf{d}}{2} \\ \mathbf{s} &= \frac{\mathbf{d} + \mathbf{a}}{2} \end{aligned}$$

(ii) Calculate $\overrightarrow{PQ}$:

$$\begin{aligned} \overrightarrow{PQ} &= \mathbf{q} - \mathbf{p} = \frac{\mathbf{b} + \mathbf{c}}{2} - \frac{\mathbf{a} + \mathbf{b}}{2} \\ &= \frac{\mathbf{b} + \mathbf{c} - \mathbf{a} - \mathbf{b}}{2} = \frac{\mathbf{c} - \mathbf{a}}{2} \end{aligned}$$

Calculate $\overrightarrow{SR}$:

$$\begin{aligned} \overrightarrow{SR} &= \mathbf{r} - \mathbf{s} = \frac{\mathbf{c} + \mathbf{d}}{2} - \frac{\mathbf{d} + \mathbf{a}}{2} \\ &= \frac{\mathbf{c} + \mathbf{d} - \mathbf{d} - \mathbf{a}}{2} = \frac{\mathbf{c} - \mathbf{a}}{2} \end{aligned}$$

Therefore $\overrightarrow{PQ} = \overrightarrow{SR}$.

(iii) Since $\overrightarrow{PQ} = \overrightarrow{SR}$, the sides PQ and SR are equal in length and parallel. By definition, a quadrilateral with one pair of opposite sides equal and parallel is a parallelogram. Therefore $PQRS$ is a parallelogram.

(iv) If $ABCD$ is a parallelogram, then $\overrightarrow{AB} = \overrightarrow{DC}$, which means $\mathbf{b} - \mathbf{a} = \mathbf{c} - \mathbf{d}$.

Calculate $\overrightarrow{PS}$:

$$\overrightarrow{PS} = \mathbf{s} - \mathbf{p} = \frac{\mathbf{d} + \mathbf{a}}{2} - \frac{\mathbf{a} + \mathbf{b}}{2} = \frac{\mathbf{d} - \mathbf{b}}{2}$$

Calculate $\overrightarrow{QR}$:

$$\overrightarrow{QR} = \mathbf{r} - \mathbf{q} = \frac{\mathbf{c} + \mathbf{d}}{2} - \frac{\mathbf{b} + \mathbf{c}}{2} = \frac{\mathbf{d} - \mathbf{b}}{2}$$

So $\overrightarrow{PS} = \overrightarrow{QR}$, meaning both pairs of opposite sides are equal and parallel.

Additionally, from (ii): $|\overrightarrow{PQ}| = \frac{1}{2}|\mathbf{c} - \mathbf{a}| = \frac{1}{2}|\overrightarrow{AC}|$ (half the diagonal AC).

Similarly: $|\overrightarrow{PS}| = \frac{1}{2}|\mathbf{d} - \mathbf{b}| = \frac{1}{2}|\overrightarrow{BD}|$ (half the diagonal BD).

If $ABCD$ is a parallelogram with equal diagonals, then $PQRS$ is a rhombus (all four sides equal). If the diagonals of $ABCD$ are perpendicular, then $PQRS$ is a rectangle.

Problem 3.20: Force Vector Analysis

A particle is subject to two forces and undergoes a displacement in three-dimensional space.

- (i) Force $\mathbf{F}_1$ has magnitude 12 newtons in the direction of vector $2\mathbf{i} - 2\mathbf{j} + \mathbf{k}$.

Show that $\mathbf{F}_1 = 8\mathbf{i} - 8\mathbf{j} + 4\mathbf{k}$ newtons.

- (ii) Force $\mathbf{F}_1$ from part (i) and a second force, $\mathbf{F}_2 = -6\mathbf{i} + 12\mathbf{j} + 4\mathbf{k}$ newtons, both act upon the particle.

Show that the resultant force acting on the particle is given by:

$$\mathbf{F}_3 = 2\mathbf{i} + 4\mathbf{j} + 8\mathbf{k} \text{ newtons}$$

- (iii) The particle moves through a displacement $\mathbf{d} = \mathbf{i} + \mathbf{j} + 2\mathbf{k}$ metres under the action of the resultant force $\mathbf{F}_3$ from part (ii).

Calculate the work done by the resultant force on the particle.

Note: Here $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ are the standard unit vectors in the x , y , and z directions respectively. Work done is calculated using $W = \mathbf{F} \cdot \mathbf{d}$ and is measured in joules (J) when force is in newtons (N) and displacement is in metres (m).

Hint: (i) Find the unit vector in the given direction by dividing by its magnitude, then multiply by 12 to get the force vector. (ii) Add the two force vectors component-wise to find the resultant. (iii) Use the dot product formula $W = \mathbf{F} \cdot \mathbf{d}$ to compute the work done.

Solution 3.20: Sketch

- (i) Direction vector: $\mathbf{v} = 2\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ has magnitude $|\mathbf{v}| = \sqrt{2^2 + (-2)^2 + 1^2} = \sqrt{4 + 4 + 1} = \sqrt{9} = 3$.
Unit vector in this direction: $\hat{\mathbf{u}} = \frac{1}{3}(2\mathbf{i} - 2\mathbf{j} + \mathbf{k})$.
Force with magnitude 12 N:

$$\mathbf{F}_1 = 12\hat{\mathbf{u}} = 12 \times \frac{1}{3}(2\mathbf{i} - 2\mathbf{j} + \mathbf{k}) = 4(2\mathbf{i} - 2\mathbf{j} + \mathbf{k}) = 8\mathbf{i} - 8\mathbf{j} + 4\mathbf{k} \text{ N}$$

- (ii) Resultant force (vector addition):

$$\begin{aligned}\mathbf{F}_3 &= \mathbf{F}_1 + \mathbf{F}_2 \\ &= (8\mathbf{i} - 8\mathbf{j} + 4\mathbf{k}) + (-6\mathbf{i} + 12\mathbf{j} + 4\mathbf{k}) \\ &= (8 - 6)\mathbf{i} + (-8 + 12)\mathbf{j} + (4 + 4)\mathbf{k} \\ &= 2\mathbf{i} + 4\mathbf{j} + 8\mathbf{k} \text{ N}\end{aligned}$$

- (iii) Work done is $W = \mathbf{F}_3 \cdot \mathbf{d}$:

$$\begin{aligned}W &= (2\mathbf{i} + 4\mathbf{j} + 8\mathbf{k}) \cdot (\mathbf{i} + \mathbf{j} + 2\mathbf{k}) \\ &= (2)(1) + (4)(1) + (8)(2) \\ &= 2 + 4 + 16 \\ &= 22 \text{ J}\end{aligned}$$

The work done by the resultant force is 22 joules.

Problem 3.21: Double Angle with Vectors

Consider the vectors $\mathbf{a} = \mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$ and $\mathbf{b} = 2\mathbf{i} - 4\mathbf{j} + 4\mathbf{k}$.

Let θ be the acute angle between these two vectors. Find the exact value of $\sin 2\theta$.

Note: Here $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ are the standard unit vectors in the x , y , and z directions respectively.

Hint: Find $\cos \theta$ from dot product, then $\sin \theta$ from Pythagorean identity, and use $\sin 2\theta = 2 \sin \theta \cos \theta$.

Solution 3.21: Sketch

$|\mathbf{a}| = 3$, $|\mathbf{b}| = 6$, $\mathbf{a} \cdot \mathbf{b} = 2$. Thus $\cos \theta = \frac{2}{18} = \frac{1}{9}$. Then $\sin \theta = \sqrt{1 - \frac{1}{81}} = \frac{4\sqrt{5}}{9}$. Therefore $\sin 2\theta = 2 \cdot \frac{4\sqrt{5}}{9} \cdot \frac{1}{9} = \frac{8\sqrt{5}}{81}$.

Problem 3.22: Vector Projection

Find the integer value of m such that the projection of $\mathbf{a} = 2\mathbf{i} - 3\mathbf{j} + \mathbf{k}$ onto $\mathbf{b} = \mathbf{i} + m\mathbf{j} - \mathbf{k}$ equals $-\frac{11}{18}\mathbf{b}$.

Note: Here $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ are the standard unit vectors in the x , y , and z directions respectively.

Hint: Use projection formula: $\text{proj}_{\mathbf{b}} \mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{b}|^2} \mathbf{b}$.

Solution 3.22: Sketch

$\mathbf{a} \cdot \mathbf{b} = 2 - 3m - 1 = 1 - 3m$, $|\mathbf{b}|^2 = 1 + m^2 + 1 = m^2 + 2$. Set $\frac{1-3m}{m^2+2} = -\frac{11}{18}$. Cross-multiply: $18(1 - 3m) = -11(m^2 + 2)$, giving $11m^2 - 54m + 40 = 0$. Factor: $(11m - 10)(m - 4) = 0$. Since m is an integer, $m = 4$.

Problem 3.23: Perpendicular Vectors Condition

For $\mathbf{u} = -2\mathbf{i} - \mathbf{j} + 3\mathbf{k}$ and $\mathbf{v} = p\mathbf{i} + \mathbf{j} + 2\mathbf{k}$, find values of p such that $\mathbf{u} - \mathbf{v}$ and $\mathbf{u} + \mathbf{v}$ are perpendicular.

Note: Here $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ are the standard unit vectors in the x , y , and z directions respectively.

Hint: Use $(\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = |\mathbf{u}|^2 - |\mathbf{v}|^2 = 0$.

Solution 3.23: Sketch

$|\mathbf{u}|^2 = 4 + 1 + 9 = 14$. $|\mathbf{v}|^2 = p^2 + 1 + 4 = p^2 + 5$. Setting equal: $14 = p^2 + 5 \Rightarrow p^2 = 9 \Rightarrow p = \pm 3$.

Problem 3.24: Parallel and Perpendicular Lines

For lines with direction vectors $\begin{pmatrix} 3 \\ -3 \\ 3 \end{pmatrix}$ and $\begin{pmatrix} a-2 \\ -7 \\ 7 \end{pmatrix}$, find a if they are: (a) parallel, (b) perpendicular.

Hint: Parallel: direction vectors are scalar multiples. Perpendicular: dot product equals zero.

Solution 3.24: Sketch

(a) For parallel: $\frac{3}{a-2} = \frac{-3}{-7} = \frac{3}{7}$. Solving: $3 \cdot 7 = 3(a-2) \Rightarrow a = 9$. (b) For perpendicular: $3(a-2) + (-3)(-7) + 3(7) = 0 \Rightarrow 3a + 36 = 0 \Rightarrow a = -12$.

Problem 3.25: Angle BCD Using Dot Product

Relative to a fixed origin, the points B , C , and D are defined respectively by the position vectors:


$$\mathbf{b} = \mathbf{i} - \mathbf{j} + 2\mathbf{k}$$

$$\mathbf{c} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$$

$$\mathbf{d} = a\mathbf{i} - 2\mathbf{j}$$

where a is a real constant.

Given that the magnitude of angle BCD is $\frac{\pi}{3}$ (i.e., $\angle BCD = \frac{\pi}{3}$), find the value of a .



(Note: The angle BCD is the angle at vertex C between rays CB and CD .)

Hint: Find vectors $\overrightarrow{CB}$ and $\overrightarrow{CD}$ emanating from vertex C , then use the dot product formula $\cos \theta = \frac{\overrightarrow{u} \cdot \overrightarrow{v}}{|\overrightarrow{u}| |\overrightarrow{v}|}$ with $\cos(\pi/3) = 1/2$. Be careful to check that your solution gives a positive value for $1 - a$.

Solution 3.25: Sketch

Step 1: Find vectors from C : $\overrightarrow{CB} = \mathbf{b} - \mathbf{c} = -\mathbf{i} + \mathbf{k}$ and $\overrightarrow{CD} = \mathbf{d} - \mathbf{c} = (a - 2)\mathbf{i} - \mathbf{j} - \mathbf{k}$.
 Step 2: Calculate magnitudes: $|\overrightarrow{CB}| = \sqrt{2}$ and $|\overrightarrow{CD}| = \sqrt{(a - 2)^2 + 2}$.
 Step 3: Dot product: $\overrightarrow{CB} \cdot \overrightarrow{CD} = -(a - 2) + 0 - 1 = 1 - a$.
 Step 4: Use formula with $\cos(\pi/3) = 1/2$: $1 - a = \sqrt{2} \cdot \sqrt{(a - 2)^2 + 2} \cdot \frac{1}{2}$.
 Step 5: Square both sides: $4(1 - a)^2 = 2[(a - 2)^2 + 2]$, which simplifies to $a^2 = 4$, giving $a = \pm 2$.
 Step 6: Check constraint: Since the RHS is positive, we need $1 - a > 0$, so $a < 1$. Therefore $a = -2$.

Problem 3.26: Direction Cosines — Compute and Verify

Let $\mathbf{r} = 6\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$. The **direction cosines** of $\mathbf{r}$ are

$$l = \cos \alpha, \quad m = \cos \beta, \quad n = \cos \gamma,$$

where α , β , and γ are the angles between $\mathbf{r}$ and the positive x -, y -, and z -axes.
 Find l , m , and n , and verify that $l^2 + m^2 + n^2 = 1$.

Hint: First find $|\mathbf{r}|$, then use $l = \frac{|\mathbf{r}|}{6}$, $m = \frac{|\mathbf{r}|}{2}$, $n = \frac{|\mathbf{r}|}{3}$ (see the direction-cosines proof in Problem 3.9).

Solution 3.26: Sketch

$$|\mathbf{r}| = \sqrt{6^2 + 2^2 + 3^2} = \sqrt{49} = 7.$$

The direction cosines are the component-to-length ratios:

$$l = \frac{6}{7}, \quad m = \frac{2}{7}, \quad n = \frac{3}{7}.$$

Check the identity:

$$l^2 + m^2 + n^2 = \frac{36 + 4 + 9}{49} = \frac{49}{49} = 1.$$

So the direction cosines are $\left(\frac{6}{7}, \frac{2}{7}, \frac{3}{7}\right)$, and they satisfy $l^2 + m^2 + n^2 = 1$ as expected from Problem 3.9.

Problem 3.27: Section Formula Proof

Prove that if C divides AB in ratio $m : n$, then $\overrightarrow{OC} = \frac{m\mathbf{a} + n\mathbf{b}}{m + n}$ where $\mathbf{a} = \overrightarrow{OA}$, $\mathbf{b} = \overrightarrow{OB}$.

Hint: Use $\overrightarrow{AC} = \frac{m}{n} \overrightarrow{AB}$ and vector addition.

Solution 3.27: Sketch

Since $\frac{CB}{AC} = \frac{m}{n}$, we have $\overrightarrow{AC} = \frac{n}{m+n}(\mathbf{b} - \mathbf{a})$. Then $\overrightarrow{OC} = \mathbf{a} + \overrightarrow{AC} = \mathbf{a} + \frac{n}{m+n}(\mathbf{b} - \mathbf{a}) = \frac{m\mathbf{a} + n\mathbf{b}}{m+n}$.

Problem 3.28: Skew or Intersecting Lines

Consider the two lines:

$$L_1 : \quad \mathbf{r} = \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 4 \\ 1 \end{pmatrix}$$

$$L_2 : \quad \mathbf{r} = \begin{pmatrix} -4 \\ 6 \\ -2 \end{pmatrix} + \mu \begin{pmatrix} 4 \\ -11 \\ 3 \end{pmatrix}$$

Determine whether lines L_1 and L_2 are skew or intersecting.

Hint: Check if direction vectors are parallel. If not, solve system to see if it's consistent.

Solution 3.28: Sketch

Direction vectors not parallel (check ratios). Solve system: from equations 1 and 3, find $\lambda = 6$, $\mu = 3$. Check equation 2: $4(6) + 11(3) = 57 \neq 9$. System inconsistent, so lines are skew.

Problem 3.29: Line Through Points, Intersection Check

A line passes through the points $A(1, 3, -2)$ and $B(2, -1, 5)$.

(i) Show that the vector equation of the line AB is given by:

$$\mathbf{r} = (\mathbf{i} + 3\mathbf{j} - 2\mathbf{k}) + \lambda_1(\mathbf{i} - 4\mathbf{j} + 7\mathbf{k}), \quad \lambda_1 \in \mathbb{R}$$

(ii) Determine if the point $C(3, 4, 9)$ lies on the line.

(iii) Consider a second line with parametric equations $x = 1 - \lambda_2$, $y = 2 + 3\lambda_2$, $z = -1 + \lambda_2$.

Assuming this line is neither parallel nor perpendicular to AB , determine whether the two lines intersect or are skew.

Hint: (i) Find the direction vector $\overrightarrow{AB}$ and use the point A to write the vector equation. (ii) Substitute the coordinates of C into the line equation and check if a consistent value of λ_1 exists for all three components. (iii) Set the two lines equal component-wise, solve for both parameters, then verify if the solution is consistent across all three equations.

Solution 3.29: Sketch

(i) Direction vector: $\overrightarrow{AB} = \mathbf{b} - \mathbf{a} = \begin{pmatrix} 2-1 \\ -1-3 \\ 5-(-2) \end{pmatrix} = \begin{pmatrix} 1 \\ -4 \\ 7 \end{pmatrix} = \mathbf{i} - 4\mathbf{j} + 7\mathbf{k}$.

Using point A as the position vector: $\mathbf{r} = (\mathbf{i} + 3\mathbf{j} - 2\mathbf{k}) + \lambda_1(\mathbf{i} - 4\mathbf{j} + 7\mathbf{k})$.

(ii) If C lies on the line: $\begin{pmatrix} 3 \\ 4 \\ 9 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ -2 \end{pmatrix} + \lambda_1 \begin{pmatrix} 1 \\ -4 \\ 7 \end{pmatrix}$.

From x -component: $3 = 1 + \lambda_1 \Rightarrow \lambda_1 = 2$.

Check y -component: $y = 3 - 4(2) = -5 \neq 4$. Since the y -coordinate doesn't match, point C does not lie on the line.

(iii) Line 1: $\mathbf{r}_1 = \begin{pmatrix} 1 \\ 3 \\ -2 \end{pmatrix} + \lambda_1 \begin{pmatrix} 1 \\ -4 \\ 7 \end{pmatrix}$. Line 2: $\mathbf{r}_2 = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + \lambda_2 \begin{pmatrix} -1 \\ 3 \\ 1 \end{pmatrix}$.

Equate components: From x : $1 + \lambda_1 = 1 - \lambda_2 \Rightarrow \lambda_1 = -\lambda_2$. From y : $3 - 4\lambda_1 = 2 + 3\lambda_2$. Substitute $\lambda_1 = -\lambda_2$: $3 + 4\lambda_2 = 2 + 3\lambda_2 \Rightarrow \lambda_2 = -1$, so $\lambda_1 = 1$.

Check z -component: LHS: $-2 + 7(1) = 5$. RHS: $-1 + (-1) = -2$. Since $5 \neq -2$, the system is inconsistent. The lines do not intersect and are not parallel, therefore they are skew.

Problem 3.30: Line-Sphere Intersection Angle

Let S be the sphere with center at the origin and equation $x^2 + y^2 + z^2 = 10$.

Let ℓ be the line with equation:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + \lambda \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$$

(i) Find A and B , the points of intersection of S and ℓ .

(ii) Find $\angle AOB$ to the nearest degree, where O is the origin.

Hint: (i) Write the line in parametric form ($x = 1, y = 2 + \lambda, z = 1 - \lambda$), substitute into the sphere equation, and solve the resulting quadratic equation for λ . (ii) Use the dot product formula $\cos \theta = \frac{\overrightarrow{OA} \cdot \overrightarrow{OB}}{|\overrightarrow{OA}| |\overrightarrow{OB}|}$ to find the angle between the two position vectors.

Solution 3.30: Sketch

(i) Parametric form: $x = 1, y = 2 + \lambda, z = 1 - \lambda$.

Substitute into $x^2 + y^2 + z^2 = 10$: $1 + (2 + \lambda)^2 + (1 - \lambda)^2 = 10$
 $1 + 4 + 4\lambda + \lambda^2 + 1 - 2\lambda + \lambda^2 = 10$
 $2\lambda^2 + 2\lambda + 6 = 10$
 $\lambda^2 + \lambda - 2 = 0$
 $(\lambda + 2)(\lambda - 1) = 0$

So $\lambda = 1$ or $\lambda = -2$.

When $\lambda = 1$: $A(1, 3, 0)$. When $\lambda = -2$: $B(1, 0, 3)$.

(ii) $\overrightarrow{OA} = \begin{pmatrix} 1 \\ 3 \\ 0 \end{pmatrix}, \overrightarrow{OB} = \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}$.

$\overrightarrow{OA} \cdot \overrightarrow{OB} = (1)(1) + (3)(0) + (0)(3) = 1$.

$|\overrightarrow{OA}| = \sqrt{1^2 + 3^2 + 0^2} = \sqrt{10}, |\overrightarrow{OB}| = \sqrt{1^2 + 0^2 + 3^2} = \sqrt{10}$.

$\cos \theta = \frac{1}{\sqrt{10} \cdot \sqrt{10}} = \frac{1}{10} = 0.1$.

Therefore $\theta = \cos^{-1}(0.1) \approx 84.26^\circ \approx 84^\circ$.

Problem 3.31: Linear Combination of Vectors

Consider the following four vectors in three-dimensional space:

$$\mathbf{a} = 2\mathbf{i} + 3\mathbf{j} + \mathbf{k}$$

$$\mathbf{b} = \mathbf{i} - \mathbf{j} + 2\mathbf{k}$$

$$\mathbf{c} = -\mathbf{i} + 2\mathbf{j} - \mathbf{k}$$

$$\mathbf{u} = 5\mathbf{i} + 5\mathbf{j} + 5\mathbf{k}$$

Find the values of the scalars λ , μ , and ν that satisfy the equation:

$$\mathbf{u} = \lambda\mathbf{a} + \mu\mathbf{b} + \nu\mathbf{c}$$

Note: This means expressing vector $\mathbf{u}$ as a linear combination of vectors $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$. Any vector in 3D space can be written as a combination of three non-coplanar (non-parallel) vectors.

Hint: Substitute the given vectors into the equation $\mathbf{u} = \lambda\mathbf{a} + \mu\mathbf{b} + \nu\mathbf{c}$, then equate the coefficients of $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ to form a system of three linear equations. Solve the system by elimination.

Solution 3.31: Sketch

Substituting: $5\mathbf{i} + 5\mathbf{j} + 5\mathbf{k} = \lambda(2\mathbf{i} + 3\mathbf{j} + \mathbf{k}) + \mu(\mathbf{i} - \mathbf{j} + 2\mathbf{k}) + \nu(-\mathbf{i} + 2\mathbf{j} - \mathbf{k})$.

Equating coefficients gives the system:

$$2\lambda + \mu - \nu = 5 \quad (1)$$

$$3\lambda - \mu + 2\nu = 5 \quad (2)$$

$$\lambda + 2\mu - \nu = 5 \quad (3)$$

Subtract (1) from (3): $-\lambda + \mu = 0 \Rightarrow \lambda = \mu$.

Substitute $\mu = \lambda$ into (2): $3\lambda - \lambda + 2\nu = 5 \Rightarrow 2\lambda + 2\nu = 5 \Rightarrow \lambda + \nu = \frac{5}{2}$.

Substitute $\mu = \lambda$ into (1): $2\lambda + \lambda - \nu = 5 \Rightarrow 3\lambda - \nu = 5$.

Add these two results: $4\lambda = \frac{5}{2} + 5 = \frac{15}{2} \Rightarrow \lambda = \frac{15}{8}$.

Since $\lambda = \mu$: $\mu = \frac{15}{8}$.

From $3\lambda - \nu = 5$: $\nu = 3 \cdot \frac{15}{8} - 5 = \frac{45}{8} - \frac{40}{8} = \frac{5}{8}$.

Therefore: $\lambda = \frac{15}{8}$, $\mu = \frac{15}{8}$, $\nu = \frac{5}{8}$.


Takeaways 3.2

Key Concepts:

- **Linear Combination of Vectors:** Any vector can potentially be expressed as a linear combination of other vectors: $\mathbf{v} = \lambda \mathbf{u}_1 + \mu \mathbf{u}_2 + \nu \mathbf{u}_3$. The scalars λ, μ, ν are the coefficients.
- **Equating Coefficients Method:** When vectors are expressed in component form (using $\mathbf{i}, \mathbf{j}, \mathbf{k}$), equate the coefficients of each unit vector separately to create a system of linear equations.
- **Solving Linear Systems:** The systematic approach involves:
 1. Look for relationships by adding/subtracting equations
 2. Find simple relationships (e.g., $\lambda = \mu$) to reduce variables
 3. Substitute to eliminate variables progressively
 4. Solve for remaining variables and back-substitute
- **Verification Strategy:** Always substitute final values back into the original equations to verify correctness. Here: $2\left(\frac{15}{8}\right) + \frac{15}{8} - \frac{5}{8} = \frac{30+15-5}{8} = \frac{40}{8} = 5 \checkmark$
- **Linear Independence:** The fact that a unique solution exists ($\lambda = \mu = \frac{15}{8}, \nu = \frac{5}{8}$) indicates that the three given vectors are **linearly independent**—no one of them can be written as a linear combination of the others—and therefore span $\mathbb{R}^3$. Any vector in 3D space can be expressed as a linear combination of these three vectors. See Appendix A.7 for the formal definition and geometric meaning.

Problem 3.32: Parallelogram Fourth Vertex

Three vertices of a parallelogram are $O(0,0,0)$, $A(2,2,1)$, $B(1,2,2)$. Find all possible positions of the fourth vertex.



Three possible parallelograms with vertices O , A , B
(shown in blue, red, and green)

Hint: Consider three cases: which point is opposite to which, giving three parallelograms.

Solution 3.32: Sketch

Given three vertices $O(0, 0, 0)$, $A(2, 2, 1)$, and $B(1, 2, 2)$, we need to find all possible fourth vertices C such that $OABC$ forms a parallelogram (in some order).

In a parallelogram, opposite sides are equal and parallel. This gives us three distinct cases depending on which vertex is opposite to which.

Case 1: C_1 opposite to O (parallelogram $OABC_1$ with $OA \parallel BC_1$ and $OB \parallel AC_1$)

Using the parallelogram property: $\overrightarrow{OC_1} = \overrightarrow{OA} + \overrightarrow{OB}$.

Calculate:

$$\begin{aligned}\overrightarrow{OA} &= (2, 2, 1) \\ \overrightarrow{OB} &= (1, 2, 2) \\ \overrightarrow{OC_1} &= (2, 2, 1) + (1, 2, 2) = (3, 4, 3)\end{aligned}$$

Therefore, $C_1 = (3, 4, 3)$.

Verification: $\overrightarrow{OA} = (2, 2, 1)$ and $\overrightarrow{BC_1} = (3, 4, 3) - (1, 2, 2) = (2, 2, 1) \checkmark$

Case 2: C_2 opposite to B (parallelogram OAC_2B with $OA \parallel C_2B$ and $OC_2 \parallel AB$)

Using the parallelogram property: $\overrightarrow{OC_2} = \overrightarrow{OA} - \overrightarrow{OB}$.

Calculate:

$$\overrightarrow{OC_2} = (2, 2, 1) - (1, 2, 2) = (1, 0, -1)$$

Therefore, $C_2 = (1, 0, -1)$.

Verification: $\overrightarrow{OA} = (2, 2, 1)$ and $\overrightarrow{C_2B} = (1, 2, 2) - (1, 0, -1) = (0, 2, 3) \dots$ Actually, $\overrightarrow{AC_2} = (1, 0, -1) - (2, 2, 1) = (-1, -2, -2)$ and $\overrightarrow{OB} = (1, 2, 2)$. Let me verify: $\overrightarrow{OC_2} + \overrightarrow{OB} = (1, 0, -1) + (1, 2, 2) = (2, 2, 1) = \overrightarrow{OA} \checkmark$

Case 3: C_3 opposite to A (parallelogram OBC_3A with $OB \parallel C_3A$ and $OC_3 \parallel BA$)

Using the parallelogram property: $\overrightarrow{OC_3} = \overrightarrow{OB} - \overrightarrow{OA}$.

Calculate:

$$\overrightarrow{OC_3} = (1, 2, 2) - (2, 2, 1) = (-1, 0, 1)$$

Therefore, $C_3 = (-1, 0, 1)$.

Verification: $\overrightarrow{OB} = (1, 2, 2)$ and $\overrightarrow{C_3A} = (2, 2, 1) - (-1, 0, 1) = (3, 2, 0) \dots$ Let me verify: $\overrightarrow{OC_3} + \overrightarrow{OA} = (-1, 0, 1) + (2, 2, 1) = (1, 2, 2) = \overrightarrow{OB} \checkmark$

Final Answer: The three possible positions for the fourth vertex are:

$$C_1 = (3, 4, 3), \quad C_2 = (1, 0, -1), \quad C_3 = (-1, 0, 1)$$

3.3 Part 2 Advanced Problems

Problem 3.33: Triangle Inequality and Cauchy-Schwarz

Let $P(x, y, z)$ be a point on the unit sphere centered at the origin, so that $x^2 + y^2 + z^2 = 1$. Prove:

(i) $|x| + |y| + |z| \geq 1$

(ii) For any two vectors $\mathbf{a} = (a_1, a_2, a_3)$ and $\mathbf{b} = (b_1, b_2, b_3)$, the Cauchy-Schwarz inequality states:

$$|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}| \cdot |\mathbf{b}|$$

Equivalently: $|a_1b_1 + a_2b_2 + a_3b_3| \leq \sqrt{a_1^2 + a_2^2 + a_3^2} \cdot \sqrt{b_1^2 + b_2^2 + b_3^2}$

Derive this inequality from the dot product formula.

(iii) $|x| + |y| + |z| \leq \sqrt{3}$


Hint: Use triangle inequality for part (i) and Cauchy-Schwarz with vector $(1, 1, 1)$ for part (iii).

Solution 3.33: Sketch

(i) By triangle inequality: $1 = |\mathbf{r}| \leq |x| + |y| + |z|$. (ii) From dot product: $|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}||\mathbf{b}|$. (iii) Apply Cauchy-Schwarz with $\mathbf{a} = (|x|, |y|, |z|)$, $\mathbf{b} = (1, 1, 1)$: $|x| + |y| + |z| \leq \sqrt{x^2 + y^2 + z^2}\sqrt{3} = \sqrt{3}$.

Problem 3.34: Bimedians of Tetrahedron

On the triangular pyramid (tetrahedron) $ABCD$, L, M, N, P, Q, R are the midpoints of AB, AC, AD, CD, BD, BC respectively.



Note: The three colored arrows show the bimedians $\overrightarrow{LP}$ (red), $\overrightarrow{MQ}$ (green), and $\overrightarrow{NR}$ (violet). Each bimedian connects the midpoint of one edge to the midpoint of the opposite edge.

Let $\mathbf{b} = \overrightarrow{AB}$, $\mathbf{c} = \overrightarrow{AC}$ and $\mathbf{d} = \overrightarrow{AD}$.

(i) Show that $\overrightarrow{LP} = \frac{1}{2}(-\mathbf{b} + \mathbf{c} + \mathbf{d})$.

(ii) It can be shown that:

$$\overrightarrow{MQ} = \frac{1}{2}(\mathbf{b} - \mathbf{c} + \mathbf{d}) \quad \text{and} \quad \overrightarrow{NR} = \frac{1}{2}(\mathbf{b} + \mathbf{c} - \mathbf{d})$$

(Do NOT prove these.)

Prove that:

$$|AB|^2 + |AC|^2 + |AD|^2 + |BC|^2 + |BD|^2 + |CD|^2 = 4(|LP|^2 + |MQ|^2 + |NR|^2)$$

Note: The segments LP , MQ , and NR are called the bimedians of the tetrahedron. This problem proves that the sum of the squares of all six edges equals four times the sum of the squares of the three bimedians.

Hint:(i) Express $\overrightarrow{AL}$ and $\overrightarrow{AP}$ in terms of $\mathbf{b}$, $\mathbf{c}$, and $\mathbf{d}$, then find $\overrightarrow{LP} = \overrightarrow{AP} - \overrightarrow{AL}$. (ii) Expand the left side as the sum of squared magnitudes of all six edges. Expand the right side by computing $4|LP|^2$, $4|MQ|^2$, and $4|NR|^2$ using dot products. Show both sides equal $3(b^2 + c^2 + d^2) - 2(\mathbf{b} \cdot \mathbf{c} + \mathbf{b} \cdot \mathbf{d} + \mathbf{c} \cdot \mathbf{d})$.

Solution 3.34: Sketch

(i) Since L is midpoint of AB : $\overrightarrow{AL} = \frac{1}{2}\mathbf{b}$. Since P is midpoint of CD : $\overrightarrow{AP} = \frac{1}{2}(\mathbf{c} + \mathbf{d})$. Therefore:
 $\overrightarrow{LP} = \overrightarrow{AP} - \overrightarrow{AL} = \frac{1}{2}(\mathbf{c} + \mathbf{d}) - \frac{1}{2}\mathbf{b} = \frac{1}{2}(-\mathbf{b} + \mathbf{c} + \mathbf{d})$.

(ii) LHS: The six edges are AB , AC , AD , $BC = \mathbf{c} - \mathbf{b}$, $BD = \mathbf{d} - \mathbf{b}$, $CD = \mathbf{d} - \mathbf{c}$.

$$\begin{aligned} LHS &= |\mathbf{b}|^2 + |\mathbf{c}|^2 + |\mathbf{d}|^2 + |\mathbf{c} - \mathbf{b}|^2 + |\mathbf{d} - \mathbf{b}|^2 + |\mathbf{d} - \mathbf{c}|^2 \\ &= 3(b^2 + c^2 + d^2) - 2(\mathbf{b} \cdot \mathbf{c} + \mathbf{b} \cdot \mathbf{d} + \mathbf{c} \cdot \mathbf{d}) \end{aligned}$$

RHS: Using part (i) and the given expressions:

$$4|LP|^2 = b^2 + c^2 + d^2 - 2\mathbf{b} \cdot \mathbf{c} - 2\mathbf{b} \cdot \mathbf{d} + 2\mathbf{c} \cdot \mathbf{d}$$

$$4|MQ|^2 = b^2 + c^2 + d^2 - 2\mathbf{b} \cdot \mathbf{c} + 2\mathbf{b} \cdot \mathbf{d} - 2\mathbf{c} \cdot \mathbf{d}$$

$$4|NR|^2 = b^2 + c^2 + d^2 + 2\mathbf{b} \cdot \mathbf{c} - 2\mathbf{b} \cdot \mathbf{d} - 2\mathbf{c} \cdot \mathbf{d}$$

Summing: $RHS = 3(b^2 + c^2 + d^2) - 2(\mathbf{b} \cdot \mathbf{c} + \mathbf{b} \cdot \mathbf{d} + \mathbf{c} \cdot \mathbf{d}) = LHS$.

Takeaways 3.3

Key Concepts and Connections:

- **Bimedians of a Tetrahedron:** The three bimedians are segments connecting midpoints of opposite edges in a tetrahedron. This problem proves a beautiful relationship: the sum of squares of all six edge lengths equals four times the sum of squares of the three bimedian lengths.
- **Connection to Apollonius's Theorem:** This result is a 3D generalization of Apollonius's Theorem. In 2D, Apollonius's Theorem states that for a triangle with sides a, b, c and median m_c to side c :

$$a^2 + b^2 = 2m_c^2 + \frac{c^2}{2}$$

Equivalently: $2(a^2 + b^2 + c^2) = 4(m_a^2 + m_b^2 + m_c^2)$ where m_a, m_b, m_c are the three medians.

The tetrahedron bimedian theorem extends this median-edge relationship from triangles (2D) to tetrahedra (3D).

- **Algebraic Technique:** The solution demonstrates a powerful technique: expanding dot products systematically and observing how cross-terms cancel when summing. Notice how the coefficients ± 2 in the dot product terms sum to -2 for each pair $(\mathbf{b} \cdot \mathbf{c})$, $(\mathbf{b} \cdot \mathbf{d})$, and $(\mathbf{c} \cdot \mathbf{d})$.
- **Symmetric Structure:** The three bimedian expressions have a beautiful symmetry:

$$\begin{aligned}\overrightarrow{LP} &= \frac{1}{2}(-\mathbf{b} + \mathbf{c} + \mathbf{d}) \\ \overrightarrow{MQ} &= \frac{1}{2}(\mathbf{b} - \mathbf{c} + \mathbf{d}) \\ \overrightarrow{NR} &= \frac{1}{2}(\mathbf{b} + \mathbf{c} - \mathbf{d})\end{aligned}$$

Each expression has exactly one negative sign, cycling through the three vectors. This symmetry guarantees that when we compute $|\overrightarrow{LP}|^2 + |\overrightarrow{MQ}|^2 + |\overrightarrow{NR}|^2$, the cross-terms will combine correctly.

- **Geometric Insight:** This theorem provides a metric relationship in tetrahedra, analogous to how Pythagoras relates sides in right triangles or how Apollonius relates medians in triangles. It's a fundamental identity in 3D geometry that can be used to prove other properties or solve optimization problems involving tetrahedra.

Problem 3.35: Concurrent Medians of a Tetrahedron

In a tetrahedron $ABCD$, define a *median* as the segment from a vertex to the centroid of the opposite face.

Let the position vectors of A, B, C, D be $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ relative to origin O . Let M_A, M_B, M_C, M_D be centroids of faces BCD, ACD, ABD, ABC respectively.

(i) Show that

$$\mathbf{m}_A = \frac{\mathbf{b} + \mathbf{c} + \mathbf{d}}{3}, \quad \mathbf{m}_B = \frac{\mathbf{a} + \mathbf{c} + \mathbf{d}}{3}, \quad \mathbf{m}_C = \frac{\mathbf{a} + \mathbf{b} + \mathbf{d}}{3}, \quad \mathbf{m}_D = \frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3}.$$

(ii) Define

$$\mathbf{g} = \frac{\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{d}}{4}.$$

Prove that G lies on all four medians AM_A, BM_B, CM_C, DM_D .

(iii) Deduce the internal division ratio on each median.

Hint: Rewrite $\mathbf{g}$ by isolating one vertex and grouping the other three. For example: $\mathbf{g} = \frac{1}{4}(\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{d}) = \frac{1}{4}\mathbf{a} + \frac{3}{4}\mathbf{m}_A$. Then repeat by symmetry for B, C, D .

Solution 3.35: Sketch

(i) Each face centroid is the average of its three vertex vectors, so the four formulas follow directly.

(ii) Start with:

$$\mathbf{g} = \frac{\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{d}}{4} = \frac{\mathbf{a} + (\mathbf{b} + \mathbf{c} + \mathbf{d})}{4} = \frac{\mathbf{a} + 3\mathbf{m}_A}{4} = \frac{1}{4}\mathbf{a} + \frac{3}{4}\mathbf{m}_A.$$

So G lies on AM_A by the section formula.

By symmetry:

$$\mathbf{g} = \frac{1}{4}\mathbf{b} + \frac{3}{4}\mathbf{m}_B = \frac{1}{4}\mathbf{c} + \frac{3}{4}\mathbf{m}_C = \frac{1}{4}\mathbf{d} + \frac{3}{4}\mathbf{m}_D,$$

so G lies on BM_B, CM_C, DM_D as well. Therefore all four medians are concurrent.

(iii) From $\mathbf{g} = \frac{1}{4}\mathbf{a} + \frac{3}{4}\mathbf{m}_A$, point G divides AM_A internally in ratio

$$AG : GM_A = 3 : 1.$$

Similarly:

$$BG : GM_B = CG : GM_C = DG : GM_D = 3 : 1.$$

Takeaways 3.4

This is the 3D analogue of triangle centroid concurrency. In 2D, averaging 3 vertices gives ratio 2 : 1 on each median; in 3D, averaging 4 vertices gives ratio 3 : 1 on each tetrahedron median. The symmetry of the averaged vector is the key idea that avoids solving multiple line equations.

Problem 3.36: Triangle Intersection Ratios

The diagram shows triangle OQA .


The point P lies on OA so that $OP : OA = 3 : 5$.

The point B lies on OQ so that $OB : OQ = 1 : 3$.

The point R is the intersection of AB and PQ .

The point T is chosen on AQ so that O, R and T are collinear.

Let $\mathbf{a} = \overrightarrow{OA}$, $\mathbf{b} = \overrightarrow{OB}$ and $\overrightarrow{PR} = k\overrightarrow{PQ}$ where k is a real number.



(i) Show that $\overrightarrow{OR} = \frac{3}{5}(1-k)\mathbf{a} + 3k\mathbf{b}$.

(ii) Writing $\overrightarrow{AR} = h\overrightarrow{AB}$, where h is a real number, it can be shown that $\overrightarrow{OR} = (1-h)\mathbf{a} + h\mathbf{b}$. (Do NOT prove this.)

Show that $k = \frac{1}{6}$.

(iii) Find $\overrightarrow{OT}$ in terms of $\mathbf{a}$ and $\mathbf{b}$.

Hint: (i) Express $\overrightarrow{OR}$ as $\overrightarrow{OP} + \overrightarrow{PR}$. Use $\overrightarrow{OP} = \frac{3}{5}\mathbf{a}$ and $\overrightarrow{PQ} = -\frac{3}{5}\mathbf{a} + 3\mathbf{b}$ where $\overrightarrow{OQ} = 3\mathbf{b}$.
(ii) Equate coefficients of $\mathbf{a}$ and $\mathbf{b}$ from the two expressions for $\overrightarrow{OR}$. Use the relationship $\overrightarrow{OT} = \lambda\overrightarrow{OR}$. Since O, R, T are collinear: $\overrightarrow{OT} = \lambda\overrightarrow{OR}$. Since T is on AQ : $\overrightarrow{OT} = (1-\mu)\mathbf{a} + 3\mu\mathbf{b}$. Equate coefficients.

Solution 3.36: Sketch

(i) $\overrightarrow{OR} = \overrightarrow{OP} + k\overrightarrow{PQ}$. Since $\overrightarrow{OP} = \frac{3}{5}\mathbf{a}$ and $\overrightarrow{OQ} = 3\mathbf{b}$, we have: $\overrightarrow{PQ} = -\frac{3}{5}\mathbf{a} + 3\mathbf{b}$. Therefore: $\overrightarrow{OR} = \frac{3}{5}\mathbf{a} + k(-\frac{3}{5}\mathbf{a} + 3\mathbf{b}) = \frac{3}{5}(1-k)\mathbf{a} + 3k\mathbf{b}$.

(ii) Equating coefficients from both expressions: For $\mathbf{b}$: $3k = h$. For $\mathbf{a}$: $\frac{3}{5}(1-k) = 1-h$. Substitute $h = 3k$ into second equation: $\frac{3}{5}(1-k) = 1-3k \Rightarrow 3(1-k) = 5(1-3k) \Rightarrow 3-3k = 5-15k \Rightarrow 12k = 2 \Rightarrow k = \frac{1}{6}$.

(iii) From part (ii): $\overrightarrow{OR} = \frac{3}{5}(1-\frac{1}{6})\mathbf{a} + 3(\frac{1}{6})\mathbf{b} = \frac{1}{2}\mathbf{a} + \frac{1}{2}\mathbf{b}$.

Since O, R, T collinear: $\overrightarrow{OT} = \lambda\overrightarrow{OR} = \frac{\lambda}{2}(\mathbf{a} + \mathbf{b})$.

Since T on AQ : $\overrightarrow{OT} = \mathbf{a} + \mu(\overrightarrow{OQ} - \mathbf{a}) = (1-\mu)\mathbf{a} + 3\mu\mathbf{b}$.

Equating coefficients: $\frac{\lambda}{2} = 1-\mu$ and $\frac{\lambda}{2} = 3\mu$. From these: $1-\mu = 3\mu \Rightarrow \mu = \frac{1}{4}$. Therefore: $\overrightarrow{OT} = (1-\frac{1}{4})\mathbf{a} + 3(\frac{1}{4})\mathbf{b} = \frac{3}{4}\mathbf{a} + \frac{3}{4}\mathbf{b} = \frac{3}{4}(\mathbf{a} + \mathbf{b})$.

Problem 3.37: Circle Intersection of Sets

Let A and B be two distinct points in three-dimensional space. Let M be the midpoint of AB .

Let S_1 be the set of all points P such that $\overrightarrow{AP} \cdot \overrightarrow{BP} = 0$.

Let S_2 be the set of all points N such that $|\overrightarrow{AN}| = |\overrightarrow{MN}|$.

The intersection of S_1 and S_2 is the circle S .

What is the radius of the circle S ?

- A. $\frac{|\overrightarrow{AB}|}{2}$
- B. $\frac{|\overrightarrow{AB}|}{4}$
- C. $\frac{\sqrt{3}|\overrightarrow{AB}|}{2}$
- D. $\frac{\sqrt{3}|\overrightarrow{AB}|}{4}$

Hint: (Step 1) Recognize that $\overrightarrow{AP} \cdot \overrightarrow{BP} = 0$ defines a sphere with diameter AB centered at M . (Step 2) The condition $|\overrightarrow{AN}| = |\overrightarrow{MN}|$ describes the perpendicular bisecting plane of segment AM . (Step 3) Use the Pythagorean relationship $r^2 + d^2 = R_{\text{sphere}}^2$ where d is the distance from the sphere center to the plane.

Solution 3.37: Sketch

Step 1: Analyze set S_1 . The condition $\overrightarrow{AP} \cdot \overrightarrow{BP} = 0$ means vectors $\overrightarrow{AP}$ and $\overrightarrow{BP}$ are perpendicular. The locus of points P that subtend a right angle to segment AB is a sphere with diameter AB . Center: M (midpoint of AB). Radius: $R_{\text{sphere}} = \frac{1}{2}|\overrightarrow{AB}|$.

Step 2: Analyze set S_2 . The condition $|\overrightarrow{AN}| = |\overrightarrow{MN}|$ describes points equidistant from A and M , which is the perpendicular bisecting plane of segment AM .

Step 3: Find intersection. The intersection of a sphere (centered at M) and a plane is a circle. Using the Pythagorean relationship: $r^2 + d^2 = R_{\text{sphere}}^2$ where d is the perpendicular distance from center M to the plane.

Step 4: Calculate distances. Since the plane passes through the midpoint of AM , and $|\overrightarrow{AM}| = \frac{1}{2}|\overrightarrow{AB}|$, we have:

$$d = \frac{1}{2}|\overrightarrow{AM}| = \frac{1}{2} \left(\frac{1}{2}|\overrightarrow{AB}| \right) = \frac{1}{4}|\overrightarrow{AB}|$$

Step 5: Solve for radius r .

$$\begin{aligned} r^2 &= R_{\text{sphere}}^2 - d^2 = \left(\frac{|\overrightarrow{AB}|}{2} \right)^2 - \left(\frac{|\overrightarrow{AB}|}{4} \right)^2 \\ &= \frac{|\overrightarrow{AB}|^2}{4} - \frac{|\overrightarrow{AB}|^2}{16} = \frac{4|\overrightarrow{AB}|^2 - |\overrightarrow{AB}|^2}{16} = \frac{3|\overrightarrow{AB}|^2}{16} \\ r &= \frac{\sqrt{3}|\overrightarrow{AB}|}{4} \end{aligned}$$

Therefore, the answer is **D**.

Problem 3.38: Complex Numbers and Centroid

The complex numbers x , w , and z are all different and all have modulus 1 (i.e., they lie on the unit circle in the complex plane).

The **centroid** (also called the center of mass) of three points is the average of their positions, given by $G = \frac{1}{3}(x + w + z)$.

Prove that $\frac{1}{3}(x + w + z)$ is never a cube root of xwz .

Note: A cube root of xwz is any complex number K satisfying $K^3 = xwz$.

Hint: Show that the centroid has modulus strictly less than 1 (using the triangle inequality with strict inequality since the points are distinct), while any cube root of xwz has modulus exactly 1.

Solution 3.38: Sketch

Given: $|x| = |w| = |z| = 1$ and x, w, z are all different.


Step 1: Find modulus of any cube root. Let K be any cube root of xwz , so $K^3 = xwz$. Taking modulus: $|K|^3 = |xwz| = |x||w||z| = 1 \cdot 1 \cdot 1 = 1$. Therefore $|K| = 1$. Thus, any cube root of xwz lies on the unit circle.

Step 2: Find modulus of centroid. The centroid is $G = \frac{1}{3}(x + w + z)$. By triangle inequality: $|x + w + z| \leq |x| + |w| + |z| = 3$. Equality holds only if x, w, z have the same argument (same direction), which means $x = w = z$. But they are given to be different, so strict inequality holds: $|x + w + z| < 3$. Therefore: $|G| = \left| \frac{x+w+z}{3} \right| < 1$.

Conclusion: Since $|G| < 1$ but $|K| = 1$ for any cube root K , we have $G \neq K$. Therefore, the centroid is never a cube root of the product.

Problem 3.39: Pyramid Centroid

Consider a pyramid $SABCD$ with square base $ABCD$ and apex S . Let H be the center of the square base (the intersection point of the diagonals AC and BD).



Note: H (red) is the center of the square base, and G (green) is the centroid of the pyramid. The blue dotted lines show vectors from H to all five vertices.

The centroid G of the pyramid is the point such that:

$$\vec{GA} + \vec{GB} + \vec{GC} + \vec{GD} + \vec{GS} = \vec{0}$$

Show that G lies on the line HS with $\vec{HG} = \frac{1}{5}\vec{HS}$.

Hint: Express each $\vec{GA_i}$ in terms of $\vec{GH}$ and $\vec{HA_i}$. Use the fact that H is the center of the square base, so $\vec{HA} + \vec{HB} + \vec{HC} + \vec{HD} = \vec{0}$ by symmetry.

Solution 3.39

Finding the centroid G

The centroid G of the five vertices satisfies:

$$\vec{GA} + \vec{GB} + \vec{GC} + \vec{GD} + \vec{GS} = \vec{0}$$

Express each vector using the triangle rule: $\vec{GA} = \vec{GH} + \vec{HA}$:

$$\begin{aligned}\vec{0} &= \vec{GA} + \vec{GB} + \vec{GC} + \vec{GD} + \vec{GS} \\ &= (\vec{GH} + \vec{HA}) + (\vec{GH} + \vec{HB}) + (\vec{GH} + \vec{HC}) + (\vec{GH} + \vec{HD}) + (\vec{GH} + \vec{HS}) \\ &= 5\vec{GH} + (\vec{HA} + \vec{HB} + \vec{HC} + \vec{HD}) + \vec{HS}\end{aligned}$$

Since H is the center of the square base, by symmetry the vectors from H to the four base vertices cancel:

$$\vec{HA} + \vec{HB} + \vec{HC} + \vec{HD} = \vec{0}$$

(This follows because H is the midpoint of both diagonals: $\vec{HA} = -\vec{HC}$ and $\vec{HB} = -\vec{HD}$.)

Therefore:

$$\vec{0} = 5\vec{GH} + \vec{HS}$$

Rearranging:

$$5\vec{GH} = -\vec{HS} = \vec{SH}$$

Therefore:

$$\vec{GH} = \frac{1}{5}\vec{SH}$$

Or equivalently:

$$\vec{HG} = \frac{1}{5}\vec{HS}$$

This shows that G lies on the line segment HS , located $\frac{1}{5}$ of the way from H to S .

Answer: The centroid G divides the segment from the center of the base to the apex in the ratio $HG : GS = 1 : 4$.

Problem 3.40: Line-Sphere Intersection Points

Find intersection points of line $\mathbf{r} = \mathbf{i} + 3\mathbf{j} - 4\mathbf{k} + t(\mathbf{i} + 2\mathbf{j} + 2\mathbf{k})$ and sphere $(x - 1)^2 + (y - 3)^2 + (z + 4)^2 = 81$.


Hint: Substitute parametric equations into sphere equation and solve resulting quadratic.

Solution 3.40: Sketch

Parametric: $x = 1 + t$, $y = 3 + 2t$, $z = -4 + 2t$. Substitute: $(t)^2 + (2t)^2 + (2t)^2 = 81 \Rightarrow 9t^2 = 81 \Rightarrow t = \pm 3$.
Points: $(4, 9, 2)$ when $t = 3$ and $(-2, -3, -10)$ when $t = -3$.

Problem 3.41: Regular Octagon Vector Sum

In regular octagon with side length 4, find magnitude of sum of vectors from midpoint of one side to all vertices.



Note: The regular octagon has vertices $P_1, P_2, \dots, P_8$, center O (blue), and A (red) is the midpoint of side P_1P_2 (shown in violet). Red arrows show vectors from A to all eight vertices. The green arrow highlights $\overrightarrow{AO}$.

Hint: Use symmetry: sum of vectors from center to vertices is zero. Express via center.

Solution 3.41

Step 1: Use vector decomposition via the center.

For any point A and the center O , we can write each vector from A to a vertex P_i as:

$$\overrightarrow{AP_i} = \overrightarrow{AO} + \overrightarrow{OP_i}$$

Summing over all 8 vertices:

$$\begin{aligned}\sum_{i=1}^8 \overrightarrow{AP_i} &= \sum_{i=1}^8 (\overrightarrow{AO} + \overrightarrow{OP_i}) \\ &= 8\overrightarrow{AO} + \sum_{i=1}^8 \overrightarrow{OP_i}\end{aligned}$$

Step 2: Apply symmetry to eliminate the sum from center.

By symmetry of a regular octagon, the 8 vertices are evenly distributed around the center O . The vectors $\overrightarrow{OP_1}, \overrightarrow{OP_2}, \dots, \overrightarrow{OP_8}$ point in directions separated by 45° and all have equal magnitude. Their vector sum is:

$$\sum_{i=1}^8 \overrightarrow{OP_i} = \vec{0}$$

Therefore:

$$\sum_{i=1}^8 \overrightarrow{AP_i} = 8\overrightarrow{AO}$$

Step 3: Calculate the apothem $|AO|$.

The apothem is the perpendicular distance from center to midpoint of a side. Consider the right triangle formed by O , midpoint A , and vertex P_1 . The central angle between adjacent vertices is 45° , so half this angle at O is 22.5° . With opposite side $|AP_1| = 2$ (half the side length) and adjacent side $|AO|$:

$$\tan(22.5^\circ) = \frac{2}{|AO|} \Rightarrow |AO| = 2 \cot(22.5^\circ)$$

Step 4: Evaluate $\cot(22.5^\circ)$.

Using the half-angle formula: $\tan(22.5^\circ) = \sqrt{2} - 1$. Therefore:

$$\cot(22.5^\circ) = \frac{1}{\sqrt{2} - 1} = \frac{\sqrt{2} + 1}{(\sqrt{2} - 1)(\sqrt{2} + 1)} = \sqrt{2} + 1$$

Thus: $|AO| = 2(\sqrt{2} + 1)$.

Step 5: Find the final magnitude.

From Step 2, we have $\sum_{i=1}^8 \overrightarrow{AP_i} = 8\overrightarrow{AO}$.

Taking magnitudes:

$$\left| \sum_{i=1}^8 \overrightarrow{AP_i} \right| = |8\overrightarrow{AO}| = 8|AO| = 8 \cdot 2(\sqrt{2} + 1) = 16(\sqrt{2} + 1)$$

Answer: The magnitude of the sum of vectors is $16(\sqrt{2} + 1)$ or approximately $16(2.414) \approx 38.63$.

4 Conclusion

Vectors unify geometry, algebra, and analytic work for Extension 1 and Extension 2. Build two habits: manipulate components reliably, and always know what the result *means* geometrically.

The hardest material here is integrative by design—it reaches into Euclidean geometry, trigonometry, and (where applicable) complex numbers and planes. That mirrors the HSC and the mathematics beyond it: vector technique earns its keep when combined with other tools, not kept in isolation.

Teachers can use these problems for extension, proof practice, and exam-style reasoning. Learners aiming past the HSC carry the same ideas into linear algebra, mechanics, and multi-

variable courses. Revisit what resisted you first; explain *why* a method works, not only *how*.
Best of luck with your studies and the HSC examination!

Contact Information:

LinkedIn: <https://www.linkedin.com/in/nguyenvuhung/>


GitHub: <https://github.com/vuhung16au/>

Repository: <https://github.com/vuhung16au/math-olympiad-ml/tree/main/HSC-Vectors>

A Appendices

A.1 Appendix 1: 2D Parametric Curves


Enrichment only. These 2D parametric curves are *outside the current NESA Mathematics Extension 1 and Extension 2 syllabuses*. They are included for students interested in university mathematics, physics, engineering, and computer graphics.



Cardioid

Heart-shaped curve from a rolling circle.


$$\begin{aligned}x &= a(2 \cos t - \cos 2t) \\y &= a(2 \sin t - \sin 2t)\end{aligned}$$



Epitrochoid

Epitrochoid with the tracing point offset from the rolling circle.


$$\begin{aligned}x &= (a + b) \cos t - c \cos\left(\frac{a}{b} + 1\right)t \\y &= (a + b) \sin t - c \sin\left(\frac{a}{b} + 1\right)t\end{aligned}$$



Hypocycloid

Curve from a circle rolling inside a fixed circle.


$$\begin{aligned}x &= (a - b) \cos t + b \cos\left(\frac{a}{b} - 1\right)t \\y &= (a - b) \sin t - b \sin\left(\frac{a}{b} - 1\right)t\end{aligned}$$



Lissajous Curves

Closed curves from harmonic motion in two perpendicular directions.


$$x(t) = a \sin(nt + c), \quad y(t) = b \sin t$$



Epicycloid

Curve traced by a point on a circle rolling outside another.

$$\begin{aligned}x &= (a + b) \cos t - b \cos\left(\frac{a}{b} + 1\right)t \\y &= (a + b) \sin t - b \sin\left(\frac{a}{b} + 1\right)t\end{aligned}$$




Folium of Descartes

A loop with an oblique asymptote.

$$\text{Cartesian: } x^3 + y^3 = 3axy$$


$$x(t) = \frac{3at}{1 + t^3}, \quad y(t) = \frac{3at^2}{1 + t^3}$$



Hypotrochoid

Hypocycloid with the tracing point offset from the rolling circle.

$$\begin{aligned}x &= (a - b) \cos t + c \cos\left(\frac{a}{b} - 1\right)t \\y &= (a - b) \sin t - c \sin\left(\frac{a}{b} - 1\right)t\end{aligned}$$




Plateau Curves

Rose-like curves from rational trigonometric combinations.

$$\begin{aligned}x(t) &= \frac{a \sin((m + n)t)}{\sin((m - n)t)} \\y(t) &= \frac{2a \sin(mt) \sin(nt)}{\sin((m - n)t)}\end{aligned}$$

A.2 Appendix 2: 3D Parametric Curves


Enrichment only. These 3D parametric curves are *outside the current NESA syllabus*. They preview ideas from multi-variable calculus, mechanics, and computer graphics.



The Helix (Slinky)

Simplest continuous spiral in three dimensions.

$$x = \cos t, \quad y = \sin t, \quad z = t$$



Torus Knot (Figure-Eight)

A tube wraps along a closed, twisting figure-eight path.


$$\begin{aligned} x &= \cos t + 2 \cos 2t \\ y &= \sin t - 2 \sin 2t, \quad z = -\sin 3t \end{aligned}$$



3D Rose (Rhodonea)

A 2D flower curve lifted into 3D by a periodic z -component.


$$x = \cos(4t) \cos t, \quad y = \cos(4t) \sin t, \quad z = \sin(4t)$$



3D Lissajous Curve

Wavelike motion on all three axes creates symmetric nodes.

$$x = \sin(3t), \quad y = \sin(4t), \quad z = \sin t$$




Heart

Classic heart profile in the xy -plane ($z = 0$).

$$x = 4R \sin^3 u$$


$$y = \frac{R}{4} (13 \cos u - 5 \cos 2u - 2 \cos 3u - \cos 4u)$$



Spring

Circular motion in xy with oscillating height in z .

$$x = R \cos t, \quad y = R \sin t, \quad z = R \cos t$$




Möbius Strip

Non-orientable surface; centre circle ($v = 0$) and edge ($v = 1$) shown.

$$\begin{aligned} x &= 2R \left(\cos u + v \cos \frac{u}{2} \cos u \right) \\ y &= 2R \left(\sin u + v \cos \frac{u}{2} \sin u \right), \quad z = 2Rv \sin \frac{u}{2} \end{aligned}$$

Tools to plot 3D curves (Last checked: Jun 2026)

- MATLAB `fplot3()` function
- Desmos 3D (<https://www.desmos.com/3d>)



Cone

Spiral path on a conical surface as radius and height grow.

$$x = t \cos t, \quad y = t \sin t, \quad z = t$$

A.3 Appendix 3: Projections of a 3D Parametric Curve


Enrichment. Orthogonal projections onto coordinate planes are a core sketching strategy for 3D parametric curves. This example uses the circular helix from part (a) of Problem 2.21 (Sketching 3D Parametric Curves).

Example curve.

$$\mathbf{r}(t) = (2 \sin t)\mathbf{i} + (2 \cos t)\mathbf{j} + \frac{t}{\pi}\mathbf{k}, \quad 0 \leq t \leq 4\pi.$$

Equivalently, $x(t) = 2 \sin t$, $y(t) = 2 \cos t$, and $z(t) = t/\pi$.


The curve in 3D.



Orthogonal projections onto the coordinate planes. Each projection keeps two components and sets the third to zero—the “shadow” of the curve when viewed along the missing axis.

Projection onto the xy -plane


Drop z : $(x, y) = (2 \sin t, 2 \cos t)$



Circle of radius 2, traversed **twice** as t runs from 0 to 4π .

Projection onto the yz -plane


Drop x : $(y, z) = (2 \cos t, t/\pi)$



Sinusoidal trace: $y = 2 \cos(\pi z)$ for $0 \leq z \leq 4$ (two full periods).

Projection onto the xz -plane

Drop y : $(x, z) = (2 \sin t, t/\pi)$



Sinusoidal trace: $x = 2 \sin(\pi z)$ for $0 \leq z \leq 4$ (two full periods).

What we find, and how to read the projections.

- **xy -projection (view from above, along $\mathbf{k}$).** Only the $\mathbf{i}$ and $\mathbf{j}$ components matter. Here $x^2 + y^2 = 4$, so the shadow is a circle of radius 2. Because t ranges over 4π , the particle winds around this circle *twice* while rising in z . The projection alone does not show the vertical motion—it only reveals that the helix sits on a vertical cylinder of radius 2.
- **yz - and xz -projections (side views).** Each pairs one horizontal coordinate with height. Substituting $t = \pi z$ gives $y = 2 \cos(\pi z)$ and $x = 2 \sin(\pi z)$ on $0 \leq z \leq 4$. Both shadows are sinusoidal waves with amplitude 2 and two complete periods—matching the two revolutions of the helix.
- **Reconstructing the 3D curve.** Read the curve component-wise: circular motion in the xy -plane combined with steady linear increase in z . The start point is $\mathbf{r}(0) = (0, 2, 0)$ and the end point is $\mathbf{r}(4\pi) = (0, 2, 4)$ —both visible on every projection at $z = 0$ and $z = 4$ with $x = 0$.
- **Sketching strategy (HSC and beyond).** When a 3D parametric equation looks unfamiliar, project onto each coordinate plane first. Each projection is a 2D curve you can sketch using familiar

methods. Then combine the information: if the xy -shadow is a circle and z grows steadily, expect a helix; if two horizontal coordinates match (as in part (b) of the same problem), the curve is confined to a plane.

Enrichment material. *The five appendices that follow explore ideas from Victorian and Queensland Specialist Mathematics, as well as NSW’s historical 1966–1980 Level 1 / 4 Unit curriculum. They are intended as enrichment for NSW students studying the NESA 2024 syllabuses, **not** as additional HSC examination requirements. They also provide a natural bridge to ideas encountered in first-year university mathematics, physics, engineering and computer science.*

Beyond HSC Vectors — Modern Exts and NSW’s 60s Heritage

Vectors are an important part of senior mathematics across Australia. While NSW Mathematics Extension 1 and Extension 2 develop vector algebra, geometry, proofs, equations of lines and curves, other states introduce some additional techniques.

Moreover, these extensions are **not foreign to NSW**. Between 1966 and 1980, the highest level of senior mathematics in NSW (originally called **First Level**, and later **4 Unit**) was a **two-year senior course** across Years 11 and 12. Section 4 of the 1965 syllabus was explicitly titled “*Plane Geometry, Geometric Algebra, Matrices, Geometry in Three Dimensions*”. It unified 3D vectors, equations of planes, determinants, and linear dependence. When 4 Unit was compressed into a 1-year Year 12 course in 1981, these topics were removed. Exploring them today reconnects students with the complete mathematical picture that once anchored NSW senior mathematics.

This section explores five ideas:

1. **Polar form of two-dimensional vectors** (QLD Specialist Mathematics — Unit 1)
2. **The cross product** (VIC Specialist Mathematics — Units 3–4; QLD Specialist Mathematics — Unit 3; HSC First Level 1966–1980; including applications to 2D polygon and parametric curve areas)
3. **Equations of planes in three dimensions** (VIC Specialist Mathematics — Units 3–4; QLD Specialist Mathematics — Unit 3; HSC 4 Unit 1966–1980)
4. **Linear dependence and independence** (VIC Specialist Mathematics — Units 3–4; HSC First Level 1966–1980)
5. **Vector-valued motion in three dimensions** (VIC Specialist Mathematics — Units 3–4; QLD Specialist Mathematics — Unit 3)

Curriculum and historical references. These appendices draw upon the following curriculum and historical sources:

- NESA — Mathematics Extension 1 (2024)
- NESA — Mathematics Extension 2 (2024)
- NSW Board of Senior School Studies — Mathematics First Level Syllabus (1965/1967)
- VCAA — VCE Specialist Mathematics Study Design
- QCAA — Specialist Mathematics 2025 v1.4

These sections are enrichment material, designed to broaden geometric intuition and prepare students for advanced tertiary STEM study.

A.4 Appendix 4: Polar Form of Two-Dimensional Vectors

Enrichment only. Polar form of vectors is part of *QLD Specialist Mathematics (Unit 1)*. It is not prescribed in the current NESA Extension 1 or Extension 2 syllabus.

Key concepts.

A vector in two dimensions has both magnitude and direction. We usually express a vector in Cartesian form:

$$\mathbf{v} = x\mathbf{i} + y\mathbf{j}.$$

An alternative is **polar form**, which represents a non-zero vector by

$$(r, \theta)$$

where:

- $r > 0$ is the magnitude (length) of the vector.
- θ is its direction, measured anticlockwise from the positive x -axis.

The two forms are related by

$$x = r \cos \theta, \quad y = r \sin \theta.$$

Conversely,

$$r = \sqrt{x^2 + y^2}, \quad \tan \theta = \frac{y}{x} \quad (x \neq 0).$$

When calculating θ , always check the quadrant. The direction is determined modulo 2π . The zero vector has no uniquely defined direction.

NSW connection. Extension 1 already teaches vector components, magnitude and direction, while Extension 2 uses trigonometric vector equations of circles. The enrichment here is the systematic use of polar-form notation and conversion.

Worked example. Express

$$\mathbf{v} = -\sqrt{3}\mathbf{i} + \mathbf{j}$$

in polar form.

Step 1: Find the magnitude.

$$r = \sqrt{(-\sqrt{3})^2 + 1^2} = 2.$$

Step 2: Find the direction. Since $x < 0$ and $y > 0$, the vector lies in Quadrant II.

$$\tan \theta = -\frac{1}{\sqrt{3}}.$$

Taking $0 \leq \theta < 2\pi$,

$$\theta = \frac{5\pi}{6}.$$

Therefore,

$$(r, \theta) = \left(2, \frac{5\pi}{6}\right).$$

Verification.

$$2 \cos \frac{5\pi}{6} = -\sqrt{3}, \quad 2 \sin \frac{5\pi}{6} = 1. \quad \checkmark$$

Key takeaways.

- Cartesian form is convenient for adding vectors and performing algebra; polar form makes magnitude and direction immediately visible.

- Always check the quadrant when converting to polar form.
- Polar form connects naturally with the modulus and argument of complex numbers.

Beyond school — University connection. Polar coordinates are widely used in physics, robotics and engineering. They also connect vector geometry with complex numbers through

$$z = r(\cos \theta + i \sin \theta).$$

A.5 Appendix 5: The Cross Product of Two Vectors

Enrichment only. The cross product is part of **VIC Specialist Mathematics (Units 3–4)** and **QLD Specialist Mathematics (Unit 3)**. It is **not** prescribed in the current **NESA Mathematics Extension 1 or Extension 2 syllabus**, so none of the core worked methods in Parts 1 and 2 of this booklet depend on it. It is included here because students may meet it later in university mathematics, physics, or engineering, and it can help connect familiar vector ideas to further study.

Key concepts.

The dot product produces a *scalar*:

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta.$$

The **cross product** produces a *vector* perpendicular to both original vectors.

For two vectors in three dimensions,

$$\mathbf{a} = (a_1, a_2, a_3), \quad \mathbf{b} = (b_1, b_2, b_3),$$

their cross product is defined by

$$\mathbf{a} \times \mathbf{b} = \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix}.$$

Determinant mnemonic. The same formula can be written using a 3×3 determinant expansion:

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \mathbf{k}.$$

Understanding the matrix notation. While matrices and determinants are not in the current NESA syllabus, they were historically integrated alongside 3D vectors in the 1966–1980 NSW Level 1 / 4 Unit syllabus Section 4. The determinant notation is used here both as an exact algebraic operator and a practical calculation mnemonic:

- **2×2 Determinant (Criss-Cross):** $\begin{vmatrix} p & q \\ r & s \end{vmatrix} = ps - qr$ (main diagonal minus off-diagonal).
- **Top-Row Expansion:** For each unit vector ($\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$), delete its row and column; the remaining four entries form a 2×2 sub-determinant.
- **Alternating Signs (+, −, +):** Notice the **minus sign** on the middle $\mathbf{j}$ component: $-(a_1 b_3 - a_3 b_1)\mathbf{j}$, which is the most frequent source of errors.

Magnitude and direction.

$$|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$$


where $0 \leq \theta \leq \pi$. The direction follows the **right-hand rule**. In particular,

$$\mathbf{a} \times \mathbf{b} = -(\mathbf{b} \times \mathbf{a}).$$

The cross product has an elegant geometric interpretation: its magnitude is the area of the parallelogram formed by the two vectors.

Properties.

- $\mathbf{a} \times \mathbf{b}$ is perpendicular to both $\mathbf{a}$ and $\mathbf{b}$.
- $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$ (area of parallelogram).
- $\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}$ (anticommutative).
- Two vectors have zero cross product precisely when they are parallel, including the case where one is the zero vector.



Left: The cross product $\mathbf{a} \times \mathbf{b}$ (green) is perpendicular to both $\mathbf{a}$ (blue) and $\mathbf{b}$ (red), using the right-hand rule for direction. *Right:* The magnitude $|\mathbf{a} \times \mathbf{b}|$ equals the area of the parallelogram formed by $\mathbf{a}$ and $\mathbf{b}$.

Worked example 1 — Cross product and triangle area.

Given

$$\mathbf{a} = (1, 0, 1), \quad \mathbf{b} = (0, 2, 0),$$

find: (a) $\mathbf{a} \times \mathbf{b}$; (b) the area of the triangle formed by the two vectors.

Solution. Using the component formula,

$$\mathbf{a} \times \mathbf{b} = (0 \cdot 0 - 1 \cdot 2, 1 \cdot 0 - 1 \cdot 0, 1 \cdot 2 - 0 \cdot 0) = \boxed{(-2, 0, 2)}.$$

Its magnitude is

$$|\mathbf{a} \times \mathbf{b}| = \sqrt{(-2)^2 + 0^2 + 2^2} = 2\sqrt{2}.$$

The triangle has half the area of the parallelogram:

$$\boxed{\text{Area of triangle} = \sqrt{2}}.$$

Verification of perpendicularity.

$$(\mathbf{a} \times \mathbf{b}) \cdot \mathbf{a} = (-2)(1) + 0(0) + 2(1) = 0, \quad (\mathbf{a} \times \mathbf{b}) \cdot \mathbf{b} = (-2)(0) + 0(2) + 2(0) = 0.$$

Thus the cross product is perpendicular to both original vectors. ✓

Worked example 2 — Distance from point to line.

Problem. Find the distance from point $P(1, 2, 0)$ to the line $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$.

Solution using Cross Product. Let $A(1, 0, 1)$ be a point on the line and $\mathbf{d} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$ be the direction

vector. The vector from A to P is $\overrightarrow{AP} = \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}$.

Using the cross product:

$$\overrightarrow{AP} \times \mathbf{d} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 2 & -1 \\ 0 & 1 & 1 \end{vmatrix} = \begin{vmatrix} 2 & -1 \\ 1 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 0 & -1 \\ 0 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 0 & 2 \\ 0 & 1 \end{vmatrix} \mathbf{k} = 3\mathbf{i} - 0\mathbf{j} + 0\mathbf{k} = \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}.$$

$$\text{Distance: } d = \frac{|\overrightarrow{AP} \times \mathbf{d}|}{|\mathbf{d}|} = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2} \text{ units.}$$

Comparison with the syllabus method. The in-syllabus method uses projection. The component of $\overrightarrow{AP}$ perpendicular to the line is found by:

$$\overrightarrow{AP}_{\perp} = \overrightarrow{AP} - \frac{\overrightarrow{AP} \cdot \mathbf{d}}{\mathbf{d} \cdot \mathbf{d}} \mathbf{d}.$$

Then $d = |\overrightarrow{AP}_{\perp}|$. Both methods yield the same result; for HSC purposes, the projection method is the syllabus-aligned approach to remember.

Worked example 3 — Flattening 3D to 2D: Triangle Area on an Ellipse.

Problem. Let $P(a \cos \theta, b \sin \theta)$ and $Q(a \cos \phi, b \sin \phi)$ be two points on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where $a, b > 0$. Find the area of triangle OPQ , where O is the origin $(0, 0)$.

Solution using the Cross Product (The “Two-Line” Shortcut). Embed the two-dimensional position vectors into three-dimensional space by setting the z -coordinate to zero:

$$\mathbf{p} = \begin{pmatrix} a \cos \theta \\ b \sin \theta \\ 0 \end{pmatrix}, \quad \mathbf{q} = \begin{pmatrix} a \cos \phi \\ b \sin \phi \\ 0 \end{pmatrix}.$$

Their cross product points strictly perpendicular to the xy -plane along $\mathbf{k}$:

$$\begin{aligned} \mathbf{p} \times \mathbf{q} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a \cos \theta & b \sin \theta & 0 \\ a \cos \phi & b \sin \phi & 0 \end{vmatrix} = 0\mathbf{i} - 0\mathbf{j} + (ab \cos \theta \sin \phi - ab \sin \theta \cos \phi)\mathbf{k} \\ &= ab \sin(\phi - \theta) \mathbf{k}. \end{aligned}$$

The area of the triangle is half the area of the parallelogram formed by $\mathbf{p}$ and $\mathbf{q}$:

$$\boxed{\text{Area}(\triangle OPQ) = \frac{1}{2}|\mathbf{p} \times \mathbf{q}| = \frac{1}{2}ab|\sin(\phi - \theta)|.}$$

The Historical 4U HSC Pain Point. Between 1966 and 1980, when NSW 4 Unit / First Level was a two-year senior course, Section 4 of the syllabus integrated 3D vectors and determinants alongside conics. When 4 Unit was compressed into a one-year Year 12 course in 1981, matrices and cross products were excised, but conic sections remained examinable until 2019. Consequently, generations of students were forced to solve triangle area questions using punishing coordinate geometry:

1. Apply the distance formula to find the base length $PQ = \sqrt{a^2(\cos \phi - \cos \theta)^2 + b^2(\sin \phi - \sin \theta)^2}$.
2. Determine the Cartesian equation of the chord PQ : $(b \cos \frac{\theta+\phi}{2})x + (a \sin \frac{\theta+\phi}{2})y = ab \cos \frac{\phi-\theta}{2}$.
3. Apply the perpendicular distance formula $h = \frac{|Ax_0 + By_0 + C|}{\sqrt{A^2 + B^2}}$ from $O(0, 0)$ to the chord.
4. Calculate $\text{Area} = \frac{1}{2} \times \text{base} \times \text{height}$, expanding pages of nested square roots and trig identities to obtain cancellations.

With the cross product, this entire algebraic marathon collapses into two elementary lines! The syllabus question was originally conceived when students had these very tools.

Connection to Matrices. Notice that the $\mathbf{k}$ -component of $\mathbf{p} \times \mathbf{q}$ is identically the 2×2 matrix determinant $\det \begin{pmatrix} a \cos \theta & b \sin \theta \\ a \cos \phi & b \sin \phi \end{pmatrix}$. For full coverage of coordinate determinants and the general 3×3 area

formula $\frac{1}{2} \left| \det \begin{pmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{pmatrix} \right|$, see our companion booklet *Matrix Practice: NSW Extension Enrichment & QLD Specialist Mathematics*.

Key takeaways.

- The dot product measures how much two vectors point in the same direction; the cross product gives a perpendicular direction and an area.
- In 2D plane geometry, embedding vectors into 3D with $z = 0$ turns triangle and polygon area calculations into a one-step cross product, bypassing tedious coordinate geometry formulas.
- Two vectors have zero cross product precisely when they are parallel, including the case where one is the zero vector.
- The order matters: $\mathbf{a} \times \mathbf{b} \neq \mathbf{b} \times \mathbf{a}$ in general.

Beyond school — University connection. The cross product appears in rotational mechanics, electromagnetism and computer graphics. For example, the torque produced by a force is

$$\boldsymbol{\tau} = \mathbf{r} \times \mathbf{F},$$

where $\mathbf{r}$ is a position vector and $\mathbf{F}$ is the applied force.

A.6 Appendix 6: Equations of Planes in Three Dimensions

Beyond the current HSC syllabus. The NESA Mathematics Extension 2 syllabus requires understanding of coordinate planes ($z = 0$, etc.) and coplanarity through vector relationships—see §1.4.8 in the Vectors Primer. This appendix covers the **full algebra of general plane equations** as taught in **VIC Specialist Mathematics (Units 3–4)** and **QLD Specialist Mathematics (Unit 3)**, and previously in the **NSW 1966–1980 First Level / 4 Unit syllabus**. It is intended as enrichment for university preparation, **not** as additional HSC examination content.

Key concepts.

In three-dimensional space, a line can be determined by a point and a direction vector:

$$\mathbf{r} = \mathbf{p} + t \mathbf{d}.$$

A **plane** can be determined by a point and a non-zero **normal vector**.

A normal vector is perpendicular to every direction lying in the plane.

Parametric form of a plane. A plane can also be described using a point and **two non-parallel direction vectors** that lie in the plane:

$$\boxed{\mathbf{r} = \mathbf{p} + s \mathbf{u} + t \mathbf{v}}, \quad s, t \in \mathbb{R}.$$

This is analogous to the parametric form of a line $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$, but with *two* independent parameters instead of one.

From parametric form to Cartesian form. Given two direction vectors $\mathbf{u}$ and $\mathbf{v}$ in the plane, a normal vector can be found using the cross product (Appendix A.5):

$$\mathbf{n} = \mathbf{u} \times \mathbf{v}.$$

Then the point-normal equation is

$$\boxed{\mathbf{n} \cdot (\mathbf{r} - \mathbf{p}) = 0}.$$

If $\mathbf{n} = (a, b, c)$ and $\mathbf{p} = (x_0, y_0, z_0)$, this becomes

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$

Equivalently, the Cartesian equation of a plane is

$$\boxed{ax + by + cz = d},$$

where a, b, c are not all zero.

Worked example — Finding a plane from three points.

Find the Cartesian equation of the plane passing through

$$A(1, 0, 0), \quad B(2, 0, 1), \quad C(1, 2, 0).$$

Step 1: Find two vectors in the plane.

$$\overrightarrow{AB} = (1, 0, 1), \quad \overrightarrow{AC} = (0, 2, 0).$$

Step 2: Find a normal vector using the cross product.

From Appendix A.5,

$$\overrightarrow{AB} \times \overrightarrow{AC} = (0 \cdot 0 - 1 \cdot 2, 1 \cdot 0 - 1 \cdot 0, 1 \cdot 2 - 0 \cdot 0) = (-2, 0, 2).$$

A simpler normal vector is therefore

$$\mathbf{n} = (1, 0, -1).$$

Step 3: Form the plane equation.

Using point $A(1, 0, 0)$,

$$(1, 0, -1) \cdot (x - 1, y, z) = 0.$$

Expanding,

$$x - 1 - z = 0.$$

Hence,

$$\boxed{x - z = 1}.$$

Verification. All three given points satisfy the equation: A : $1 - 0 = 1$ ✓; B : $2 - 1 = 1$ ✓; C : $1 - 0 = 1$ ✓.

Key takeaways.

- A line is described by one direction vector; a plane can be described by two non-parallel direction vectors.
- The cross product provides a quick method of finding a plane's normal vector.
- Coordinate planes such as $z = 0$ are special cases of general plane equations.

NSW HSC Heritage (1975 Q8). In the 1975 NSW HSC 4 Unit examination (Question 8), students were required to find the equation of the tangent plane to the sphere $x^2 + y^2 + z^2 - 2x - 6y + 2z + 2 = 0$ at the point $(3, 4, 1)$, and prove geometric properties of planes intersecting spheres in three dimensions. General plane equations were standard examinable content in NSW prior to the 1981 curriculum truncation.

Beyond school — University connection. Plane equations appear in linear algebra, computer-aided design, 3D graphics and optimisation. They also form the foundation for studying intersections of surfaces in multivariable calculus.

A.7 Appendix 7: Linear Dependence and Independence

Enrichment only. Linear dependence and independence are part of *VIC Specialist Mathematics (Units 3–4)* and were an integral component of the *NSW 1966–1980 First Level / 4 Unit curriculum*. The current *NESA syllabus* uses these concepts informally in geometric proofs (e.g. showing coplanarity via linear combinations); this appendix provides the formal definitions and matrix connections.

Key concepts.

A **linear combination** of vectors is an expression formed by multiplying vectors by scalars and adding them. For example,

$$\mathbf{w} = 2\mathbf{a} - 3\mathbf{b}$$

is a linear combination of $\mathbf{a}$ and $\mathbf{b}$.

Vectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ are **linearly independent** if

$$c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n = \mathbf{0}$$

is possible **only** when

$$c_1 = c_2 = \dots = c_n = 0.$$

Otherwise, they are **linearly dependent**.

Informally, dependence means at least one vector can be constructed from the others: it does not provide a genuinely new direction.

Worked example.

Consider the three vectors

$$\mathbf{a} = (1, 0, 1), \quad \mathbf{b} = (0, 2, 0), \quad \mathbf{c} = (1, 2, 1).$$

Determine whether they are linearly independent.

Solution. Observe that

$$\mathbf{a} + \mathbf{b} = (1, 2, 1) = \mathbf{c}.$$

Therefore,

$$\mathbf{a} + \mathbf{b} - \mathbf{c} = \mathbf{0}.$$

The coefficients 1, 1, -1 are not all zero. Hence, the three vectors are **linearly dependent**.

Geometric interpretation. All three vectors lie in the plane $x - z = 0$ through the origin.

However, $\mathbf{a}$ and $\mathbf{b}$ themselves are independent. Indeed,

$$s\mathbf{a} + t\mathbf{b} = (s, 2t, s) = \mathbf{0}$$

implies $s = t = 0$.

Thus, two independent vectors can describe the whole plane, while the third adds no new direction.

The Matrix Determinant Connection (1966–1980 NSW Heritage). In three dimensions, whether three vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ are linearly independent is directly determined by the determinant of the 3×3 matrix containing them as columns:

$$\mathbf{u}, \mathbf{v}, \mathbf{w} \text{ are linearly independent} \iff \det \begin{bmatrix} \mathbf{u} & \mathbf{v} & \mathbf{w} \end{bmatrix} \neq 0 \iff \mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) \neq 0.$$

If the determinant is zero, the scalar triple product vanishes, meaning the volume of the parallelepiped they span is zero and the vectors are coplanar (linearly dependent). For the worked example vectors above:

$$\det \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 2 \\ 1 & 0 & 1 \end{bmatrix} = 1(2 - 0) - 0 + 1(0 - 2) = 2 - 2 = 0,$$

confirming their linear dependence in one calculation without needing to solve simultaneous equations!

Key takeaways.

- Linear dependence means some vector information is redundant.
- Two non-zero vectors are independent precisely when they are **not parallel**.
- In three-dimensional space, three independent vectors can describe every vector through linear combinations. Any set of four or more 3D vectors is necessarily dependent.

Beyond school — University connection. Linear independence leads to the concept of a *basis* in linear algebra. It is fundamental to matrix rank, solving simultaneous equations, computer graphics and data science.

A.8 Appendix 8: Vector-Valued Motion in Three Dimensions

Enrichment only. Three-dimensional vector calculus is part of **VIC Specialist Mathematics (Units 3–4)** and **QLD Specialist Mathematics (Unit 3)**. NSW Extension 1 already covers vector functions for motion in a plane; this appendix extends those ideas by adding a third component.

Key concepts.

The position vector of a moving particle is

$$\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}.$$

The **velocity** vector is the derivative of position:

$$\mathbf{v}(t) = \mathbf{r}'(t).$$

The **acceleration** vector is the derivative of velocity:

$$\mathbf{a}(t) = \mathbf{r}''(t).$$

The particle's **speed** is the magnitude of its velocity:

$$|\mathbf{v}(t)| = \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2}.$$

The same differentiation rules used in two dimensions apply component by component in three dimensions.

Worked example.

A particle moves through space with position

$$\mathbf{r}(t) = t\mathbf{i} + t^2\mathbf{j} + t^3\mathbf{k}, \quad t \geq 0.$$

Find: (a) its velocity and acceleration; (b) its speed at $t = 1$; (c) the equation of the tangent line to its path at $t = 1$.

(a) Velocity and acceleration.

Differentiating each component,

$$\mathbf{v}(t) = \mathbf{i} + 2t\mathbf{j} + 3t^2\mathbf{k}.$$

Differentiating again,

$$\mathbf{a}(t) = 2\mathbf{j} + 6t\mathbf{k}.$$

(b) Speed at $t = 1$.

The velocity at $t = 1$ is

$$\mathbf{v}(1) = (1, 2, 3).$$

Therefore,

$$|\mathbf{v}(1)| = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}.$$

(c) Tangent line.

At $t = 1$, the particle is at

$$\mathbf{r}(1) = (1, 1, 1).$$

The velocity vector $(1, 2, 3)$ gives the direction of the tangent. Hence the tangent line is

$$\mathbf{R} = (1, 1, 1) + \lambda(1, 2, 3), \quad \lambda \in \mathbb{R}.$$

This connects the familiar NSW vector equation of a line with calculus: the velocity gives the tangent direction to the particle's path.

Key takeaways.

- Velocity is a vector; speed is its magnitude. They are not interchangeable.
- Differentiation of vector functions works component by component.
- The velocity vector gives the instantaneous tangent direction to a particle's path, even in three-dimensional space.

Beyond school — University connection. Three-dimensional vector calculus introduces space curves, tangent vectors, arc length and curvature (see also Problem 2.20 in Part 1 of this booklet). These ideas are used in robotics, aerospace engineering, mechanics and computer animation.

Final Thoughts: Three Operations, Three Geometric Ideas

The dot product, cross product and linear independence describe different aspects of vector geometry.

Concept	What it tells us
Dot product $\mathbf{a} \cdot \mathbf{b}$	Angles, projections and perpendicularity
Cross product $\mathbf{a} \times \mathbf{b}$	A perpendicular vector and an area
Linear independence	Whether vectors provide genuinely different directions

Together, these concepts explain how vectors describe directions, lines, planes and motion.

The underlying message is simple:

Vectors are more than arrows. They provide a common mathematical language for geometry, motion, linear algebra and many problems encountered in university STEM courses.