

How do trees grow in a non-stationary climate?

EM Wolkovich (University of British Columbia, Forest & Conservation Sciences)

Janneke Hille Ris Lambers (ETH Zurich, Environmental Systems Science)

Michael Betancourt (Symplectomorphic, LLC, NA)

Victor van der Meersch (CNRS, Center for Functional and Evolutionary Ecology)

9–16 March 2025

1 Overview of the Field

What determines how much trees grow and when? Answering this question has become increasingly difficult as climate change has warmed up forests, shifted rainfall patterns increased extreme weather. This research in teams (RIT) aimed to bridge across multiple fields and build better tree growth models leveraging new modeling opportunities and data streams.

Tree growth is influenced by a complex interplay of external events and size-dependent developmental stages that are mediated by environmental factors [1, 4]. Different fields have addressed this complexity with varying foci. Foresters and ecologists have generally used data that spans a wide spatial range, but is coarse temporally and short-term (10-30 years) and focused on models that partition growth variation across populations, sites and species [e.g., 5]. In contrast, dendrochronology, a subfield of climatology has used ring width extracted from internal tree cores to build extensive time-series (centuries to millennia) to estimate relationships between annual tree growth and climate—especially temperature and drought [2].

2 Recent Developments and Open Problems

The diverging approaches to modeling tree growth—across forestry, ecology and dendrochronology—may have worked well enough in the past, but recent forecasting attempts have uncovered major issues [e.g., the ‘divergence problem’ in dendrochronology, 3]. Alongside this, global forecasts increasingly need accurate models of not only how trees grow, but also how that growth varies within and across ecosystems and their climates. Both fields are therefore increasingly recognizing the need for new models that handle shifting temperature and drought regimes. To date, however, few have attempted to merge these disciplines’ data-streams and modeling approaches—this challenge was the focus of our RIT.

3 Scientific Progress Made

Our aim over one week was to improve models of how tree growth interacts with a changing climate by combining ecological and dendrochronological domain expertise and state-of-the-art statistical methodology to develop principled mathematical models and fit them. The team had met over video-conference several times in advance and thus arrived with data in hand and modeling ideas to begin work.

One major outcome of work over the first days at BIRS was developing models of how to combine the temporally coarse but spatially dense data from forestry (tree diameter measurements taken every several years) with annual tree-ring data from tree cores. We quickly realized the best way to combine these data would be to build from data using each measurement approach on the same trees. Towards this aim, we are currently collecting this unique multi-method data type at Mount Rainier by coring trees (to gather annual ring data) that have been monitored by diameter measurements for over 40 years as part of the Permanent Sample Plot program. We hope to have these data within a year and will return to this modeling challenge then.

Moving on from this we focused on modeling annual tree ring data, which is far more powerful for addressing questions of how climate affects growth. While our pre-BIRS meetings had focused on using a Gompertz growth model, we found at BIRS that it was too inflexible to capture the variability in growth observed across our data and thus developed a novel hierarchical Gaussian process model for tree growth, which combines flexible but domain-expertise-informed functions for long-term and short-term variable growth with three climatic predictors. We detail this model below.

3.1 Gaussian process

From a simplified point of view, a (finite- N -dimensional projection) of a Gaussian process f_N is defined as:

$$f_N \sim \text{multinormal}(m, K)$$

with a certain mean function m and a covariance matrix K . In particular, the elements of K are defined by a covariance function. In our model, we use an *exponentiated quadratic* function:

$$\forall i, j \in \llbracket 1, N \rrbracket \quad k_{i,j} = \gamma^2 \exp \left(-\frac{1}{2} \left(\frac{|x_j - x_i|}{\rho} \right)^2 \right)$$

This covariance function capture the "link" between observations along a continuous variable, such as time or space. The function is characterized by two parameters:

- a marginal deviation γ , which controls the variability around the mean
- a length scale ρ , which controls how long/far the correlations persist

3.2 Application to tree growth

Now, assume that we have ring width measurements r_w for one tree, for each year x_n . Using one Gaussian process, a simple model would look like this:

$$\forall x_n \quad \log r_w(x_n) \sim \text{normal}(\mu + f(x_n), \sigma^2)$$

with:

$$\vec{f} \sim \text{multinormal}(\vec{0}, K)$$

The two equations are equivalent to:

$$\log \vec{r_w} \sim \text{multinormal}(\vec{\mu}, \Sigma + K)$$

where:

$$\vec{\mu} = \begin{bmatrix} \mu \\ \mu \\ \vdots \\ \mu \end{bmatrix} \quad \Sigma = \begin{bmatrix} \sigma^2 & 0 & \dots & 0 \\ 0 & \sigma^2 & \dots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \dots & \dots & \sigma^2 \end{bmatrix}$$

In practice, we choose to model the log of ringwidth with two Gaussian processes:

- one **long-term scale** Gaussian process f_{long} , to capture the long-term variation of tree growth, likely related to allometric constraints
- one **short-term scale** Gaussian process $f_{\text{short},p}$ per stand p ('stand' is a common forestry term that refers to a set of trees in one geographical location), to capture climatic variations that span several years (while our climate metrics, below, occur at the annual-scale)

The long-term scale GP is characterized by one pair of parameters $\{\gamma_{\text{sp}}, \rho_{\text{sp}}\}$ per species sp . The short-term scale GP is characterized by two parameters $\{\gamma_{\text{short}}, \rho_{\text{short}}\}$, common across all stands. The short-term scale does not depend on species, but its effect is scaled by a factor κ_{sp} (see below). We chose two different prior models based on domaine-expertise for these two Gaussian processes, to have the prior mass of ρ_{short} between 3 and 10 years, and the prior mass of ρ_{sp} between 20 and 60 years.

The full model is, for n observations for one tree k of the species sp on the stand p :

$$\log \vec{r\mathbf{w}}_{sp,p,k} \sim \text{multinormal}(\mu_{sp,p}, \Sigma + \mathbf{K}_{\text{sp}})$$

with:

$$\log \vec{r\mathbf{w}}_{sp,k} = \log \circ \begin{bmatrix} \text{rw}_{sp,p,k}(t_1) \\ \text{rw}_{sp,p,k}(t_2) \\ \vdots \\ \text{rw}_{sp,k}(t_n) \end{bmatrix} \quad \log \vec{\mu}_{sp,p} = \log \circ \begin{bmatrix} \mu_{sp,p}(t_1) \\ \mu_{sp,p}(t_2) \\ \vdots \\ \mu_{sp,p}(t_n) \end{bmatrix}$$

$$\forall i \in \llbracket 1, n \rrbracket \quad \mu_{sp,p}(t_i) = \alpha + \beta_{\text{GSL}} * \text{GSL}(p, t_i) + \beta_{\text{GDD}} * \text{GDD}(p, t_i) + \beta_{\text{SM}} * \text{SM}(p, t_i) + \kappa_{\text{sp}} * f_{\text{short},p}(t_i)$$

$$\vec{f}_{\text{short},p} \sim \text{multinormal}(\vec{0}, \mathbf{K}_{\text{short}})$$


We currently are considering three different climate predictors, at the stand scale:

- the growing season length $\text{GSL}(p, t_i)$, in days
- the growing degree days $\text{GDD}(p, t_i)$ accumulated during the growing season, in $\times 10^\circ \text{C}$
- the mean soil moisture $\text{SM}(p, t_i)$ (weighted average across soil horizons) during the growing season, in %


3.3 Retrodictive check

We can examine how the model fit some trees in one particular stand (from a Pacific Northwest forest):


StandAE10, TreeAE10_F1



StandAE10, TreeAE10_F10



StandAE10, TreeAE10_F12



4 Outcome of the Meeting

Having developed a new model of tree growth that yields remarkable flexibility in capturing the long-term variability seen in tree growth across individuals, stands and species—with a hierarchical form that captures these levels—we are excited to share this model with the research community in a publication. These are hierarchical levels of variability that are generally ignored by the dendrochronological community (most dendrochronological models focus on one species and average all trees across a site), while foresters rarely have data that includes annual individual-tree variation. Thus, we hope our models will help bridge diverging perspectives across these fields and provide improved forecasting. First, however, we need to finalize how to robustly add in climate drivers.

Work since the meeting has focused on finding the best-available climate data to merge with our data on annual tree growth. Given that our tree growth data was often collected across complex topologies with the goal to span climate variability this is no easy task, as weather stations are often quite far from locations of sampled trees. We are addressing this challenge by working with a climatologist from NASA GISS, but may need to consider modeling some climate metrics (e.g., soil moisture) as latent variables informed by local weather data. We expect progress on this challenge by the end of the summer.

Taken together, we have advanced on our goal of improving models of how tree growth interacts with a changing climate far more than we expected was possible in such a short time. While work began before our meeting at BIRS and continues on after, the one week to focus on the challenge and work intensively with experts from ecology, advanced modeling and forestry was pivotal to our progress.

References

- [1] F. S. Chapin, P. A. Matson, and H. A. Mooney. *Principles of Terrestrial Ecosystem Ecology*. Springer-Verlag, New York, 2002.
- [2] Edward R Cook and Leonardas A Kairiukstis. *Methods of dendrochronology: applications in the environmental sciences*. Springer Science & Business Media, 2013.
- [3] Rosanne D’Arrigo, Rob Wilson, Beate Liepert, and Paolo Cherubini. On the ‘divergence problem’ in northern forests: a review of the tree-ring evidence and possible causes. *Global and planetary change*, 60(3-4):289–305, 2008.
- [4] W. Larcher. Plants under stress. In *Physiological Plant Ecology: Ecophysiology and Stress Physiology of Functional Groups*, chapter 6, pages 345–450. Springer, Heidelberg, 4th edition, 2003.
- [5] Zhaochen Zhang, Michael J Papaik, Xugao Wang, Zhanqing Hao, Ji Ye, Fei Lin, and Zuoqiang Yuan. The effect of tree size, neighborhood competition and environment on tree growth in an old-growth temperate forest. *Journal of Plant Ecology*, 10(6):970–980, 2017.