

Overlooked model uncertainties may misinform forest management strategies

Victor, Jérôme, Isabelle

Abstract—still need work: Forests play a major role in mitigating climate change, but increasing threats to forests from climate change have heightened the importance of managing these systems. Robust forecasts of forest composition with increasing climate change are critical to this aim, but are currently highly variable. To help guide management in the face of this variability and understand where we can most rapidly reduce uncertainty through improved models, we compare over 1350 ecological models and climate scenarios in forecasts for forests across Europe. Our approach considers a gradient of more mechanistic (‘process-based’) to correlative models of species distributions to find that uncertainty in ecological models can drive more variation than vastly different climate scenarios (e.g., SSP2 vs. SSP5), but also areas with relatively consistent projections [give overview of these and say that this could reduce uncertainty in how to manage for these areas]. [Maybe something on using existing range data leads to more pessimistic forecasts?] Our results highlight a new way to approach ecological forecasting that better identifies areas of higher certainty and, conversely, the areas where managers will need more diversified approaches and where more ecological study may be most useful.

1 Main

2 Forests are key to pursuing climate change mitigation policies and achieving carbon neutrality^{1,2}.
3 Yet, forests are increasingly under pressure. In Europe, temperatures are rising twice as fast as
4 the global average³, and unprecedented pulses of tree mortality have been reported in the last
5 decade⁴. As a result, some European forests are becoming net CO₂ sources^{5,6}, due to decreased
6 growth^{5,7}, larger burned areas^{8,9}, and increased pest- and drought-induced dieback^{6,10,11}. Forest
7 managers are facing unprecedented challenges, as they must address current threats while also
8 promoting long-term adaptation to climate change. In this context of high uncertainty, better
9 guidance is needed to implement successful strategies.

10 Given the diversity of predictive ecological models, the challenge of providing practical in-
11 sights for forest management is even greater. Different models, ranging from correlative to more
12 mechanistic approaches, may provide highly divergent projections^{12–15}. While it remains un-
13 clear under which conditions one approach is more reliable than another¹⁶, most projections still
14 rely on a limited set of models^{17–20}, ultimately increasing the risk of policy and management
15 failures²¹. We often lack a robust estimation of uncertainty and a comprehensive understanding
16 of what drives differences between projections²². Given the urgency of climate change, we must
17 incorporate the diversity of models and merge across ecological and climatological models to
18 provide a complete picture of both the threats and opportunities for forests.

19 Gaining a better understanding of where uncertainties originate and how they relate is crucial
20 to identify opportunities to address policy-relevant questions^{23–25}. Species shifts are predicted
21 to have major impacts on timber production and on the forest economic sector^{18,19}. Forest
22 managers need to know whether the current species will be able to tolerate future climate
23 conditions, whether they can rely on its natural regeneration, or whether they should capitalize
24 on new species opportunities. If the main driver of variation across projections is the different




Figure 1: Ecological models represent the main source of uncertainty in future projections of species climatic suitability across European biomes (2080-2100).

ecological models, even more than different global emissions scenarios, it becomes critical to encompass the a wide range of models. Failing to do so could lead to overly confident predictions about which species will or will not be able to survive in future climates, ultimately leading to counterproductive or even detrimental forest management decisions.

To understand the level of confidence we can place in predictions requires a framework that account for all the various components of climatological and biological uncertainties, including socioeconomic scenarios, global climate models, ecological models, down to the species level. To this aim, we combined over 1350 projections of forest tree species distributions, incorporating a wide range of models, from more mechanistic (‘process-based’) to correlative models. Fully accounting for our current level of knowledge about future climate states and species functioning allows us to quantify the contribution of each component to the total variation across projections. This approach represents a significant advancement over previous studies, which overlooked large portions of uncertainty, and will lead to better informed decision-making to improve the resilience of forests.

Results and discussion

Our dataset included 9 tree species, both deciduous and coniferous, adapted to diverse climatic conditions across Europe. We simulated their suitability from 1970 to 2100, at a 0.1° spatial resolution, using a diverse set of ecological models spanning different hypotheses and calibration methods. We used 10 different future climate simulations, based on 2 forcing scenarios and 5 global climate models with different climate sensitivities.

Across species, ecological models drive more variation than vastly different climate scenarios (Figure 1). They consistently represent the main source of uncertainty across major European biomes (explaining between 42.9% and 63.9% of the variation between projections), with the exception of the Alpine biome. At the species-level, across all biomes, the differences between ecological models is also the main source of uncertainty for all the species considered here, and represents between 40% and 62% of the total uncertainty on average. One of the striking example is the climatic suitability change of sessile oak in the Atlantic region, where this species represents an important cultural and economic value, and for which more than 80% of the uncertainty in




Figure 2: Considering a broad range of models provides a more comprehensive view of possible future scenarios.

climate change impact projections was due to variations among ecological models.

Failing to account for a broad range of ecological models severely underestimate projection uncertainty, and ultimately lead to incorrect trend estimates (Figure 2). Considering only correlative models would have misled to an overestimation of the contribution of climate projections (forcing scenarios, climate models, and their two-way interaction) to the total projection uncertainty in all regions, except the Mediterranean. In particular, divergence between climate models would have appeared to contribute as much as ecological models to projection uncertainty (on average, 36.6% and 37.5%, respectively). When accounting for more diverse ecological models, as done in our study, the uncertainty introduced by different ecological models (51.0% on average for all species) becomes greater than that introduced by climate models (19.9%). One of the key challenges for reducing uncertainty thus remains at the biological and ecological levels, even before considering the broad variations across future climate projections.

Our results also revealed that the divergence between ecological model projections followed a regular pattern. Models calibrated using current species range data consistently predict greater extinctions than models calibrated using experimental data (Figure 3). Current distribution data may capture only a portion of the climatic niche of a species, underestimating the range of conditions where it could survive^{26,27}. These discrepancies between models can significantly alter country-level projections, and impact national strategies derived from them. For example, by the end of the 21st century, beech showed an average suitability decrease of -0.19 (± 0.14) across its historical distribution when considering only models entirely calibrated with current species distribution data, leading to an average loss of 30.5% of its historical distribution (Figure 3). But this decreasing trend vanish once a broad range of models is accounted for (-0.028 ± 0.17). Relying on a narrow set of models—especially derived from the same calibration process—undermines the robustness of projections²¹, and may ultimately bias decisions towards intensive intervention strategies (e.g. introduction of species outside their native range), potentially overlooking alternative strategies.

Despite large uncertainties, comparing diverse models improve prediction robustness and enable to identify areas with relatively consistent projections that differ in terms of future climate risks and levers of action to address them (Figure 4). Around the Mediterranean Basin, the models consistently predict less favorable climatic conditions for the species we considered here. In areas where most species are threatened, forest managers may thus consider introducing more drought-tolerant species. Along the Atlantic margin, the suitability of most species is also projected to decrease, except for the two Mediterranean species—pubescent and evergreen oaks—which could replace less adapted temperate species such as beech²⁸. Mechanistic model




Figure 3: Models build on current species range data project higher extinctions.

projections are less pessimistic for deciduous oaks and beech (Figure 3), suggesting that some better-adapted populations could survive if the existing standing genetic variation is maintained and promoted by forest managers²⁹. Additionally, adapting management practices, such as decreasing stand density to limit competition for water, could support their long-term survival to drought events³⁰.

A large part of the Continental biome exhibit less clearer trends (Figure 4), as well as mountainous regions at the transition between Mediterranean and Continental/Atlantic climates (Pyrenees, Massif Central, Balkans). Yet, looking at divergence across projections still provide important insights into species and locations most likely to experience significant changes in future climatic conditions. Models agree on a lower suitability for Scots pines (with uncertainty driven more by climatic models and scenarios, 45.7%, than by ecological models, 30.8%), which is a commercially very important species [cite] in several countries of Central Europe (such as Poland, Eastern Germany, Czech Republic, Belarus). Projections also suggest that temperate deciduous species (e.g. beech, deciduous oaks) will be less affected by climate change, despite high uncertainties due to high divergence among ecological models (between 45.7 and 75.4% of the total uncertainty). A key lever of action in this region is the diversification of tree species, as well as increasing genetic diversity within populations, to mitigate the risks associated with uncertain future conditions^{31–34}. Promoting uneven-aged stands could also enhance forest stability by improving structural complexity and buffering against climate extremes^{35,36}.

Finally, boreal biomes in Scandinavian and Baltic countries are projected to get an overall increase of climatic suitability (Figure 4). These include Finland and Sweden, two very important forestry countries in terms of wood stock, added value, and forest-based workforce [cite]. Forests in these regions are dominated by two conifers species, Scots pine and spruce, favoured by commercial forest management. The lack of experimental data prevented us from making mechanistic model projections for spruce, but uncertainty for Scot pine future suitability was very high. Models consistently project that temperate deciduous trees will become more competitive at the northern margin of their range (Figure 3), and the extending growing season could offer an opportunity to convert pure coniferous stands into mixed forest to increase their resilience³⁷.




Figure 4: Accounting for uncertainties supports evidence-based forest management.

The implications of these results extend beyond European forests. The large uncertainties we found raise questions about the robustness of existing projections, and highlight that further advances are necessary to provide policy-relevant projections. Given the rapid pace of climate change, there is no time for fruitless debates over which modeling approach to favor. Instead, efforts should focus on bridging diverse methodologies to generate practical insights and support evidence-based decision making. Interdisciplinary synergies will be essential to developing models that are both grounded in mechanistic understanding and statistically robust. Correlative models may be refined by integrating experimental data³⁸—yet, with some caution³⁹. Mechanistic model improvements should go beyond mere sophistication—adding new processes and new parameters—toward robust statistical inference. Too often, mechanistic models often rely on a single immutable best-parameter set per species⁴⁰, preventing any proper propagation of uncertainty. Ultimately, as scientists, we need to be transparent about these uncertainties if we expect forest managers to acknowledge and integrate them²⁴. This is how ecological modeling can become truly relevant to decision making in the face of a changing climate.

References

- [1] Anu Korosuo, Roberto Pilli, Raúl Abad Viñas, Viorel N. B. Blujdea, Rene R. Colditz, Giulia Fiorese, Simone Rossi, Matteo Vizzarri, and Giacomo Grassi. The role of forests in the EU climate policy: are we on the right track? *Carbon Balance and Management*, 18(1):15, July 2023. ISSN 1750-0680. doi: 10.1186/s13021-023-00234-0. URL <https://doi.org/10.1186/s13021-023-00234-0>.
- [2] Matti Hyrynen, Markku Ollikainen, and Jyri Seppälä. European forest sinks and climate targets: past trends, main drivers, and future forecasts. *European Journal of Forest Research*, 142(5):1207–1224, October 2023. ISSN 1612-4677. doi: 10.1007/s10342-023-01587-4. URL <https://doi.org/10.1007/s10342-023-01587-4>.
- [3] Copernicus Climate Change Service. European state of the climate 2023. Technical report, 2024. URL <https://climate.copernicus.eu/esotc/2023>.
- [4] Cornelius Senf, Allan Buras, Christian S. Zang, Anja Rammig, and Rupert Seidl. Excess forest mortality is consistently linked to drought across Europe. *Nature Communications*, 11(1):6200, December 2020. ISSN 2041-1723. doi: 10.1038/s41467-020-19924-1. URL <https://www.nature.com/articles/s41467-020-19924-1>.
- [5] David Hadden and Achim Grelle. Changing temperature response of respiration turns boreal forest from carbon sink into carbon source. *Agricultural and Forest Meteorology*, 223:30–38, June 2016. ISSN 0168-1923. doi: 10.1016/j.agrformet.2016.03.020. URL <https://www.sciencedirect.com/science/article/pii/S0168192316302131>.
- [6] D. V. Karelin, D. G. Zamolodchikov, A. V. Shilkin, S. Yu. Popov, A. S. Kumanyaev, V. O. Lopes de Gerenyu, N. O. Tel’nova, and Michael L. Gitarskiy. The effect of tree mortality on CO₂ fluxes in an old-growth spruce forest. *European Journal of Forest Research*, 140(2):287–305, April 2021. ISSN 1612-4677. doi: 10.1007/s10342-020-01330-3. URL <https://doi.org/10.1007/s10342-020-01330-3>.
- [7] Auke M. van der Woude, Wouter Peters, Emilie Joetzjer, Sébastien Lafont, Gerbrand Koren, Philippe Ciais, Michel Ramonet, Yidi Xu, Ana Bastos, Santiago Botía, Stephen Sitch, Remco de Kok, Tobias Kneuer, Dagmar Kubistin, Adrien Jacotot, Benjamin Loubet, Pedro-Henrique Herig-Coimbra, Denis Loustau, and Ingrid T. Luijkx. Temperature extremes of 2022 reduced carbon uptake by forests in Europe. *Nature Communications*, 14(1):6218, October 2023. ISSN 2041-1723. doi: 10.1038/s41467-023-41851-0. URL <https://www.nature.com/articles/s41467-023-41851-0>.
- [8] Jofre Carnicer, Andrés Alegria, Christos Giannakopoulos, Francesca Di Giuseppe, Anna Karali, Nikos Koutsias, Piero Lionello, Mark Parrington, and Claudia Vitolo. Global warming is shifting the relationships between fire weather and realized fire-induced CO₂ emissions in Europe. *Scientific Reports*, 12(1):10365, June 2022. ISSN 2045-2322. doi: 10.1038/s41598-022-14480-8. URL <https://www.nature.com/articles/s41598-022-14480-8>.
- [9] Julia Kelly, Natascha Kljun, Zhanzhang Cai, Stefan H. Doerr, Claudio D’Onofrio, Thomas Holst, Irene Lehner, Anders Lindroth, Shangharsha Thapa, Patrik Vestin, and Cristina Santín. Wildfire impacts on the carbon budget of a managed Nordic boreal forest. *Agricultural and Forest Meteorology*, 351:110016, May 2024. ISSN 0168-1923. doi: 10.1016/j.agrformet.2024.110016. URL <https://www.sciencedirect.com/science/article/pii/S016819232400131X>.
- [10] Emil Cienciala and Jan Melichar. Forest carbon stock development following extreme drought-induced dieback of coniferous stands in Central Europe: a CBM-CFS3 model ap-

plication. *Carbon Balance and Management*, 19(1):1, January 2024. ISSN 1750-0680. doi: 10.1186/s13021-023-00246-w. URL <https://doi.org/10.1186/s13021-023-00246-w>.

[11] Lejla Latifovic and M. Altaf Arain. The impact of spongy moth (*Lymantria dispar*) defoliation on carbon balance of a temperate deciduous forest in North America. *Agricultural and Forest Meteorology*, 354:110076, July 2024. ISSN 0168-1923. doi: 10.1016/j.agrformet.2024.110076. URL <https://www.sciencedirect.com/science/article/pii/S0168192324001916>.

[12] Xavier Morin and Wilfried Thuiller. Comparing niche- and process-based models to reduce prediction uncertainty in species range shifts under climate change. *Ecology*, 90(5):1301–1313, 2009. ISSN 1939-9170. doi: 10.1890/08-0134.1. URL <https://onlinelibrary.wiley.com/doi/abs/10.1890/08-0134.1>.

[13] Trevor Keenan, Josep Maria Serra, Francisco Lloret, Miquel Ninyerola, and Santiago Sabate. Predicting the future of forests in the Mediterranean under climate change, with niche- and process-based models: CO2 matters! *Global Change Biology*, 17(1):565–579, 2011. ISSN 1365-2486. doi: 10.1111/j.1365-2486.2010.02254.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2486.2010.02254.x>.

[14] Alissar Cheaib, Vincent Badeau, Julien Boe, Isabelle Chuine, Christine Delire, Eric Dufrêne, Christophe François, Emmanuel S. Gritti, Myriam Legay, Christian Pagé, Wilfried Thuiller, Nicolas Viovy, and Paul Leadley. Climate change impacts on tree ranges: model intercomparison facilitates understanding and quantification of uncertainty. *Ecology Letters*, 15(6):533–544, 2012. ISSN 1461-0248. doi: 10.1111/j.1461-0248.2012.01764.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1461-0248.2012.01764.x>.

[15] Antti Takolander, Thomas Hickler, Laura Meller, and Mar Cabeza. Comparing future shifts in tree species distributions across Europe projected by statistical and dynamic process-based models. *Regional Environmental Change*, 19(1):251–266, January 2019. ISSN 1436-378X. doi: 10.1007/s10113-018-1403-x. URL <https://doi.org/10.1007/s10113-018-1403-x>.

[16] Victor Van der Meersch, Edward Armstrong, Florent Mouillot, Anne Duputié, Hendrik Davi, Frédérik Saltré, and Isabelle Chuine. Biological mechanisms are necessary to improve projections of species range shifts. *bioRxiv*, 2024. doi: 10.1101/2024.05.06.592679. URL <https://www.biorxiv.org/content/early/2024/05/08/2024.05.06.592679>.

[17] Marcin K. Dyderski, Sonia Paz, Lee E. Frelich, and Andrzej M. Jagodzinski. How much does climate change threaten European forest tree species distributions? *Global Change Biology*, 24(3):1150–1163, 2018. ISSN 1365-2486. doi: 10.1111/gcb.13925. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.13925>.

[18] Johannes Wessely, Franz Essl, Konrad Fiedler, Andreas Gattringer, Bernhard Hülber, Olesia Ignateva, Dietmar Moser, Werner Rammer, Stefan Dullinger, and Rupert Seidl. A climate-induced tree species bottleneck for forest management in Europe. *Nature Ecology & Evolution*, 8(6):1109–1117, June 2024. ISSN 2397-334X. doi: 10.1038/s41559-024-02406-8. URL <https://www-nature-com.inee.bib.cnrs.fr/articles/s41559-024-02406-8>.

[19] Marc Hanewinkel, Dominik A. Cullmann, Mart-Jan Schelhaas, Gert-Jan Nabuurs, and Niklaus E. Zimmermann. Climate change may cause severe loss in the economic value of European forest land. *Nature Climate Change*, 3(3):203–207, March 2013. ISSN 1758-6798. doi: 10.1038/nclimate1687. URL <https://www.nature.com/articles/nclimate1687>.

[20] Silvio Schueler, Wolfgang Falk, Jarkko Koskela, François Lefèvre, Michele Bozzano, Jason Hubert, Hojka Kraigher, Roman Longauer, and Ditte C. Olrik. Vulnerability of dynamic

- genetic conservation units of forest trees in Europe to climate change. *Global Change Biology*, 20(5):1498–1511, 2014. ISSN 1365-2486. doi: 10.1111/gcb.12476. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.12476>.
- [21] Terence P. Dawson, Stephen T. Jackson, Joanna I. House, Iain Colin Prentice, and Georgina M. Mace. Beyond Predictions: Biodiversity Conservation in a Changing Climate. *Science*, 332(6025):53–58, April 2011. doi: 10.1126/science.1200303. URL <https://www.science.org/doi/10.1126/science.1200303>.
- [22] Emily G. Simmonds, Kwaku P. Adjei, Benjamin Cretois, Lisa Dickel, Ricardo González-Gil, Jack H. Laverick, Caitlin P. Mandeville, Elizabeth G. Mandeville, Otso Ovaskainen, Jorge Sicacha-Parada, Emma S. Skarstein, and Bob O’Hara. Recommendations for quantitative uncertainty consideration in ecology and evolution. *Trends in Ecology & Evolution*, 39(4):328–337, April 2024. ISSN 0169-5347. doi: 10.1016/j.tree.2023.10.012. URL <https://www.sciencedirect.com/science/article/pii/S0169534723002793>.
- [23] M. C. Urban, G. Bocedi, A. P. Hendry, J.-B. Mihoub, G. Pe’er, A. Singer, J. R. Bridle, L. G. Crozier, L. De Meester, W. Godsoe, A. Gonzalez, J. J. Hellmann, R. D. Holt, A. Huth, K. Johst, C. B. Krug, P. W. Leadley, S. C. F. Palmer, J. H. Pantel, A. Schmitz, P. A. Zollner, and J. M. J. Travis. Improving the forecast for biodiversity under climate change. *Science*, 353(6304):aad8466, September 2016. doi: 10.1126/science.aad8466. URL <https://www.science.org/doi/10.1126/science.aad8466>.
- [24] Andrea Saltelli, Gabriele Bammer, Isabelle Bruno, Erica Charters, Monica Di Fiore, Emmanuel Didier, Wendy Nelson Espeland, John Kay, Samuele Lo Piano, Deborah Mayo, Roger Pielke Jr, Tommaso Portaluri, Theodore M. Porter, Arnald Puy, Ismael Rafols, Jerome R. Ravetz, Erik Reinert, Daniel Sarewitz, Philip B. Stark, Andrew Stirling, Jeroen van der Sluijs, and Paolo Vineis. Five ways to ensure that models serve society: a manifesto. *Nature*, 582(7813):482–484, June 2020. doi: 10.1038/d41586-020-01812-9. URL <https://www.nature.com/articles/d41586-020-01812-9>.
- [25] T. F. Johnson, A. P. Beckerman, D. Z. Childs, T. J. Webb, K. L. Evans, C. A. Griffiths, P. Capdevila, C. F. Clements, M. Besson, R. D. Gregory, G. H. Thomas, E. Delmas, and R. P. Freckleton. Revealing uncertainty in the status of biodiversity change. *Nature*, 628(8009):788–794, March 2024. ISSN 1476-4687. doi: 10.1038/s41586-024-07236-z.
- [26] Mathieu Chevalier, Olivier Broennimann, and Antoine Guisan. Climate change may reveal currently unavailable parts of species’ ecological niches. *Nature Ecology & Evolution*, pages 1–13, May 2024. ISSN 2397-334X. doi: 10.1038/s41559-024-02426-4. URL <https://www.nature.com/articles/s41559-024-02426-4>.
- [27] D. Nogués-Bravo, S. Veloz, B. G. Holt, J. Singarayer, P. Valdes, B. Davis, S. C. Brewer, J. W. Williams, and C. Rahbek. Amplified plant turnover in response to climate change forecast by Late Quaternary records. *Nature Climate Change*, 6(12):1115–1119, December 2016. ISSN 1758-6798. doi: 10.1038/nclimate3146. URL <https://www.nature.com/articles/nclimate3146>.
- [28] Josep Peñuelas and Martí Boada. A global change-induced biome shift in the Montseny mountains (NE Spain). *Global Change Biology*, 9(2):131–140, 2003. ISSN 1365-2486. doi: 10.1046/j.1365-2486.2003.00566.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2486.2003.00566.x>.
- [29] Peter Brang, Peter Spathelf, J. Bo Larsen, Jürgen Bauhus, Andrej Bonccina, Christophe Chauvin, Lars Drössler, Carlos García-Güemes, Caroline Heiri, Gary Kerr, Manfred J. Lexer, Bill Mason, Frits Mohren, Urs Mühlethaler, Susanna Nocentini, and Miroslav

Svoboda. Suitability of close-to-nature silviculture for adapting temperate European forests to climate change. *Forestry: An International Journal of Forest Research*, 87 (4):492–503, October 2014. ISSN 0015-752X. doi: 10.1093/forestry/cpu018. URL <https://doi.org/10.1093/forestry/cpu018>.

[30] Derek J. N. Young, Becky L. Estes, Shana Gross, Amarina Wuenschel, Christina Restaino, and Marc D. Meyer. Effectiveness of forest density reduction treatments for increasing drought resistance of ponderosa pine growth. *Ecological Applications*, 33(4), April 2023. ISSN 1939-5582. doi: 10.1002/eap.2854.

[31] Xavier Morin, Lorenz Fahse, Claire de Mazancourt, Michael Scherer-Lorenzen, and Harald Bugmann. Temporal stability in forest productivity increases with tree diversity due to asynchrony in species dynamics. *Ecology Letters*, 17(12):1526–1535, 2014. ISSN 1461-0248. doi: 10.1111/ele.12357. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.12357>.

[32] Christian Ammer. Diversity and forest productivity in a changing climate. *New Phytologist*, 221(1):50–66, 2019. ISSN 1469-8137. doi: 10.1111/nph.15263. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/nph.15263>.

[33] Hans Pretzsch. Genetic diversity reduces competition and increases tree growth on a Norway spruce (*Picea abies* [L.] Karst.) provenance mixing experiment. *Forest Ecology and Management*, 497:119498, October 2021. ISSN 0378-1127. doi: 10.1016/j.foreco.2021.119498. URL <https://www.sciencedirect.com/science/article/pii/S0378112721005880>.

[34] Sonja Vospernik, Carl Vigren, Xavier Morin, Maude Toïgo, Kamil Bielak, Gediminas Brazaitis, Felipe Bravo, Michael Heym, Miren del Río, Aris Jansons, Magnus Löf, Arne Nothdurft, Marta Pardos, Maciej Pach, Quentin Ponette, and Hans Pretzsch. Can mixing *Quercus robur* and *Quercus petraea* with *Pinus sylvestris* compensate for productivity losses due to climate change? *Science of The Total Environment*, 942:173342, September 2024. ISSN 0048-9697. doi: 10.1016/j.scitotenv.2024.173342. URL <https://www.sciencedirect.com/science/article/pii/S0048969724034892>.

[35] Elia Vangi, Daniela Dalmonech, Elisa Cioccolo, Gina Marano, Leonardo Bianchini, Paulina F. Puchi, Elisa Grieco, Alessandro Cescatti, Andrea Colantoni, Gherardo Chirici, and Alessio Collalti. Stand age diversity (and more than climate change) affects forests’ resilience and stability, although unevenly. *Journal of Environmental Management*, 366: 121822, August 2024. ISSN 0301-4797. doi: 10.1016/j.jenvman.2024.121822.

[36] Shengmin Zhang, Jörgen Sjögren, and Mari Jönsson. Retention forestry amplifies microclimate buffering in boreal forests. *Agricultural and Forest Meteorology*, 350:109973, May 2024. ISSN 0168-1923. doi: 10.1016/j.agrformet.2024.109973.

[37] Bastian Schauer, Simon Thorn, Markus Blaschke, and Thomas Kudernatsch. Conversion of pure spruce to mixed spruce beech stands: Effects on alpha and beta diversity of multiple taxonomic groups. *Forest Ecology and Management*, 545:121297, October 2023. ISSN 0378-1127. doi: 10.1016/j.foreco.2023.121297. URL <https://www.sciencedirect.com/science/article/pii/S0378112723005315>.

[38] Chi-Yun Kuo, Chia-Ying Ko, and Yin-Zheng Lai. Assessing warming impacts on marine fishes by integrating physiology-guided distribution projections, life-history changes and food web dynamics. *Methods in Ecology and Evolution*, 13(6):1343–1357, 2022. ISSN 2041-210X. doi: 10.1111/2041-210X.13846. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13846>.

- [39] Mathieu Chevalier, Vincent Pignard, Olivier Broennimann, and Antoine Guisan. A cautionary message on combining physiological thermal limits with macroclimatic data to predict species distribution. *Ecosphere*, 15(7), July 2024. ISSN 2150-8925. doi: 10.1002/ecs2.4931.
- [40] Sandy P. Harrison, Wolfgang Cramer, Oskar Franklin, Iain Colin Prentice, Han Wang, Ake Brännström, Hugo de Boer, Ulf Dieckmann, Jaideep Joshi, Trevor F. Keenan, Aliénor Lavergne, Stefano Manzoni, Giulia Mengoli, Catherine Morfopoulos, Josep Peñuelas, Stephan Pietsch, Karin T. Rebel, Youngryel Ryu, Nicholas G. Smith, Benjamin D. Stocker, and Ian J. Wright. Eco-evolutionary optimality as a means to improve vegetation and land-surface models. *New Phytologist*, 231(6):2125–2141, 2021. ISSN 1469-8137. doi: 10.1111/nph.17558. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/nph.17558>.