

Overlooked model uncertainties may misinform forest management strategies

Victor, Jérôme, Isabelle, and more?

Abstract: Forests play a major role in mitigating climate change, but increasing threats to forests from climate change have heightened the importance of managing these systems. Robust forecasts of forest composition with increasing climate change are critical to this aim, but are currently highly variable. To help guide management in the face of this variability and understand where we can most rapidly reduce uncertainty through improved models, we compare over XX ecological models and climate scenarios in forecasts for forests across Europe. Our approach considers a gradient of more mechanistic (‘process-based’) to correlative models of species distributions to find that uncertainty in ecological models can drive more variation than vastly different climate scenarios (e.g., SSP2 vs. SSP5), but also areas with relatively consistent projections [give overview of these and say that this could reduce uncertainty in how to manage for these areas]. [Maybe something on using existing range data leads to more pessimistic forecasts?] Our results highlight a new way to approach ecological forecasting that better identifies areas of higher certainty and, conversely, the areas where managers will need more diversified approaches and where more ecological study may be most useful.

1 Main

Forests are key to pursuing climate change mitigation policies and achieving carbon neutrality^{1,2}. Yet, forests are increasingly under pressure. In Europe, temperatures are rising twice as fast as the global average³, and unprecedented pulses of tree mortality have been reported in the last decade⁴. As a result, some European forests are becoming net CO₂ sources^{5,6}, due to decreased growth^{5,7}, larger burned areas^{8,9}, and increased pest- and drought-induced dieback^{6,10,11}. Forest managers are facing unprecedented challenges, as they must address current threats while also promoting long-term adaptation to climate change. In this context of high uncertainty, better guidance is needed to implement successful strategies.

Given the diversity of predictive ecological models, the challenge of providing practical insights for forest management is even greater. Different models, ranging from correlative to more mechanistic approaches, may provide highly divergent projections^{12–15}. While it remains unclear under which conditions one approach is more reliable than another¹⁶, most forecasting studies still rely on a limited set of models^{17–20}. We thus often lack a comprehensive understanding of what drives differences between projections²¹. Given the urgency of climate change, we must incorporate this diversity and merge across ecological and climatological models to provide a complete picture of both the threats and opportunities for forests.

Gaining a better understanding of where uncertainties originate and how they relate is crucial to identify opportunities to address policy-relevant questions²³. Species shifts are predicted to have major impacts on timber production and on the forest economic sector^{18,19}. Forest managers need to know whether the current species will be able to tolerate future climate conditions, whether they can rely on its natural regeneration, or whether they should capitalize

on new species opportunities. If the main driver of variation across projections is the different ecological models, even more than different global emissions scenarios, it becomes critical to encompass the full range of models. Failing to do so could lead to overly confident predictions about which species will or will not be able to survive in future climates, ultimately leading to counterproductive or even detrimental forest management decisions.

To understand the level of confidence we can place in predictions requires methods that account for all the various components of climatological and biological uncertainties, including socio-economic scenarios, global climate models, ecological models, down to the species level. To this aim, we combined over 2,000 projections of forest tree species distributions, considering in particular a range of models, from more mechanistic (‘process-based’) to correlative models. Fully accounting for our current level of knowledge about future climate states and species functioning allows us to quantify the contribution of each component to the total variation across projections. This approach represents a significant advancement over previous studies, which overlooked large portions of uncertainty, and will lead to better informed decision-making to improve the resilience of our forests.

Results and discussion

References

- [1] Anu Korosuo, Roberto Pilli, Raúl Abad Viñas, Viorel N. B. Blujdea, Rene R. Colditz, Giulia Fiorese, Simone Rossi, Matteo Vizzarri, and Giacomo Grassi. The role of forests in the EU climate policy: are we on the right track? *Carbon Balance and Management*, 18(1):15, July 2023. ISSN 1750-0680. doi: 10.1186/s13021-023-00234-0. URL <https://doi.org/10.1186/s13021-023-00234-0>.
- [2] Matti Hyyrynen, Markku Ollikainen, and Jyri Seppälä. European forest sinks and climate targets: past trends, main drivers, and future forecasts. *European Journal of Forest Research*, 142(5):1207–1224, October 2023. ISSN 1612-4677. doi: 10.1007/s10342-023-01587-4. URL <https://doi.org/10.1007/s10342-023-01587-4>.
- [3] Copernicus Climate Change Service. European state of the climate 2023. Technical report, 2024. URL <https://climate.copernicus.eu/esotc/2023>.
- [4] Cornelius Senf, Allan Buras, Christian S. Zang, Anja Rammig, and Rupert Seidl. Excess forest mortality is consistently linked to drought across Europe. *Nature Communications*, 11(1):6200, December 2020. ISSN 2041-1723. doi: 10.1038/s41467-020-19924-1. URL <https://www.nature.com/articles/s41467-020-19924-1>.
- [5] David Hadden and Achim Grelle. Changing temperature response of respiration turns boreal forest from carbon sink into carbon source. *Agricultural and Forest Meteorology*, 223:30–38, June 2016. ISSN 0168-1923. doi: 10.1016/j.agrformet.2016.03.020. URL <https://www.sciencedirect.com/science/article/pii/S0168192316302131>.
- [6] D. V. Karelin, D. G. Zamolodchikov, A. V. Shilkin, S. Yu. Popov, A. S. Kumanyaev, V. O. Lopes de Gerenyu, N. O. Tel’nova, and Michael L. Ginarskiy. The effect of tree mortality on CO₂ fluxes in an old-growth spruce forest. *European Journal of Forest Research*, 140(2):287–305, April 2021. ISSN 1612-4677. doi: 10.1007/s10342-020-01330-3. URL <https://doi.org/10.1007/s10342-020-01330-3>.
- [7] Auke M. van der Woude, Wouter Peters, Emilie Joetzjer, Sébastien Lafont, Gerbrand Koren, Philippe Ciais, Michel Ramonet, Yidi Xu, Ana Bastos, Santiago Botía, Stephen Sitch, Remco de Kok, Tobias Kneuer, Dagmar Kubistin, Adrien Jacotot, Benjamin Loubet, Pedro-Henrique Herig-Coimbra, Denis Loustau, and Ingrid T. Luijkx. Temperature extremes of 2022 reduced carbon uptake by forests in Europe. *Nature Communications*, 14(1):6218, October 2023. ISSN 2041-1723. doi: 10.1038/s41467-023-41851-0. URL <https://www.nature.com/articles/s41467-023-41851-0>.
- [8] Jofre Carnicer, Andrés Alegria, Christos Giannakopoulos, Francesca Di Giuseppe, Anna Karali, Nikos Koutsias, Piero Lionello, Mark Parrington, and Claudia Vitolo. Global warming is shifting the relationships between fire weather and realized fire-induced CO₂ emissions in Europe. *Scientific Reports*, 12(1):10365, June 2022. ISSN 2045-2322. doi: 10.1038/s41598-022-14480-8. URL <https://www.nature.com/articles/s41598-022-14480-8>.
- [9] Julia Kelly, Natascha Kljun, Zhanzhang Cai, Stefan H. Doerr, Claudio D’Onofrio, Thomas Holst, Irene Lehner, Anders Lindroth, Shangharsha Thapa, Patrik Vestin, and Cristina Santín. Wildfire impacts on the carbon budget of a managed Nordic boreal forest. *Agricultural and Forest Meteorology*, 351:110016, May 2024. ISSN 0168-1923. doi: 10.1016/j.agrformet.2024.110016. URL <https://www.sciencedirect.com/science/article/pii/S016819232400131X>.

- [10] Emil Cienciala and Jan Melichar. Forest carbon stock development following extreme drought-induced dieback of coniferous stands in Central Europe: a CBM-CFS3 model application. *Carbon Balance and Management*, 19(1):1, January 2024. ISSN 1750-0680. doi: 10.1186/s13021-023-00246-w. URL <https://doi.org/10.1186/s13021-023-00246-w>.
- [11] Lejla Latifovic and M. Altaf Arain. The impact of spongy moth (*Lymantria dispar*) defoliation on carbon balance of a temperate deciduous forest in North America. *Agricultural and Forest Meteorology*, 354:110076, July 2024. ISSN 0168-1923. doi: 10.1016/j.agrformet.2024.110076. URL <https://www.sciencedirect.com/science/article/pii/S0168192324001916>.
- [12] Xavier Morin and Wilfried Thuiller. Comparing niche- and process-based models to reduce prediction uncertainty in species range shifts under climate change. *Ecology*, 90(5):1301–1313, 2009. ISSN 1939-9170. doi: 10.1890/08-0134.1. URL <https://onlinelibrary.wiley.com/doi/abs/10.1890/08-0134.1>.
- [13] Trevor Keenan, Josep Maria Serra, Francisco Lloret, Miquel Ninyerola, and Santiago Sabate. Predicting the future of forests in the Mediterranean under climate change, with niche- and process-based models: CO2 matters! *Global Change Biology*, 17(1):565–579, 2011. ISSN 1365-2486. doi: 10.1111/j.1365-2486.2010.02254.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2486.2010.02254.x>.
- [14] Alissar Cheaib, Vincent Badeau, Julien Boe, Isabelle Chuine, Christine Delire, Eric Dufrêne, Christophe François, Emmanuel S. Gritti, Myriam Legay, Christian Pagé, Wilfried Thuiller, Nicolas Viovy, and Paul Leadley. Climate change impacts on tree ranges: model intercomparison facilitates understanding and quantification of uncertainty. *Ecology Letters*, 15(6):533–544, 2012. ISSN 1461-0248. doi: 10.1111/j.1461-0248.2012.01764.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1461-0248.2012.01764.x>.
- [15] Antti Takolander, Thomas Hickler, Laura Meller, and Mar Cabeza. Comparing future shifts in tree species distributions across Europe projected by statistical and dynamic process-based models. *Regional Environmental Change*, 19(1):251–266, January 2019. ISSN 1436-378X. doi: 10.1007/s10113-018-1403-x. URL <https://doi.org/10.1007/s10113-018-1403-x>.
- [16] Victor Van der Meersch, Edward Armstrong, Florent Mouillot, Anne Duputié, Hendrik Davi, Frédéric Saltré, and Isabelle Chuine. Biological mechanisms are necessary to improve projections of species range shifts. *bioRxiv*, 2024. doi: 10.1101/2024.05.06.592679. URL <https://www.biorxiv.org/content/early/2024/05/08/2024.05.06.592679>.
- [17] Marcin K. Dyderski, Sonia Paz, Lee E. Frelich, and Andrzej M. Jagodzinski. How much does climate change threaten European forest tree species distributions? *Global Change Biology*, 24(3):1150–1163, 2018. ISSN 1365-2486. doi: 10.1111/gcb.13925. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.13925>.
- [18] Johannes Wessely, Franz Essl, Konrad Fiedler, Andreas Gattringer, Bernhard Hülber, Oleksia Ignateva, Dietmar Moser, Werner Rammer, Stefan Dullinger, and Rupert Seidl. A climate-induced tree species bottleneck for forest management in Europe. *Nature Ecology & Evolution*, 8(6):1109–1117, June 2024. ISSN 2397-334X. doi: 10.1038/s41559-024-02406-8. URL <https://www-nature-com.inee.bib.cnrs.fr/articles/s41559-024-02406-8>.
- [19] Marc Hanewinkel, Dominik A. Cullmann, Mart-Jan Schelhaas, Gert-Jan Nabuurs, and Niklaus E. Zimmermann. Climate change may cause severe loss in the economic value of

- European forest land. *Nature Climate Change*, 3(3):203–207, March 2013. ISSN 1758-6798. doi: 10.1038/nclimate1687. URL <https://www.nature.com/articles/nclimate1687>.
- [20] Silvio Schueler, Wolfgang Falk, Jarkko Koskela, François Lefèvre, Michele Bozzano, Jason Hubert, Hojka Kraigher, Roman Longauer, and Ditte C. Olrik. Vulnerability of dynamic genetic conservation units of forest trees in Europe to climate change. *Global Change Biology*, 20(5):1498–1511, 2014. ISSN 1365-2486. doi: 10.1111/gcb.12476. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.12476>.
- [21] Emily G. Simmonds, Kwaku P. Adjei, Benjamin Cretois, Lisa Dickel, Ricardo González-Gil, Jack H. Laverick, Caitlin P. Mandeville, Elizabeth G. Mandeville, Otso Ovaskainen, Jorge Sicacha-Parada, Emma S. Skarstein, and Bob O’Hara. Recommendations for quantitative uncertainty consideration in ecology and evolution. *Trends in Ecology & Evolution*, 39(4):328–337, April 2024. ISSN 0169-5347. doi: 10.1016/j.tree.2023.10.012. URL <https://www.sciencedirect.com/science/article/pii/S0169534723002793>.
- [22] Owen L. Petchey, Mikael Pontarp, Thomas M. Massie, Sonia Kéfi, Arpat Ozgul, Maja Weilenmann, Gian Marco Palamara, Florian Altermatt, Blake Matthews, Jonathan M. Levine, Dylan Z. Childs, Brian J. McGill, Michael E. Schaepman, Bernhard Schmid, Piet Spaak, Andrew P. Beckerman, Frank Pennekamp, and Ian S. Pearse. The ecological forecast horizon, and examples of its uses and determinants. *Ecology Letters*, 18(7):597–611, 2015. ISSN 1461-0248. doi: 10.1111/ele.12443. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.12443>.
- [23] M. C. Urban, G. Bocedi, A. P. Hendry, J.-B. Mihoub, G. Pe’er, A. Singer, J. R. Bridle, L. G. Crozier, L. De Meester, W. Godsoe, A. Gonzalez, J. J. Hellmann, R. D. Holt, A. Huth, K. Johst, C. B. Krug, P. W. Leadley, S. C. F. Palmer, J. H. Pantel, A. Schmitz, P. A. Zollner, and J. M. J. Travis. Improving the forecast for biodiversity under climate change. *Science*, 353(6304):aad8466, September 2016. doi: 10.1126/science.aad8466. URL <https://www.science.org/doi/10.1126/science.aad8466>.